

ON SOME INTEGRALS INVOLVING $\Delta_2(x)$ AND $\Delta_3(x)$

WENGUANG ZHAI

In memory of Professor Aleksandar Ivić

(Presented at the 2nd Meeting, held on March 26, 2021)

A b s t r a c t. Let $k \geq 2$ be a fixed natural number and $d_k(n)$ denote the number of ways n can be written as a product of k positive integers. Let $\Delta_k(x)$ denote the error term in the asymptotic formula of the summatory function of $d_k(n)$. In this paper we study some integrals involving $\Delta_2(x)$ and $\Delta_3(x)$.

AMS Mathematics Subject Classification (2020): 11N37.

Key Words: General divisor problem, Dirichlet divisor problem, Power moment.

1. Introduction and statements of results

Let $k \geq 2$ be a fixed natural number and $d_k(n)$ denote the number of ways n can be written as a product of k positive integers. The general divisor problem is to study the asymptotic behaviour of the sum

$$D_k(x) := \sum_{n \leq x} d_k(n)$$

as x tends to infinity. It is well-known that the asymptotic formula of $D_k(x)$ is of the form

$$D_k(x) = xP_k(\log x) + \Delta_k(x), \quad (1.1)$$

where $P_k(u)$ is a polynomial in u of degree $k - 1$, and $\Delta_k(x)$ is the error term. It is a classical and important problem in the analytic number theory to study properties of $\Delta_k(x)$.

When $k = 2$, this is usually called the Dirichlet divisor problem since Dirichlet first proved that $\Delta_2(x) \ll \sqrt{x}$. Dirichlet's result was improved by many authors. The best upper bound up to date is $\Delta_2(x) \ll x^{517/1648+\varepsilon}$ proved in Bourgain and Watt [1]. It is conjectured that the estimate $\Delta_2(x) \ll x^{1/4+\varepsilon}$ holds, which is supported by the mean square result(see Cramér [3])

$$\int_1^T (\Delta_2(x))^2 dx = C_2^{(2)} T^{3/2} + O(T^{5/4+\varepsilon}) \quad (1.2)$$

and the estimate (see Ivić [5], Thm 13.9)

$$\int_1^T |\Delta_2(x)|^{A_0} dx \ll T^{1+A_0/4+\varepsilon} \quad (A_0 = 8.75), \quad (1.3)$$

where

$$C_2^{(2)} := \frac{1}{6\pi^2} \sum_{n=1}^{\infty} \frac{d_2^2(n)}{n^{3/2}}.$$

The estimate $T^{5/4+\varepsilon}$ was improved in Tong[14], Preissmann[12], Lau and Tsang [10], respectively. For any integer $3 \leq l \leq 9$, we have the asymptotic formula

$$\int_1^T (\Delta_2(x))^l dx = C_2^{(l)} T^{1+l/4} + O(T^{1+l/4-\delta_l+\varepsilon}), \quad (1.4)$$

where $C_2^{(l)}$ and $\delta_l > 0$ are explicit constants. Tsang [15] first proved the asymptotic formula (1.4) for $l = 3$ and $l = 4$ with $\delta_3 = 1/14$ and $\delta_4 = 1/23$. Ivić and Sargos [6] proved the asymptotic formula (1.4) for $l = 3$ and $l = 4$ with $\delta_3 = 7/20$ and $\delta_4 = 1/12$. Zhai [17] proved (1.4) for any integer $3 \leq l \leq 9$ by a unified approach. For more results about (1.4), see [9, 11, 16, 19, 20].

When $k = 3$, this is usually called the Piltz divisor problem. The best estimate of $\Delta_3(x)$ up to date is $\Delta_3(x) \ll x^{43/96+\varepsilon}$ proved in Kolesnik [8]. Tong [14] proved the mean square result

$$\int_1^T (\Delta_3(x))^2 dx = C_3^{(2)} T^{5/3} + O(T^{14/9+\varepsilon}), \quad (1.5)$$

where

$$C_3^{(2)} := \frac{1}{10\pi^2} \sum_{n=1}^{\infty} \frac{d_3^2(n)}{n^{4/3}}.$$

Heath-Brown[4] proved that the estimate

$$\int_1^T |\Delta_3(x)|^3 dx \ll T^{2+\varepsilon} \quad (1.6)$$

holds.

Ivić and Zhai [7] first studied the hybrid moments of $\Delta_2(x)$ and $\Delta_3(x)$. They proved that

$$\int_1^T \Delta_2(x)\Delta_3(x) dx \ll T^{13/9} \log^{10/3} T. \quad (1.7)$$

In this paper we shall study the upper bound of the integral

$$\int_1^T \Delta_2^k(x)\Delta_3(x) dx$$

with $k = 2$ and $k = 3$. We have the following theorems.

Theorem 1.1. *We have the estimate*

$$\int_1^T \Delta_2^2(x)\Delta_3(x) dx \ll T^{16/9+\varepsilon}. \quad (1.8)$$

Theorem 1.2. *We have the estimate*

$$\int_1^T \Delta_2^3(x)\Delta_3(x) dx \ll T^{73/36+\varepsilon}. \quad (1.9)$$

Remark 1.1. From (1.3), (1.5) and Cauchy's inequality we get

$$\int_1^T \Delta_2^2(x)\Delta_3(x) dx \ll T^{11/6+\varepsilon}, \quad \int_1^T \Delta_2^3(x)\Delta_3(x) dx \ll T^{25/12+\varepsilon}.$$

Note that $16/9 = 11/6 - 1/18$, $73/36 = 25/12 - 1/18$. So Theorem 1.1 and Theorem 1.2 are non-trivial.

Notation. For any $k \geq 2$, $d_k(n)$ denotes the number of ways n can be written as a product of k natural numbers. $n \sim N$ means $N < n \leq 2N$ and $n \asymp N$ means $c_1N < n \leq c_2N$ for absolute positive constants $c_2 > c_1 > 0$. ε always denotes a small positive constant which may be different at different places. For any complex number α , we define $e(\alpha) := \exp(2\pi i\alpha)$.

2. Some preliminary lemmas

In order to prove our theorems, we need the following lemmas.

Lemma 2.1. *Let $c > 0$ be a non-integer real number, $M \geq 2$ be a large parameter, $\delta > 0$ be any real number. Let $\mathcal{A}(M, \delta; c)$ denote the number of solutions of the inequality*

$$|m_1^c + m_2^c - m_3^c - m_4^c| \leq \delta M^c, \quad M < m_1, m_2, m_3, m_4 \leq 2M.$$

Then we have

$$\mathcal{A}(M, \delta) \ll M^\varepsilon (M^2 + \delta M^4).$$

PROOF. This is Theorem 2 of [13]. □

Lemma 2.2. *Let $T \geq 10$ be a large parameter and y be a real number such that $T^\varepsilon \ll y \ll T$. For any $T \leq x \leq 2T$ define*

$$\Delta_{31}(x; y) := \frac{x^{1/3}}{\sqrt{3}\pi} \sum_{n \leq y} \frac{d_3(n)}{n^{2/3}} \cos(6\pi(nx)^{1/3}).$$

Then uniformly for $y \ll T^{1/3}$ we have

$$\int_T^{2T} |\Delta_{31}(x; y)|^4 dx \ll T^{7/3+\varepsilon}. \quad (2.1)$$

PROOF. By a splitting argument and Hölder's inequality we get for some $1 \ll N \ll y$ that

$$\begin{aligned} \int_T^{2T} |\Delta_{31}(x; y)|^4 dx &= \int_T^{2T} \left| \frac{x^{1/3}}{\sqrt{3}\pi} \sum_j \sum_{\frac{y}{2^{j+1}} < n \leq \frac{y}{2^j}} \frac{d_3(n)}{n^{2/3}} \cos(6\pi(nx)^{1/3}) \right|^4 dx \\ &\ll T^{4/3} \log^3 T \sum_j \int_T^{2T} \left| \sum_{\frac{y}{2^{j+1}} < n \leq \frac{y}{2^j}} \frac{d_3(n)}{n^{2/3}} \cos(6\pi(nx)^{1/3}) \right|^4 dx \\ &\ll T^{4/3} \log^4 T \sum_{n_1, n_2, n_3, n_4 \sim N} \frac{\prod_{j=1}^4 d_3(n_j)}{\prod_{j=1}^4 n_j^{2/3}} \int_T^{2T} e(3x^{1/3}\eta) dx, \end{aligned}$$

where $\eta := n_1^{1/3} + n_2^{1/3} - n_3^{1/3} - n_4^{1/3}$. By the first derivative test we have

$$\int_T^{2T} e(3x^{1/3}\eta) dx \ll \min \left(T, \frac{T^{2/3}}{|\eta|} \right).$$

So from the above two estimates we have

$$\int_T^{2T} |\Delta_{31}(x; y)|^4 dx \ll \frac{T^{2+\varepsilon}}{N^{8/3}} \sum_{\substack{n_j \sim N \\ j=1,2,3,4}} \min\left(T^{1/3}, \frac{1}{|\eta|}\right). \quad (2.2)$$

According to Lemma 2.1, the contribution of $T^{1/3}$ (in this case $|\eta| \leq T^{-1/3}$) is

$$\ll \frac{T^{7/3+\varepsilon}}{N^{8/3}} (N^2 + T^{-1/3} N^{-1/3} N^4) \ll T^{7/3+\varepsilon} N^{-2/3} + T^{2+\varepsilon} N \ll T^{7/3+\varepsilon}$$

if $y \ll T^{1/3}$.

Now we consider the contribution of $1/|\eta|$ for which $T^{-1/3} \ll |\eta| \ll y^{1/3}$. We divide the range of η into $O(\log T)$ subcases of the form $\xi < |\eta| \leq 2\xi$. By Lemma 2.1 again we get the the contribution of $1/|\eta|$ is

$$\begin{aligned} &\ll \frac{T^{2+\varepsilon}}{N^{8/3}} \max_{T^{-1/3} \ll \xi \ll y^{1/3}} \sum_{\xi < |\eta| \leq 2\xi} \frac{1}{|\eta|} \\ &\ll \frac{T^{2+\varepsilon}}{N^{8/3}} \max_{T^{-1/3} \ll \xi \ll y^{1/3}} \frac{1}{\xi} (N^2 + \xi N^{-1/3} N^4) \\ &\ll T^{7/3+\varepsilon} N^{-2/3} + T^{2+\varepsilon} N \ll T^{7/3+\varepsilon}. \end{aligned}$$

Whence Lemma 2.2 follows. \square

Lemma 2.3. *Let $T \geq 10$ be a large parameter and y be a real number such that $T^\varepsilon \ll y \ll T$. Define for any $T \leq x \leq 2T$ that*

$$\Delta_{32}(x; y) := \Delta_3(x) - \Delta_{31}(x; y).$$

Then we have

$$\int_T^{2T} |\Delta_{32}(x; y)|^2 dx \ll T^{5/3+\varepsilon} y^{-1/3} \quad (y \ll T^{1/3}). \quad (2.3)$$

PROOF. We give a very short proof (2.3). We take $d = 3$, $a(n) = d_3(n)$, $N = [T^{5-\varepsilon}]$, $\sigma^* = 7/12$, and $M = [T^{2/3}]$. As in the proof of Theorem 1 in [2], we can write

$$\Delta_{32}(x; y) = R_1^*(x; y) + \sum_{j=2}^7 R_j(x),$$

where

$$R_1^*(x; y) := \frac{x^{1/3}}{\sqrt{3}\pi} \sum_{y < n \leq M} \frac{d_3(n)}{n^{2/3}} \cos(6\pi(nx)^{1/3})$$

and $R_j(x)$ ($j = 2, 3, 4, 5, 6, 7$) were defined in Page 2129 of [2]. Similar to the formula (8.11) of [2], we have the estimate (noting that $y \ll T^{1/3}$)

$$\begin{aligned} & \int_T^{2T} (R_1^*(x; y) + R_2(x))^2 dx \\ & \ll \sum_{y < n < M} \frac{d_3^2(n)}{n^{4/3}} \int_T^{2T} x^{2/3} dx + T^{5/3+\varepsilon} M^{-1/6} + T^{4/3+\varepsilon} M^{1/3} \\ & \ll T^{5/3+\varepsilon} y^{-1/3} + T^{14/9+\varepsilon} \ll T^{5/3+\varepsilon} y^{-1/3}, \end{aligned}$$

which combining (8.17) of [2] gives (2.3). \square

Lemma 2.4. *Let $T \geq 10$ be a large parameter and Y be a real number such that $T^\varepsilon \ll Y \ll T$. For any $T \leq x \leq 2T$ define*

$$\Delta_{21}(x; Y) := \frac{x^{1/4}}{\sqrt{2}\pi} \sum_{n \leq Y} \frac{d_2(n)}{n^{3/4}} \cos \left[4\pi(nx)^{1/2} - \frac{\pi}{4} \right].$$

Then uniformly for $Y \ll T$ we have

$$\int_T^{2T} |\Delta_{21}(x; Y)|^4 dx \ll T^{2+\varepsilon}. \quad (2.4)$$

If $Y \ll T^{1/3}$, then we have

$$\int_T^{2T} |\Delta_{21}(x; Y)|^8 dx \ll T^{3+\varepsilon}. \quad (2.5)$$

PROOF. For any $1 \ll N \ll x$, we have (see, for example, [5])

$$\Delta_2(x) = \Delta_{21}(x; N) + O\left(x^{1/2+\varepsilon} N^{-1/2}\right). \quad (2.6)$$

We first show (2.4). If $Y \ll T^{1/2}$, then by the same approach as the proof of Lemma 2.2 we get (2.4). If $T \gg T^{1/2}$, from (2.6) we have

$$\Delta_{21}(x; Y) = \Delta_{21}(x; \sqrt{T}) + O(x^{1/4+\varepsilon}).$$

So (2.4) follows from the case $y = \sqrt{T}$.

Using the approaches of Theorems 13.8 and 13.9 directly to $\Delta_{21}(x; Y)$ we can get (2.5) easily. So, we omit the details. \square

Lemma 2.5. *Let $T \geq 10$ be a large parameter and Y be a real number such that $T^\varepsilon \ll Y \ll T$. For any $T \leq x \leq 2T$ define*

$$\Delta_{22}(x; Y) := \Delta_2(x) - \Delta_{21}(x; Y).$$

Then for $Y \ll T$ we have

$$\int_T^{2T} |\Delta_{22}(x; Y)|^2 dx \ll T^{3/2+\varepsilon} Y^{-1/2}. \quad (2.7)$$

If $Y \ll T^{1/3}$, then we have

$$\int_T^{2T} |\Delta_{22}(x; Y)|^4 dx \ll T^{2+\varepsilon} Y^{-1/3}. \quad (2.8)$$

PROOF. The estimate (2.7) is Lemma 2.2 of Ivić and Zhai [7].

From (2.5) and (1.3) with $A_0 = 8$ we have

$$\begin{aligned} \int_T^{2T} |\Delta_{22}(x; Y)|^8 dx &= \int_T^{2T} |\Delta_2(x) - \Delta_{21}(x; Y)|^8 dx \\ &\ll \int_T^{2T} |\Delta_{21}(x; Y)|^8 dx + \int_T^{2T} |\Delta_2(x)|^8 dx \ll T^{3+\varepsilon}. \end{aligned} \quad (2.9)$$

Now from (2.7), (2.9) and Hölder's inequality we get

$$\begin{aligned} \int_T^{2T} |\Delta_{22}(x; Y)|^4 dx &\ll \left(\int_T^{2T} |\Delta_{22}(x; Y)|^2 \right)^{2/3} \left(\int_T^{2T} |\Delta_{22}(x; Y)|^8 \right)^{1/3} \\ &\ll T^{2+\varepsilon} Y^{-1/3}. \end{aligned} \quad \square$$

Lemma 2.6. *Suppose $(i_1, i_2) \in \{0, 1\}^2$ and $Y \geq 10$ is a real number. For $(n_1, n_2, n_3) \in \mathbb{N}^3$, define*

$$\alpha_3 := \sqrt{n_1} + (-1)^{i_1} \sqrt{n_2} + (-1)^{i_2} \sqrt{n_3},$$

$$H(Y; i_1, i_2) := \sum_{\substack{n_j \leq Y, 1 \leq j \leq 3 \\ \alpha_3 \neq 0}} \frac{d_2(n_1) d_2(n_2) d_2(n_3)}{(n_1 n_2 n_3)^{3/4} |\alpha_3|}.$$

Then we have

$$H(Y; i_1, i_2) \ll Y^{1/4+\varepsilon}.$$

PROOF. This is Lemma 2.6 of the author [18]. \square

Lemma 2.7. Let $N, M, K \geq 10$, $D = \max(N, M, K)$, $0 < \Delta \ll D^{1/2}$. Let

$$\mathcal{A}(N, M, K; \Delta) := \sum_{\substack{n \sim N, m \sim M, k \sim K \\ |\sqrt{n} + \sqrt{m} - \sqrt{k}| \leq \Delta}} 1.$$

Then we have

$$D^{-\varepsilon} \mathcal{A}(N, M, K; \Delta) \ll \Delta D^{-1/2} N M K + D^{-1/2} (N M K)^{1/2}.$$

PROOF. This is Lemma 2.4 of the author [18]. \square

Lemma 2.8. Suppose $f(x)$ is real-valued such that $|f'(x)| \gg \Delta > 0$ in $[a, b]$, then we have

$$\int_a^b e(f(x)) dx \ll \frac{1}{\Delta}.$$

PROOF. See, for example, Ivić [5]. \square

Lemma 2.9. We have the estimates

$$\sum_{n \leq x} d_2(n) \ll x \log x, \quad \sum_{n \leq x} d_3(n) \ll x \log^2 x.$$

PROOF. See, for example, Ivić [5]. \square

Lemma 2.10. Suppose $T \geq 3$ is a large parameter. Then we have the estimate

$$\sum_{\substack{n, m \leq T \\ n \neq m}} \frac{d_2(n) d_2(m)}{m^{3/4} n^{3/4} |n^{1/2} - m^{1/2}|} \ll \log^4 T.$$

PROOF. See, for example, Ivić and Zhai [7]. \square

Lemma 2.11. If $\sqrt{n} + \sqrt{m} - \sqrt{k} \neq 0$, then we have

$$|\sqrt{n} + \sqrt{m} - \sqrt{k}| \gg \frac{1}{\sqrt{nmk}}.$$

PROOF. This is Lemma 2.5 of the author [18]. \square

3. Proof of Theorem 1.1

Let $T \geq 10$ be a large parameter. We only need to estimate the integral

$$P(T) := \int_T^{2T} \Delta_3(x) \Delta_2^2(x) dx.$$

Suppose $T^\varepsilon \ll y \ll T^{1/3}$ is a parameter to be determined later. Write

$$P(T) = P_1(T) + P_2(T) \tag{3.1}$$

where

$$P_1(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_2^2(x) dx,$$

and

$$P_2(T) := \int_T^{2T} \Delta_{32}(x; y) \Delta_2^2(x) dx.$$

By Lemma 2.3 and (1.3) with $A_0 = 4$ we get

$$P_2(T) \ll T^{11/6+\varepsilon} y^{-1/6}. \tag{3.2}$$

It suffices for us to bound $P_1(T)$. Suppose $T^\varepsilon \ll Y \ll T^{1/3}$ is a parameter to be determined later. Write

$$P_1(T) = P_{11}(T) + 2P_{12}(T) + P_{13}(T), \tag{3.3}$$

where

$$P_{11}(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_{21}^2(x; Y) dx,$$

$$P_{12}(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_{21}(x; Y) \Delta_{22}(x; Y) dx,$$

$$P_{13}(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_{22}^2(x; Y) dx.$$

From Lemma 2.2, (2.4) of Lemma 2.4, (2.7) of Lemma 2.5, and using Hölder's inequality, we get

$$\begin{aligned} P_{12}(T) &\ll \left(\int_T^{2T} \Delta_{31}^4(x; y) dx \right)^{1/4} \left(\int_T^{2T} \Delta_{21}^4(x; Y) dx \right)^{1/4} \left(\int_T^{2T} \Delta_{22}^2(x; Y) dx \right)^{1/2} \\ &\ll T^{11/6+\varepsilon} Y^{-1/4}. \end{aligned} \tag{3.4}$$

By Lemma 2.2, (2.8) of Lemma 2.5 and Hölder's inequality we get

$$\begin{aligned} P_{13}(T) &\ll \left(\int_T^{2T} \Delta_{31}^4(x; y) dx \right)^{1/4} \left(\int_T^{2T} 1 dt \right)^{1/4} \left(\int_T^{2T} \Delta_{22}^4(x; Y) dx \right)^{1/2} \\ &\ll T^{11/6+\varepsilon} Y^{-1/6}. \end{aligned} \quad (3.5)$$

We now bound $P_{11}(T)$. By the elementary formula

$$\cos \alpha_1 \cos \alpha_2 \cos \alpha_3 = \frac{1}{4} \sum_{(j_1, j_2) \in \{0, 1\}^2} \cos [\alpha_1 + (-1)^{j_1} \alpha_2 + (-1)^{j_2} \alpha_3]$$

we can write

$$\begin{aligned} &\Delta_{31}(x; y)(\Delta_{21}(x; Y))^2 \\ &= \frac{x^{5/6}}{2\sqrt{3}\pi^3} \sum_{(j_1, j_2) \in \{0, 1\}^2} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}} m_1^{\frac{3}{4}} m_2^{\frac{3}{4}}} \\ &\quad \times \cos \left[6\pi(nx)^{1/3} + (-1)^{j_1} 4\pi(xm_1)^{1/2} \right. \\ &\quad \left. + (-1)^{j_2} 4\pi(xm_2)^{1/2} - \frac{\pi}{4}((-1)^{j_1} + (-1)^{j_2}) \right] \\ &= S_1(x) + S_2(x) - S_3(x), \end{aligned}$$

where

$$\begin{aligned} S_1(x) &:= \frac{x^{5/6}}{2\sqrt{3}\pi^3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}} m_1^{\frac{3}{4}} m_2^{\frac{3}{4}}} \\ &\quad \times \sin \left[6\pi(nx)^{1/3} + 4\pi(xm_1)^{1/2} + 4\pi(xm_2)^{1/2} \right], \\ S_2(x) &:= \frac{x^{5/6}}{2\sqrt{3}\pi^3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}} m_1^{\frac{3}{4}} m_2^{\frac{3}{4}}} \\ &\quad \times \sin \left[6\pi(nx)^{1/3} - 4\pi(xm_1)^{1/2} - 4\pi(xm_2)^{1/2} \right], \\ S_3(x) &:= \frac{x^{5/6}}{\sqrt{3}\pi^3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}} m_1^{\frac{3}{4}} m_2^{\frac{3}{4}}} \\ &\quad \times \cos \left[6\pi(nx)^{1/3} - 4\pi(xm_1)^{1/2} + 4\pi(xm_2)^{1/2} \right]. \end{aligned}$$

Let

$$f_1(x) = 6\pi(nx)^{1/3} + 4\pi(xm_1)^{1/2} + 4\pi(xm_2)^{1/2}.$$

Then for any $T \leq x \leq 2T$, we have

$$f_1'(x) \gg \frac{m_1^{1/2} + m_2^{1/2}}{T^{1/2}}.$$

By Lemma 2.8, Lemma 2.9 and the inequality $a^2 + b^2 \geq 2ab$ we have

$$\begin{aligned} \int_T^{2T} S_1(x) dx &\ll T^{4/3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{2/3}m_1^{3/4}m_2^{3/4}(m_1^{1/2} + m_2^{1/2})} \\ &\ll T^{4/3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{2/3}m_1m_2} \\ &\ll T^{4/3}y^{1/3} \log^6 T. \end{aligned} \quad (3.6)$$

Let

$$f_2(x) = 6\pi(nx)^{1/3} - 4\pi(xm_1)^{1/2} - 4\pi(xm_2)^{1/2}.$$

Then

$$f_2'(x) = 2\pi \left(\frac{n^{1/3}}{x^{2/3}} - \frac{m_1^{1/2} + m_2^{1/2}}{x^{1/2}} \right).$$

Since $y \ll T^{1/3}$, it is easy to see that for any $n \leq y$, $m_1 \leq Y$, $m_2 \leq Y$, and $T \leq x \leq 2T$ we have

$$|f_2'(x)| \gg \frac{m_1^{1/2} + m_2^{1/2}}{x^{1/2}}.$$

So similar to $S_1(x)$, by Lemma 2.8 and Lemma 2.9 we have

$$\begin{aligned} \int_T^{2T} S_2(x) dx &\ll T^{4/3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{2/3}m_1^{3/4}m_2^{3/4}(m_1^{1/2} + m_2^{1/2})} \\ &\ll T^{4/3}y^{1/3} \log^6 T. \end{aligned} \quad (3.7)$$

Now we consider the contribution of $S_3(x)$. We write

$$S_3(x) = S_{31}(x) + S_{32}(x) + S_{33}(x), \quad (3.8)$$

where

$$\begin{aligned}
S_{31}(x) &:= \frac{x^{5/6}}{\sqrt{3}\pi^3} \sum_{n \leq y} \sum_{m \leq Y} \frac{d_3(n)d_2^2(m)}{n^{\frac{2}{3}}m^{\frac{3}{2}}} \cos \left[6\pi(nx)^{1/3} \right], \\
S_{32}(x) &:= \frac{x^{5/6}}{\sqrt{3}\pi^3} \sum_{n \leq y} \sum_{\substack{m_1 \leq Y, m_2 \leq Y \\ m_1 < m_2}} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}}m_1^{\frac{3}{4}}m_2^{\frac{3}{4}}} \\
&\quad \times \cos \left[6\pi(nx)^{1/3} - 4\pi(xm_1)^{1/2} + 4\pi(xm_2)^{1/2} \right], \\
S_{33}(x) &:= \frac{x^{5/6}}{\sqrt{3}\pi^3} \sum_{n \leq y} \sum_{\substack{m_1 \leq Y, m_2 \leq Y \\ m_1 > m_2}} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}}m_1^{\frac{3}{4}}m_2^{\frac{3}{4}}} \\
&\quad \times \cos \left[6\pi(nx)^{1/3} - 4\pi(xm_1)^{1/2} + 4\pi(xm_2)^{1/2} \right].
\end{aligned}$$

Obviously, by Lemma 2.8 and Lemma 2.9 we have

$$\int_T^{2T} S_{31}(x) dx \ll T^{3/2} \log^3 T. \quad (3.9)$$

Let

$$f_3(x) = 6\pi(nx)^{1/3} - 4\pi(xm_1)^{1/2} + 4\pi(xm_2)^{1/2}.$$

Then

$$f'_3(x) = 2\pi \left(\frac{n^{1/3}}{x^{2/3}} - \frac{m_1^{1/2} - m_2^{1/2}}{x^{1/2}} \right).$$

If $m_1 < m_2$, it is easy to see that for any $T \leq x \leq 2T$ we have

$$|f'_3(x)| \gg \frac{|m_2^{1/2} - m_1^{1/2}|}{T^{1/2}}.$$

So by Lemma 2.8 and Lemma 2.10 we have

$$\begin{aligned}
\int_T^{2T} S_{32}(x) dx &\ll T^{4/3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{2}{3}}m_1^{\frac{3}{4}}m_2^{\frac{3}{4}}|m_1^{1/2} - m_2^{1/2}|} \\
&\ll T^{4/3} y^{1/3} \log^6 T.
\end{aligned} \quad (3.10)$$

Finally, we consider the contribution of $S_{33}(x)$. By a splitting argument, $S_{33}(x)$ can be written as a sum of $O(\log^3 T)$ terms of the form

$$J(x; N, M_1, M_2) := \frac{x^{5/6}}{\sqrt{3}\pi^3} \sum_{n \sim N} \sum_{\substack{m_1 \sim M_1, m_2 \sim M_2 \\ m_1 > m_2}} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{2/3}m_1^{3/4}m_2^{3/4}} \cos f_3(x).$$

We write further that

$$J(x; N, M_1, M_2) = J_1(x; N, M_1, M_2) + J_2(x; N, M_1, M_2) + J_3(x; N, M_1, M_2),$$

where

$$\begin{aligned} SC(J_1) : n \sim N, \quad m_1 \sim M_1, \quad m_2 \sim M_2, \\ m_1 > m_2, \quad n^{1/3} > 10T^{1/6}(m_1^{1/2} - m_2^{1/2}), \\ SC(J_2) : n \sim N, \quad m_1 \sim M_1, \quad m_2 \sim M_2, \\ m_1 > m_2, \quad n^{1/3} < \frac{1}{10}T^{1/6}(m_1^{1/2} - m_2^{1/2}), \\ SC(J_3) : n \sim N, \quad m_1 \sim M_1, \quad m_2 \sim M_2, \\ m_1 > m_2, \quad n^{1/3} \asymp T^{1/6}(m_1^{1/2} - m_2^{1/2}). \end{aligned}$$

For J_1 , we have $|f_3(x)| \gg n^{1/3}T^{-2/3}$. So by Lemma 2.8 and Lemma 2.9 we get

$$\begin{aligned} \int_T^{2T} J_1(x; N, M_1, M_2) dx &\ll T^{3/2} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{nm_1^{3/4}m_2^{3/4}} \\ &\ll T^{3/2}Y^{1/2} \log^6 T. \end{aligned} \quad (3.11)$$

For J_2 , we have $|f_3(x)| \gg |m_1^{1/2} - m_2^{1/2}|/T^{1/2}$. By Lemma 2.8 and Lemma 2.10 we get

$$\begin{aligned} \int_T^{2T} J_2(x; N, M_1, M_2) dx &\ll T^{4/3} \sum_{n \leq y} \sum_{m_1 \leq Y} \sum_{m_2 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{2/3}m_1^{3/4}m_2^{3/4} |m_1^{1/2} - m_2^{1/2}|} \\ &\ll T^{4/3}y^{1/3} \log^6 T. \end{aligned} \quad (3.12)$$

Now we consider the contribution of $J_3(x; N, M_1, M_2)$. We only need to bound the integral $\int_T^{2T} x^{5/6} \cos f_3(x) dx$. Suppose $\eta > 0$ is a small parameter to be determined later. Let

$$I_1 = \{T \leq x \leq 2T : |f_3(x)| > \eta\}, \quad I_2 = \{T \leq x \leq 2T : |f_3(x)| \leq \eta\}.$$

Without loss of generality, we suppose that both I_1 and I_2 are not empty.

Suppose $x \in I_2$. Then we have $|n^{1/3} - x^{1/6}(m_1^{1/2} - m_2^{1/2})| \leq \eta(2T)^{2/3}$, which implies that

$$x^{1/6} = \frac{n^{1/3}}{m_1^{1/2} - m_2^{1/2}} + O\left(\frac{\eta T^{2/3}}{m_1^{1/2} - m_2^{1/2}}\right).$$

Obviously I_2 is a closed interval. Let $I_2 = [x_1, x_2]$. By the Lagrange theorem, we get

$$x_2 - x_1 \ll \frac{\eta T^{3/2}}{(m_1^{1/2} - m_2^{1/2})},$$

which implies that

$$\int_{I_2} x^{5/6} \cos f_3(x) dx \ll T^{5/6}(x_2 - x_1) \ll \frac{\eta T^{14/6}}{m_1^{1/2} - m_2^{1/2}}.$$

By Lemma 2.8 we have

$$\int_{I_1} x^{5/6} \cos f_3(x) dx \ll \frac{T^{5/6}}{\eta}.$$

So we get by choosing $\eta = |m_1^{1/2} - m_2^{1/2}|^{1/2}/T^{3/4}$ that

$$\int_T^{2T} x^{5/6} \cos f_3(x) dx \ll \frac{T^{19/12}}{(m_1^{1/2} - m_2^{1/2})^{1/2}}.$$

Since $n^{1/3} \asymp T^{1/6}|m_1^{1/2} - m_2^{1/2}|$, the above bound becomes

$$\int_T^{2T} x^{5/6} \cos f_3(x) dx \ll \frac{T^{5/3}}{n^{1/6}}, \quad (3.13)$$

which implies that

$$\begin{aligned} \int_T^{2T} J_3(x; N, M_1, M_2) dx &\ll T^{5/3} \sum_{n \sim N} \sum_{\substack{m_1 \sim M_1, m_2 \sim M_2 \\ |m_1^{1/2} - m_2^{1/2}| \asymp \frac{n^{1/3}}{T^{1/6}}}} \frac{d_3(n)d_2(m_1)d_2(m_2)}{n^{\frac{5}{6}}m_1^{\frac{3}{4}}m_2^{\frac{3}{4}}} \\ &\ll \frac{T^{5/3}}{N^{5/6}M_1^{3/2}} \sum_{n \sim N} d_3(n) \sum_{\substack{m_1 \sim M_1, m_2 \sim M_2 \\ |m_1^{1/2} - m_2^{1/2}| \asymp \frac{n^{1/3}}{T^{1/6}}}} d_2(m_1)d_2(m_2). \end{aligned} \quad (3.14)$$

It suffices for us to bound

$$U = \sum_{n \sim N} d_3(n) \sum_{\substack{m_1 \sim M_1, m_2 \sim M_2 \\ |m_1^{1/2} - m_2^{1/2}| \asymp \frac{n^{1/3}}{T^{1/6}}}} d_2(m_1) d_2(m_2).$$

It is easy to see that if $|m_1^{1/2} - m_2^{1/2}| \asymp \frac{n^{1/3}}{T^{1/6}}$, then for fixed m_1 , the number of m_2 's is $\ll 1 + \frac{n^{1/3} m_1^{1/2}}{T^{1/6}}$. By the symmetry of m_1 and m_2 , Cauchy's inequality and Lemma 2.9 we have

$$\begin{aligned} U &\ll \sum_{n \sim N} d_3(n) \sum_{m_1 \sim M_1} d_2^2(m_1) \sum_{\substack{m_2 \sim M_2 \\ |m_1^{1/2} - m_2^{1/2}| \asymp \frac{n^{1/3}}{T^{1/6}}}} 1 \\ &\ll \sum_{n \sim N} d_3(n) \sum_{m_1 \sim M_1} d_2^2(m_1) \left(1 + \frac{n^{1/3} m_1^{1/2}}{T^{1/6}} \right) \\ &\ll N M_1 \log^5 T + \frac{N^{4/3}}{T^{1/6}} M_1^{3/2} \log^5 T. \end{aligned} \quad (3.15)$$

Inserting (3.15) into (3.14) we get

$$\begin{aligned} \int_T^{2T} J_3(x; N, M_1, M_2) dx &\ll \frac{T^{5/3}}{N^{5/6} M_1^{3/2}} \left(N M_1 \log^5 T + \frac{N^{4/3}}{T^{1/6}} M_1^{3/2} \log^5 T \right) \\ &\ll T^{5/3} y^{1/6} \log^5 T + T^{3/2} y^{1/2} \log^5 T. \end{aligned} \quad (3.16)$$

From (3.11), (3.12) and (3.16) we have

$$\int_T^{2T} S_{33}(x) dx \ll K(T, y, Y),$$

which combining with (3.8)-(3.10) gives

$$\int_T^{2T} S_3(x) dx \ll K(T, y, Y), \quad (3.17)$$

where

$$\begin{aligned} K(T, y, Y) &:= T^{5/3} y^{1/6} \log^9 T + T^{3/2} y^{1/2} \log^9 T \\ &\quad + T^{4/3} y^{1/3} \log^6 T + T^{3/2} Y^{1/2} \log^4 T. \end{aligned}$$

From (3.6), (3.7) and (3.17) we obtain

$$P_{11}(T) \ll K(T, y, Y). \quad (3.18)$$

From (3.1), (3.2) and (3.18) we get by choosing $y = Y = T^{1/3}$ that

$$P(T) \ll K(T, y, Y) + T^{11/6+\varepsilon}Y^{-1/6} + T^{11/6+\varepsilon}y^{-1/6} \ll T^{16/9+\varepsilon}, \quad (3.19)$$

which combining with a splitting argument proves Theorem 1.1.

4. Proof of Theorem 1.2

We shall estimate the integral

$$Q(T) := \int_T^{2T} \Delta_3(x) \Delta_2^3(x) dx.$$

Suppose $T^\varepsilon \ll y \ll T^{1/3}$ is a parameter to be determined later. Write

$$Q(T) = Q_1(T) + Q_2(T), \quad (4.1)$$

where

$$Q_1(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_2^3(x) dx, \quad Q_2(T) := \int_T^{2T} \Delta_{32}(x; y) \Delta_2^3(x) dx.$$

By Lemma 2.3 and (1.3) with $A_0 = 6$ we have

$$Q_2(T) \ll T^{25/12+\varepsilon}y^{-1/6}. \quad (4.2)$$

It suffices for us to bound $Q_1(T)$. Suppose $T^\varepsilon \ll Y \ll T^{1/3}$ is a parameter to be determined later. We write

$$Q_1(T) = Q_{11}(T) + 3Q_{12}(T) + 3Q_{13}(T) + Q_{14}(T), \quad (4.3)$$

where

$$Q_{11}(T) := \int_T^{2T} \Delta_{31}(x; y) (\Delta_{21}(x; Y))^3 dx,$$

$$Q_{12}(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_{21}^2(x; Y) \Delta_{22}(x; Y) dx,$$

$$Q_{13}(T) := \int_T^{2T} \Delta_{31}(x; y) \Delta_{21}(x; Y) (\Delta_{22}(x; Y))^2 dx,$$

$$Q_{14}(T) := \int_T^{2T} \Delta_{31}(x; y) (\Delta_{22}(x; Y))^3 dx.$$

By Lemma 2.2, (2.5) of Lemma 2.4, (2.7) of Lemma 2.5 and Hölder's inequality, we have

$$\begin{aligned} Q_{12}(T) &\ll \left(\int_T^{2T} \Delta_{31}^4(x; y) dx \right)^{\frac{1}{4}} \left(\int_T^{2T} \Delta_{21}^8(x; Y) dx \right)^{\frac{1}{4}} \left(\int_T^{2T} \Delta_{22}^2(x; Y) dx \right)^{\frac{1}{2}} \\ &\ll T^{25/12+\varepsilon} Y^{-1/4}. \end{aligned} \quad (4.4)$$

By Lemma 2.2, (2.4) of Lemma 2.4, (2.8) of Lemma 2.5, as well as Hölder's inequality, we have

$$\begin{aligned} Q_{13}(T) &\ll \left(\int_T^{2T} \Delta_{31}^4(x; y) dx \right)^{\frac{1}{4}} \left(\int_T^{2T} \Delta_{21}^4(x; Y) dx \right)^{\frac{1}{4}} \left(\int_T^{2T} \Delta_{22}^4(x; Y) dx \right)^{\frac{1}{2}} \\ &\ll T^{25/12+\varepsilon} Y^{-1/6}. \end{aligned} \quad (4.5)$$

By Lemma 2.2, (2.8) of Lemma 2.5 and Hölder's inequality, we have

$$\begin{aligned} Q_{14}(T) &\ll \left(\int_T^{2T} \Delta_{31}^4(x; y) dx \right)^{\frac{1}{4}} \left(\int_T^{2T} \Delta_{22}^4(x; Y) dx \right)^{\frac{3}{4}} \\ &\ll T^{25/12+\varepsilon} Y^{-1/4}. \end{aligned} \quad (4.6)$$

It suffices to bound $Q_1(T)$. By the elementary formula

$$\prod_{l=1}^4 \cos \alpha_l = \frac{1}{8} \sum_{(j_1, j_2, j_3) \in \{0,1\}^3} \cos [(-1)^{j_1} \alpha_1 + (-1)^{j_2} \alpha_2 + (-1)^{j_3} \alpha_3 + \alpha_4],$$

we have

$$\begin{aligned} &\Delta_{31}(x; y) (\Delta_{21}(x; Y))^3 \\ &= \frac{x^{13}}{2\sqrt{6} \pi^4} \sum_{n \leq y} \sum_{m_1, m_2, m_3 \leq Y} \frac{d_3(n) d_2(m_1) d_2(m_2) d_2(m_3)}{n^{\frac{2}{3}} m_1^{\frac{3}{4}} m_2^{\frac{3}{4}} m_3^{\frac{3}{4}}} \cos(6\pi(xn)^{1/3}) \\ &\quad \times \cos \left[4\pi(xm_1)^{1/2} - \frac{\pi}{4} \right] \cos \left[4\pi(xm_2)^{1/2} - \frac{\pi}{4} \right] \cos \left[4\pi(xm_3)^{1/2} - \frac{\pi}{4} \right] \\ &= \frac{x^{13/12}}{16\sqrt{6} \pi^4} \sum_{(j_1, j_2, j_3) \in \{0,1\}^3} \sum_{n \leq y} \sum_{m_1, m_2, m_3 \leq Y} \frac{d_3(n) d_2(m_1) d_2(m_2) d_2(m_3)}{n^{\frac{2}{3}} m_1^{\frac{3}{4}} m_2^{\frac{3}{4}} m_3^{\frac{3}{4}}} \\ &\quad \times \cos f(x; n, \mathbf{m}, \mathbf{j}), \end{aligned} \quad (4.7)$$

where $\mathbf{m} = (m_1, m_2, m_3)$, $\mathbf{j} = (j_1, j_2, j_3)$, and

$$\begin{aligned} f(x; n, \mathbf{m}, \mathbf{j}) &= 6\pi(xn)^{1/3} + (-1)^{j_1}4\pi(xm_1)^{1/2} \\ &\quad + (-1)^{j_2}4\pi(xm_2)^{1/2} + (-1)^{j_3}4\pi(xm_3)^{1/2} \\ &\quad + \frac{\pi}{4}((-1)^{j_1+1} + (-1)^{j_2+1} + (-1)^{j_3+1}). \end{aligned} \quad (4.8)$$

For each \mathbf{j} , we shall estimate the integral

$$H(T; \mathbf{j}) := \int_T^{2T} x^{13/12} \sum_{n \leq y} \sum_{\mathbf{m}} \frac{d_3(n)d_2(m_1)d_2(m_2)d_2(m_3)}{n^{\frac{2}{3}}m_1^{\frac{3}{4}}m_2^{\frac{3}{4}}m_3^{\frac{3}{4}}} \cos f(x; n, \mathbf{m}, \mathbf{j}) dx.$$

Since $y \ll T^{1/3}$, it is easy to see that if $\mathbf{j} = (0, 0, 0)$ or $\mathbf{j} = (1, 1, 1)$, we have

$$|f'(x; n, \mathbf{m}, \mathbf{j})| \gg m_1^{1/2} + m_2^{1/2} + m_3^{1/2}.$$

By Lemma 2.8 and Lemma 2.6 we get

$$\begin{aligned} H(T; \mathbf{j}) &\ll T^{19/12} \sum_{n \leq y} \sum_{m_1, m_2, m_3 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)d_2(m_3)}{n^{\frac{2}{3}}m_1^{\frac{3}{4}}m_2^{\frac{3}{4}}m_3^{\frac{3}{4}}(m_1^{1/2} + m_2^{1/2} + m_3^{1/2})} \\ &\ll T^{19/12}y^{1/3}Y^{1/4} \log^6 T. \end{aligned} \quad (4.9)$$

Now we suppose $\mathbf{j} \neq (0, 0, 0)$ and $\mathbf{j} \neq (1, 1, 1)$. Let

$$\alpha = (-1)^{j_1}(m_1)^{1/2} + (-1)^{j_2}(m_2)^{1/2} + (-1)^{j_3}(m_3)^{1/2}.$$

Then $f'(x; n, \mathbf{m}, \mathbf{j}) = 2\pi(n^{1/3}/x^{2/3} + \alpha/x^{1/2})$.

If $n^{1/3} \geq 10T^{1/6}|\alpha|$, then by Lemma 2.8 we have

$$\int_T^{2T} x^{13/12} \cos f(x; n, \mathbf{m}, \mathbf{j}) dx \ll T^{7/4}n^{-1/3},$$

whose contribution to $H(T; \mathbf{j})$ is

$$\begin{aligned} &\ll T^{7/4} \sum_{n \leq y} \sum_{m_1, m_2, m_3 \leq Y} \frac{d_3(n)d_2(m_1)d_2(m_2)d_2(m_3)}{nm_1^{\frac{3}{4}}m_2^{\frac{3}{4}}m_3^{\frac{3}{4}}} \\ &\ll T^{7/4}Y^{3/4} \log^6 T. \end{aligned} \quad (4.10)$$

If $n^{1/3} \leq \frac{1}{10}T^{1/6}|\alpha|$, then by Lemma 2.8 again we have

$$\int_T^{2T} x^{13/12} \cos f(x; n, \mathbf{m}, \mathbf{j}) dx \ll T^{19/12}/|\alpha|.$$

whose contribution to $H(T; \mathbf{j})$ is (via Lemma 2.6)

$$\begin{aligned} &\ll T^{19/12} \sum_{n \leq y} \sum_{m_1, m_2, m_3 \leq Y} \frac{d_3(n) d_2(m_1) d_2(m_2) d_2(m_3)}{n^{2/3} m_1^{3/4} m_2^{3/4} m_3^{3/4} |\alpha|} \\ &\ll T^{19/12+\varepsilon} y^{1/3} Y^{1/4}. \end{aligned} \quad (4.11)$$

Now suppose $n^{1/3} \asymp T^{1/6}|\alpha|$, then similarly to (3.13) we have the estimate

$$\int_T^{2T} x^{13/12} \cos f(x; n, \mathbf{m}, \mathbf{j}) dx \ll \frac{T^{23/12}}{|\alpha|^{1/6}},$$

whose contribution of $H(T; \mathbf{j})$ is

$$\ll T^{23/12} \sum_{n \leq y} \frac{d_3(n)}{n^{5/6}} G(T, n), \quad (4.12)$$

where

$$G(T, n) := \sum_{\substack{m_1, m_2, m_3 \leq Y \\ |\alpha| \leq n^{1/3} T^{-1/6}}} \frac{d_2(m_1) d_2(m_2) d_2(m_3)}{m_1^{3/4} m_2^{3/4} m_3^{3/4}}.$$

By a splitting argument and Lemma 2.7 we get (suppose $M_1 \ll M_2 \ll M_3 \ll Y$)

$$\begin{aligned} G(T, n) &\ll \log^3 Y \sum_{\substack{m_1 \sim M_1, m_2 \sim M_2, m_3 \sim M_3 \\ |\alpha| \leq n^{1/3} T^{-1/6}}} \frac{d_2(m_1) d_2(m_2) d_2(m_3)}{m_1^{3/4} m_2^{3/4} m_3^{3/4}} \\ &\ll \frac{Y^\varepsilon}{M_1^{3/4} M_2^{3/4} M_3^{3/4}} \times \left(\frac{n^{1/3}}{T^{1/6}} M_1 M_2 M_3^{1/2} + M_1^{1/2} M_2^{1/2} \right) \\ &\ll \frac{n^{1/3} Y^{1/4+\varepsilon}}{T^{1/6}} + \frac{Y^\varepsilon}{M_1^{1/4} M_2^{1/4} M_3^{3/4}} \\ &\ll \frac{n^{1/3} Y^{1/4+\varepsilon}}{T^{1/6}} + \frac{Y^\varepsilon}{(M_1 M_2 M_3)^{5/12}}. \end{aligned}$$

By Lemma 2.11 we have $|\alpha| \gg 1/\sqrt{M_1 M_2 M_3}$, which combining with $|\alpha| \leq n^{1/3} T^{-1/6}$ implies that

$$\frac{1}{(M_1 M_2 M_3)^{5/12}} \ll \frac{n^{5/18}}{T^{5/36}}.$$

So we get

$$G(T, n) \ll \frac{n^{1/3} Y^{1/4+\varepsilon}}{T^{1/6}} + \frac{n^{5/18} Y^\varepsilon}{T^{5/36}}.$$

Inserting the above bound into (4.12) we get via Lemma 2.9 that the contribution of the case $n^{1/3} \asymp T^{1/6} |\alpha|$ is

$$\ll T^{7/4+\varepsilon} y^{1/2} Y^{1/4} + T^{16/9+\varepsilon} y^{4/9}. \quad (4.13)$$

From (4.10), (4.11) and (4.13) we see that if $\mathbf{j} \neq (0, 0, 0)$ and $\mathbf{j} \neq (1, 1, 1)$, then we have

$$H(T; \mathbf{j}) \ll L(T, y, Y) T^\varepsilon \quad (4.14)$$

by noting that $T^{19/12} y^{1/3} T^{1/4} \ll T^{7/4} y^{1/2} Y^{1/4}$, where

$$L(T, y, Y) := T^{7/4} y^{1/2} Y^{1/4} + T^{16/9} y^{4/9} + T^{7/4} Y^{3/4}.$$

From (4.7), (4.9) and (4.14) we get

$$Q_{11}(T) \ll L(T, y, Y) T^\varepsilon,$$

which combining with (4.4)–(4.6) gives

$$Q_1(T) \ll L(T, y, Y) T^\varepsilon + T^{25/12+\varepsilon} Y^{-1/6}. \quad (4.15)$$

From (4.1), (4.2) and (4.15) we get by choosing $y = Y = T^{1/3}$ that

$$\begin{aligned} Q(T) &\ll L(T, y, Y) T^\varepsilon + T^{25/12+\varepsilon} Y^{-1/6} + T^{25/12+\varepsilon} y^{-1/6} \\ &\ll T^{73/36+\varepsilon}, \end{aligned}$$

which combining with a splitting argument proves Theorem 1.2.

Acknowledgement. This work is supported by the National Natural Science Foundation of China(Grant No. 11971476).

REFERENCES

- [1] J. Bourgain, N. Watt, *Mean square of zeta function, circle problem and divisor problem revisited*, preprint available at arXiv:1709.04340.
- [2] X. Cao, Y. Tanigawa, W. Zhai, *Tong-type identity and the mean square of the error term for an extended Selberg class*, Science China Mathematics **59** (2016), No. 11, 2103–2144.
- [3] H. Cramér, *Über zwei Sätze von Herrn G. H. Hardy*, Math. Z. **15** (1922), 201–210.
- [4] D.R. Heath-Brown, *The distribution and moments of the error term in the Dirichlet divisor problem*, Acta Arith. **60**(1992), 389–415.
- [5] A. Ivić, *The Riemann Zeta-Function*, John Wiley and Sons, 1985.
- [6] A. Ivić, P. Sargos, *On the higher moments of the error term in the divisor problem*, Illinois J Math, **51** (2007), 353–377.
- [7] A. Ivić, W. Zhai, *On certain integrals involving the Dirichlet divisor problem*, Functiones et Approximatio **62.2** (2020), 247–267.
- [8] G. Kolesnik, *On the estimation of multiple exponential sums*, Recent Progress in Analytic Number Theory, Vol. 1 (Durham, 1979), pp. 231–246, Academic Press, London-New York, 1981.
- [9] K. L. Kong, *Some mean value theorems for certain error terms in analytic number theory*, Master Degree Thesis, The University of Hong Kong, 2014.
- [10] Y. K. Lau, K-M Tsang, *On the mean square formula of the error term in the Dirichlet divisor problem*, Mathematical Proceedings of the Cambridge Philosophical Society, 2009, 146: 277–287.
- [11] J. Li, *On the seventh power moment of $\Delta(x)$* , Int. J. Number Theory **13** (2017), no. 3, 571–591.
- [12] E. Preissmann, *Sur la moyenne quadratique du terme de reste du problème du cercle*, C R Acad Sci Paris Sér I, 1988, 306: 151–154.
- [13] O. Robert, P. Sargos, *Three-dimensional exponential sums with monomials*, J. Reine Angew. Math. **591** (2006), 1–20.
- [14] K. C. Tong, *On divisor problem III*, Acta Math. Sinica **6** (1956), 515–541.
- [15] K.-M. Tsang, *Higher-power moments of $\Delta(x)$, $E(t)$ and $P(x)$* , Proc. London Math. Soc. (3) **65** (1992), 65–84.

- [16] J. Wang, *On the sixth-power moment of $\Delta(x)$* , Master Degree Thesis, Shandong Normal University, 2010.
- [17] W. Zhai, *On higher-power moments of $\Delta(x)$ (II)*, Acta Arith. **114** (2004), 35–54.
- [18] W. Zhai, *On higher-power moments of $E(t)$* , Acta Arith. **115** (2004), no. 4, 329–348.
- [19] W. Zhai, *On higher-power moments of $\Delta(x)$ (III)*, Acta Arith. **118** (2005), 263–281.
- [20] D. Zhang, W. Zhai, *On the fifth-power moment of $\Delta(x)$* , Int. J. Number Theory **7** (2011), no. 1, 71–86.

Department of Mathematics
China University of Mining and Technology
Beijing 100083, P. R. China
e-mail: zhaiwg@hotmail.com