

## SEMI-FREDHOLM OPERATORS AND PERTURBATION FUNCTIONS

Snežana Živković–Zlatanović

**Abstract.** In this paper we give several remarks on [3] and a different proof of the inequalities in [8]. Also we introduce the concept of a lower perturbation function, and prove that some usual measures of noncompactness of operators and also some measures of non-strict-cosingularity of operators are lower perturbation functions. For each lower perturbation function  $\delta$ , we define functions  $\nabla_\delta$  and  $K_\delta$ , and give the connection with lower semi-Fredholm operators.

### 1. Introduction and preliminaries

In this paper  $X$ ,  $Y$  and  $Z$  are complex Banach spaces,  $B(X, Y)$  (respectively  $K(X, Y)$ ) the set of all bounded (respectively compact) linear operators from  $X$  into  $Y$ . We shall write  $B(X)$  ( $K(X)$ ) instead of  $B(X, X)$  ( $K(X, X)$ ).

An operator  $T \in B(X, Y)$  is in  $\Phi_+(X, Y)$  ( $\Phi_-(X, Y)$ ) if the range  $R(T)$  is closed and the dimension  $\alpha(T)$  of the null space  $N(T)$  of  $T$  is finite (the codimension  $\beta(T)$  of  $R(T)$  in  $Y$  is finite). Operators in  $\Phi_+(X, Y) \cup \Phi_-(X, Y)$  are called semi-Fredholm operators. For such operators the index,  $i(T)$ , is defined by  $i(T) = \alpha(T) - \beta(T)$ . We set  $\Phi(X, Y) = \Phi_+(X, Y) \cap \Phi_-(X, Y)$ . The operators in  $\Phi(X, Y)$  are called Fredholm. We shall write  $\Phi_+(X)$  (resp.  $\Phi_-(X)$ ,  $\Phi(X)$ ) instead of  $\Phi_+(X, X)$  (resp.  $\Phi_-(X, X)$ ,  $\Phi(X, X)$ ).

Recall that the essential spectral radius of  $T \in B(X)$ ,  $r_e(T)$ , is defined by

$$r_e(T) = \max\{|\lambda| : T - \lambda I \notin \Phi(X)\}.$$

Let  $B_X$  denote the closed unit ball of  $X$ . Let  $T \in B(X, Y)$  and

$$m(T) = \inf\{\|Tx\| : \|x\| = 1\}$$

be the *minimum modulus* of  $T$ , and let

$$n(T) = \sup\{\epsilon \geq 0 : \epsilon B_Y \subset TB_X\}$$

---

Received May 7, 1997

1991 *Mathematics Subject Classification*: 47A10, 47A53, 47A55.

Supported by Grant 04M03 of RFNS through Math. Inst. SANU

be the *surjection modulus* of  $T$ . Recall that  $m(T^*) = n(T)$ , where  $T^* \in B(Y^*, X^*)$  is the adjoint operator.

If  $M$  is a subspace of  $X$ , then  $J_M$  will denote the embedding map of  $M$  into  $X$ , and if  $V$  is a subspace of  $Y$ , then  $Q_V$  will denote the canonical map of  $Y$  onto the quotient space  $Y/V$ .

An operator  $T \in B(X, Y)$  is *strictly singular* ( $T \in S(X, Y)$ ) if, for every infinite dimensional (closed) subspace  $M$  of  $X$ , the restriction of  $T$  to  $M$ ,  $T|_M$ , is not a homeomorphism, i.e.  $m(T|_M) = 0$ . An operator  $T \in B(X, Y)$  is *strictly cosingular* ( $T \in CS(X, Y)$ ) if, for every infinite codimensional closed subspace  $V$  of  $Y$  the composition  $Q_V T$  is not surjective.

For  $A \in B(X, Y)$ , set

$$\|A\|_C = \inf\{\|A + K\| : K \in K(X, Y)\}.$$

If  $\Omega$  is a non-empty bounded subset of  $X$ , then the Kuratowski measure of noncompactness of  $\Omega$  is denoted by  $\alpha(\Omega)$ , and

$$\alpha(\Omega) = \inf\{\epsilon > 0 : \Omega \subset \cup_{i=1}^n D_i, D_i \subset X, \text{diam} D_i \leq \epsilon\}.$$

For  $A \in B(X, Y)$  the Kuratowski measure of noncompactness of  $A$  is denoted by  $\|A\|_\alpha$  and defined by

$$\|A\|_\alpha = \inf\{k \geq 0 : \alpha(A\Omega) \leq k\alpha(\Omega), \Omega \subset X \text{ is bounded}\}.$$

It is easy to see that  $\|A\|_\alpha = \sup\{\alpha(A\Omega) : \Omega \subset X, \alpha(\Omega) = 1\}$ .

If  $\Omega$  is a non-empty bounded subset of  $X$ , then the Hausdorff measure of noncompactness of  $\Omega$ , is denoted by  $q(\Omega)$ , and

$$q(\Omega) = \inf\{\epsilon > 0 : \Omega \text{ has a finite } \epsilon\text{-net in } X\}.$$

For  $A \in B(X, Y)$  the Hausdorff measure of noncompactness of  $A$  is denoted by  $\|A\|_q$  and defined by

$$\|A\|_q = \inf\{k \geq 0 : q_Y(A\Omega) \leq kq_X(\Omega), \Omega \subset X \text{ is bounded}\}.$$

Recall that  $\|A\|_q = q_Y(AB_X)$ .

For  $A \in B(X, Y)$ , set (see [6])

$$\|A\|_\mu = \inf\{\|A|_L\| : L \text{ subspace of } X, \text{codim} L < \infty\},$$

and

$$\begin{aligned} \Gamma_M(A) &= \inf_{N \subset M} \|A|_N\|, & \Gamma(A) &= \Gamma_X(A), \\ \Delta_M(A) &= \sup_{N \subset M} \Gamma_N(A), & \Delta(A) &= \Delta_X(A), \end{aligned}$$

where  $M, N$  denote infinite dimensional subspaces of  $X$  (see [9]). Schechter [9] proved that  $\Delta$  is a submultiplicative seminorm and

$$(1.1) \quad \Delta(A) = 0 \iff A \in S(X, Y).$$

For  $A \in B(X, Y)$ , set (see [12])

$$\begin{aligned} K_V(A) &= \inf_{W \supset V} \|Q_W A\|, & K(A) &= K_{\{0\}}(A), \\ \nabla_V(A) &= \sup_{W \supset V} K_W(A), & \nabla(A) &= \nabla_{\{0\}}(A), \end{aligned}$$

where  $V, W$  denote closed infinite codimensional subspaces of  $Y$ .

Recall that [12],

$$(1.2) \quad \begin{aligned} K(A) > 0 &\iff A \in \Phi_-(X, Y), \\ \nabla(A) = 0 &\iff A \in CS(X, Y). \end{aligned}$$

Zemánek [13] considered the following functions

$$\begin{aligned} u(A) &= \sup\{m(A|_W) : W \text{ is closed subspace of } X \text{ with } \dim W = \infty\}, \\ v(A) &= \sup\{n(Q_V A) : V \text{ is closed subspace of } Y \text{ with } \operatorname{codim} V = \infty\}. \end{aligned}$$

From the definition of the strictly singular and strictly cosingular operators it is obvious that

$$(1.3) \quad u(A) = 0 \iff A \in S(X, Y),$$

$$(1.4) \quad v(A) = 0 \iff A \in CS(X, Y).$$

## 2. Results

Recall that F. Galaz-Fontes [3] introduced and investigated a perturbation function.

**Definition 2.1.** A perturbation function is a function  $\gamma$ , assigning to each pair of complex Banach spaces  $X, Y$ , and  $T \in B(X, Y)$  a nonnegative number  $\gamma(T)$ , with the following properties:

$$(2.1.1) \quad \gamma(\lambda T) = |\lambda| \gamma(T), \quad \lambda \in \mathbb{C},$$

$$(2.1.2) \quad \gamma(T + K) = \gamma(T), \quad K \in K(X, Y),$$

$$(2.1.3) \quad \gamma(T) \leq \|T\|,$$

$$(2.1.4) \quad m(T) \leq \gamma(T),$$

$$(2.1.5) \quad \gamma(T|_M) \leq \gamma(T),$$

where  $M$  denotes an infinite dimensional subspace of  $X$ .

He proved that the quantities  $\|\cdot\|_C$ ,  $\|\cdot\|_\alpha$ ,  $\|\cdot\|_q$ ,  $\|\cdot\|_\mu$ ,  $\Delta$  and  $u$  are perturbation functions.

Let us note that the following quantity is a measure of non-strict-singularity and we shall show that it is a perturbation function.

**Example 2.2.** For  $T \in B(X, Y)$ , set

$$\|T\|_S = \inf\{\|T + S\| : S \in S(X, Y)\}.$$

(2.1.1)-(2.1.3) follow easily from the definition.

Since  $\Delta$  is a seminorm which annihilates precisely on the strictly singular operators [9] and such that  $\Delta(T) \leq \|T\|$ ,  $T \in B(X, Y)$ , it follows that

$$\Delta(T) = \Delta(T + S) \leq \|T + S\|, \text{ for each } S \in S(X, Y).$$

Therefore

$$\Delta(T) \leq \|T\|_S.$$

Since  $m(T) \leq u(T) \leq \Delta(T)$ , we get  $m(T) \leq \|T\|_S$ . (2.1.5) follows from the fact that a restriction of a strictly singular operator to an infinite dimensional subspace of  $X$  is a strictly singular operators also.

Recall that for a given perturbation function  $\gamma$ , and  $T \in B(X, Y)$ , F. Galaz-Fontes [3] defined

$$\begin{aligned} \Gamma_{\gamma, M}(T) &= \inf\{\gamma(T|_V) : V \subset M\}, & \Gamma_\gamma(T) &= \Gamma_{\gamma, X}(T); \\ \Delta_{\gamma, M}(T) &= \sup\{\Gamma_{\gamma, V}(T) : V \subset M\}, & \Delta_\gamma(T) &= \Delta_{\gamma, X}(T), \end{aligned}$$

where  $M, V$  denote closed infinite dimensional subspaces of  $X$ .

In the following  $\gamma$  is a perturbation function. F. Galaz-Fontes proved that if  $T \in S(X, Y)$ , then  $\Delta_\gamma(T) = 0$  [3, Proposition 9]. Actually, we shall prove that the equivalence holds.

**Proposition 2.3.** *Let  $T \in B(X, Y)$ . Then*

$$(2.3.1) \quad T \in S(X, Y) \iff \Delta_\gamma(T) = 0.$$

*Proof.* First we show:

$$(2.3.2) \quad u(T) \leq \Delta_\gamma(T) \leq \Delta(T).$$

From (2.1.3) and the definitions of  $\Delta$  and  $\Delta_\gamma$  it follows that  $\Delta_\gamma(T) \leq \Delta(T)$ .

Let  $M$  be a closed infinite dimensional subspace of  $X$ . We have

$$\gamma(T|_V) \geq m(T|_V) \geq m(T|_M),$$

for each closed infinite dimensional subspace  $V$  of  $M$ . It implies

$$\Gamma_{\gamma, M}(T) = \inf_{V \subset M} \gamma(T|_V) \geq m(T|_M).$$

Hence

$$\Delta_\gamma(T) = \sup_M \Gamma_{\gamma, M}(T) \geq \sup_M m(T|_M) = u(T).$$

(2.3.1) follows directly from (2.3.2), (1.1) and (1.3).  $\square$

Let us remark that if  $\gamma$  is a submultiplicative seminorm then it can be proved (analogously as in [9]) that  $\Delta_\gamma$  is a submultiplicative seminorm and

$$\Gamma_\gamma(T + S) \leq \Gamma_\gamma(T) + \Delta_\gamma(S), \quad T, S \in B(X, Y).$$

If  $\gamma = \|\cdot\|_q$  we shall write  $\Gamma_q$  ( $\Delta_q$ ) instead of  $\Gamma_\gamma$  ( $\Delta_\gamma$ ). Analogously, we introduce  $\Gamma_\alpha$  ( $\Delta_\alpha$ ),  $\Gamma_\mu$  ( $\Delta_\mu$ ),  $\Gamma_C$  ( $\Delta_C$ ),  $\Gamma_u$  ( $\Delta_u$ ) and  $\Gamma_\Delta$  ( $\Delta_\Delta$ ). By [3, Lemma 8] it follows that  $\Gamma_\Delta = \Gamma$  and  $\Delta_\Delta = \Delta$ .

**Lemma 2.4.** *If there exists a constant  $c > 0$  such that  $\gamma(T) \geq c\Delta(T)$ , for each  $T \in B(X, Y)$ , then*

$$(2.4.1) \quad \Gamma_\gamma(T) \leq \Gamma(T) \leq c^{-1}\Gamma_\gamma(T),$$

$$(2.4.2) \quad \Delta_\gamma(T) \leq \Delta(T) \leq c^{-1}\Delta_\gamma(T), \quad T \in B(X, Y).$$

*Proof.* (2.4.1). From (2.1.3) and the definitions of  $\Gamma$  and  $\Gamma_\gamma$  it follows that  $\Gamma_\gamma(T) \leq \Gamma(T)$ . Further from the hypothesis and [3, Lemma 8] we get

$$\Gamma_\gamma(T) \geq c\Gamma_\Delta(T) = c\Gamma(T).$$

(2.4.2) can be proved analogously to (2.4.1).  $\square$

Recall that

$$\begin{aligned} \|T\|_\mu &\geq \Delta(T) \text{ [9, Theorem 2.10]}, \\ \|T\|_q &\geq 2^{-1}\|T\|_\mu \geq 2^{-1}\Delta(T) \text{ [6, Theorem 3.1]}, \\ \|T\|_\alpha &\geq 2^{-1}\|T\|_q \geq 4^{-1}\Delta(T), \\ \|T\|_C &\geq \|T\|_\mu \geq \Delta(T), \\ \|T\|_S &\geq \Delta(T), \quad T \in B(X, Y). \end{aligned}$$

By Lemma 2.4 we get

$$(2.5) \quad \begin{aligned} \Gamma_\mu(T) &= \Gamma(T), \quad \Delta_\mu(T) = \Delta(T), \\ \Gamma_q(T) &\leq \Gamma(T) \leq 2\Gamma_q(T), \quad \Delta_q(T) \leq \Delta(T) \leq 2\Delta_q(T), \\ \Gamma_\alpha(T) &\leq \Gamma(T) \leq 4\Gamma_\alpha(T), \quad \Delta_\alpha(T) \leq \Delta(T) \leq 4\Delta_\alpha(T), \\ \Gamma_C(T) &= \Gamma(T), \quad \Delta_C(T) = \Delta(T), \\ \Gamma_S(T) &= \Gamma(T), \quad \Delta_S(T) = \Delta(T). \end{aligned}$$

The inequalities in (2.5) were proved in [8] and [5, Proposition 3.9]. However, our proof is different from the proofs in [8] and [5].

**Remark 2.6.** Clearly,  $\Delta_\gamma \leq \gamma$ , and by (2.3.2), for  $\gamma = u$ , we get

$$\Delta_u = u.$$

Recall that [3, Lemma 11]

$$B \leq \Gamma_u \leq \Gamma,$$

where  $B(T) = \sup\{m(T|_V) : V \text{ closed subspace of } X, \text{codim}V < \infty\}$ . M. González and A. Martínón proved that the quantities  $B$  and  $\Gamma_u$  are not equivalent and also that the quantities  $\Gamma_u$  and  $\Gamma$  are not equivalent [5, Theorem 2.7 and Corollary 3.5].

In the terminology of [10], by [3, Theorem 6] it results that  $u/\Gamma_u$  is a perturbation function for  $\Phi_+(X, Y)$ . Since  $\Gamma_u \geq B$  it follows that the perturbation function  $u/\Gamma_u$  is better than the perturbation function  $u/B$ . Thus, Theorem 2.14 in [9] is a consequence of the fact that  $u/\Gamma_u$  is a perturbation function for  $\Phi_+(X, Y)$ .

The above fact can be established from the inequality [7, Proposicion 25.8.4]

$$\Gamma_u(T + S) \leq u(T) + \Gamma(S), \quad T, S \in B(X, Y).$$

Indeed, let  $u(S) < \Gamma_u(T)$ . Then

$$\Gamma_u(T) = \Gamma_u(T + S + (-S)) \leq \Gamma(T + S) + u(S) < \Gamma(T + S) + \Gamma_u(T),$$

and

$$\Gamma(T + S) > 0 \implies T + S \in \Phi_+(X, Y).$$

Now we introduce and investigate a new function connected with the lower semi-Fredholm operators.

**Definition 2.7.** A lower perturbation function is a function  $\delta$ , which assigns to each pair of complex Banach spaces  $X, Y$ , and  $T \in B(X, Y)$  a nonnegative number  $\delta(T)$ , with the following properties:

$$(2.7.1) \quad \delta(\lambda T) = |\lambda|\delta(T),$$

$$(2.7.2) \quad \delta(T + K) = \delta(T), \quad K \in K(X, Y),$$

$$(2.7.3) \quad \delta(T) \leq \|T\|,$$

$$(2.7.4) \quad n(T) \leq \delta(T),$$

$$(2.7.5) \quad U \subset V \implies \delta(Q_U T) \geq \delta(Q_V T),$$

where  $U, V$  denote closed infinite codimensional subspaces of  $Y$ .

We shall give several examples.

**Example 2.8.** The quantity  $\|\cdot\|_\mu$  is a lower perturbation function.

We shall prove only (2.7.4) and (2.7.5). Recall that  $\|T\|_\mu = \|T^*\|_q$  [1, Teorema 2.5.2]. Now from [3, Example 3] it follows:

$$\|T\|_\mu = \|T^*\|_q \geq m(T^*) = n(T).$$

Let  $U$  and  $V$  denote closed infinite codimensional subspaces of  $Y$  and  $U \subset V$ . Let be  $\phi : Y/U \rightarrow Y/V$  a map defined by  $\phi(y + U) = y + V$ . Then, by [6, Lemma 3.2] we have

$$\|Q_V T\|_\mu = \|\phi Q_U T\|_\mu \leq \|\phi\|_\mu \|Q_U T\|_\mu \leq \|\phi\| \|Q_U T\|_\mu \leq \|Q_U T\|_\mu.$$

**Example 2.9.** The quantity  $\|\cdot\|_q$  is a lower perturbation function.

We shall prove only (2.7.4) and (2.7.5). Recall that

$$\|T\|_q \geq \|T^*\|_\mu$$

(see [4, Theorem 1 (ii), Proposition 6 (ii)] or [1, Corollary 2.5.4]). Since  $\|T^*\|_\mu \geq m(T^*) = n(T)$ , we get

$$\|T\|_q \geq n(T).$$

Analogously as in Example 2.8, it can be proved that  $\|\cdot\|_q$  has the property (2.7.5).

**Example 2.10.** The quantity  $\|\cdot\|_C$  is a lower perturbation function.

Indeed, since  $\|T\|_C \geq \|T\|_q$ , we get  $\|T\|_C \geq n(T)$ . Since  $B(Y, Z)K(X, Y) \subset K(X, Z)$ , it follows that  $\|BA\|_C \leq \|B\| \|A\|_C$ ,  $A \in B(X, Y)$ ,  $B \in B(Y, Z)$ . In an analogous way as in Example 2.8, we obtain (2.7.5).

**Example 2.11.** The quantity  $\nabla$  is a lower perturbation function.

Since  $\nabla$  is a seminorm on  $B(X, Y)$  which annihilates precisely on the set  $CS(X, Y)$  and  $K(X, Y) \subset CS(X, Y)$  we obtain properties (2.7.1) and (2.7.2). (2.7.3) is obvious.

To prove the property (2.7.4), suppose that  $V$  is a closed subspace of  $Y$  with  $\text{codim} V \geq 1$ . Then  $V^\circ$  is a subspace of  $Y^*$ ,  $\dim V^\circ = \text{codim} V \geq 1$  and  $\|Q_V T\| = \|T^* J_{V^\circ}\|$ . Thus

$$\begin{aligned} & \{\|Q_V T\| : V \text{ closed subspace of } Y, \text{codim} V \geq 1\} \\ & \subset \{\|T^* J_M\| : M \text{ subspace of } Y^*, \dim M \geq 1\}. \end{aligned}$$

Hence

$$\begin{aligned} \nabla(T) \geq K(T) &= \inf\{\|Q_V T\| : V \text{ closed subspace of } Y, \text{codim} V = \infty\} \\ &\geq \inf\{\|Q_V T\| : V \text{ closed subspace of } Y, \text{codim} V \geq 1\} \\ &\geq \inf\{\|T^* J_M\| : M \text{ subspace of } Y^*, \dim M \geq 1\} \\ &= \inf\{\|T^* J_M\| : M \text{ subspace of } Y^*, \dim M = 1\} \\ &= \inf\{\|T^* f\| : f \in Y^*, \|f\| = 1\} = m(T^*) = n(T). \end{aligned}$$

Let us remark that

$$n(T) \leq K(T) \leq \nabla(T) \leq \|T\|_q.$$

Since  $\nabla(BA) \leq \nabla(B)\nabla(A)$ ,  $A \in B(X, Y)$ ,  $B \in B(Y, Z)$ , in an analogous way as in Example 2.8, we obtain the property (2.7.5).

**Example 2.12.** For  $T \in B(X, Y)$ , set

$$\|T\|_{CS} = \inf\{\|T + C\| : C \in CS(X, Y)\}.$$

The quantity  $\|\cdot\|_{CS}$  is a lower perturbation function.

Really, as in the previous example, we obtain that  $\|\cdot\|_{CS}$  has properties (2.7.1)-(2.7.3). It is easy to see that  $\nabla(T) \leq \|T\|_{CS}$  and since  $n(T) \leq \nabla(T)$ , we get  $n(T) \leq \|T\|_{CS}$ . Since  $B(Y, Z)CS(X, Y) \subset CS(X, Z)$ , it follows that  $\|BA\|_{CS} \leq \|B\|\|A\|_{CS}$ ,  $A \in B(X, Y)$ ,  $B \in B(Y, Z)$ . Now, in an analogous way as in Example 2.8, we obtain the property (2.7.5).

**Problem:** The quantity  $v$  has properties (2.7.1)-(2.7.4), but we do not know whether it has property (2.7.5).

From Definition 2.7 we have

**Lemma 2.13.** *If  $\delta$  is a lower perturbation function, then:*

$$(2.13.1) \quad \delta(K) = 0, \quad K \in K(X, Y),$$

$$(2.13.2) \quad \delta(Q_V) = 1,$$

*$V$  closed infinite codimensional subspace of  $Y$ .*

*Proof.* (2.13.2) follows from the following inequalities:

$$1 = m(J_{V^\circ}) = n(Q_V) \leq \delta(Q_V) \leq \|Q_V\| = 1. \quad \square$$

In the following  $\delta$  is a lower perturbation function.

**Lemma 2.14.** *Let  $P \in B(X)$ . If  $\delta(P) < 1$ , then  $I + P \in \Phi(X)$  and  $i(I + P) = 0$ .*

*Proof.* Assume that  $I + P \notin \Phi_-(X)$ . By [6, Lemma 5.4] there exists  $K \in K(X)$  such that  $\text{codim} \overline{R(I + P - K)} = \infty$ . Set  $U = \overline{R(I + P - K)}$ . From  $Q_U(I + P - K) = 0$  we get  $Q_U = Q_U(K - P)$ . Hence, by (2.13.2), (2.7.2) and (2.7.5), it follows that

$$1 = \delta(Q_U) = \delta(Q_U(K - P)) = \delta(Q_U P) \leq \delta(P).$$

This contradicts the hypothesis. Thus,  $I + P \in \Phi_-(X)$ .

Let  $0 \leq \lambda \leq 1$ . Then  $\delta(\lambda P) < 1$  and therefore  $I + \lambda P \in \Phi_-(X)$ . Since the index is locally constant it follows that  $i(I + P) = i(I) = 0$ . Consequently,  $I + P \in \Phi(X)$ .  $\square$

The following proposition can be proved (see [3, Theorem 5]).



**Proposition 2.15.**  $r_\epsilon(T) = \lim_{n \rightarrow \infty} (\delta(T^n))^{\frac{1}{n}}, \quad T \in B(X).$

**Definition 2.16.** For  $T \in B(X, Y)$ , set

$$\begin{aligned} K_{\delta, V}(T) &= \inf\{\delta(Q_W T) : W \supset V\}, & K_\delta(T) &= K_{\delta, \{0\}}(T), \\ \nabla_{\delta, V}(T) &= \sup\{K_{\delta, W}(T) : W \supset V\}, & \nabla_\delta(T) &= \nabla_{\delta, \{0\}}(T), \end{aligned}$$

where  $V, W$  denote closed infinite codimensional subspaces of  $Y$ .

**Theorem 2.17.** *Let  $S, T \in B(X, Y)$ . If  $\nabla_\delta(T) < K_\delta(T)$ , then  $T, T + S \in \Phi_-(X, Y)$  and  $i(T + S) = i(T)$ .*

*Proof.* Suppose that  $T + S \notin \Phi_-(X, Y)$ . Then, by [6, Lemma 5.4], there is  $K \in K(X, Y)$  such that  $\text{codim}R(T + S - K) = \infty$ . Set  $U = R(T + S - K)$ . Let  $V$  be a closed infinite codimensional subspace of  $Y$  such that  $V \supset U$ . Then  $Q_V(T + S - K) = 0$ , and  $Q_V T = Q_V(K - S)$ . Therefore

$$\begin{aligned} K_\delta(T) &\leq K_{\delta, U}(T) = \inf\{\delta(Q_V T) : V \supset U\} = \inf\{\delta(Q_V(K - S)) : V \supset U\} \\ &= \inf\{\delta(Q_V S) : V \supset U\} = K_{\delta, U}(S) \leq \nabla_\delta(S). \end{aligned}$$

This contradicts the hypothesis.

Let  $0 \leq \lambda \leq 1$ . Then  $\nabla_\delta(\lambda S) < K_\delta(T)$ . It follows that  $T + \lambda S \in \Phi_-(X, Y)$ . Thus  $T, T + S \in \Phi_-(X, Y)$  and, since the index is locally constant, we get  $i(T) = i(T + S)$ .  $\square$

**Theorem 2.18.** *Let  $T \in B(X, Y)$ . Then  $T \in \Phi_-(X, Y) \iff K_\delta(T) > 0$ .*

*Proof.* Let  $K_\delta(T) > 0$ . If  $S = 0$  then  $\nabla_\delta(S) = 0 < K_\delta(T)$ , and by Theorem 2.17 we get  $T \in \Phi_-(X, Y)$ .

Let  $T \in \Phi_-(X, Y)$ . Then  $\text{codim}R(T) < \infty$ , and we can express  $Y$  as a direct sum  $Y = R(T) \oplus V$  where  $V$  is a subspace of  $Y$  with  $\dim V < \infty$ . This implies that  $Q_V T$  is surjective, i.e.  $n(Q_V T) > 0$ . Let  $W$  be a closed infinite codimensional subspace of  $Y$ . Clearly,  $\text{codim}(V + W) = \infty$ . Now, by (2.7.4) and (2.7.5), we have:

$$n(Q_V T) \leq n(Q_{V+W} T) \leq \delta(Q_{V+W} T) \leq \delta(Q_W T).$$

Consequently,

$$\begin{aligned} K_\delta(T) &= \inf\{\delta(Q_W T) : W \text{ closed subspace of } Y, \text{codim}W = \infty\} \\ &\geq n(Q_V T) > 0. \quad \square \end{aligned}$$

**Proposition 2.19.**  $T \in B(X, Y)$  is strictly cosingular if and only if  $\nabla_\delta(T) = 0$ .

*Proof.* First we shall prove the following inequality

$$(2.19.1) \quad v(T) \leq \nabla_\delta(T) \leq \nabla(T).$$

The right side of the above inequality follows from (2.7.3) and the definitions of  $\nabla$  and  $\nabla_\delta$ . To prove the left side, let  $V$  be a closed infinite codimensional subspace of  $Y$ . We have

$$\delta(Q_W T) \geq n(Q_W T) \geq n(Q_V T),$$

for each closed infinite codimensional subspace  $W$ , such that  $W \supset V$ . It implies

$$K_{\delta, V}(T) = \inf_{W \supset V} \delta(Q_W T) \geq n(Q_V T).$$

Hence

$$\nabla_\delta(T) = \sup_V K_{\delta, V}(T) \geq \sup_V n(Q_V T) = v(T).$$

Now the assertion of Proposition follows from (2.19.1), (1.2) and (1.4).  $\square$

Let us remark that if  $\delta$  is a submultiplicative seminorm then it can be proved that  $\nabla_\delta$  is a submultiplicative seminorm and

$$K_\delta(T + S) \leq K_\delta(T) + \nabla_\delta(S), \quad T, S \in B(X, Y).$$

Also in this case we can show that  $\nabla_\delta$  is a lower perturbation function with  $\nabla_\delta(T) \leq \delta(T)$  (the property (2.7.4) follows from the inequality  $n(T) \leq v(T) \leq \nabla_\delta(T)$  and the property (2.7.5) can be proved analogously as in Example 2.8). Since

$$M(T) \leq K_\delta(T) \leq K(T),$$

where  $M(T) = \sup\{n(Q_V T) : \dim V < \infty\}$  [13], from [13, Theorem 8.1] it follows:

$$s_-(T) = \lim_{n \rightarrow \infty} (K_\delta(T^n))^{\frac{1}{n}},$$

where  $s_-(T) = \inf\{|\lambda| : \lambda I - T \notin \Phi_-(X)\}$ .

The next lemma implies that  $\nabla(\nabla_\delta) = \nabla_\delta$ .

**Lemma 2.20.**  $K_{\delta, V}(T) = \inf\{\nabla_{\delta, W}(T) : W \supset V\}$ , where  $V, W$  denote closed infinite codimensional subspaces of  $Y$ .

*Proof.* Since

$$\nabla_{\delta, W}(T) \geq K_{\delta, W}(T) \geq K_{\delta, V}(T),$$

for each  $W$  with  $W \supset V$ , we get

$$\inf\{\nabla_{\delta, W}(T) : W \supset V\} \geq K_{\delta, V}(T).$$

In the following  $M, N$  are closed infinite codimensional subspace of  $Y$ . Let  $M \supset W$ . From (2.7.5) it follows that

$$K_{\delta, M}(T) = \inf_{N \supset M} \delta(Q_N T) \leq \delta(Q_M T) \leq \delta(Q_W T).$$

Consequently

$$\nabla_{\delta, W}(T) = \sup_{M \supset W} K_{\delta, M}(T) \leq \delta(Q_W T).$$

This implies

$$\inf_{W \supset V} \nabla_{\delta, W}(T) \leq \inf_{W \supset V} \delta(Q_W T) = K_{\delta, V}(T). \quad \square$$

If  $\delta = \|\cdot\|_q$  we shall write  $K_q$  ( $\nabla_q$ ) instead of  $K_\delta$  ( $\nabla_\delta$ ). Analogously, we introduce  $K_\mu$  ( $\nabla_\mu$ ),  $K_C$  ( $\nabla_C$ ),  $K_\nabla$  ( $\nabla_\nabla$ ) i  $K_{CS}$  ( $\nabla_{CS}$ ). From Lemma 2.20 it follows that  $K_\nabla = K$  and  $\nabla_\nabla = \nabla$ .

In an analogous way as Lemma 2.4 the next lemma can be proved.

**Lemma 2.21.** *If there exists a constant  $c > 0$  such that  $\delta(T) \geq c\nabla(T)$ , for each  $T \in B(X, Y)$ , then*

$$\begin{aligned} K_\delta(T) &\leq K(T) \leq \frac{1}{c}K_\delta(T), \\ \nabla_\delta(T) &\leq \nabla(T) \leq \frac{1}{c}\nabla_\delta(T), \quad T \in B(X, Y). \end{aligned}$$

Recall that

$$\begin{aligned} \|T\|_q &\geq \nabla(T), \\ \|T\|_\mu &\geq \frac{1}{2}\|T\|_q \geq \frac{1}{2}\nabla(T), \\ \|T\|_C &\geq \|T\|_q \geq \nabla(T), \\ \|T\|_{CS} &\geq \nabla(T), \quad T \in B(X, Y). \end{aligned}$$

Now, by Lemma 2.21 we obtain

$$(2.22) \quad \begin{aligned} K_q(T) &= K(T), \quad \nabla_q(T) = \nabla(T), \\ K_\mu(T) &\leq K(T) \leq 2K_\mu(T), \quad \nabla_\mu(T) \leq \nabla(T) \leq 2\nabla_\mu(T), \\ K_C(T) &= K(T), \quad \nabla_C(T) = \nabla(T), \\ K_{CS}(T) &= K(T), \quad \nabla_{CS}(T) = \nabla(T). \end{aligned}$$

The equalities in (2.22) were proved in [11, Summary and discussion, Remark 2]. However, our proof is different from this one.

**Acknowledgment.** I am grateful to Professor Vladimir Rakočević for helpful comments and suggestions.

## References

- [1] P. P. Ахмеров, М. И. Каменский, А. С. Потапов и др., *Меры некомпактности и уплотняющие операторы*, Наука, Новосибирск, 1986.
- [2] A. S. Fainstein, *On measures of noncompactness of linear operators and analogous of the minimum modulus for semi-Fredholm operators (Russian)*, Spektral'naja Teorija Operatorov i Prilož., Baku **6** (1985), 182–195..
- [3] F. Galaz-Fontes, *Measures of noncompactness and upper semi-Fredholm perturbation theorems*, Proc. Amer. Math. Soc. **118** (1993), 891–897.
- [4] M. González and A. Martínón, *Operational quantities derived from the norm and measures of non-compactness*, Proc. R. Ir. Acad. **91A** (1991), 63–70.
- [5] M. González and A. Martínón, *Operational quantities characterizing semi-Fredholm operators*, Studia Math. **114** (1995), 13–27.
- [6] A. Lebow and M. Schechter, *Semigroups of operators and measures of noncompactness*, J. Funct. Anal. **7** (1971), 1–26.
- [7] A. Martínón, *Cantidades operacionales en teoria de Fredholm*, Doctoral thesis, University of La Laguna, 1989.
- [8] V. Rakočević, *Measures of non-strict singularity of operators*, Mat. Vesnik **35** (1983), 79–82.
- [9] M. Schechter, *Quantities related to strictly singular operators*, Indiana Univ. Math. J. **21** (11) (1972), 1061–1071.
- [10] M. Schechter and R. Whitley, *Best Fredholm perturbation theorems*, Studia Math. **90** (1988).
- [11] H.-O. Tylli, *On semifredholm operators, Calkin algebras and some related quantities*, Dissertation, Helsinki, 1986.
- [12] L. Weist, *Über strikt singuläre und strikt cosinguläre Operatoren in Banachräumen*, Dissertation, Bonn, 1974.
- [13] J. Zemánek, *Geometric characteristics of semi-Fredholm operators and asymptotic behaviour*, Studia Math. **80** (1984), 219–234.

UNIVERSITY OF NIŠ, FACULTY OF PHILOSOPHY, DEPARTMENT OF MATHEMATICS, ĆIRILA AND METODIJA 2, 18000 NIŠ, YUGOSLAVIJA, SERBIA