



Statistical Convergence of Bivariate Generalized Bernstein Operators via Four-Dimensional Infinite Matrices

Faruk Özger^a, Khursheed J. Ansari^b

^aDepartment of Engineering Sciences, İzmir Katip Çelebi University, 35620, İzmir, Turkey

^bDepartment of Mathematics, College of Science, King Khalid University, 61413, Abha, Saudi Arabia

Abstract. Our main aim in this work is to construct an original extension of bivariate Bernstein type operators based on multiple shape parameters to give an application of four-dimensional infinite matrices to approximation theory, and prove some Korovkin theorems using two summability methods: a statistical convergence method which is stronger than the classical case and a power series method. We obtain the rate of generalized statistical convergence, and the rate of convergence for the power series method. Moreover, we provide some computer graphics to numerically analyze the efficiency and accuracy of convergence of our operators and obtain corresponding error plots. All the results that have been obtained in the present paper can be extended to the case of n -variate functions.

1. Introduction and Preliminaries

Bernstein opened a new way [1, 2] by giving the most well-known proof of Weierstrass approximation theorem (see [42]). He constructed a sequence of approximating polynomials and many researchers have successfully extended this idea to approximate functions (see [22–24, 32, 33]). Korovkin-type theorems provide a process to decide whether a given sequence of positive linear operators converges strongly. Using certain types of statistical convergences instead of the classical convergence in Korovkin type approximation theory gives us many advantages. Applications of Korovkin type approximation on positive linear operators can be seen in [6–8]. We also note that classical convergence is also used in Korovkin type approximation theory in many papers [10–12, 32–39].

In this study, we construct an original extension of bivariate Bernstein type operators based on multiple shape parameters and prove certain Korovkin theorems using a four-dimensional summability method, and a power series method. We obtain the rate of D -statistical convergence, and the rate of convergence for the power series method (PSM) with the help of the modulus of continuity. Finally, we demonstrate some computer graphics to numerically see the efficiency and accuracy of convergence of proposed operators, and obtain corresponding error plots.

First we provide standard notations, notions and auxiliary results.

2020 *Mathematics Subject Classification.* Primary 41A10; Secondary 41A25, 41A36, 26A16, 40C05

Keywords. Multiple shape parameters, power series method, four-dimensional summability method, Robinson-Hamilton conditions, bivariate Bernstein operators, computer graphics.

Received: 3 June 2021; Accepted: 13 August 2021

Communicated by Eberhard Malkowsky

The authors extend their appreciation to the Deanship of Scientific Research at King Khalid University for funding this work through research groups program under Grant number R.G.P.1/195/42.

Email addresses: farukozger@gmail.com (Faruk Özger), ansari.jkhursheed@gmail.com (Khursheed J. Ansari)

Assume that there is $N = N(\tau) \in \mathbb{N}$ for each $\tau > 0$, so that $|\varrho_{u,v} - Q| < \tau$ whenever $u, v > N$, in this case double sequence $\varrho = (\varrho_{u,v})$ is said to be convergent to Q in Pringsheim’s sense (or simply Π -convergent), and it is denoted by $\Pi - \lim_{u,v} \varrho_{u,v} = Q$ (see [17]). When there is a positive number E so that $|\varrho_{u,v}| \leq E$ for all $(u, v) \in \mathbb{N}^2 = \mathbb{N} \times \mathbb{N}$, double sequence is said to be bounded. As it is well known, a convergent single sequence is bounded whereas a convergent double sequence need not to be bounded.

Assume that $D = (d_{l,o,\mu,\nu})$ is a four-dimensional summability method. Given a double sequence $\varrho = (\varrho_{u,v})$, D transform of ϱ , denoted by $D\varrho := ((D\varrho)_{l,o})$, is defined as

$$(D\varrho)_{l,o} = \sum_{u,v=1}^{\infty} d_{l,o,\mu,\nu} \varrho_{u,v},$$

and the double series is Π -convergent for $(l, o) \in \mathbb{N}^2$. When four-dimensional matrix $D = (d_{l,o,\mu,\nu})$ maps every bounded Π -convergent sequence into a Π -convergent sequence with the same Π -limit, it is called *RH-regular* (shortly *RHR*). A four-dimensional matrix $D = (d_{l,o,\mu,\nu})$ is *RHR* if and only if

- (a) $\Pi - \lim_{l,o} d_{l,o,\mu,\nu} = 0$,
- (b) $\Pi - \lim_{l,o} \sum_{u,v=1}^{\infty} d_{l,o,\mu,\nu} = 1$,
- (c) $\Pi - \lim_{l,o} \sum_{u=1}^{\infty} |d_{l,o,\mu,\nu}| = 0 \ (\forall v \in \mathbb{N})$,
- (d) $\Pi - \lim_{l,o} \sum_{v=1}^{\infty} |d_{l,o,\mu,\nu}| = 0 \ (\forall u \in \mathbb{N})$,
- (e) $\sum_{u,v=1}^{\infty} |d_{l,o,\mu,\nu}|$ is Π -convergent,
- (f) The inequality $\sum_{u,v>E_2} |d_{l,o,\mu,\nu}| < E_1$ is satisfied for finite positive integers E_1 and E_2 and for each $(l, o) \in \mathbb{N}^2$.

These conditions are called Robison-Hamilton conditions [18]. Assume that $D = (d_{l,o,\mu,\nu})$ is a nonnegative *RHR* matrix, and $S \subset \mathbb{N}^2$, then D -density of S is defined as

$$\delta_D^2(S) := \Pi - \lim_{l,o} \sum_{(u,v) \in S} d_{l,o,\mu,\nu}$$

provided that the limit on the right-hand side exists in the Pringsheim sense. A real double sequence $\varrho = (\varrho_{u,v})$ is called D -statistically convergent to Q and denoted by $st_D^2 - \lim_{u,v} \varrho_{u,v} = Q$ if, for every $\tau > 0$,

$$\delta_D^2(\{(u, v) \in \mathbb{N}^2 : |\varrho_{u,v} - Q| \geq \tau\}) = 0$$

(see also [16, 19]). A Π -convergent double sequence is D -statistically convergent to the same number even if converse statement may not be true. When $D = C(1, 1)$, $C(1, 1)$ -statistical convergence becomes statistical convergence for double sequences (see also [15]), where $C(1, 1) = (c_{l,o,\mu,\nu})$ is double Cesàro matrix, defined by $c_{l,o,\mu,\nu} = 1/lo$ if $1 \leq u \leq o, 1 \leq v \leq l$, and $c_{l,o,\mu,\nu} = 0$ otherwise. Suppose that $(\xi_{u,v})$ is a double sequence of nonnegative numbers with condition $\xi_{0,0} > 0$, then power series

$$\xi(a, b) := \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v$$

has radius of convergence Θ , where $\Theta \in (0, \infty]$ and $a, b \in (0, \Theta)$. When following equality is satisfied

$$\lim_{a,b \rightarrow \Theta^-} \frac{1}{\xi(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \varrho_{u,v} = Q$$

for each $a, b \in (0, \Theta)$, then double sequence $\rho = (\rho_{u,v})$ is said to be convergent to Q in the sense of PSM [27]. PSM for double sequences is regular if and only if

$$\lim_{a,b \rightarrow \Theta^-} \frac{\sum_{r=0}^{\infty} \xi_{r,v} a^r}{\xi(a,b)} = 0; \quad \lim_{a,b \rightarrow \Theta^-} \frac{\sum_{s=0}^{\infty} \xi_{\mu,s} b^s}{\xi(a,b)} = 0$$

are satisfied for any μ, v [27]. In this work, we assume that PSM is regular. When $\Theta = 1$ and $\xi_{u,v} = 1$ PSM becomes Abel summability method, and it becomes logarithmic summability method if $\xi_{u,v} = \frac{1}{(u+1)(v+1)}$. Also, PSM becomes Borel summability method when $\Theta = \infty$ and $\xi_{u,v} = \frac{1}{u!v!}$. Some properties of modified Szász-Mirakyan, Baskakov-Schurer-Szasz, and generalized Szasz operators in polynomial weight spaces were studied by power summability methods in [3–5]. We also note that applications of various statistical summability methods in approximation theory can be seen in the papers [13, 14, 29–31]. Finally, one can see more information about double sequences in [15, 16, 20, 21, 40, 41], and application of double sequences in approximation theory in [6, 27].

2. Bivariate Operators and Statistical Convergence

In this part, we construct an original extension of bivariate Bernstein type operators based on multiple shape parameters and prove some Korovkin theorems using statistical convergence four-dimensional matrices and power series method. The following polynomial functions

$$\begin{aligned} a_{u,0}(\rho; x) &= (1-x)^u (1-\rho_1 x), \\ a_{u,i}(\rho; x) &= x^i (1-x)^{u-i} \left(\binom{u}{v} + \rho_i - \rho_i x - \rho_{i+1} x \right), \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor - 1, \\ a_{u, \lfloor \frac{u}{2} \rfloor}(\rho; x) &= x^{\lfloor \frac{u}{2} \rfloor} (1-x)^{u-\lfloor \frac{u}{2} \rfloor} \left(\binom{u}{\lfloor \frac{u}{2} \rfloor} + \rho^{\lfloor \frac{u}{2} \rfloor} - \rho^{\lfloor \frac{u}{2} \rfloor} x + \rho^{\lfloor \frac{u}{2} \rfloor + 1} x \right), \\ a_{u,i}(\rho; x) &= x^i (1-x)^{u-i} \left(\binom{u}{v} - \rho_i + \rho_i x + \rho_{i+1} x \right), \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u-1, \\ a_{u,u}(\rho; x) &= x^u (1-\rho_u + \rho_u x) \end{aligned} \tag{1}$$

are called generalized Bernstein polynomials of degree u ($u \geq 2$) and for $x \in [0, 1]$ with shape parameters $\rho_i, i = 1, 2, \dots, u$, where

$$\begin{cases} \rho_i \in [-(\binom{u}{i}), (\binom{u}{i-1})] & ; i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor \\ \rho_i \in [-(\binom{u}{i-1}), (\binom{u}{i})] & ; i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u \end{cases} \quad \text{with} \quad \begin{cases} \lfloor \frac{u}{2} \rfloor = \frac{u}{2} & ; \text{if } u \text{ is even} \\ \lfloor \frac{u}{2} \rfloor = \frac{u-1}{2} & ; \text{if } u \text{ is odd.} \end{cases} \tag{2}$$

These polynomials were introduced by Han et al. in [26] and they are reduced to classical Bernstein basis functions $b_{u,i}(x)$ of degree u on $x \in [0, 1]$ which is defined as

$$b_{u,i}(x) = \binom{u}{v} x^i (1-x)^{u-i}, \quad i = 0, \dots, u$$

when $\rho_i = 0$ ($i = 1, 2, \dots, u$). Generalized Bernstein basis functions with parameters ρ_i ($i = 1, 2, \dots, u$) are linearly independent (see [25]) and these basis functions are effectively and flexibly used in designing parametric curves and surfaces (see [25, 26]). These functions also have partition of unity, symmetry and nonnegativity properties (see [26]). In 2017, Hu et al. [25] have obtained the following equations to convert classical Bernstein polynomials of degree u to generalized Bernstein polynomials of degree u associated

with shape parameters ρ_i :

$$\begin{aligned}
 a_{u,0}(\rho; x) &= b_{u+1,0}(x) + \frac{\binom{u}{0} - \rho_1}{\binom{u+1}{1}} b_{u+1,1}(x), \\
 a_{u,i}(\rho; x) &= \frac{\binom{u}{i} + \rho_i}{\binom{u+1}{i}} b_{u+1,i}(x) + \frac{\binom{u}{i} - \rho_{i+1}}{\binom{u+1}{i+1}} b_{u+1,i+1}(x), \quad i = 1, 2, \dots, \left\lfloor \frac{u}{2} \right\rfloor - 1, \\
 a_{u,i}(\rho; x) &= \frac{\binom{u}{i} + \rho_v}{\binom{u+1}{i}} b_{u+1,i}(x) + \frac{\binom{u}{i} + \rho_{i+1}}{\binom{u+1}{i+1}} b_{u+1,i+1}(x), \quad i = \left\lfloor \frac{u}{2} \right\rfloor, \\
 a_{u,i}(\rho; x) &= \frac{\binom{u}{i} - \rho_i}{\binom{u+1}{i}} b_{u+1,i}(x) + \frac{\binom{u}{i} + \rho_{i+1}}{\binom{u+1}{i+1}} b_{u+1,i+1}(x), \quad i = \left\lfloor \frac{u}{2} \right\rfloor + 1, \dots, u - 1, \\
 a_{u,u}(\rho; x) &= \frac{\binom{u}{u} - \rho_u}{\binom{u+1}{u}} b_{u+1,u}(x) + b_{u+1,u+1}(x).
 \end{aligned} \tag{3}$$

Let $C[0, 1] = \mathbf{C}$ be the space of all continuous functions on unit interval $[0, 1]$ and $C([0, 1] \times [0, 1]) = \bar{\mathbf{C}}$. The operators $\mathcal{B}_u^v, \mathcal{B}_v^u : \mathbf{C} \rightarrow \mathbf{C}$ for any $u, v \in \mathbb{N}$ are given as follows, respectively,

$$\mathcal{B}_u^v(f; y) = \sum_{i=0}^u f\left(\frac{i}{u}\right) a_{u,i}(v; y), \tag{4}$$

$$\mathcal{B}_v^u(g; z) = \sum_{j=0}^v g\left(\frac{j}{v}\right) a_{v,j}(u; z), \tag{5}$$

where polynomials $a_{u,i}(v; y)$ and $a_{v,j}(u; z)$ are given in (3). The parametric extension of (4) and (5) for $u, v \in \mathbb{N}$ and $h \in \bar{\mathbf{C}}$ are the operators

$$\mathcal{B}_u^{v,y}, \mathcal{B}_v^{u,z} : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}},$$

where

$$\mathcal{B}_u^{v,y}(h; y, z) = \sum_{i=0}^u a_{u,i}(v; y) h\left(\frac{i}{u}, \frac{i}{u}\right), \tag{6}$$

$$\mathcal{B}_v^{u,z}(h; y, z) = \sum_{j=0}^v a_{v,j}(u; z) h\left(\frac{j}{v}, \frac{j}{v}\right). \tag{7}$$

Lemma 2.1. *The parametric extension of operators defined in (6) and (7) are linear and positive.*

Proof. The assertion follows from the definitions of $\mathcal{B}_u^{v,y}$ and $\mathcal{B}_v^{u,z}$. \square

Lemma 2.2. *The parametric extensions of bivariate operators commute on $\bar{\mathbf{C}}$. Their product establishes bivariate operators $\mathcal{B}_{u,v}^{v,u} : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ defined for any $u, v \in \mathbb{N}$ and any $h \in \bar{\mathbf{C}}$ by the relation*

$$\mathcal{B}_{u,v}^{v,u}(h; y, z) = \sum_{i=0}^u \sum_{j=0}^v a_{u,i}(v; y) a_{v,j}(u; z) h\left(\frac{i}{u}, \frac{j}{v}\right). \tag{8}$$

Proof. We get the desired result by direct computation, taking into account definitions (6), (7) and Lemma 2.1. \square

Lemma 2.3. *The bivariate operators (8) are linear and positive.*

Proof. Using the fact that the product of linear and positive operators are also linear and positive, and applying Lemma 2.1 we obtain desired result. \square

Lemma 2.4. [13] For any natural number u ($u \geq 2$) the following equalities hold:

$$\begin{aligned} \mathcal{B}_u^p(1; x) &= 1, \mathcal{B}_u^p(t; x) = x + \frac{1-x}{u} \mathcal{R}_0(x), \mathcal{B}_u^p(t^2; x) = \frac{u-1}{u} x^2 + \frac{x}{u} + \frac{1-x}{u^2} [2\mathcal{R}_1(x) - \mathcal{R}_0(x)], \\ \mathcal{B}_u^p(t^3; x) &= \frac{(u-1)(u-2)}{u^2} x^3 + \frac{3(u-1)}{u^2} x^2 + \frac{x}{u^2} + \frac{1-x}{u^3} [3\mathcal{R}_2(x) - 3\mathcal{R}_1(x) + \mathcal{R}_0(x)], \\ \mathcal{B}_u^p(t^4; x) &= \frac{(u-1)(u-2)(u-3)}{u^3} x^4 + \frac{6(u-1)(u-2)}{u^3} x^3 + \frac{7(u-1)}{u^3} x^2 + \frac{x}{u^3} \\ &\quad + \frac{1-x}{u^4} [4\mathcal{R}_3(x) - 6\mathcal{R}_2(x) + 4\mathcal{R}_1(x) - \mathcal{R}_0(x)], \end{aligned}$$

where

$$\mathcal{R}_p(x) = \sum_{i=1}^{\lfloor \frac{u}{2} \rfloor} \binom{u}{i} i^p \rho_i b_{u,i}(x) - \sum_{i=\lfloor \frac{u}{2} \rfloor + 1}^u \binom{u}{i} i^p \rho_i b_{u,i}(x), \quad p = 0, 1, 2, \dots, u, \quad x \in [0, 1].$$

Lemma 2.5. The parametric extension $\mathcal{B}_u^{v,y}$ satisfies the identities in Lemma 2.1.

Proof. By using the definition (6) of $\mathcal{B}_u^{v,y}$ and Lemma 2.1, we get the result. \square

Remark 2.6. The parametric extension $\mathcal{B}_v^{u,z}$ satisfies identities similar to the identities in Lemma 2.1.

Lemma 2.7. Let $e_{uv}(y, z) = y^u z^v$, $u, v \in \mathbb{N}$, $y, z \in \mathbb{R}$ be the two-dimensional test functions. The bivariate operators defined in (8) satisfy

$$\begin{aligned} \mathcal{B}_{u,v}^{v,\mu}(e_{00}; y, z) &= 1, \mathcal{B}_{u,v}^{v,\mu}(e_{10}; y, z) = y + \frac{1-y}{u} \mathcal{R}_0(y), \mathcal{B}_{u,v}^{v,\mu}(e_{01}; y, z) = z + \frac{1-z}{v} \mathcal{R}_0(z), \\ \mathcal{B}_{u,v}^{v,\mu}(e_{20}; y, z) &= \frac{u-1}{u} y^2 + \frac{y}{u} + \frac{1-y}{u^2} [2\mathcal{R}_1(y) - \mathcal{R}_0(y)], \\ \mathcal{B}_{u,v}^{v,\mu}(e_{02}; y, z) &= \frac{v-1}{v} z^2 + \frac{z}{v} + \frac{1-z}{v^2} [2\mathcal{R}_1(z) - \mathcal{R}_0(z)], \\ \mathcal{B}_{u,v}^{v,\mu}(e_{30}; y, z) &= \frac{(u-1)(u-2)(u-3)}{u^3} y^4 + \frac{6(u-1)(u-2)}{u^3} y^3 + \frac{7(u-1)}{u^3} y^2 + \frac{y}{u^3} \\ &\quad + \frac{1-y}{u^4} [4\mathcal{R}_3(y) - 6\mathcal{R}_2(y) + 4\mathcal{R}_1(y) - \mathcal{R}_0(y)], \\ \mathcal{B}_{u,v}^{v,\mu}(e_{03}; y, z) &= \frac{(v-1)(v-2)(v-3)}{v^3} z^4 + \frac{6(v-1)(v-2)}{v^3} z^3 + \frac{7(v-1)}{v^3} z^2 + \frac{z}{v^3} \\ &\quad + \frac{1-z}{v^4} [4\mathcal{R}_3(z) - 6\mathcal{R}_2(z) + 4\mathcal{R}_1(z) - \mathcal{R}_0(z)], \\ \mathcal{B}_{u,v}^{v,\mu}(e_{40}; y, z) &= \frac{(u-1)(u-2)(u-3)}{u^3} y^4 + \frac{6(u-1)(u-2)}{u^3} y^3 + \frac{7(u-1)}{u^3} y^2 + \frac{y}{u^3} \\ &\quad + \frac{1-y}{u^4} [4\mathcal{R}_3(y) - 6\mathcal{R}_2(y) + 4\mathcal{R}_1(y) - \mathcal{R}_0(y)], \\ \mathcal{B}_{u,v}^{v,\mu}(e_{04}; y, z) &= \frac{(v-1)(v-2)(v-3)}{v^3} z^4 + \frac{6(v-1)(v-2)}{v^3} z^3 + \frac{7(v-1)}{v^3} z^2 + \frac{z}{v^3} \\ &\quad + \frac{1-z}{v^4} [4\mathcal{R}_3(z) - 6\mathcal{R}_2(z) + 4\mathcal{R}_1(z) - \mathcal{R}_0(z)]. \end{aligned}$$

Proof. Taking into account definition (8) and Lemma 2.4, we complete the proof. \square

Lemma 2.8. *The bivariate operators (8) satisfy the relations*

$$\mathcal{B}_{u,v}^{\nu,\mu}((e_{10} - y)^2; y, z) = \frac{y(1-y)}{u}(1 - 2\mathcal{R}_0(y)) + \frac{1-y}{u^2} [2\mathcal{R}_1(y) - \mathcal{R}_0(y)], \tag{9}$$

$$\mathcal{B}_{u,v}^{\nu,\mu}((e_{01} - z)^2; y, z) = \frac{z(1-z)}{v}(1 - 2\mathcal{R}_0(z)) + \frac{1-z}{v^2} [2\mathcal{R}_1(z) - \mathcal{R}_0(z)]. \tag{10}$$

Proof. Since $\mathcal{B}_{u,v}^{\nu,\mu}$ is linear, we have

$$\begin{aligned} \mathcal{B}_{u,v}^{\nu,\mu}((e_{10} - y)^2; y, z) &= \mathcal{B}_{u,v}^{\nu,\mu}(e_{20}; y, z) \\ &\quad - 2y\mathcal{B}_{u,v}^{\nu,\mu}(e_{10}; y, z) + y^2\mathcal{B}_{u,v}^{\nu,\mu}(e_{10}; y, z). \end{aligned}$$

By applying Lemma 2.7, we get relation (9). Similarly we have the equality (10). \square

Following theorem gives Korovkin type approximation for statistical convergence by four-dimensional matrices:

Theorem 2.9. [19] *Let $D = (d_{l,o,u,v})$ be a nonnegative RHR matrix. Let $(Q_{u,v})$ be a double sequence of operators acting from $C([a, b] \times [c, d])$ into itself. So, for each $h \in C([a, b] \times [c, d])$,*

$$st_D^2 - \lim_{u,v} \|Q_{u,v}(h) - h\|_{C([a,b] \times [c,d])} = 0$$

if and only if

$$st_D^2 - \lim_{u,v} \|Q_{u,v}(h_u) - h_u\|_{C([a,b] \times [c,d])} = 0,$$

where $h_0(y, z) = 1, h_1(y, z) = y, h_2(y, z) = z$ and $h_3(y, z) = y^2 + z^2$.

Theorem 2.9 provides next result.

Theorem 2.10. *Let $h \in \tilde{C}$, then*

$$st_D^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h) - h\|_{\tilde{C}} = 0.$$

Proof. Assume that

$$st_D^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h_u) - h_u\|_{\tilde{C}} = 0. \tag{11}$$

We have the following result for h_0 using Lemma 2.7 (a):

$$st_D^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h_0) - h_0\|_{\tilde{C}} = 0.$$

By Lemma 2.7, we obtain

$$\|\mathcal{B}_{u,v}^{\nu,\mu}(h_1) - h_1\|_{\tilde{C}} = \sup_{(y,z) \in [0,1] \times [0,1]} \left| y + \frac{1-y}{u} \mathcal{R}_0(y) - y \right| \leq \frac{1}{u}. \tag{12}$$

For a given $\epsilon' > 0$, we choose a number $\epsilon > 0$ such that $\epsilon < \epsilon'$. Let us define the following sets:

$$\begin{aligned} \mathcal{S} &: = \{(u, v) : \|\mathcal{B}_{u,v}^{\nu,\mu}(h_1) - h_1\|_{C([0,1] \times [0,1])} \geq \epsilon'\}, \\ \mathcal{S}_1 &: = \{(u, v) : \frac{1}{u} \geq \epsilon - \epsilon'\}. \end{aligned}$$

We observe that $\mathcal{S} \subseteq \mathcal{S}_1$. So $\delta_D^2(\mathcal{S}) \leq \delta_D^2(\mathcal{S}_1)$ is satisfied and the following relation is satisfied for h_1 :

$$st_D^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h_1) - h_1\|_{\bar{C}} = 0.$$

Using the same manner we have

$$st_D^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h_2) - h_2\|_{\bar{C}} = 0,$$

that is (11) is satisfied for h_2 . Considering the following relations

$$\|\mathcal{B}_{u,v}^{\nu,\mu}(h_3) - h_3\|_{\bar{C}} \tag{13}$$

$$\leq \|\mathcal{B}_{u,v}^{\nu,\mu}(e_{20}) - e_{20}\|_{\bar{C}} + \|\mathcal{B}_{u,v}^{\nu,\mu}(e_{02}) - e_{02}\|_{\bar{C}}$$

$$= \sup_{(y,z) \in [0,1] \times [0,1]} \left| \frac{y(1-y)}{u} + \frac{1-y}{u^2} [2\mathcal{R}_1(y) - \mathcal{R}_0(y)] \right|$$

$$+ \sup_{(y,z) \in [0,1] \times [0,1]} \left| \frac{z(1-z)}{v} + \frac{1-z}{v^2} [2\mathcal{R}_1(z) - \mathcal{R}_0(z)] \right|$$

$$\leq \sup_{(y,z) \in [0,1] \times [0,1]} \left| \frac{y(1-y)}{u} + \frac{2(1+u)}{u^2} \left[y^2 + \frac{y(1-y)}{u} \right] \right| \tag{14}$$

$$+ \sup_{(y,z) \in [0,1] \times [0,1]} \left| \frac{z(1-z)}{v} + \frac{2(1+v)}{v^2} \left[z^2 + \frac{z(1-z)}{v} \right] \right|$$

and taking D -statistical limit in both-sides of last inequality, we obtain

$$st_D^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h_3) - h_3\|_{\bar{C}} = 0,$$

that is (11) is satisfied for h_3 . \square

The following result is obtained by replacing the matrix D in Theorem 2.9 with double identity matrix.

Corollary 2.11. Assume that $h \in \bar{C}$, then

$$\Pi - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h) - h\|_{\bar{C}} = 0.$$

The $C(1,1)$ -statistical convergence becomes statistical convergence for double sequences if $D = C(1,1)$ is chosen, and following result is satisfied:

Corollary 2.12. Assume that $h \in \bar{C}$, then

$$st_{C(1,1)}^2 - \lim_{u,v} \|\mathcal{B}_{u,v}^{\nu,\mu}(h) - h\|_{\bar{C}} = 0.$$

3. Power series method for operators $\mathcal{B}_{u,v}^{\nu,\mu}$

Assume throughout the section, $\Psi := [a, b] \times [c, d]$ and $(Q_{u,v})$ be a double sequence of positive linear operators acting from $C(\Psi)$ into itself such that

$$\sup_{0 < a, b < \Theta} \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \|Q_{u,v}(1)\|_{C(\Psi)} < \infty. \tag{15}$$

Set

$$S_{a,b}(h; y, z) = \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v Q_{u,v}(h; y, z), \quad a, b \in (0, \Theta)$$

and

$$T_{a,b}(h; y, z) = \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \mathcal{B}_{u,v}^{\nu, \mu}(h; y, z), \quad a, b \in (0, \Theta).$$

The proof of following theorem was given in [23], for the readers convince we also give it here.

Theorem 3.1. *Let $h \in C(\Psi)$, then*

$$\lim_{a,b \rightarrow \Theta^-} \|S_{a,b}(h) - h\|_{C(\Psi)} = 0 \tag{16}$$

if and only if

$$\lim_{a,b \rightarrow \Theta^-} \|S_{a,b}(h_u) - h_u\|_{C(\Psi)} = 0, \tag{17}$$

where $h_0(y, z) = 1, h_1(y, z) = y, h_2(y, z) = z$ and $h_3(y, z) = y^2 + z^2$.

Proof. The implication (16) \Rightarrow (17) is clear since $h_u \in C(\Psi)$ for each $u = 0, 1, 2, 3$. Let $h \in C(\Psi)$ and $(y, z) \in \Psi$ be fixed. Since function h is continuous on Ψ , following inequality is satisfied:

$$|h(y, z)| \leq M_h.$$

Therefore

$$|h(s, t) - h(y, z)| \leq 2M_h.$$

Also, since h is continuous on Ψ , there is a number $\rho > 0$ so that $|h(s, t) - h(y, z)| < \tau$ holds for each $\tau > 0$ and $(s, t) \in \Psi$ satisfying $|s - y| < \rho$ and $|t - z| < \rho$. Hence, we get

$$|h(s, t) - h(y, z)| < \tau + \frac{2M_h}{\rho^2} \{(s - y)^2 + (t - z)^2\}.$$

This means

$$-\tau - \frac{2M_h}{\rho^2} \{(s - y)^2 + (t - z)^2\} < h(s, t) - h(y, z) < \tau + \frac{2M_h}{\rho^2} \{(s - y)^2 + (t - z)^2\}.$$

So, we can write

$$\begin{aligned} & |S_{a,b}(h; y, z) - h(y, z)| \\ &= \left| \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v Q_{u,v}(h; y, z) - h(y, z) \right| \\ &\leq \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v Q_{u,v}(|h(s, t) - h(y, z)|) \\ &\quad + |h(y, z)| \left| \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v Q_{u,v}(h_0; y, z) - h_0(y, z) \right| \\ &\leq \tau + \left(\tau + M_h + \frac{2M_h \|h_3\|_{C(\Psi)}}{\rho^2} \right) |S_{a,b}(h_0; y, z) - h_0(y, z)| \\ &\quad + \frac{4M_h \|h_1\|_{C(\Psi)}}{\rho^2} |S_{a,b}(h_1; y, z) - h_1(y, z)| \\ &\quad + \frac{4M_h \|h_2\|_{C(\Psi)}}{\rho^2} |S_{a,b}(h_2; y, z) - h_2(y, z)| \\ &\quad + \frac{2M_h}{\rho^2} |S_{a,b}(h_3; y, z) - h_3(y, z)|. \end{aligned}$$

Then taking supremum over $(y, z) \in \Psi$, we have

$$\begin{aligned} & \|S_{a,b}(h) - h\|_{C(\Psi)} \\ & \leq \tau + N \left\{ \sum_{u=0}^3 \|S_{a,b}(h_u; y, z) - h_u(y, z)\|_{C(\Psi)} \right\}, \end{aligned}$$

where

$$N := \max \left\{ \tau + M_h + \frac{2M_h \|h_3\|_{C(\Psi)}}{\rho^2}, \frac{4M_h \|h_1\|_{C(\Psi)}}{\rho^2}, \frac{4M_h \|h_2\|_{C(\Psi)}}{\rho^2}, \frac{2M_h}{\rho^2} \right\}.$$

By the aid of relation (17), following result, which completes the proof, is obtained:

$$\lim_{a,b \rightarrow \Theta^-} \|S_{a,b}(h) - h\|_{C(\Psi)} = 0.$$

□

Theorem 3.2. Let $h \in \tilde{C}$, then

$$\lim_{a,b \rightarrow \Theta^-} \|T_{a,b}(h) - h\|_{\tilde{C}} = 0.$$

Proof. Since $\mathcal{B}_{u,v}^{\nu,\mu}(e_{00}; y, z) = 1$, we see that (15) holds. Considering Lemma 2.7 and the inequalities (12) and (13), the proof is completed. □

4. Rate of statistical convergence, and rate of convergence for PSM

In this section, with the aid of modulus of continuity, we calculate the rate of statistical convergence by four-dimensional matrices and the rate of convergence for PSM, here modulus of continuity is expressed as

$$\omega(h, \rho) = \sup_{\sqrt{(s-y)^2 + (t-z)^2} \leq \rho} |h(s, t) - h(y, z)| \quad (\rho > 0), \quad h \in \tilde{C}.$$

We know that, for any $\rho > 0$ and for all $h \in \tilde{C}$

$$\omega(h, \rho\rho) \leq (1 + [\rho]) \omega(h, \rho),$$

where $[\rho]$ is greatest integer less than or equal to ρ .

Theorem 4.1. Assume that $h \in \tilde{C}$, $u, v \in \mathbb{Z}_+$, $D = (d_{i,o,u,v})$ is a nonnegative RHR matrix and $(\alpha_{u,v})$ is a positive non-increasing double sequence so that $\omega(h, \rho_{u,v}) = st_D^2 - o(\alpha_{u,v})$, then

$$\|\mathcal{B}_{u,v}^{\nu,\mu}(h) - h\|_{\tilde{C}} = st_D^2 - o(\alpha_{u,v}),$$

where

$$\rho_{u,v} := \left\{ \frac{2}{u} + \frac{3}{u^2} + \frac{4(u+1)}{u^3} + \frac{2}{v} + \frac{3}{v^2} + \frac{4(v+1)}{v^3} \right\}^{\frac{1}{2}}.$$

Proof. Assume that the hypotheses are satisfied. Using positivity and monotonicity of operators $\mathcal{B}_{u,v}^{\nu,\mu}$ we get

$$\begin{aligned} |\mathcal{B}_{u,v}^{\nu,\mu}(h; y, z) - h(y, z)| & \leq \mathcal{B}_{u,v}^{\nu,\mu}(|h(s, t) - h(y, z)|; y, z) \\ & \leq \mathcal{B}_{u,v}^{\nu,\mu} \left(\left(1 + \frac{(s-y)^2 + (t-z)^2}{\rho^2} \right) \omega(h, \rho); y, z \right) \\ & = \omega(h, \rho) + \frac{\omega(h, \rho)}{\rho^2} \mathcal{B}_{u,v}^{\nu,\mu}((s-y)^2 + (t-z)^2; y, z). \end{aligned}$$

Then taking supremum over $(y, z) \in [0, 1] \times [0, 1]$, we have

$$\begin{aligned} & \left\| \mathcal{B}_{u,v}^{\nu,\mu}(h) - h \right\|_{\bar{C}} \\ & \leq \omega(h, \rho) + \frac{\omega(h, \rho)}{\rho^2} \left\{ \left\| \mathcal{B}_{u,v}^{\nu,\mu}((s - \cdot)^2) \right\|_{\bar{C}} + \left\| \mathcal{B}_{u,v}^{\nu,\mu}((t - \cdot)^2) \right\|_{\bar{C}} \right\} \\ & \leq \omega(h, \rho) + \frac{\omega(h, \rho)}{\rho^2} \left\{ \frac{y(1-y)}{u} + \frac{2y^2-y}{u^2} + \frac{2(u+1)y(1-y)}{u^3} \right\} \\ & \quad + \frac{\omega(h, \rho)}{\rho^2} \left\{ \frac{z(1-z)}{v} + \frac{2z^2-z}{v^2} + \frac{2(v+1)z(1-z)}{v^3} \right\} \\ & \leq \omega(h, \rho) + \frac{\omega(h, \rho)}{\rho^2} \left\{ \frac{2}{u} + \frac{3}{u^2} + \frac{4(u+1)}{u^3} + \frac{2}{v} + \frac{3}{v^2} + \frac{4(v+1)}{v^3} \right\}. \end{aligned}$$

By the following chose of ρ

$$\rho = \rho_{u,v} := \left\{ \frac{2}{u} + \frac{3}{u^2} + \frac{4(u+1)}{u^3} + \frac{2}{v} + \frac{3}{v^2} + \frac{4(v+1)}{v^3} \right\}^{\frac{1}{2}}$$

the following inequality is satisfied for any positive integers u, v :

$$\left\| \mathcal{B}_{u,v}^{\nu,\mu}(h) - h \right\|_{\bar{C}} \leq 2\omega(h, \rho_{u,v}).$$

Hence the following relation is satisfied for any $\tau > 0$:

$$\frac{1}{\alpha_{u,v}} \sum_{\left\| \mathcal{B}_{u,v}^{\nu,\mu}(h) - h \right\|_{\bar{C}} \geq \tau} d_{l,o,u,v} \leq \frac{1}{\alpha_{u,v}} \sum_{\omega(h, \rho_{u,v}) \geq \frac{\tau}{2}} d_{l,o,u,v}$$

and the following final step is obtained from the hypothesis

$$\left\| \mathcal{B}_{u,v}^{\nu,\mu}(h) - h \right\|_{\bar{C}} = st_D^2 - o(\alpha_{u,v}).$$

□

We give rate of convergence for PSM by the following theorem.

Theorem 4.2. Let $h \in \bar{C}$ and ζ be a positive real function defined on $(0, \Theta) \times (0, \Theta)$. If $\omega(h, \psi) = o(\zeta)$, as $a, b \rightarrow \Theta^-$, then

$$\left\| T_{a,b}(h) - h \right\|_{\bar{C}} = o(\zeta)$$

as $a, b \rightarrow \Theta^-$, where $\psi : (0, \Theta) \times (0, \Theta) \rightarrow \mathbb{R}$ is given as

$$\begin{aligned} & \psi(a, b) \\ & : = \left\{ \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \left\| \mathcal{B}_{u,v}^{\nu,\mu}((s - \cdot)^2 + (t - \cdot)^2) \right\|_{\bar{C}} \right\}^{\frac{1}{2}}. \end{aligned}$$

Proof. Since the operators are linear and the positive the following relations are satisfied

$$\begin{aligned} & |T_{a,b}(h; y, z) - h(y, z)| \\ & = \left| \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \mathcal{B}_{u,v}^{\nu,\mu}(h; y, z) - h(y, z) \right| \\ & \leq \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \mathcal{B}_{u,v}^{\nu,\mu}(|h(s, t) - h(y, z)|; y, z) \\ & \leq \omega(h, \rho) + \frac{\omega(h, \rho)}{\rho^2} \left\{ \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \mathcal{B}_{u,v}^{\nu,\mu}((s - y)^2 + (t - z)^2; y, z) \right\} \end{aligned}$$

for any $a, b \in (0, \Theta)$ and $(y, z) \in [0, 1] \times [0, 1]$. Then taking supremum, we have

$$\|\mathcal{T}_{u,v}^{\nu,\mu}(h) - h\|_{\mathcal{C}} \leq 2 \omega(h, \psi),$$

where

$$\begin{aligned} \rho &= \psi(a, b) \\ &: = \left\{ \frac{1}{\varrho(a, b)} \sum_{u,v=0}^{\infty} \xi_{u,v} a^u b^v \left\| \mathcal{B}_{u,v}^{\nu,\mu}((s - \cdot)^2 + (t - \cdot)^2) \right\|_{\mathcal{C}} \right\}^{\frac{1}{2}}. \end{aligned}$$

□

5. Convergence by graphics

In this part, we first focus on the behavior of generalized basis polynomials $a_{u,v}(v; y)a_{u,v}(\mu; z)$ and give several illustrative examples with the help of Mathematica to verify the convergence behavior, computational efficiency and consistency of the generalized bivariate Bernstein operators.

In order to see the effect of shape parameters ν and μ to behaviour of polynomials $a_{u,v}(v; y)a_{u,v}(\mu; z)$ we choose certain parameters ν and μ . In Figure 1 (a), we present behaviour of polynomials $b_{u,v}(y)b_{u,v}(z)$ to see the difference. In Figure 1 (b)-(f), we choose the following shape parameters, respectively:

$$\begin{aligned} \text{(b)} \quad v_i &= \binom{u}{i-1} - \binom{u}{i}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = \binom{v}{j-1} - \binom{v}{j}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ v_i &= \binom{u}{i} - \binom{u}{i-1}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = \binom{v}{j} - \binom{v}{j-1}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v; \\ \text{(c)} \quad v_i &= \binom{u}{i-1}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = \binom{v}{j-1}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ v_i &= \binom{u}{i}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = \binom{v}{j}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v; \\ \text{(d)} \quad v_i &= -\binom{u}{i}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = -\binom{v}{j}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ v_i &= \binom{u}{i}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = \binom{v}{j}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v; \\ \text{(e)} \quad v_i &= \binom{u}{i-1}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = \binom{v}{j-1}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ v_{i, \mu_j} &= -\binom{u}{v-1}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v; \\ \text{(f)} \quad v_i &= -\binom{u}{i}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = -\binom{v}{j}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ v_i &= -\binom{u}{i-1}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = -\binom{v}{j-1}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v. \end{aligned}$$

Example 5.1. Consider the following function

$$h(y, z) = \frac{(z^3 - \frac{1}{2}) \sin(\pi y) \cos(\pi y)}{(y^7 + 2)(z^7 + \frac{1}{2})}$$

u	v	Absolute Error	y	z
4	4	1.4204	1	0.8635
10	10	0.7281	1	0.8617
20	20	0.3976	1	0.8589
30	30	0.2731	1	0.8579
40	40	0.2079	1	0.8575
50	50	0.1679	1	0.8573
5	50	0.5915	0.8349	0.8490
5	100	0.5291	0.8346	0.8476
100	100	0.0855	1	0.8569

Table 1: Absolute errors of approximation for certain u and v values and related occurring points (y, z) for Example 5.1

u	v	GB	B
4	4	0.1507	0.1596
6	6	0.1148	0.1216
8	8	0.0936	0.0984
10	10	0.0793	0.0826
12	12	0.0688	0.0712
20	4	0.0552	0.0658
4	20	0.1482	0.1565
50	10	0.0261	0.0313

Table 2: Comparison of bivariate Bernstein and $\mathcal{B}_{u,v}^{\nu,\mu}$ operators by absolute errors for Example 5.3

on $(y, z) \in [0, 1] \times [0, 1]$. We choose the following shape parameters for this example:

$$\nu_i = \frac{1}{10000} \binom{u}{i-1}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = \frac{1}{10000} \binom{v}{j-1}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor,$$

$$\nu_i = \binom{u}{i}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = \binom{v}{j}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v.$$

In Table 1, we give maximum error of approximation for $\mathcal{B}_{u,v}^{\nu,\mu}$ for certain u and v values and related occurring (y, z) points.




Figure 1: Behaviour of polynomials $a_{u,v}(v;y)a_{u,v}(\mu;z)$ and $b_{u,v}(y)b_{u,v}(z)$




Figure 2: Approximations of $\mathcal{B}_{u,v}^{v,u}$ operators with different values of u and v for Example 5.2

Example 5.2. Consider the following function

$$h(y, z) = \frac{|5z^3 - 1|(y^3 - 22)(z + 1) \cos(1.5\pi z)}{|y^3 - 15|}$$

on $(y, z) \in [0, 1] \times [0, 1]$. We choose the following shape parameters for this example:

$$\begin{aligned} \nu_i &= \frac{7}{ui} \binom{u}{i-1}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = \frac{7}{vj} \binom{v}{j-1}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ \nu_i &= \frac{1}{10i} \binom{u}{i}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = \frac{1}{10j} \binom{v}{j}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v. \end{aligned}$$

In Figures 2-3, we give approximations of $\mathcal{B}_{u,v}^{v,u}$ operators with $u = v = 10$ and related error of approximation, respectively. In Figure 4, we extensively examine errors with different values of u and v .

Example 5.3. Consider the following function

$$h(y, z) = \frac{(z^3 - 2z^3y^2) \sin(2\pi z)}{2(y^3 + 1)}$$

on $(y, z) \in [0, 1] \times [0, 1]$. We choose the following shape parameters for this example:

$$\begin{aligned} \nu_i &= -0.000005 \binom{u}{i}, \quad i = 1, 2, \dots, \lfloor \frac{u}{2} \rfloor, \quad \mu_j = -0.000005 \binom{v}{j}, \quad j = 1, 2, \dots, \lfloor \frac{v}{2} \rfloor, \\ \nu_i &= -0.000001 \binom{u}{i-1}, \quad i = \lfloor \frac{u}{2} \rfloor + 1, \dots, u, \quad \mu_j = -0.000001 \binom{v}{j-1}, \quad j = \lfloor \frac{v}{2} \rfloor + 1, \dots, v. \end{aligned}$$

In Figures 5-6, we provide convergence of $\mathcal{B}_{u,v}^{v,u}$ and related errors of approximation for the given function. In Figure 7, we extensively examine errors with different values of u and v . We also compare the maximum error of $\mathcal{B}_{u,v}^{v,u}$ with the classical Bernstein operators in Table 2.




Figure 3: Errors of approximation for Example 5.2




Figure 7: Approximation errors of $\mathcal{B}_{u,v}^{\nu,\mu}$ to the function $h(y, z)$ for different ν and ν values




Figure 4: Approximation errors of $\mathcal{B}_{u,v}^{v,H}$ to the function $h(y, z)$ for different u and v values




Figure 5: Approximation of $\mathcal{B}_{u,v}^{\nu,\mu}$ operators for $u = v = 30$




Figure 6: Error of approximation

References

- [1] SN. Bernstein, Démonstration du théorème de Weierstrass fondée sur le calcul des probabilités, *Comm. Soc. Math. Kharkow*, (1912) 13, 1-2.
- [2] SN. Bernstein, Sur les recherches récentes relatives à la meilleure approximation des fonctions continues par les polynômes, *Proc. of 5th Inter. Math. Congress*, (1912) 1, 256-266.
- [3] NL. Braha, Some properties of new modified Szász-Mirakyan operators in polynomial weight spaces via power summability methods, *Bull Math Anal Appl.* 10 (3) (2018), 53-65.
- [4] NL. Braha, Some properties of Baskakov-Schurer-Szasz operators via power summability methods, *Quaestiones Mathematicae* 42(10) (2019), 1411-1426.
- [5] NL. Braha and U. Kadak, Approximation properties of the generalized Szasz operators by multiple Appell polynomials via power summability method, *Mathematical Methods in the Applied Sciences* 43(5) (2020), 2337-2356.
- [6] E. Yavuz, Ö. Talo, Approximation of continuous periodic functions of two variables via power series methods of summability. *Bull. Malays. Math. Sci. Soc.* 2019, 42, 1709–1717.
- [7] K. Demirci, S.Yıldız, and F. Dirik, Approximation via power series method in two-dimensional weighted spaces, *Bull. Malays. Math. Sci. Soc.* (2020), 1-13.
- [8] D. Söylemez and M. Ünver, Korovkin type theorems for Cheney–Sharma Operators via summability methods, *Results in Mathematics*, 72 (3) (2017), 1601-1612.
- [9] F. Özger, H. M. Srivastava, S. A. Mohiuddine, Approximation of functions by a new class of generalized Bernstein-Schurer operators, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM* (2020) 114:173.
- [10] S.A. Mohiuddine, N. Ahmad, F. Özger, et al. Approximation by the Parametric Generalization of Baskakov-Kantorovich Operators Linking with Stancu Operators. *Iran J Sci Technol Trans Sci* (2021). <https://doi.org/10.1007/s40995-020-01024-w>
- [11] A. Alotaibi, F. Özger, S.A. Mohiuddine et al. Approximation of functions by a class of Durrmeyer–Stancu type operators which includes Euler’s beta function. *Adv Differ Equ* 2021, 13 (2021). DOI: 10.1186/s13662-020-03164-0
- [12] F. Özger, Applications of generalized weighted statistical convergence to approximation theorems for functions of one and two variables, *Numerical Functional Analysis and Optimization*, 41(16): 1990-2006, (2020). DOI: 10.1080/01630563.2020.1868503
- [13] U. Kadak, F. Özger, Generalized Bernstein operators associated with multiple shape parameters, under review.
- [14] U. Kadak, F. Özger, A numerical comparative study of generalized Bernstein-Kantorovich operators, *Mathematical Foundations of Computing*, (2021), 4(2).
- [15] F. Moricz, Statistical convergence of multiple sequences, *Arch. Math.* 81 no.1, 82-89 (2003).
- [16] HI. Miller, A -statistical convergence of subsequence of double sequences, *Boll. U.M.I.* 8 (2007), 727-739.
- [17] A. Pringsheir, Zur theorie der zweifach unendlichen zahlenfolgen, *Math. Ann.* 53, 289-321 (1900).
- [18] GM. Robison, Divergent double sequences and series, *Amer. Math. Soc. Transl.* 28, 50-73 (1926).
- [19] F. Dirik and K. Demirci, Korovkin type approximation theorem for functions of two variables in statistical sense, *Turkish J. Math.*, 34(2010), 73-83.
- [20] M. Malkowsky, V. Velickovic, The duals of certain matrix domains of factorable triangles and some related visualisations, *Filomat* 27:5 (2013), 821–829.
- [21] M. Malkowsky, V. Velickovic, Some new sequence spaces, their duals and a connection with Wulff’s crystal, *MATCH Commun. Math. Comput. Chem.* 67 (2012) 589-607
- [22] H. M. Srivastava, F. Özger, S. A. Mohiuddine, Construction of Stancu-type Bernstein operators based on Bézier bases with shape parameter λ , *Symmetry* 11(3) (2019), Article 316.
- [23] H. M. Srivastava, K. J. Ansari, F. Özger, Z. Ödemiş Özger. A link between approximation theory and summability methods via four-dimensional infinite matrices. *Mathematics* 2021, 9(16), 1895. <https://doi.org/10.3390/math9161895>
- [24] F. Özger, Weighted statistical approximation properties of univariate and bivariate λ -Kantorovich operators, *Filomat*, 33(11), (2019) 473-486.
- [25] G. Hu, C. Bo, X. Qin, Continuity conditions for q -Bezier curves of degree n , *J. Inequal. Appl.* (2017) 2017:115.
- [26] X. Han, Y.C. Ma, X.L. Huang, A novel generalization of Bezier curve and surface, *J. Comput. Appl. Math.* 217 (2008) 180–193.
- [27] S. Baron, U. Stadtmüller, Tauberian theorems for power series methods applied to double sequences, *J. Math. Anal. Appl.* 211(2) (1997), 574-589.
- [28] X. Qin, G. Hu, N. Zhang, X. Shen, Y. Yang, A novel extension to the polynomial basis functions describing Bezier curves and surfaces of degree n with multiple shape parameters, *Appl. Math. Comput.* 223 (2013) 1–16.
- [29] U. Kadak, Modularly weighted four dimensional matrix summability with application to Korovkin type approximation theorem, *J. Math. Anal. Appl.* 468, 1(1) 2018, 38-63.
- [30] U. Kadak, N.L. Braha, H.M. Srivastava, Statistical weighted B-summability and its applications to approximation theorems, *Appl. Math. Comput.* 302 (2017) 80-96.
- [31] S. A. Mohiuddine, Statistical weighted A -summability with application to Korovkins type approximation theorem, *J. Inequal. Appl.* 2016 (2016) Article ID 101.
- [32] F. Özger, On new Bézier bases with Schurer polynomials and corresponding results in approximation theory, *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 69(1) (2020) 376-393.
- [33] F. Özger, K. Demirci, S. Yıldız, Approximation by Kantorovich variant of λ -Schurer operators and related numerical results, *Topics in Contemporary Mathematical Analysis and Applications*, Boca Raton, USA: CRC Press, ISBN 9781003081197, 77-94, 2020.
- [34] S. A. Mohiuddine, F. Özger, Approximation of functions by Stancu variant of Bernstein–Kantorovich operators based on shape parameter α , *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM* 114, 70 (2020).

- [35] O.H.H. Edelya, M. Mursaleen, A. Khan, Approximation for periodic functions via weighted statistical convergence. *Appl. Math. Comput.* 2013, 219, 8231–8236.
- [36] S. Karateke, Ç. Atakut, İ. Büyükyazıcı, Approximation by generalized integral Favard-Szász type operators involving Sheffer polynomials, *Filomat*, (2019) 33(7), 1921-1935.
- [37] A. Alotaibi, M. Nasiruzzaman, M. Mursaleen, Approximation by Phillips operators via q -Dunkl generalization based on a new parameter, *Journal of King Saud University-Science*, (2021) 33(4), 101413.
- [38] S. A. Mohiuddine, A. Kajla, M. Mursaleen, M. A. Alghamdi, Blending type approximation by τ -Baskakov-Durrmeyer type hybrid operators, *Adv Differ Equ*, (2020) 2020(1), 1-12.
- [39] R. Aslan, A. İzgi, Approximation by one and two variables of the Bernstein-Schurer-type operators and associated GBS operators on symmetrical mobile interval. *Journal of Function Spaces*, 2021 (2021), 9979286. <https://doi.org/10.1155/2021/9979286>
- [40] M. Yesilkayagil, F. Başar, Comparison of four dimensional summability methods. *Aligarh Bull. Math.*, (2015) 33(1-2), 1-11.
- [41] M. Yesilkayagil, F. Başar, Note on Abel summability of double series. *Numerical Functional Analysis and Optimization*, (2017) 38(8), 1069-1076.
- [42] V. K. Weierstrass, Ueber die analytische Darstellbarkeit sogenannter willkürlicher Functionen einer reellen Veränderlichen, in: *Sitzungsberichte der Akademie zu Berlin*, 1885, pp.633-639, 789-805.