



Approximation and existence of fixed points via interpolative enriched contractions

Shivam Rawat^{a,*}, Ayush Bartwal^b, R.C. Dimri^a

^aDepartment of Mathematics, H.N.B. Garhwal University, Uttarakhand 246174, India

^bDepartment of Mathematics, Himwant Kavi Chandra Kunwar Bartwal Govt. P.G. College, Nagnath Pokhari, Uttarakhand 246473, India

Abstract. In this paper, we define interpolative enriched contractions of Kannan type, Hardy-Rogers type and Matkowski type, by enriching existing interpolative contractions, in the setting of convex metric space. For these newly introduced contractions, we prove existence of fixed points and approximation results using Krasnoselskij iteration. Examples are also given to indicate the relevance of our results in comparison to some of the existing ones in the literature.

1. Introduction

Banach developed the Banach contraction principle [6], a fundamental consequence of fixed point theory on metric spaces, in 1922. This approach has been developed and extended in numerous ways as a result of its applications in diverse domains of nonlinear analysis and applied mathematical analysis. As the map following the Banach contraction principle is a continuous map, it was logical to ask whether a discontinuous map with similar contractive criteria has a fixed point or not. Kannan [17] gave an affirmative answer to this question by defining a contractive condition for a discontinuous map T as follows:

$$d(Tx, Ty) \leq c[d(x, Tx) + d(y, Ty)],$$

for all $x, y \in X$ and $c \in [0, \frac{1}{2})$. Moreover, he proved the existence and uniqueness of fixed points in the setting of complete metric spaces.

Takahashi [37] defined a convex structure in a metric space and referred to it as a convex metric space. He also investigated numerous features of this space to verify the presence of a fixed point for nonexpansive mappings in the context of convex metric space. Introducing convexity provides a basic tool for the building of different fixed point iterative algorithms ([2],[7]-[9],[32]-[36],[38]). Recently, Berinde [11] incorporated enrichment to contractive type mappings in the setting of Banach spaces in order to generalize the literature. A self-mapping T on X is called an enriched contraction or (b, θ) -enriched contraction if there exist two constants, $b \in [0, \infty)$ and $\theta \in [0, b + 1)$ such that for all $x, y \in X$

$$\|b(x - y) + Tx - Ty\| \leq \theta\|x - y\|.$$

2020 *Mathematics Subject Classification.* 47H10, 41A65, 46L52.

Keywords. Convex metric space; Interpolative enriched contractions; Fixed point.

Received: 28 September 2022; Accepted: 13 November 2022

Communicated by Erdal Karapinar

The first author would like to thank CSIR-HRDG Fund, under grant 09/386(0064)/2019-EMR-I, for financial support.

* Corresponding author: Shivam Rawat

Email addresses: rawat.shivam09@gmail.com (Shivam Rawat), ayushbartwal@gmail.com (Ayush Bartwal), dimrirc@gmail.com (R.C. Dimri)

The fact that enriched contraction is an extension of the Banach contraction class has supported the idea that in the context of a Banach space, Hilbert space and a convex metric space, a fixed point x^* will exist and Krasnoselskij iteration can approximate that point [10, 12, 13].

In 2018, Karapinar [18] used the technique of interpolation to revisit the Kannan type contraction. However, prior to Karapinar, interpolative theory was a vital technique in functional analysis. In 1982, Krein et al. [27] defined the notion of interpolative triples in the following way. Two Banach spaces A and B are known as a Banach couple denoted as (A, B) , when they are algebraically and topologically embedded in different topological linear space. If the embedding $A \cap B \subset E \subset A + B$ holds, the Banach space E is said to be an intermediate for the spaces of the Banach pair (A, B) . Let (A, B) and (C, D) be two Banach couples. If the limitations of T to the spaces A and B are, respectively bounded operators from A to C and B to D , then the linear mapping T operating from the space $A + B$ to $C + D$ is referred to as a bounded operator from (A, B) to (C, D) . The linear space of all bounded operators from (A, B) to (C, D) is denoted by $L(AB, CD)$. This is a Banach space in the norm $\|T\|_{L(AB, CD)} = \max\{\|T\|_{A \rightarrow B}, \|T\|_{C \rightarrow D}\}$. Let (A, B) and (C, D) be two Banach couples, and E (respectively F) be an intermediate for the spaces of the Banach couple (A, B) (respectively (C, D)). The triple (A, B, E) is called an interpolation triple, relative to (C, D, F) , if every bounded operator from (A, B) to (C, D) maps E to F . A triple (A, B, E) is said to be an interpolation triple of type α ($0 \leq \alpha \leq 1$), relative to (C, D, F) , if it is an interpolation triple and the following inequality holds: $\|T\|_{E \rightarrow F} \leq c\|T\|^\alpha \cdot \|T\|^{1-\alpha}$. Karapinar defined the interpolative Kannan type contraction as follows:

$$d(Tx, Ty) \leq \lambda([d(x, Tx)]^\alpha \cdot [d(y, Ty)]^{1-\alpha}),$$

for all $x, y \in X \setminus \text{Fix}(T)$, where $\text{Fix}(T) = \{x \in X, Tx = x\}$ and $\lambda \in [0, 1)$. Some recent works in the direction of interpolative Ćirić-Reich-Rus type contraction [4, 5, 31], Meir-Keeler type contraction [20, 25] may be referred to. Many authors followed this new dimension in the study of fixed point theory [1, 3, 14–16, 19, 22–24, 26, 30, 39].

Motivated by the recent results, in this paper, we define interpolative enriched contractions of Kannan type, Hardy-Rogers type and Matkowski type to prove existence of fixed points and approximation results using Krasnoselskij iteration in a convex metric space. Examples are also given in order to indicate the relevance of our newly obtained results in comparison to some of the existing ones in the literature.

2. Preliminaries

Definition 2.1. [18] Let (X, d) be a metric space. A mapping $T : X \rightarrow X$ is said to be an interpolative Kannan contraction mapping if there exist some $c \in [0, 1)$ and $\alpha \in (0, 1)$ such that

$$d(Tx, Ty) \leq c[d(x, Tx)]^\alpha \cdot [d(y, Ty)]^{1-\alpha}$$

for all $x, y \in X$ with $x \neq Tx$.

The main result in [18] is the following.

Theorem 2.2. [18] Let (X, d) be a complete metric space and T be an interpolative Kannan type contraction. Then T has a unique fixed point in X .

Karapinar, Agarwal and Aydi [21] gave a counter-example to Theorem 2.2, showing that the fixed point may be not unique. The revised version of Theorem 2.2 is the following.

Theorem 2.3. [21] Let (X, d) be a complete metric space. Let $T : X \rightarrow X$ be a given mapping such that

$$d(Tx, Ty) \leq c[d(x, Tx)]^\alpha \cdot [d(y, Ty)]^{1-\alpha},$$

for all $x, y \in X \setminus \text{Fix}(T)$. Then T has a fixed point in X .

Definition 2.4. [22] Let (X, d) be a metric space. We say that the self-mapping $T : X \rightarrow X$ is an interpolative Hardy-Rogers type contraction if there exist $c \in [0, 1)$ and $\alpha, \beta, \gamma \in (0, 1)$ with $\alpha + \beta + \gamma < 1$, such that

$$d(Tx, Ty) \leq c[d(x, y)]^\beta \cdot [d(x, Tx)]^\alpha \cdot [d(y, Ty)]^\gamma \cdot \left[\frac{1}{2}(d(x, Ty) + d(y, Tx))\right]^{1-\alpha-\beta-\gamma},$$

for all $x, y \in X \setminus \text{Fix}(T)$.

Theorem 2.5. [22] Let (X, d) be a complete metric space and T be an interpolative Hardy-Rogers type contraction. Then, T has a fixed point in X .

Definition 2.6. [28] Let Φ be a set of functions, $\phi : [0, \infty) \rightarrow [0, \infty)$ such that

- (i) ϕ is non decreasing,
- (ii) $\lim_{n \rightarrow \infty} \phi^n(t) = 0$ for each $t > 0$.

Lemma 2.7. [29] Let $\phi \in \Phi$, then $\phi(t) < t$ for all $t > 0$ and $\phi(0) = 0$.

Definition 2.8. [23] Let (X, d) be a metric space. We say that the self mapping $T : X \rightarrow X$ is an interpolative Matkowski type contraction, if there exist $\alpha, \beta, \gamma \in (0, 1)$ and $\phi \in \Phi$ such that

$$d(Tx, Ty) \leq \phi([d(x, y)]^\beta \cdot [d(x, Tx)]^\alpha \cdot [d(y, Ty)]^\gamma \cdot \left[\frac{1}{2}(d(x, Ty) + d(y, Tx))\right]^{1-\alpha-\beta-\gamma}),$$

for all $x, y \in X \setminus \text{Fix}(T)$.

Theorem 2.9. [23] Let (X, d) be a complete metric space and T be an interpolative Matkowski type contraction. Then T has a fixed point in X .

Now, we give some basic preliminaries regarding convex metric spaces.

Definition 2.10. [37] Let X be a metric space. A continuous function $W : X \times X \times [0, 1] \rightarrow X$ is said to be a convex structure on X , if for all $x, y \in X$ and $\lambda \in [0, 1]$ the following inequality holds

$$d(u, W(x, y; \lambda)) \leq \lambda d(u, x) + (1 - \lambda)d(u, y), \text{ for any } u \in X. \quad (1)$$

A metric space X endowed with a convex structure W is called a Takahashi convex metric space or simply a convex metric space and is usually denoted by (X, d, W) .

Example 2.11. [37] Let I be the unit interval $[0, 1]$ and X be the family of closed intervals $[a_i, b_i]$, such that $0 \leq a_i \leq b_i \leq 1$. For $I_i = [a_i, b_i]$, $I_j = [a_j, b_j]$ and $\lambda (0 \leq \lambda \leq 1)$, we define a mapping W by $W(I_i, I_j; \lambda) = [\lambda a_i + (1 - \lambda)a_j, \lambda b_i + (1 - \lambda)b_j]$ and define a metric d on X by the Hausdorff distance, i.e.

$$d(I_i, I_j) = \sup_{a \in I_i} \{ \inf_{b \in I_j} \{|a - b|\} - \inf_{c \in I_j} \{|a - c|\} \}.$$

(X, d, W) is a convex metric space.

The next lemmas present some fundamental properties of a convex metric space.

Lemma 2.12. [37] Let (X, d, W) be a convex metric space. For all $x, y \in X$ and any $\lambda \in [0, 1]$, we have

$$d(x, y) = d(x, W(x, y; \lambda)) + d(W(x, y; \lambda), y).$$

Lemma 2.13. [37] Let (X, d, W) be a convex metric space. For all $x, y \in X$ and any $\lambda \in [0, 1]$, we have

$$d(x, W(x, y; \lambda)) = (1 - \lambda)d(x, y) \text{ and } d(W(x, y; \lambda), y) = \lambda d(x, y).$$

Lemma 2.14. [2] Let (X, d, W) be a convex metric space. For each $x, y \in X$ and $\lambda, \lambda_1, \lambda_2 \in [0, 1]$, we have the following:

1. $W(x, x; \lambda) = x; W(x, y; 0) = y$ and $W(x, y; 1) = x$.
2. $|\lambda_1 - \lambda_2|d(x, y) \leq d(W(x, y; \lambda_1), W(x, y; \lambda_2))$.

Lemma 2.15. [2] Let (X, d, W) be a convex metric space and $T : X \rightarrow X$ be a self mapping. Define the mapping $T_\lambda : X \rightarrow X$ by

$$T_\lambda x = W(x, Tx; \lambda), \quad x \in X. \quad (2)$$

Then, for any $\lambda \in [0, 1)$, we have $\text{Fix}(T) = \text{Fix}(T_\lambda)$.

In 2021, Berinde and Păcurar [13] introduced the notion of enriched contraction in convex metric spaces and proved some fixed point results.

Definition 2.16. [13] Let (X, d, W) be a convex metric space. A mapping $T : X \rightarrow X$ is said to be an enriched contraction if there exist $c \in [0, 1)$ and $\lambda \in [0, 1)$ such that

$$d(W(x, Tx; \lambda), W(y, Ty; \lambda)) \leq cd(x, y),$$

for all $x, y \in X$.

Theorem 2.17. [13] Let (X, d, W) be a complete convex metric space and let $T : X \rightarrow X$ be an enriched contraction. Then,

1. $\text{Fix}(T) = \{p\}$ for some $p \in X$.
2. The sequence $\{x_n\}_{n=0}^\infty$ obtained from the iterative process

$$x_{n+1} = W(x_n, Tx_n; \lambda) \rightarrow p, \quad \text{for any } x_0 \in X \text{ and } n \geq 0. \quad (3)$$

3. The following estimate holds

$$d(x_{n+i-1}, p) \leq \frac{c^i}{1-c} \cdot d(x_n, x_{n-1}), \quad n = 1, 2, \dots; \quad i = 1, 2, \dots \quad (4)$$

3. Main results

We enrich the interpolative Kannan type contraction as follows:

Definition 3.1. Let (X, d, W) be a convex metric space. A self-mapping $T : X \rightarrow X$ is an interpolative enriched Kannan type contraction if there exist $\lambda \in [0, 1)$, $c \in [0, 1)$ and $\alpha \in (0, 1)$, such that

$$d(W(x, Tx; \lambda), W(y, Ty; \lambda)) \leq c[d(x, W(x, Tx; \lambda))]^\alpha \cdot [d(y, W(y, Ty; \lambda))]^{1-\alpha} \quad (5)$$

for all $x, y \in X \setminus \text{Fix}(T)$. We also call T an interpolative (λ, k) -enriched Kannan type contraction.

It is easily seen that an interpolative $(0, k)$ -enriched Kannan type contraction is a usual interpolative Kannan type contraction.

Theorem 3.2. Let (X, d, W) be a complete convex metric space. $T : X \rightarrow X$ be a continuous interpolative enriched Kannan type contraction. Then,

1. $\text{Fix}(T) \neq \emptyset$.
2. The following estimate holds

$$d(x_{n+i-1}, p) \leq \frac{c^i}{1-c} \cdot d(x_n, x_{n-1}), \quad n = 1, 2, \dots; \quad i = 1, 2, \dots \quad (6)$$

Proof. Using the interpolative enriched Kannan type contraction condition (5), the mapping $T_\lambda : X \rightarrow X$ defined by (2) satisfies

$$d(T_\lambda x, T_\lambda y) \leq c[d(x, T_\lambda x)]^\alpha \cdot [d(y, T_\lambda y)]^{1-\alpha} \quad (7)$$

that is, T_λ is an interpolative Kannan type contraction. The Picard iteration associated with T_λ is actually the Krasnoselskij iterative process $\{x_n\}_{n=0}^\infty$ associated to T , which is defined by $x_{n+1} = W(x_n, Tx_n; \lambda)$, i.e.,

$$x_{n+1} = T_\lambda x_n, \quad n \geq 0. \quad (8)$$

Without loss of generality, we assume that $x_n \neq x_{n+1}$ for each nonnegative integer n . Indeed, if there exists a nonnegative integer n_0 , such that $x_{n_0} = x_{n_0+1} = T_\lambda x_{n_0}$. Then x_{n_0} is a fixed point. Thus, we have

$$d(x_n, T_\lambda x_n) = d(x_n, x_{n+1}) > 0, \quad \text{for each nonnegative integer } n.$$

Taking $x = x_n$ and $y = x_{n-1}$ in (7), we obtain

$$d(x_{n+1}, x_n) \leq c[d(x_n, x_{n+1})]^\alpha \cdot [d(x_{n-1}, x_n)]^{1-\alpha},$$

which further gives

$$[d(x_n, x_{n+1})]^{1-\alpha} \leq c[d(x_{n-1}, x_n)]^{1-\alpha}.$$

We deduce that

$$d(x_n, x_{n+1}) \leq c d(x_{n-1}, x_n) \leq c^n d(x_0, x_1). \quad (9)$$

Since, $c < 1$ and letting $n \rightarrow \infty$ in the above inequality, we get $d(x_n, x_{n+1}) \rightarrow 0$. Now, we will prove that $\{x_n\}$ is a Cauchy sequence. Using the triangular inequality, we obtain

$$\begin{aligned} d(x_n, x_{n+r}) &\leq d(x_n, x_{n+1}) + \dots + d(x_{n+r-1}, x_{n+r}) \\ &\leq c^n d(x_0, x_1) + \dots + c^{n+r-1} d(x_0, x_1) \\ &\leq \frac{c^n}{1-c} (1-c^r) d(x_0, x_1). \end{aligned} \quad (10)$$

Letting $n \rightarrow \infty$ in the above inequality, we get that $\{x_n\}$ is a Cauchy sequence. Since, (X, d, W) is a complete convex metric space, there exists a $x^* \in X$ such that $\lim_{n \rightarrow \infty} d(x_n, x^*) = 0$.

On substituting $x = x_n$ and $y = x^*$ in (7), we obtain

$$d(T_\lambda x_n, T_\lambda x^*) \leq c[d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x^*, T_\lambda x^*)]^{1-\alpha}.$$

Letting $n \rightarrow \infty$, we obtain $T_\lambda x^* = x^*$, i.e., $x^* \in \text{Fix}(T_\lambda)$.

To prove the last statement of our theorem, we first deduce the following result from (9),

$$d(x_{n+r}, x_n) \leq \frac{c}{1-c} d(x_n, x_{n-1}). \quad (11)$$

Now, letting $r \rightarrow \infty$ in (10) and (11), we get

$$d(x_n, x^*) \leq \frac{c^n}{1-c} d(x_0, x_1), \quad n \geq 1, \quad (12)$$

and

$$d(x_n, x^*) \leq \frac{c}{1-c} d(x_n, x_{n-1}), \quad n \geq 1, \quad (13)$$

respectively. From (12) and (13), we get the unifying error estimate (6). \square

Example 3.3. Let $X = [0, 5]$ and define the distance function $d : X \times X \rightarrow [0, \infty)$ as $d(x, y) = |x - y|$, for any $x, y \in X$. The mapping $W : X \times X \times [0, 1] \rightarrow X$ is defined as $W(x, y; \lambda) = \lambda x + (1 - \lambda)y$, for all $x, y \in X$ and let $T : X \rightarrow X$ be given by

$$Tx = \frac{5 - x}{9}. \quad (14)$$

For any $x, y, z \in X$, we have

$$\begin{aligned} d(z, W(x, y, \lambda)) &= |\lambda(z - x) + (1 - \lambda)(z - y)| \\ &\leq \lambda|z - x| + (1 - \lambda)|z - y| \\ &= \lambda d(z, x) + (1 - \lambda)d(z, y). \end{aligned}$$

Hence, (X, d, W) is a convex metric space. Also, for $\lambda = \frac{1}{10}$

$$W(x, Tx; \lambda) = T_{\lambda}x = \frac{1}{2}. \quad (15)$$

So, we have $d(T_{\lambda}x, T_{\lambda}y) = 0$, where $x, y \in [0, 5] \setminus \text{Fix}(T)$. Therefore, for any $\alpha \in (0, 1)$, T is an interpolative enriched Kannan type contraction. Thus, by Theorem 3.2, T has a fixed point. In this case, $\text{Fix}(T) = \{\frac{1}{2}\}$.

Corollary 3.4. Let (X, d) be a complete metric space and T be an interpolative Kannan type contraction. Then T has a fixed point in X .

Proof. Taking $\lambda = 0$ in the above theorem, the result holds. \square

Now, we obtain fixed point results by enriching the interpolative Hardy-Rogers type contraction as given in the following definition.

Definition 3.5. Let (X, d, W) be a convex metric space. A self-mapping $T : X \rightarrow X$ is an interpolative enriched Hardy-Rogers type contraction if there exist $\lambda \in [0, 1)$, $c \in [0, 1)$ and $\alpha, \beta, \gamma \in (0, 1)$ with $\alpha + \beta + \gamma < 1$, such that

$$\begin{aligned} d(W(x, Tx; \lambda), W(y, Ty; \lambda)) &\leq c[d(x, y)]^{\beta} \cdot [d(x, W(x, Tx; \lambda))]^{\alpha} \cdot [d(y, W(y, Ty; \lambda))]^{\gamma} \\ &\quad \cdot \left[\frac{1}{2}(d(x, W(y, Ty; \lambda)) + d(y, W(x, Tx; \lambda))) \right]^{1-\alpha-\beta-\gamma} \quad (16) \end{aligned}$$

for all $x, y \in X \setminus \text{Fix}(T)$. We also call T an interpolative (λ, k) -enriched Hardy-Rogers type contraction.

It is easily seen that an interpolative $(0, k)$ -enriched Hardy-Rogers type contraction is a usual interpolative Hardy-Rogers type contraction.

Theorem 3.6. Let (X, d, W) be a complete convex metric space. $T : X \rightarrow X$ be a continuous interpolative enriched Hardy-Rogers type contraction. Then,

1. $\text{Fix}(T) \neq \emptyset$.
2. The following estimate holds

$$d(x_{n+i-1}, p) \leq \frac{c^i}{1-c} \cdot d(x_n, x_{n-1}), \quad n = 1, 2, \dots; \quad i = 1, 2, \dots \quad (17)$$

Proof. Using the interpolative enriched Hardy-Rogers type contraction condition (16), we have that the mapping $T_{\lambda} : X \rightarrow X$ defined by (2) satisfies

$$d(T_{\lambda}x, T_{\lambda}y) \leq c[d(x, y)]^{\beta} \cdot [d(x, T_{\lambda}x)]^{\alpha} \cdot [d(y, T_{\lambda}y)]^{\gamma} \cdot \left[\frac{1}{2}(d(x, T_{\lambda}y) + d(y, T_{\lambda}x)) \right]^{1-\alpha-\beta-\gamma}, \quad (18)$$

that is, T_λ is an interpolative Hardy-Rogers type contraction. The Picard iteration associated with T_λ is actually the Krasnoselskij iterative process $\{x_n\}_{n=0}^\infty$ associated to T , which is defined by $x_{n+1} = W(x_n, Tx_n; \lambda)$, i.e.,

$$x_{n+1} = T_\lambda x_n, \quad n \geq 0. \quad (19)$$

Without loss of generality, we assume that $x_n \neq x_{n+1}$ for each nonnegative integer n . Indeed, if there exist a nonnegative integer n_0 , such that $x_{n_0} = x_{n_0+1} = T_\lambda x_{n_0}$. Then x_{n_0} is a fixed point. Thus, we have

$$d(x_n, T_\lambda x_n) = d(x_n, x_{n+1}) > 0, \quad \text{for each nonnegative integer } n.$$

Taking $x = x_n$ and $y = x_{n-1}$ in (18), we obtain

$$\begin{aligned} d(x_{n+1}, x_n) &\leq c[d(x_n, x_{n-1})]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x_{n-1}, T_\lambda x_{n-1})]^\gamma \cdot \left[\frac{1}{2}(d(x_n, T_\lambda x_{n-1}) + d(x_{n-1}, T_\lambda x_n))\right]^{1-\alpha-\beta-\gamma} \\ &\leq c[d(x_n, x_{n-1})]^\beta \cdot [d(x_n, x_{n+1})]^\alpha \cdot [d(x_{n-1}, x_n)]^\gamma \cdot \left[\frac{1}{2}(d(x_n, x_n) + d(x_{n-1}, x_{n+1}))\right]^{1-\alpha-\beta-\gamma} \\ &\leq c[d(x_n, x_{n-1})]^\beta \cdot [d(x_n, x_{n+1})]^\alpha \cdot [d(x_{n-1}, x_n)]^\gamma \cdot \left[\frac{1}{2}(d(x_{n-1}, x_n) + d(x_n, x_{n+1}))\right]^{1-\alpha-\beta-\gamma}. \end{aligned} \quad (20)$$

Suppose that $d(x_{n-1}, x_n) < d(x_n, x_{n+1})$ for some $n \geq 1$. Thus,

$$\frac{1}{2}(d(x_{n-1}, x_n) + d(x_n, x_{n+1})) \leq d(x_n, x_{n+1}).$$

Consequently, the inequality (20) yields that

$$[d(x_n, x_{n+1})]^{\beta+\gamma} \leq c[d(x_{n-1}, x_n)]^{\beta+\gamma},$$

which implies that

$$d(x_n, x_{n+1}) \leq d(x_{n-1}, x_n),$$

which is a contradiction. Thus, we have $d(x_n, x_{n+1}) \leq d(x_{n-1}, x_n)$ for all $n \geq 1$. Hence, $\{d(x_{n-1}, x_n)\}$ is a non-increasing sequence with positive terms. Set $l = \lim_{n \rightarrow \infty} d(x_{n-1}, x_n)$. We have

$$\frac{1}{2}(d(x_{n-1}, x_n) + d(x_n, x_{n+1})) \leq d(x_n, x_{n+1}), \quad \text{for all } n \geq 1.$$

By a simple elimination, the inequality (20) implies that

$$[d(x_n, x_{n+1})]^{1-\alpha} \leq c[d(x_{n-1}, x_n)]^{1-\alpha}, \quad \text{for all } n \geq 1.$$

We deduce that

$$d(x_n, x_{n+1}) \leq c d(x_{n-1}, x_n) \leq c^n d(x_0, x_1). \quad (21)$$

Since, $c < 1$ and letting $n \rightarrow \infty$ in the above inequality, we get $l = 0$. Now, we will prove that $\{x_n\}$ is a Cauchy sequence. Using the triangular inequality, we obtain

$$\begin{aligned} d(x_n, x_{n+r}) &\leq d(x_n, x_{n+1}) + \dots + d(x_{n+r-1}, x_{n+r}) \\ &\leq c^n d(x_0, x_1) + \dots + c^{n+r-1} d(x_0, x_1) \\ &\leq \frac{c^n}{1-c} (1-c^r) d(x_0, x_1). \end{aligned} \quad (22)$$

Letting $n \rightarrow \infty$ in the above inequality, we get that $\{x_n\}$ is a Cauchy sequence. Since, (X, d, W) is a complete convex metric space, there exists a $x^* \in X$ such that $\lim_{n \rightarrow \infty} d(x_n, x^*) = 0$.

On substituting $x = x_n$ and $y = x^*$ in (18), we obtain

$$d(T_\lambda x_n, T_\lambda x^*) \leq c[d(x_n, x^*)]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x^*, T_\lambda x^*)]^\gamma \cdot \left[\frac{1}{2}(d(x_n, T_\lambda x^*) + d(x^*, T_\lambda x_n))\right]^{1-\alpha-\beta-\gamma}.$$

Letting $n \rightarrow \infty$, we obtain $d(x^*, T_\lambda x^*) = 0$, i.e., $x^* \in \text{Fix}(T_\lambda)$.

To prove the last statement of our theorem, we first deduce the following result from (21),

$$d(x_{n+r}, x_n) \leq \frac{c}{1-c} d(x_n, x_{n-1}). \quad (23)$$

Now, letting $r \rightarrow \infty$ in (22) and (23), we get

$$d(x_n, x^*) \leq \frac{c^n}{1-c} d(x_0, x_1), n \geq 1, \quad (24)$$

and

$$d(x_n, x^*) \leq \frac{c}{1-c} d(x_n, x_{n-1}), n \geq 1, \quad (25)$$

respectively. From (24) and (25) we get the unifying error estimate (17). \square

Example 3.7. Let $X = [-1, 2]$ and define the distance function $d : X \times X \rightarrow [0, \infty)$ as $d(x, y) = |x - y|$, for any $x, y \in X$. The mapping $W : X \times X \times [0, 1] \rightarrow X$ is defined as $W(x, y; \lambda) = \lambda x + (1 - \lambda)y$, for all $x, y \in X$ and let $T : X \rightarrow X$ be given by

$$Tx = \begin{cases} \frac{35-2x}{38} & \text{if } x \in [-1, 0), \\ \frac{35}{38} & \text{if } x \in [0, 2]. \end{cases} \quad (26)$$

For any $x, y, z \in X$, we have

$$\begin{aligned} d(z, W(x, y, \lambda)) &= |\lambda(z - x) + (1 - \lambda)(z - y)| \\ &\leq \lambda|z - x| + (1 - \lambda)|z - y| \\ &= \lambda d(z, x) + (1 - \lambda)d(z, y). \end{aligned}$$

Hence, (X, d, W) is a convex metric space. Also, for $\lambda = \frac{1}{20}$

$$W(x, Tx; \lambda) = T_\lambda x = \begin{cases} \frac{7}{8} & \text{if } x \in [-1, 0), \\ \frac{x}{20} + \frac{7}{8} & \text{if } x \in [0, 2]. \end{cases} \quad (27)$$

Choose $c = \frac{1}{2}$, $\alpha = \frac{1}{3}$, $\beta = \frac{1}{2}$ and $\gamma = \frac{1}{7}$. Then, we have to check that (16) holds.

Case-I: For $x, y \in [-1, 0)$, we obtain $d(T_\lambda x, T_\lambda y) = 0$. Thus, (16) holds.

Case-II: For $x \in [-1, 0)$ and $y \in [0, 2]$, the left hand side of equation (16) becomes

$$d(T_\lambda x, T_\lambda y) = d\left(\frac{7}{8}, \frac{y}{20} + \frac{7}{8}\right) = \left|\frac{y}{20}\right| \leq \frac{1}{10},$$

where the maximum value on the left is attained for $y = 2$. Thus, for $y = 2$, the right hand side of the equation is

$$\begin{aligned} c \cdot [d(x, y)]^{\frac{1}{2}} \cdot [d(x, T_\lambda x)]^{\frac{1}{3}} \cdot [d(y, T_\lambda y)]^{\frac{1}{7}} \cdot \left[\frac{1}{2}(d(x, T_\lambda y) + d(y, T_\lambda x))\right]^{\frac{1}{42}} &> \frac{1}{2} \cdot (2)^{\frac{1}{2}} \cdot \left(\frac{7}{8}\right)^{\frac{1}{3}} \cdot \left(\frac{82}{80}\right)^{\frac{1}{7}} \cdot \left[\frac{1}{2}\left(\frac{78}{80} + \frac{9}{8}\right)\right]^{\frac{1}{42}} \\ &= (0.5) \cdot (2)^{\frac{1}{2}} \cdot (0.875)^{\frac{1}{3}} \cdot (1.025)^{\frac{1}{7}} \cdot (1.05)^{\frac{1}{42}} \\ &= 0.679502 > \frac{1}{10}. \end{aligned}$$

Therefore, (16) holds.

Case-III: For $x, y \in [0, 2]$, the left hand side of equation (16) becomes

$$d(T_\lambda x, T_\lambda y) = d\left(\frac{x}{20} + \frac{7}{8}, \frac{y}{20} + \frac{7}{8}\right) = \left|\frac{x-y}{20}\right| \leq \frac{1}{10},$$

where the maximum value on the left is attained when $x = 0$ and $y = 2$ or when $x = 2$ and $y = 0$. Thus, for $x = 0$ and $y = 2$, the right hand side of the equation is

$$\begin{aligned} c.[d(x, y)]^{\frac{1}{2}}.[d(x, T_\lambda x)]^{\frac{1}{3}}.[d(y, T_\lambda y)]^{\frac{1}{7}}.[\frac{1}{2}(d(x, T_\lambda y) + d(y, T_\lambda x))]^{\frac{1}{42}} &> \frac{1}{2} \cdot (2)^{\frac{1}{2}} \cdot \left(\frac{7}{8}\right)^{\frac{1}{3}} \cdot \left(\frac{82}{80}\right)^{\frac{1}{7}} \cdot \left[\frac{1}{2}\left(\frac{78}{80} + \frac{9}{8}\right)\right]^{\frac{1}{42}} \\ &= (0.5) \cdot (2)^{\frac{1}{2}} \cdot (0.875)^{\frac{1}{3}} \cdot (1.025)^{\frac{1}{7}} \cdot (1.05)^{\frac{1}{42}} \\ &= 0.679502 > \frac{1}{10}. \end{aligned}$$

Similarly, when $x = 2$ and $y = 0$, the above equation holds. Thus, in both the possibilities, (16) holds.

From all the above three cases, we obtain that T is an interpolative enriched Hardy-Rogers type contraction. Thus, by Theorem 3.6, T has a fixed point which is $\frac{35}{38}$.

Corollary 3.8. Let (X, d) be a complete metric space and T be an interpolative Hardy-Rogers type contraction. Then T has a fixed point in X .

Proof. Taking $\lambda = 0$ in the above theorem, the result holds. \square

Now, we obtain fixed point results by enriching the interpolative Matkowski type contraction as follows.

Definition 3.9. Let (X, d, W) be a convex metric space. A self mapping $T : X \rightarrow X$ is an interpolative enriched Matkowski type contraction if there exist $\lambda \in [0, 1)$, $\phi \in \Phi$ and $\alpha, \beta, \gamma \in (0, 1)$ such that

$$d(W(x, Tx; \lambda), W(y, Ty; \lambda)) \leq \phi([d(x, y)]^\beta \cdot [d(x, W(x, Tx; \lambda))]^\alpha \cdot [d(y, W(y, Ty; \lambda))]^\gamma \cdot [\frac{1}{2}(d(x, W(y, Ty; \lambda)) + d(y, W(x, Tx; \lambda)))]^{1-\alpha-\beta-\gamma}) \quad (28)$$

It is easily seen that an interpolative $(0, k)$ -enriched Matkowski type contraction is a usual interpolative Matkowski type contraction.

Theorem 3.10. Let (X, d, W) be a complete convex metric space. $T : X \rightarrow X$ be a continuous interpolative enriched Matkowski type contraction. Then T has a fixed point in X .

Proof. Using the interpolative enriched Matkowski type contraction condition (28), we have that the mapping $T_\lambda : X \rightarrow X$ defined by (2) satisfies

$$d(T_\lambda x, T_\lambda y) \leq \phi([d(x, y)]^\beta \cdot [d(x, T_\lambda x)]^\alpha \cdot [d(y, T_\lambda y)]^\gamma \cdot [\frac{1}{2}(d(x, T_\lambda y) + d(y, T_\lambda x))]^{1-\alpha-\beta-\gamma}), \quad (29)$$

that is, T_λ is an interpolative Matkowski type contraction. The Picard iteration associated with T_λ is actually the Krasnoselskij iterative process $\{x_n\}_{n=0}^\infty$ associated to T , which is defined by $x_{n+1} = W(x_n, Tx_n; \lambda)$, i.e.

$$x_{n+1} = T_\lambda x_n, n \geq 0. \quad (30)$$

Without loss of generality, we assume that $x_n \neq x_{n+1}$ for each nonnegative integer n . Indeed, if there exist a nonnegative integer n_0 , such that $x_{n_0} = x_{n_0+1} = T_\lambda x_{n_0}$. Then x_{n_0} is a fixed point. Thus, we have

$$d(x_n, T_\lambda x_n) = d(x_n, x_{n+1}) > 0, \text{ for each nonnegative integer } n.$$

Taking $x = x_n$ and $y = x_{n-1}$ in (29), we obtain

$$\begin{aligned} d(x_{n+1}, x_n) &\leq \phi([d(x_n, x_{n-1})]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x_{n-1}, T_\lambda x_{n-1})]^\gamma \cdot [\frac{1}{2}(d(x_n, T_\lambda x_{n-1}) + d(x_{n-1}, T_\lambda x_n))]^{1-\alpha-\beta-\gamma}) \\ &= \phi([d(x_n, x_{n-1})]^\beta \cdot [d(x_n, x_{n+1})]^\alpha \cdot [d(x_{n-1}, x_n)]^\gamma \cdot [\frac{1}{2}(d(x_n, x_n) + d(x_{n-1}, x_{n+1}))]^{1-\alpha-\beta-\gamma}) \\ &\leq \phi([d(x_n, x_{n-1})]^\beta \cdot [d(x_n, x_{n+1})]^\alpha \cdot [d(x_{n-1}, x_n)]^\gamma \cdot [\frac{1}{2}(d(x_{n-1}, x_n) + d(x_n, x_{n+1}))]^{1-\alpha-\beta-\gamma}). \end{aligned} \quad (31)$$

Suppose that

$$d(x_{n-1}, x_n) < d(x_n, x_{n+1}), \text{ for some } n \geq 1.$$

Then

$$\frac{1}{2}(d(x_{n-1}, x_n) + d(x_n, x_{n+1})) \leq d(x_n, x_{n+1}).$$

Consequently, the inequality (31) yields that

$$0 < d(x_n, x_{n+1}) \leq \phi([d(x_n, x_{n-1})]^{\beta+\gamma} \cdot [d(x_n, x_{n+1})]^{1-\beta-\gamma}) \leq [d(x_n, x_{n-1})]^{\beta+\gamma} \cdot [d(x_n, x_{n+1})]^{1-\beta-\gamma}, \quad (32)$$

which implies that

$$d(x_n, x_{n+1}) \leq d(x_n, x_{n-1}),$$

which is a contradiction. Thus we have $d(x_n, x_{n+1}) \leq d(x_n, x_{n-1})$, for all $n \geq 1$. Hence, $\{d(x_n, x_{n-1})\}$ is a non-increasing sequence of positive terms. Using (32), we have

$$d(x_n, x_{n+1}) \leq \phi([d(x_n, x_{n-1})]^{\beta+\gamma} [d(x_n, x_{n+1})]^{1-\beta-\gamma}) \leq \phi(d(x_{n-1}, x_n)).$$

On repeating the same argument, we have

$$d(x_n, x_{n+1}) \leq \phi(d(x_{n-1}, x_n)) \leq \cdots \leq \phi^n(d(x_0, x_1)). \quad (33)$$

As $\phi \in \Phi$, we have $\lim_{n \rightarrow \infty} \phi^n(d(x_0, x_1)) = 0$. Therefore, (33) implies that $\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0$. Now, we shall show that sequence $\{x_n\}$ is Cauchy. For any $\epsilon > 0$, there exists some $n_0 \in \mathbb{N}$, such that for $n \geq n_0$

$$\phi^n(d(x_0, x_1)) < \epsilon - \phi(\epsilon).$$

From (33), we have

$$0 < d(x_n, x_{n+1}) < \epsilon - \phi(\epsilon). \quad (34)$$

We claim that

$$0 < d(x_n, x_m) < \epsilon, \text{ for all } m \geq n \geq n_0. \quad (35)$$

We prove the above inequality by mathematical induction. Since $\epsilon - \phi(\epsilon) < \epsilon$, from (34), we conclude that (35) holds for $m = n + 1$. Now, suppose that (35) holds for $m = k$. For $m = k + 1$, we have

$$\begin{aligned} d(x_n, x_{k+1}) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{k+1}) \\ &\leq \epsilon - \phi(\epsilon) + d(T_\lambda x_n, T_\lambda x_k) \\ &\leq \epsilon - \phi(\epsilon) + \phi([d(x_n, x_k)]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x_k, T_\lambda x_k)]^\gamma \cdot [\frac{1}{2}(d(x_n, T_\lambda x_k) + d(x_k, T_\lambda x_n))]^{1-\alpha-\beta-\gamma}). \end{aligned}$$

Since $k \geq n \geq n_0$, we have

$$[d(x_n, x_k)]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x_k, T_\lambda x_k)]^\gamma \cdot \left[\frac{1}{2}(d(x_n, T_\lambda x_k) + d(x_k, T_\lambda x_n)) \right]^{1-\alpha-\beta-\gamma} \leq \epsilon.$$

Thus, we have

$$d(x_n, x_{k+1}) \leq \epsilon - \phi(\epsilon) + \phi(\epsilon) = \epsilon.$$

So, (35) holds for $m = k + 1$, which implies that the sequence $\{x_n\}$ is Cauchy. Therefore, there exists $x^* \in X$, such that $\lim_{n \rightarrow \infty} d(x_n, x^*) = 0$. Suppose that $x^* \neq T_\lambda x^*$. Since $x_n \neq T_\lambda x_n$ for each n , by (29), we have

$$d(x_{n+1}, T_\lambda x^*) = d(T_\lambda x_n, T_\lambda x^*) \leq \phi([d(x_n, x^*)]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x^*, T_\lambda x^*)]^\gamma \cdot \left[\frac{1}{2}(d(x_n, T_\lambda x^*) + d(x^*, T_\lambda x_{n+1})) \right]^{1-\alpha-\beta-\gamma}). \quad (36)$$

It is obvious that there exists $N \in \mathbb{N}$, such that for each $n \geq N$,

$$[d(x_n, x^*)]^\beta \cdot [d(x_n, T_\lambda x_n)]^\alpha \cdot [d(x^*, T_\lambda x^*)]^\gamma \cdot \left[\frac{1}{2}(d(x_n, T_\lambda x^*) + d(x^*, T_\lambda x_{n+1})) \right]^{1-\alpha-\beta-\gamma} < d(x^*, T_\lambda x^*).$$

Since ϕ is non decreasing, by insertion of inequality (36), we have

$$d(x_{n+1}, T_\lambda x^*) \leq \phi(d(x^*, T_\lambda x^*)), \text{ for all } n \geq N. \quad (37)$$

Letting $n \rightarrow \infty$, we obtain that

$$0 < d(x^*, T_\lambda x^*) \leq \phi(d(x^*, T_\lambda x^*)),$$

which is a contradiction. Thus $x^* = T_\lambda x^*$. So from Lemma 2.15, T has a fixed point. \square

The following example justifies the previous theorem.

Example 3.11. Let $X = [-1, 1]$ and define the distance function $d : X \times X \rightarrow [0, \infty)$ as $d(x, y) = |x - y|$, for any $x, y \in X$. The mapping $W : X \times X \times [0, 1] \rightarrow X$ is defined as $W(x, y; \lambda) = \lambda x + (1 - \lambda)y$, for all $x, y \in X$ and let $T : X \rightarrow X$ be given by

$$Tx = \begin{cases} -x & \text{if } x \in [-1, 0), \\ x & \text{if } x \in [0, 1]. \end{cases} \quad (38)$$

For any $x, y, z \in X$, we have

$$\begin{aligned} d(z, W(x, y, \lambda)) &= |\lambda(z - x) + (1 - \lambda)(z - y)| \\ &\leq \lambda|z - x| + (1 - \lambda)|z - y| \\ &= \lambda d(z, x) + (1 - \lambda)d(z, y) \end{aligned}$$

Hence, (X, d, W) is a convex metric space. Also, for $\lambda = \frac{1}{2}$

$$W(x, Tx; \lambda) = T_\lambda x = \begin{cases} 0 & \text{if } x \in [-1, 0), \\ x & \text{if } x \in [0, 1]. \end{cases} \quad (39)$$

So, we have $d(T_\lambda x, T_\lambda y) = 0$, where $x, y \in [-1, 1] \setminus \text{Fix}(T)$. Therefore, for any $\phi \in \Phi$ and any $\alpha, \beta, \gamma \in (0, 1)$, T is an interpolative enriched Matkowski type contraction. Thus, by Theorem 3.10, T has a fixed point. In this case, $\text{Fix}(T) = [0, 1]$, i.e. T has infinitely many fixed point.

Corollary 3.12. Let (X, d) be a complete metric space and T be an interpolative Matkowski type contraction. Then T has a fixed point in X .

Proof. Taking $\lambda = 0$ in the above theorem, the result holds. \square

4. Conclusion

In this paper, we introduced larger classes of contractive mappings, namely, interpolative enriched Kannan type contraction, interpolative enriched Hardy-Rogers type contraction and interpolative enriched Matkowski type contraction, by enriching usual interpolative Kannan type contraction, interpolative Hardy-Rogers type contraction and interpolative Matkowski type contraction, respectively, in the Takahashi convex metric space setting. For the given contractions, we proved existence of fixed points and approximation results using Krasnoselskij iteration. Our results generalized the results presented in [18], [22], [23] and some other results in the existing literature. Examples are also given for validating the usefulness of our results. The results presented in this paper will enlighten the way for researchers to enrich the results given in [5], [19]-[21] and [24]-[26].

References

- [1] M. Abbas, R. Anjum, S. Riasat, A new type of fixed point theorem via interpolation of operators with application in homotopy theory, Arab. J. Math. (2022) 1-12.
- [2] R. P. Agarwal, D. O'Regan, D. R. Sahu, Fixed point theory for Lipschitzian-type mappings with application, Topological Fixed Point Theory and Its Applications, Springer: New York, USA, (2009) p.6.
- [3] M. U. Ali, H. Aydi, M. Alansari, New generalizations of set valued interpolative Hardy-Rogers type contractions in b-metric spaces, J. Funct. Spaces (2021) Article ID 6641342.
- [4] H. Aydi, C. M. Chen, E. Karapınar, Interpolative Ćirić-Reich-Rus type contractions via the Branciari distance, Mathematics, 7(1) (2019) 84.
- [5] H. Aydi, E. Karapınar, A. F. Roldán López de Hierro, ω -interpolative Ćirić-Reich-Rus-type contractions, Mathematics 7(1) (2019) 57.
- [6] S. Banach, Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales, Fund. Math. 3(1) (1922) 133-181.
- [7] I. Beg, An iteration scheme for asymptotically nonexpansive mappings on uniformly convex metric spaces, Nonlinear Anal. Forum 6(1) (2001) 27-34.
- [8] I. Beg, M. Abbas, Common fixed points and best approximation in convex metric spaces, Soochow J. Math. 33(4) (2007) 729-738.
- [9] I. Beg, M. Abbas, J. K. Kim, Convergence theorems of the iterative schemes in convex metric spaces, Nonlinear Funct. Anal. Appl. 11(3) (2006) 421-436.
- [10] V. Berinde, Approximating fixed points of enriched nonexpansive mappings by Krasnoselskij iteration in Hilbert spaces, Carpathian J. Math. 35(3) (2019) 293-304.
- [11] V. Berinde, M. Păcurar, Approximating fixed points of enriched contractions in Banach spaces, J. Fixed Point Theory Appl. 22 (2020) 38.
- [12] V. Berinde, M. Păcurar, Approximating fixed points of enriched Chatterjea contractions by Krasnoselskij iterative algorithm in Banach spaces, J. Fixed Point Theory Appl. 23 (2021) 66.
- [13] V. Berinde, M. Păcurar, Existence and approximation of fixed points of enriched contractions and enriched ϕ -contractions, Symmetry 13(3) (2021) 498.
- [14] P. Debnath, Z. D. Mitrovic, S. Radenovic, Interpolative Hardy-Rogers and Reich-Rus-Ćirić type contractions in b-metric spaces and rectangular b-metric spaces, Math. Vesnik 72 (2020) 368-374.
- [15] Y. U. Gaba, M. Aphane, H. Aydi, Interpolative Kannan contractions in T_0 -quasi-metric spaces, J. Math. (2021) Article ID 6685638.
- [16] P. Gautam, S. Kumar, S. Verma, G. Gupta, Nonunique fixed point results via Kannan-contraction on quasi-partial-metric space, J. Funct. Spaces (2021) Article ID 2163108.
- [17] R. Kannan, Some results on fixed points, Bull. Calcutta Math. Soc. 60 (1968) 71-76.
- [18] E. Karapınar, Revisiting the Kannan type contractions via interpolation, Adv. Theory Nonlinear Anal. Appl. 2(2) (2018) 85-87.
- [19] E. Karapınar, A survey on interpolative and hybrid contractions, In Mathematical Analysis in Interdisciplinary Research Springer, Cham (2021) 431-475.
- [20] E. Karapınar, Interpolative Kannan-Meir-Keeler type contraction, Adv. Theory Nonlinear Anal. Appl. 5(4) (2021) 611-614.
- [21] E. Karapınar, R. P. Agarwal, H. Aydi, Interpolative Reich-Rus-Ćirić type contractions on partial metric spaces, Mathematics 6(11) (2018) 256.
- [22] E. Karapınar, O. Alqahtani, H. Aydi, On interpolative Hardy-Rogers type contractions, Symmetry 11(1) (2018) 8.
- [23] E. Karapınar, H. Aydi, Z. D. Mitrovic, On interpolative Boyd-Wong and Matkowski type contractions, TWMS J. Pure Appl. Math. 11(2) (2020) 204-212.
- [24] E. Karapınar, A. Fulga, S. S. Yesilkaya, New results on Perov-interpolative contractions of Suzuki type mappings, J. Funct. Spaces (2021) Article ID 9587604.
- [25] E. Karapınar, A. Fulga, S. S. Yeşilkaya, Interpolative Meir-Keeler mappings in modular metric spaces. Mathematics, 10(16) (2022) 2986.
- [26] E. Karapınar, A. Fulga, A. F. Roldán López de Hierro, Fixed point theory in the setting of $(\alpha, \beta, \psi, \phi)$ -interpolative contractions, Adv. Differ. Equ. (2021) 339.

- [27] S. G. Krein, J. I. Petunin, E. M. Semenov, *Interpolation of linear operators*, American Mathematical Society: Providence, RI, USA, 1982.
- [28] J. Matkowski, *Integrable solutions of functional equations: dissertationes mathematicae* (1975).
- [29] J. Matkowski, Fixed point theorems for mappings with a contractive iterate at a point, *Proc. Am. Math. Soc.* 62(2) (1977) 344-348.
- [30] V. N. Mishra, L. M. Sánchez Ruiz, P. Gautam, S. Verma, Interpolative Reich-Rus-Ćirić and Hardy- Rogers contraction on quasi-partial b-metric space and related fixed point results, *Mathematics* 8(9) (2020) 1598.
- [31] B. Mohammadi, V. Parvaneh, H. Aydi, On extended interpolative Ćirić-Reich-Rus type F-contractions and an application, *J. Inequal. Appl.* (2019) 290.
- [32] S. A. Naimpally, K. L. Singh, J. H. M. Whitfield, Fixed and common fixed points for nonexpansive mappings in convex metric spaces, *Math. Sem. Notes Kobe Univ.* 11 (1983) 239–248.
- [33] S. A. Naimpally, K. L. Singh, J. H. M. Whitfield, Fixed points in convex metric spaces, *Math. Jpn.* 29 (1984) 585–597.
- [34] V. Popa, Fixed point theorems in convex metric spaces for mappings satisfying an implicit relation, *Bul. Ştiinţ. Univ. Politeh. Timiş. Ser. Mat. Fiz.* 45 (2000) 1–10.
- [35] S. Rawat, S. Kukreti, R. C. Dimri, Fixed point results for enriched ordered contractions in noncommutative Banach spaces, *J. Anal.* 30 (2022) 1555-1566.
- [36] T. Shimizu, W. Takahashi, Fixed points of multivalued mappings in certain convex metric spaces, *Topol. Methods Nonlinear Anal.* 8(1) (1996) 197–203.
- [37] W. Takahashi, A convexity in metric space and nonexpansive mappings, *I. Kodai Math. Sem. Rep.* 22(2) (1970) 142-149.
- [38] L. A. Talman, Fixed points for condensing multifunctions in metric spaces with convex structure, *Kodai Math. Sem. Rep.* 29(1-2) (1977) 62–70.
- [39] S. S. Yeşilkaya, On interpolative Hardy-Rogers contractive of Suzuki type mappings, *Topol. Algebra Appl.* 9(1) (2021) 13–19.