



## Existence and $L^\infty$ -estimates for non-uniformly elliptic equations with non-polynomial growths

Omar Benslimane<sup>a</sup>, Ahmed Aberqi<sup>b</sup>, Mhamed Elmassoudi<sup>a</sup>

<sup>a</sup>Sidi Mohamed Ben Abdellah University, Faculty of Sciences Dhar Al Mahraz, Department of Mathematics, B.P 1796 Atlas Fez, Morocco

<sup>b</sup>Sidi Mohamed Ben Abdellah University, National School of Applied Sciences Fez, Morocco

**Abstract.** In the current paper, we investigate the existence and regularity of weak solutions to a class of non-uniformly elliptic equations with degenerate coercivity and non-polynomial growth. The model case is given as follows:

$$\operatorname{div}\left(\frac{\exp(1+|Du|)}{(1+|u|)^2}Du\right) + \frac{M(|Du|)}{(1+|u|)^2} \cdot u = f \quad \text{in } \omega.$$

An  $L^\infty$ -estimate of solutions is also obtained for an  $L^1$ -datum  $f$ .

### 1. Introduction

Let  $\omega$  be a bounded open set in  $\mathbb{R}^d$  that satisfies the segment property, ( $d \geq 2$ ). The goal of the current research is to prove the existence and  $L^\infty$ -estimates of weak solutions to the nonlinear and non-degenerate equations with non-polynomial growth equations:

$$\begin{cases} -\operatorname{div}(\Gamma(x, u, Du)) + B(x, u, Du) = f & \text{on } \omega, \\ u = 0 & \text{on } \partial\omega. \end{cases} \quad (1)$$

Here,  $\Gamma : \omega \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  is a Carathéodory function that satisfies the assumptions below: for a.e.  $x \in \omega$  and for all  $s \in \mathbb{R}$ ,  $\xi, \xi^* \in \mathbb{R}^d$ ,  $\xi \neq \xi^*$ , there exist two N-functions  $M$  and  $P$  (See Definition below) such that:

$$\Gamma(x, s, \xi) \cdot \xi \geq g(|s|)M(|\xi|), \quad (2)$$

where  $g : \mathbb{R}^+ \rightarrow \mathbb{R}_*^+$  is a continuous decreasing function with  $g(0) = 1$ , and set the primitive  $G(s) = \int_0^s \frac{1}{g(t)} dt$ .

$$|\Gamma(x, s, \xi)| \leq v(a_0(x) + \overline{M}^{-1}P(k_1|s|)) + \overline{M}^{-1}M(k_2|\xi|), \quad (3)$$

2020 Mathematics Subject Classification. Primary 35J62, 35J70 ; Secondary 35J20.

Keywords. Degenerate coercivity; Weak solution; Existence; Bounded solution; Elliptic Equations; Orlicz spaces;  $\Delta_2$ -condition.

Received: 21 October 2022; Accepted: 19 November 2022

Communicated by Maria Alessandra Ragusa

Email addresses: [omar.benslimane@usmba.ac.ma](mailto:omar.benslimane@usmba.ac.ma) (Omar Benslimane), [aberqi\\_ahmed@yahoo.fr](mailto:aberqi_ahmed@yahoo.fr) (Ahmed Aberqi), [e1massoudi09@gmail.com](mailto:e1massoudi09@gmail.com) (Mhamed Elmassoudi)

where  $\nu > 0, k_1 > 0, k_2 > 0, a_0(\cdot) \in E_{\overline{M}}(\Omega)$ .

$$(\Gamma(x, s, \xi) - \Gamma(x, s, \xi^*))(\xi - \xi^*) > 0, \tag{4}$$

and  $B : \omega \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function such that

$$|B(x, s, \xi)| \leq h(s)M(|\xi|), \tag{5}$$

where  $h : \mathbb{R} \rightarrow \mathbb{R}^+$  is a continuous function,

$$B(x, s, \xi) \cdot s \geq 0, \tag{6}$$

We suppose that  $t \mapsto \frac{h(t)}{g(|t|)}$  belongs to  $L^1(\mathbb{R})$  and defining  $\psi(r) = \int_0^r \frac{h(t)}{g(|t|)} dt$  for all  $r \in \mathbb{R}$ , this implies that,

$$\psi \text{ is bounded.} \tag{7}$$

and

$$f \in L^1(\omega) \text{ and } \int_r^{+\infty} \left(\frac{t}{M(t)}\right)^p dt < \infty, \text{ with } p = \frac{1}{d-1} \text{ and } r > 0. \tag{8}$$

In the case of uniform ellipticity, i.e.,  $g(s) = \text{const}$ , the existence of bounded solutions of equation (1) has been the subject of several papers in functional frameworks of classical Sobolev spaces, as well as in general functional frameworks, see for example [1–3, 6, 13–15, 21], and their references. However, due to assumption 2, the operator degenerates as soon as the solution  $u$  is unbounded. Indeed, for large values of the solution  $u$ , a slow but steady diffusion effect can occur. The function  $\Gamma(x, s, \xi)$  strongly degenerates when  $|s|$  grows to infinity because when  $|s|$  is large,  $g(|s|)$  vanishes. This lack of coercivity prohibits us from using classical approaches.

For the results dealing with the non-coercivity case, we give the following overview of the pioneering work of Boccardo L. et al. in [9, 17], who studied (1) with  $\Gamma(x, u, Du) \geq \frac{g}{(1+|u|)^\theta} Du$ ,  $B = 0$  and  $f \in L^m(\omega)$  with  $m \geq 1$  and  $\theta \in ]0; 1]$ . After that, Croce G. in [29] introduced the term  $B(x, u, \nabla u) = |u|^{p-1}u$ , which has a regulating effect on the solution  $u$ . In the case of weighted Sobolev spaces  $W_0^{1,p}(\omega, \nu)$ , Ammar K. in [10] established the existence of a renormalized solution in the  $L^1$ -frame under the condition  $\Gamma(x, s, \xi)\xi \geq g\nu(x)|\xi|^p$  and  $B$  satisfies the sign condition. Aharouch L. et al. [7] investigated problem (1) in the presence of an obstacle,

where the right-hand side  $f \in L^1(\omega)$  and the lower-order term  $B$  satisfies  $|B(x, s, \xi)| \leq \gamma(s) + g(s) \sum_{i=1}^N \nu_i(x) |\xi_i|^p$ .

Later, in [11], the authors showed the same results using the following conditions:

$$\Gamma(x, s, \xi) \cdot \xi \geq b(s)^{p-1} \sum_{i=1}^N \nu_i(x) |\xi_i|^p, \int_{-\infty}^{+\infty} b(s) ds < +\infty \text{ and } B = 0.$$

For further results, we suggest that the reader consult [3, 4, 8, 9, 13, 16, 25] and the references therein.

This research intends to generalize the previous results (See also [1, 2, 18–20, 26, 27]) in the framework of Orlicz spaces. Moreover, to prove the existence and  $L^\infty$ -estimates of the solutions of (1) by assuming the coercivity condition (2). Therefore, we use rearrangement techniques to surmount this task, approximate problems, and choose suitable test functions.

The paper’s layout is as follows. Section 2 gives some preliminaries and technical lemmas in Orlicz Spaces. In Section 3, we prove the existence of weak solutions. In the appendix, we establish an  $L^\infty$ -estimate of the solution to (1).

## 2. Auxiliary Outcomes and Mathematical Context

This section shows the notation, goes over some basic definitions, and collects the propositions and facts we need to show our main result.

**Definition 2.1.** [2] Let  $M : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be an  $N$ -function, that is,  $M$  is continuous, convex, with  $M(s) > 0$  for  $s > 0$ ,  $\frac{M(s)}{s} \rightarrow 0$  as  $s \rightarrow 0$ , and  $\frac{M(s)}{s} \rightarrow +\infty$  as  $s \rightarrow +\infty$ . The  $N$ -function  $\bar{M}$  conjugate to  $M$  is defined by  $\bar{M}(s) = \sup_{t>0} (st - M(t))$ .

We will extend these  $N$ -functions into even functions on all  $\mathbb{R}$ .

Let  $P$  and  $Q$  be two  $N$ -functions.  $P \ll Q$  means that  $P$  grows essentially less rapidly than  $Q$ , that is, for each  $\varepsilon > 0$ ,  $\lim_{s \rightarrow +\infty} \frac{P(s)}{Q(\varepsilon s)} = 0$ .

**Definition 2.2.** [4] We define the Orlicz class  $K_M(\omega)$  (resp. the Orlicz space  $L_M(\omega)$ ) as the set of (equivalence classes of) real valued measurable functions  $u$  on  $\omega$  such that

$$\int_{\Omega} M(v(x))dx < +\infty \quad (\text{resp.} \quad \int_{\omega} M\left(\frac{v(x)}{\alpha}\right)dx < +\infty \quad \text{for some } \alpha > 0).$$

The set  $L_M(\omega)$  is Banach space under the norm

$$\|v\|_{M,\omega} = \inf\left\{\alpha > 0 : \int_{\Omega} M\left(\frac{v(x)}{\alpha}\right)dx \leq 1\right\},$$

and  $K_M(\omega)$  is a convex subset of  $L_M(\omega)$ .

- The closure in  $L_M(\omega)$  of the set of bounded measurable functions with compact support in  $\bar{\omega}$  is denoted by  $E_M(\omega)$ .
- The dual  $E_M(\omega)$  can be identified with  $L_{\bar{M}}(\omega)$  by means of the pairing  $\int_{\omega} uv dx$  and the dual norm of  $L_{\bar{M}}(\omega)$  is equivalent to  $\|v\|_{\bar{M},\omega}$ .
- The Orlicz-Sobolev space,  $W^1L_M(\omega)$  (resp.  $W^1E_M(\omega)$ ) is the space of all functions  $v$  such that  $v$  and its distributional derivatives up to order 1 lie in  $L_M(\omega)$  (resp.  $E_M(\omega)$ ). It is a Banach space under the norm

$$\|v\|_{1,M} = \sum_{|\alpha| \leq 1} \|D^{\alpha}v\|_{M,\omega}.$$

- Thus,  $W^1L_M(\omega)$  and  $W^1E_M(\omega)$  can be identified with subspaces of product of  $N + 1$  copies of  $L_M(\omega)$ . Denoting this product by  $\Pi L_M$ . We will use the weak topologies  $\sigma(\Pi L_M, \Pi E_{\bar{M}})$  and  $\sigma(\Pi L_M, \Pi L_{\bar{M}})$ .

**Definition 2.3.** [6] We define the space  $W^1_0E_M(\omega)$  as the (norm) closure of the Schwartz space  $\mathcal{D}(\omega)$  in  $W^1E_M(\omega)$  and the space  $W^1_0L_M(\omega)$  as the  $\sigma(\Pi L_M, \Pi E_{\bar{M}})$  closure of  $\mathcal{D}(\omega)$  in  $W^1L_M(\omega)$ .

We denote by  $W^{-1}L_{\bar{M}}(\omega)$  (resp.  $W^{-1}E_{\bar{M}}(\omega)$ ) the space of distributions on  $\omega$  which can be written as sums of derivatives of order  $\leq 1$  of functions in  $L_{\bar{M}}(\omega)$  (resp.  $E_{\bar{M}}(\omega)$ ). It is also a Banach space under the usual quotient norm. For more details, we refer the reader to [6].

### 2.1. Rearrangement

Denote by  $|\omega|$  the Lebesgue measure of  $\omega$ . Assume that  $v$  is a measurable function from  $\omega$  into  $\mathbb{R}$ . The distribution function  $\mu_v$  of  $v$  is defined as follows:

$$\mu_v(t) = |\{x \in \omega; |v(x)| > t\}|, t \geq 0.$$

The decreasing rearrangement  $v_*$  of  $v$  defined on  $]0, |\omega|[$  by

$$v_*(s) = \inf\{t \geq 0; \mu_v(t) \leq s\}$$

$$v_*(0) = \text{ess sup}|v|. \tag{9}$$

Furthermore, for all  $t \geq 0$ , we have

$$v_*(\mu_v(t)) \leq t. \tag{10}$$

Finally, let  $\Theta(t) = te^{\sigma t^2}$ ,  $\sigma > 0$ . It's obvious that when  $\sigma = (\frac{\lambda_1}{\lambda_2})^2$ ,  $\lambda_1 > 0$ ,  $\lambda_2 > 0$ , one has

$$\Theta'(t) - \frac{\lambda_1}{\lambda_2}|\Theta(t)| \geq \frac{1}{2} \text{ for all } t \in \mathbb{R}. \tag{11}$$

### 3. Main results

In what follows, we will assume that  $M$  and  $P$  are two N-functions such that  $H(s) = \frac{M(s)}{s}$  is a convex function.

**Definition 3.1.** A measurable function  $u \in W_0^1L_M(\omega)$  is called a weak solution to problem (1), if  $\Gamma(x, u, Du) \in (L_{\overline{M}}(\omega))^d$  and

$$\int_{\omega} \Gamma(x, u, Du)\nabla\varphi dx + \int_{\omega} B(x, u, \nabla u)\varphi dx = \int_{\Omega} f\varphi dx, \forall \varphi \in \mathcal{D}(\omega). \tag{12}$$

**Theorem 3.2.** Assume that (3)-(5) hold. Given  $f \in L^1(\omega)$  with the condition (8), then there exists a bounded weak solution  $u \in W_0^1L_M(\omega) \cap L^\infty(\omega)$  to problem (1).

#### Step 1: Approximate problems

For every  $n > 0$ , we define the following approximations:

$$\Gamma_n(x, s, \xi) = \Gamma(x, T_n(s), \xi), B_n(x, s, \xi) = \frac{B(x, s, \xi)}{1 + \frac{1}{n}|B(x, s, \xi)|}, \text{ a.e. } x \in \omega, \text{ for all } s \in \mathbb{R} \text{ and } \xi \in \mathbb{R}^d, \text{ where } T_n(s) = \max(-n, \min(n, s)).$$

Denoting by  $(f_n)_n$  the sequence of smooth functions such that  $f_n \rightarrow f$  strongly in  $L^1(\omega)$ , and

$$\|f_n\|_{L^1(\omega)} \leq \|f\|_{L^1(\omega)}. \tag{13}$$

and consider the approximated equations

$$\int_{\omega} \Gamma_n(x, T_n(u_n), Du_n)\nabla\varphi dx + \int_{\omega} B_n(x, u_n, Du_n)\varphi dx = \int_{\omega} f_n\varphi dx, \forall \varphi \in W_0^1L_M(\omega) \tag{14}$$

Now, since  $g(\cdot)$  is decreasing and by (2), we have

$$\Gamma_n(x, T_n(s), \xi) \cdot \xi \geq g(|T_n(s)|)M(|\xi|) \geq g(n)M(|\xi|).$$

We have also  $|B_n(x, s, \xi)| \leq |B(x, s, \xi)|$ ,  $|B_n(x, s, \xi)| \leq n$  and  $B_n(x, s, \xi)s \geq 0$ .

As a consequence of the [23], since  $\Gamma_n(x, s, \xi) + B_n(x, s, \xi)$  verify the assumption  $(A_4)$  of Proposition 5, there exists  $u_n \in W_0^1 L_M(\omega)$  solution of the problem (14).

**Step 2: A priori Estimates**

According to (13) and (43) (see appendix), there exists a constant still denoted  $c_0$  such that

$$\|u_n\|_{L^\infty(\omega)} \leq c_0. \tag{15}$$

Let  $n > c_0$ , then  $T_n(u_n) = u_n$ . Choosing  $\varphi = \Theta(u_n)$  as a test function of (14), by (2) and (5), we have

$$\int_\omega g(|u_n|)M(|\nabla u_n|)\Theta'(u_n)dx \leq \int_\omega h(u_n)M(|\nabla u_n|)|\Theta(u_n)|dx + \int_\omega |f_n|\Theta(u_n)dx,$$

using (13) and Dominated Convergence Theorem, we have

$$\int_\omega (g(|u_n|)\Theta'(u_n) - h(u_n)|\Theta(u_n)|)M(|\nabla u_n|)dx \leq \Theta(c_0)\|f_n\|_{L^1(\omega)},$$

using (11) with  $\sigma = \left(\frac{h(u_n)}{g(|u_n|)}\right)^2$ , we obtain

$$\int_\omega M(|\nabla u_n|)dx \leq \frac{2\Theta(c_0)}{g(c_0)}\|f\|_{L^1(\omega)}. \tag{16}$$

As a result, one has  $\{u_n\}_n$  is bounded in  $W_0^1 L_M(\omega)$ . If required, we go to a subsequence and suppose that

$$u_n \xrightarrow{weakly} u \text{ in } W_0^1 L_M(\omega) \text{ for } \sigma(\Pi L_M, \Pi E_{\overline{M}}), \text{ strongly in } E_M(\omega), \text{ and a.e. in } \omega. \tag{17}$$

and usnig the compact embedding of  $W_0^1 L_M(\omega)$  in  $E_M(\omega)$ , we have also

$$u_n \longrightarrow u \text{ strongly in } E_M(\omega) \text{ and a.e. in } \omega. \tag{18}$$

We will demonstrate that  $\{\Gamma(x, T_n(u_n), Du_n)\}_n$  is bounded in  $(L_{\overline{M}}(\omega))^d$ .

For this, we take  $v \in (E_M(\omega))^d$ , and by (4) we get,

$$(\Gamma(x, T_n(u_n), Du_n) - \Gamma(x, T_n(u_n), \frac{v}{k_2}))(Du_n - \frac{v}{k_2}) > 0,$$

then

$$\int_\omega \Gamma(x, T_n(u_n), Du_n)dx \leq I + J,$$

where

$$I = k_2 \int_\omega \Gamma(x, T_n(u_n), Du_n)\nabla u_n dx,$$

and

$$J = \int_\omega \Gamma(x, T_n(u_n), \frac{v}{k_2})v dx - k_2 \int_\omega \Gamma(x, T_n(u_n), \frac{v}{k_2})Du_n dx.$$

From (2), (14) and using the same previous techniques to establish that

$$\int_\omega \Gamma(x, T_n(u_n), Du_n)Du_n dx \leq 2|\Theta(c_0)|\|f\|_{L^1(\omega)}, \tag{19}$$

and then,

$$I \leq C_I, \tag{20}$$

where  $C_J$  is a positive constant independent of  $n$ .

By (3), the convexity of  $\overline{M}$ , and the fact that  $P \ll M$ , we have

$$\int_{\omega} \overline{M}\left(\frac{A(x, T_n(u_n), \frac{v}{k_2})}{3v}\right) dx \leq \frac{1}{3} \int_{\omega} (\overline{M}(a_0(x)) + M(k_1|T_n(u_n)|) + M(|v|)) dx + C,$$

thus  $\{\Gamma(x, T_n(u_n), \frac{v}{k_2})\}_n$  is bounded in  $(L_{\overline{M}}(\omega))^d$ .

Returning to  $J$ , we have

$$J \leq 2\|\Gamma(x, T_n(u_n), \frac{v}{k_2})\|_{\overline{M}, \omega} \|v\|_{M, \omega} + 2k_2\|\Gamma(x, T_n(u_n), \frac{v}{k_2})\|_{\overline{M}, \omega} \|Du_n\|_{M, \omega},$$

and by (16), we obtain

$$J \leq C_J,$$

where  $C_J$  is a positive constant independent of  $n$ .

So,

$$\int_{\omega} \Gamma(x, T_n(u_n), Du_n) \omega dx \leq C, \tag{21}$$

with  $C$  is a positive constant that is independent of  $n$ .

Finally, according to the Banach-Steinhaus Theorem,  $\{\Gamma(x, T_n(u_n), Du_n)\}_n$  remains bounded in  $(L_{\overline{M}}(\omega))^d$ . Hence

$$\Gamma_n(x, T_n(u_n), Du_n) \xrightarrow{weakly} \xi, \quad \text{in } (L_{\overline{M}}(\omega))^d, \tag{22}$$

for  $\sigma(\Pi L_{\overline{M}}, \Pi E_M)$ .

**Step 3 : Almost everywhere convergence of  $Du_n$**

Let  $v_j \in \mathcal{D}(\omega) \xrightarrow{modular} u$ , in  $W_0^1 L_M(\Omega)$  (cf. [24]). Let  $W_n^j = u_n - v_j$  and  $W^j = u - v_j$ .

Plug the test function  $\Theta(W_n^j)$  in (14), we get,

$$\int_{\omega} \Gamma(x, u_n, Du_n).D\Theta(W_n^j) dx + \int_{\omega} B_n(x, u_n, Du_n)\Theta(W_n^j) dx = \int_{\omega} f_n \Theta(W_n^j) dx. \tag{23}$$

For  $i \geq 1$ , we denote by  $\varepsilon_i(n, j)$  the various sequences of real numbers which satisfy

$$\lim_{j \rightarrow +\infty} \lim_{n \rightarrow +\infty} \varepsilon_i(n, j) = 0.$$

The first term in (23) is written as follows

$$\begin{aligned} \int_{\omega} \Gamma(x, u_n, Du_n).D\Theta(W_n^j) dx &= \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Dv_j \chi_j^s)] \\ &\quad \times [Du_n - Dv_j \chi_j^s] \Theta'(W_n^j) dx \\ &+ \int_{\omega} \Gamma(x, u_n, Dv_j \chi_j^s).[Du_n - Dv_j \chi_j^s] \Theta'(W_n^j) dx \\ &- \int_{\omega \setminus \omega_j^s} \Gamma(x, u_n, Du_n).Dv_j \Theta(W_n^j)' dx, \end{aligned}$$

where  $\chi_j^s$  denotes the characteristic function of the subset  $\omega_j^s = \{x \in \omega : |Dv_j| \leq s\}$ .

Starting with the third term, since  $Dv_j \chi_{\omega \setminus \omega_j^s} \in (E_M(\omega))^d$ , (17) and (22), we have

$$\int_{\omega \setminus \omega_j^s} \Gamma(x, u_n, Du_n).Dv_j \Theta'(W_n^j) dx \rightarrow \int_{\omega \setminus \omega_j^s} \xi.Dv_j \Theta'(W^j) dx \quad \text{as } n \rightarrow \infty,$$

using the modular convergence of  $\{v_j\}$ , we get

$$\int_{\omega \setminus \omega_j^s} \xi \cdot Dv_j \Theta'(W^j) dx \rightarrow \int_{\omega \setminus \omega_j^s} \xi \cdot D u dx \text{ as } j \rightarrow \infty,$$

it will allow us to write

$$\int_{\omega \setminus \omega_j^s} \Gamma(x, u_n, Du_n) \cdot Dv_j \Theta'(W_n^j) dx = \int_{\omega \setminus \omega_j^s} \xi \cdot D u dx + \varepsilon_1(n, j). \tag{24}$$

For the second term of (23), remark that

$$\begin{aligned} & \int_{\omega} \Gamma(x, u_n, \Gamma v_j \chi_j^s) \cdot [Du_n - Dv_j \chi_j^s] \Theta'(W_n^j) dx \\ & \rightarrow \int_{\omega} \Gamma(x, u, Dv_j \chi_j^s) \cdot [Du - Dv_j \chi_j^s] \Theta'(W^j) dx \end{aligned}$$

as  $n \rightarrow +\infty$ , since  $\Gamma(x, u_n, Dv_j \chi_j^s) \Theta'(W_n^j) \rightarrow \Gamma(x, u, Dv_j \chi_j^s) \Theta'(W^j)$  strongly in  $(E_M(\omega))^d$  as  $n \rightarrow \infty$  (because of lemma 1, page 405 in [12], and (18)), while  $Du_n \rightarrow Du$  weakly in  $(L_M(\omega))^d$ .

We have also,  $Dv_j \chi_j^s \rightarrow Du \chi^s$  strongly in  $(E_M(\omega))^d$  as  $j \rightarrow +\infty$ , then it is easy to see that

$$\int_{\omega} \Gamma(x, u, Dv_j \chi_j^s) \cdot [Du - Dv_j \chi_j^s] \Theta'(W^j) dx \rightarrow 0 \text{ as } j \rightarrow \infty,$$

and

$$\int_{\omega} \Gamma(x, u_n, Dv_j \chi_j^s) \cdot [Du_n - Dv_j \chi_j^s] \Theta'(W_n^j) dx = \varepsilon_2(n, j), \tag{25}$$

where  $\omega^s = \{x \in \omega : |Du| \leq s\}$ .

Now combing (23), (24) and (25), we obtain

$$\begin{aligned} \int_{\omega} \Gamma(x, u_n, Du_n) \cdot D\Theta(W_n^j) dx &= \varepsilon_3(n, j) - \int_{\omega \setminus \omega^s} \xi_k \cdot D u dx \\ &+ \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Dv_j \chi_j^s)] \cdot [Du_n - Dv_j \chi_j^s] \Theta'(W_n^j) dx. \end{aligned} \tag{26}$$

Returning to the second term on the left-hand side of (23). We have

$$\begin{aligned} \left| \int_{\omega} B_n(x, u_n, Du_n) \Theta(W_n^j) dx \right| &\leq \int_{\omega} h(u_n) M(|Du_n|) |\Theta(W_n^j)| dx \\ &\leq \int_{\omega} \frac{h(u_n)}{g(|u_n|)} \Gamma(x, u_n, Du_n) \cdot Du_n |\Theta(W_n^j)| dx \\ &\leq \frac{\|h(u_n)\|_{L^\infty}}{g(c_0)} \int_{\omega} \Gamma(x, u_n, Du_n) \cdot Du_n |\Theta(W_n^j)| dx \\ &\leq g_0 \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Dv_j \chi_j^s)] \\ &\quad \times [Du_n - Dv_j \chi_j^s] |\Theta(W_n^j)| dx \\ &+ g_0 \int_{\omega} \Gamma(x, u_n, Du_n) \cdot Dv_j \chi_j^s |\Theta(W_n^j)| dx \\ &+ g_0 \int_{\omega} \Gamma(x, u_n, Dv_j \chi_j^s) \cdot [Du_n - Dv_j \chi_j^s] |\Theta(W_n^j)| dx, \end{aligned} \tag{27}$$

where  $g_0 = \frac{\|h(u_n)\|_{L^\infty}}{g(c_0)}$ .

In a similar way as above, we have

$$g_0 \int_{\omega} \Gamma(x, u_n, Du_n) \cdot Dv_j \chi_j^s |\Theta(W_n^j)| dx = \varepsilon_4(n, j),$$

$$g_0 \int_{\omega} \Gamma(x, u_n, Dv_j \chi_j^s) \cdot [Du_n - Dv_j \chi_j^s] |\Theta(W_n^j)| dx = \varepsilon_5(n, j).$$

Hence

$$\left| \int_{\omega} B_n(x, u_n, Du_n) \Theta(W_n^j) dx \right| \leq g_0 \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Dv_j \chi_j^s)] \times [Du_n - Dv_j \chi_j^s] |\Theta(W_n^j)| dx + \varepsilon_6(n, j). \tag{28}$$

Regarding the term on the right side of (23), since  $\Theta(W_n^j) \xrightarrow{weakly^*} \Theta(W^j)$ , in  $L^\infty(\omega)$  for  $\sigma(L^\infty, L^1)$  as  $n \rightarrow \infty$ , one has

$$\int_{\omega} f_n \Theta(W_n^j) dx \rightarrow \int_{\omega} f \Theta(W^j) dx,$$

we have also  $v_j \xrightarrow{weakly^*} u$ , in  $L^\infty(\omega)$  for  $\sigma(L^\infty, L^1)$  as  $j \rightarrow \infty$ , we get

$$\int_{\omega} f_n \Theta(W_n^j) dx = \varepsilon_7(n, j). \tag{29}$$

Finally, by (23), (26), (28) and (29), we obtain

$$\int_{\omega} [\Theta'(W_n^j) - g_0 |\Theta(W_n^j)|] [a(x, u_n, \nabla u_n) - a(x, u_n, \nabla v_j \chi_j^s)] \cdot [\nabla u_n - \nabla v_j \chi_j^s] dx \leq \int_{\omega \setminus \omega^s} \xi \cdot \nabla u dx + \varepsilon_8(n, j),$$

and then

$$\int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Dv_j \chi_j^s)] \cdot [Du_n - Dv_j \chi_j^s] dx \leq 2 \int_{\omega \setminus \omega^s} \xi \cdot Du dx + 2\varepsilon_8(n, j).$$

On the other hand

$$\begin{aligned} & \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du \chi^s)] [Du_n - Du] \chi^s dx \\ &= \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Dv_j \chi_j^s)] [Du_n - Dv_j \chi_j^s] dx \\ &+ \int_{\omega} \Gamma(x, u_n, Du_n) \cdot [Dv_j \chi_j^s - Du \chi^s] dx \\ &- \int_{\omega} \Gamma(x, u_n, Du \chi^s) \cdot [Du_n - Du \chi^s] dx \\ &+ \int_{\omega} \Gamma(x, u_n, Dv_j \chi_j^s) \cdot [Du_n - Dv_j \chi_j^s] dx. \end{aligned} \tag{30}$$

We will pass to the limit in  $n$  and  $j$  in the last three terms on the right side of the above equality. Tools similar to those in (24) give

$$\int_{\omega} \Gamma(x, u_n, Du_n) \cdot [Dv_j \chi_j^s - Du \chi^s] dx = \varepsilon_9(n, j),$$



$$\int_{\omega} \Gamma(x, u_n, Du\chi^s) \cdot [Du_n - Du\chi^s] dx = \varepsilon_{10}(n, j),$$

and

$$\int_{\omega} \Gamma(x, u_n, Dv_j\chi_j^s) \cdot [Du_n - Dv_j\chi_j^s] dx = \varepsilon_{11}(n, j), \tag{31}$$

which imply that

$$\int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du\chi^s)] \cdot [Du_n - Du\chi^s] dx \leq 2 \int_{\omega \setminus \omega^s} \xi Du dx + 2\varepsilon_{12}(n, j).$$

For  $r \leq s$ , one has

$$\begin{aligned} 0 &\leq \int_{\omega_r} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du)] \cdot [Du_n - Du] dx \\ &= \int_{\omega_s} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du\chi^s)] \cdot [Du_n - Du\chi^s] dx \\ &\leq \int_{\omega} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du\chi^s)] \cdot [Du_n - Du\chi^s] dx \\ &\leq 2 \int_{\omega \setminus \omega_s} \xi \cdot Du dx + \varepsilon_{13}(n, j). \end{aligned}$$

Using the fact that  $\xi \cdot Du \in L^1(\omega)$  and  $|\omega \setminus \omega_s| \rightarrow 0$  as  $s \rightarrow +\infty$ , we get

$$\int_{\omega_r} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du)] \cdot [Du_n - Du] dx \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

As a result, we conclude that there exists a subsequence still denoted by  $u_n$  such that

$$[\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du)] \cdot [Du_n - Du] \rightarrow 0 \quad \text{a.e. in } \omega_r.$$

On the other hand, for every  $x \in \omega^r \setminus Z$  with  $|Z| = 0$ , one has by (3) and (2),

$$\begin{aligned} [\Gamma(x, u_n, Du_n) - \Gamma(x, u_n, Du)] \cdot [Du_n - Du] &\geq g(c_0)M(|Du_n|) \\ &\quad - \Gamma(x, u_n, Du_n) \cdot Du - \Gamma(x, u_n, Du) \cdot Du_n \\ &\geq g(c_0)M(|Du_n|) \\ &\quad - C(1 + |Du_n| + \overline{M}^{-1}M(|Du_n|)), \end{aligned} \tag{32}$$

where  $C$  is a constant not depend on  $n$ .

Following all the previous results,  $\{Du_n\}$  is bounded in  $\mathbb{R}^N$ , and for a subsequence of  $u_n$ , there exists  $\xi \in \mathbb{R}^d$  such that

$$\nabla u_n \rightarrow \xi \quad \text{in } \mathbb{R}^d,$$

and

$$[\Gamma(x, u, \xi) - \Gamma(x, u, \nabla u)] \cdot [\xi - Du] = 0.$$

Thus  $\xi = Du$  and  $Du_n \rightarrow Du$  a.e. in  $\omega^r$ .

Since  $r$  is arbitrary, we construct a subsequence such that

$$Du_n \rightarrow Du \quad \text{a.e. in } \omega. \tag{33}$$

From (22), (17), and (33), it follows that

$$\Gamma(x, T_n(u_n), Du_n) \xrightarrow{\text{weakly}} \Gamma(x, u, Du) \in (L_M(\omega))^d, \quad \text{for } \sigma(\Pi L_{\overline{M}}, \Pi E_M) \tag{34}$$

**Step 4:**  $Du_n \xrightarrow{\text{modular}} Du$

Let  $n > c_0$ , by (4), we obtain

$$\begin{aligned} \int_{\omega} \Gamma(x, u_n, Du_n).Du_n dx &\leq \int_{\omega} \Gamma(x, u_n, Du_n).Dv_j \chi_j^s dx \\ &+ \int_{\omega} \Gamma(x, u_n, Dv_j \chi_j^s).[Du_n - Dv_j \chi_j^s] dx \\ &+ 2 \int_{\omega \setminus \omega^s} \Gamma(x, u, Du).Dudx + 2\varepsilon_8(n, j). \end{aligned} \tag{35}$$

We return to (31) to have

$$\begin{aligned} \int_{\omega} \Gamma(x, u_n, Du_n).Du_n dx &\leq \int_{\omega} \Gamma(x, u_n, Du_n).Dv_j \chi_j^s dx \\ &+ 2 \int_{\omega \setminus \omega^s} \Gamma(x, u, Du).Dudx + \varepsilon_{15}(n, j), \end{aligned} \tag{36}$$

letting  $n \rightarrow \infty$  and  $j \rightarrow \infty$ , we get

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{\omega} \Gamma(x, u_n, Du_n).Du_n dx &\leq \int_{\omega} \Gamma(x, u, Du).Du \chi^s dx \\ &+ 2 \int_{\omega \setminus \omega^s} \Gamma(x, u, Du).Dudx. \end{aligned} \tag{37}$$

Passing to the limit as  $s \rightarrow \infty$ , we get

$$\limsup_{n \rightarrow \infty} \int_{\omega} \Gamma(x, u_n, Du_n).Du_n dx \leq \int_{\omega} \Gamma(x, u, Du).Dudx,$$

and by Fatou’s Lemma, we deduce that

$$\lim_{n \rightarrow \infty} \int_{\omega} \Gamma(x, u_n, Du_n).Du_n dx = \int_{\omega} \Gamma(x, u, Du).Dudx.$$

Using Lemma 4 page 164 in [22], we get

$$\Gamma(x, u_n, Du_n).Du_n \rightarrow \Gamma(x, u, Du).Du \text{ strongly in } L^1(\omega). \tag{38}$$

On the other hand, since  $g(c_0) \leq g(|u_n|)$  and using Young inequality, one has

$$\begin{aligned} M\left(\frac{|Du_n - Du|}{2}\right) &\leq \frac{g(|u_n|)}{2g(c_0)}M(|Du_n|) + \frac{g(|u|)}{2g(c_0)}M(|Du|) \\ &\leq \frac{1}{2g(c_0)}\Gamma(x, u_n, Du_n).Du_n + \frac{1}{2g(c_0)}\Gamma(x, u, Du).Du. \end{aligned} \tag{39}$$

As a result of (33) and Lebesgue Theorem, we reach to our result. Hence, according to (33) and Dominated Convergence Theorem, we deduce our result.

**Step 5: Equi-integrability of  $\{B(x, u_n, Du_n)\}_n$**

We aim to establish that

$$B(x, u_n, Du_n) \rightarrow B(x, u, Du) \text{ strongly in } L^1(\omega). \tag{40}$$

Indeed, by (17) and (33), one gets  $B(x, u_n, Du_n) \rightarrow B(x, u, Du)$  a.e. in  $\omega$ . However, because  $u_n$  is bounded and  $h$  is continuous, choosing  $h_0 = \|h(u_n)\|_{L^\infty(\omega)}$ , and by (5), we have

$$|B(x, u_n, Du_n)| \leq h(|u_n|)M(|Du_n|) \leq h_0M(|Du_n|).$$

Now, let  $E \subset \omega$ , then

$$\int_E |B(x, u_n, Du_n)| dx \leq \frac{h_0}{g(c_0)} \int_E \Gamma(x, u_n, Du_n) \cdot Du_n dx$$

As we have (38), we can apply the equi-integrability of  $\{\Gamma(x, u_n, Du_n)\}_n$  and finish our proof with Vitali’s theorem.

**Remark 3.3.** we can find the same result if we replaced (8) by

$$f \in L^r(\omega) \text{ such that } r = \frac{pd}{p+1} \text{ and } p > \frac{1}{d-1}.$$

**Step 6: Passing to the limit.**

Taking  $v \in \mathcal{D}(\omega)$  as a test function in (14) yields

$$\int_{\omega} \Gamma_n(x, u_n, Du_n) \cdot Dv dx + \int_{\omega} B_n(x, u_n, Du_n) v dx = \int_{\omega} f_n v dx. \tag{41}$$

By (34), (40) and (42) respectively, we get

$$\begin{aligned} \int_{\omega} \Gamma_n(x, u_n, Du_n) \cdot Dv dx &\rightarrow \int_{\omega} \Gamma(x, u, Du) \cdot Dv dx. \\ \int_{\omega} B_n(x, u_n, Du_n) v dx &\rightarrow \int_{\omega} B(x, u, Du) v dx \end{aligned}$$

and

$$\int_{\omega} f_n v dx \rightarrow \int_{\omega} f v dx$$

This completes the proof.

**Appendix**

**Theorem 3.4.** Assume that (3) - (7) hold. Given  $f \in L^1(\omega)$  with the condition (8), then any weak solution  $u$  to problem (1) (in the sense of Definition 3.1) satisfied

$$\|u\|_{L^\infty(\omega)} \leq c_0, \tag{42}$$

where  $c_0$  is a constant depending only on  $d$ .

**Proof of Theorem 3.4**

We define a decreasing and convex function  $K(\cdot)$  as  $K(s) = \frac{1}{H^{-1}(s)}$  where  $H^{-1}(s) = \sup\{r \geq 0, H(r) \leq s\}$ . Using Jensen’s inequality, the definition of  $H$  and the fact that  $g(\cdot)$  is decreasing function such that  $g(0) = 1$ , we have

$$\begin{aligned} K\left(\int_{\{t \leq |u| \leq t+h\}} \frac{g(|u|)M(|\nabla u|)}{\int_{\{t \leq |u| \leq t+h\}} |\nabla u| ds} ds\right) &= K\left(\int_{\{t \leq |u| \leq t+h\}} \frac{g(|u|)H(|\nabla u|)|\nabla u|}{\int_{\{t \leq |u| \leq t+h\}} |\nabla u| ds} ds\right) \\ &\leq \int_{\{t \leq |u| \leq t+h\}} \frac{K(g(|u|)H(|\nabla u|))|\nabla u|}{\int_{\{t \leq |u| \leq t+h\}} |\nabla u| ds} ds \\ &\leq \frac{g(t)(\mu(t) - \mu(t+h))}{\int_{\{t \leq |u| \leq t+h\}} |\nabla u| ds}. \end{aligned}$$

Letting  $h \rightarrow 0$ , we get

$$K\left(-\frac{d}{dt} \int_{\{|u|>t\}} \frac{g(|u|)M(|\nabla u|)}{-\frac{d}{dt} \int_{\{|u|>t\}} |\nabla u| ds} ds\right) \leq \frac{-g(|t|)\mu'(t)}{-\frac{d}{dt} \int_{\{|u|>t\}} |\nabla u| ds}.$$

From Lemma (See [24], Lemma 2, page 72), we have

$$-\frac{d}{dt} \int_{\{|u|>t\}} |\nabla u| dx \geq dC_d^{\frac{1}{d}} \mu(t)^{1-\frac{1}{d}},$$

where  $C_d$  is the measure of the unit ball of  $\mathbb{R}^d$ . By the same arguments in Lemma 3.3 in [5] we have

$$\begin{aligned} \frac{1}{g(|t|)} &\leq \frac{-\mu'(t)}{dC_d^{\frac{1}{d}} \mu(t)^{1-\frac{1}{d}}} H^{-1}\left(\frac{-\frac{d}{dt} \int_{\{|u|>t\}} g(|u|)M(|\nabla u|) ds}{dC_d^{\frac{1}{d}} \mu(t)^{1-\frac{1}{d}}}\right) \\ &\leq \frac{-\mu'(t)}{dC_d^{\frac{1}{d}} \mu(t)^{1-\frac{1}{d}}} H^{-1}\left(\frac{c_1 \int_{\{|u|>t\}} |f| ds}{dC_d^{\frac{1}{d}} \mu(t)^{1-\frac{1}{d}}}\right). \end{aligned}$$

By integrating between 0 and  $r$ , we obtain

$$G(r) \leq \frac{1}{dC_d^{1/d}} \int_0^r \frac{-\mu'(t)}{\mu(t)^{1-\frac{1}{d}}} H^{-1}\left(\frac{c_1 \|f\|_{L^1(\omega)}}{dC_d^{1/d} \mu(t)^{1-\frac{1}{d}}}\right) dt,$$

a change of variables gives

$$G(r) \leq \frac{1}{dC_d^{1/d}} \int_{\mu(r)}^{|\omega|} H^{-1}\left(\frac{c_1 \|f\|_{L^1(\omega)}}{dC_d^{1/d} s^{1-\frac{1}{d}}}\right) \frac{ds}{s^{1-\frac{1}{d}}},$$

as above, taking  $r = u^*(t)$  gives

$$G(u^*(t)) \leq \frac{1}{dC_d^{1/d}} \int_t^{|\omega|} H^{-1}\left(\frac{c_1 \|f\|_{L^1(\omega)}}{dC_d^{1/d} s^{1-\frac{1}{d}}}\right) \frac{ds}{s^{1-\frac{1}{d}}}.$$

Then, we have

$$G(\|u\|_\infty) \leq \frac{1}{dC_d^{1/d}} \int_0^{|\omega|} H^{-1}\left(c_1 \frac{\|f\|_{L^1(\omega)}}{dC_d^{1/d} s^{1-\frac{1}{d}}}\right) \frac{ds}{s^{1-\frac{1}{d}}},$$

a change of variables gives

$$G(\|u\|_\infty) \leq \frac{(c_1 \|f\|_{L^1(\omega)})^p}{d^p C_d^{\frac{p+1}{d}}} \int_{c_0}^{+\infty} p t^{-p-1} H^{-1}(t) dt,$$

where  $c_0 = \frac{c_1 \|f\|_{L^1(\omega)}}{dC_d^{1/d} |\omega|^{1-\frac{1}{d}}}$ . And using integration by parts we get

$$G(\|u\|_{L^\infty(\omega)}) \leq \frac{(c_1 \|f\|_{L^1(\omega)})^p}{d^p C_d^{\frac{p+1}{d}}} \left( \frac{H^{-1}(c_0)}{c_0^p} + \int_{H^{-1}(c_0)}^{+\infty} \left(\frac{r}{M(r)}\right)^p dr \right).$$

Thus

$$\|u\|_{L^\infty(\omega)} \leq G^{-1} \left( \frac{(c_1 \|f\|_{L^1(\omega)})^p}{d^p C_d^{\frac{p+1}{d}}} \left( \frac{H^{-1}(c_0)}{c_0^p} + \int_{H^{-1}(c_0)}^{+\infty} \left( \frac{r}{M(r)} \right)^p dr \right) \right). \quad (43)$$

Then, we get the  $L^\infty$ -estimates of  $u$ .

**Example 3.5.** Taking  $M(t) = t^2 \exp(t)$ , and  $g(u) = \frac{1}{(1+|u|)^2}$ .

$$\Gamma(x, u, Du) = \frac{\exp(1 + |Du|)}{(1 + |u|)^2} Du; \quad B(x, u, Du) = g(u)M(|Du|).$$

## References

- [1] A. Aberqi, J. Bennouna, M. Elmassoudi, Nonlinear Elliptic Equations with Some Measure Data in Musielak-Orlicz Spaces, *Nonlinear Dynamics and Systems Theory*, 19 (2019), 227–242.
- [2] A. Aberqi, J. Bennouna, M. Elmassoudi, Sub-supersolution Method for Nonlinear Elliptic Equation with non-coercivity in divergentiel form in Orlicz Spaces, *AIP Conference Proceedings* 2074, 020004 (2019). <https://doi.org/10.1063/1.5090621>.
- [3] A. Aberqi, J. Bennouna, M. Elmassoudi, M. Hammoumi, Existence and uniqueness of a renormalized solution of parabolic problems in Orlicz spaces, *Monatshefte für Mathematik*, 189 (2019), 195–219.
- [4] A. Aberqi, J. Bennouna, M. Elmassoudi, On some doubly nonlinear system in inhomogeneous Orlicz Spaces, *Electronic Journal of Mathematical Analysis and Applications*, 6 (2018), 156–173.
- [5] A. Aberqi, A. Benkirane and M. Elmassoudi, On some nonlinear degenerate elliptic equations having a lower term in Musielak spaces, *Advances in Operator Theory*, 7 (2022), 1–20.
- [6] R. A. Adams, *Sobolev spaces*, Pres New York, (1975).
- [7] L. Aharouch, Y. Akdim, E. Azroul, Quasilinear degenerate elliptic unilateral problems, *Abstract and Applied Analysis*, 2005 (2005), 11–31.
- [8] B. Aharrouch, J. Bennouna, B. El hamdaoui, Existence of weak and renormalized solutions of degenerated elliptic equation, *Afrika Matematika*, 30 (2019), 755–776.
- [9] A. Alvino, L. Boccardo, V. Ferone, L. Orsina, G. Trombetti, Existence results for nonlinear elliptic equations with degenerate coercivity, *Annali di Matematica Pura ed Applicata*, 182 (2003), 53–79.
- [10] K. Ammar, Renormalized solutions of degenerate elliptic problems, *Journal of Differential Equations*, 234 (2007), 1–25.
- [11] K. Ammar, H. Redwane, Renormalized Solutions for Nonlinear Degenerate Elliptic Problems with  $L^1$  Data, *Revista Matemática Complutense*, 22 (2009), 37–52.
- [12] A. Benkirane, A. Elmahi, A strongly nonlinear elliptic equation having natural growth terms and  $L^1$  data, *Nonlinear Analysis: Theory, Methods & Applications*, 39 (2000), 403–411.
- [13] O. Benslimane, A. Aberqi, J. Bennouna, The existence and uniqueness of an entropy solution to unilateral Orlicz anisotropic equations in an unbounded domain. *Axioms*, 9(3), 109. <https://doi.org/10.3390/axioms9030109>.
- [14] O. Benslimane, A. Aberqi, J. Bennouna, Existence and uniqueness of entropy solution of a nonlinear elliptic equation in anisotropic Sobolev–Orlicz space, *Rendiconti del Circolo Matematico di Palermo Series 2*, 70 (2021), 1579–1608.
- [15] O. Benslimane, A. Aberqi, J. Bennouna, Study of some nonlinear elliptic equation with non-polynomial anisotropic growth, *Advances in Operator Theory*, 7 (2022), 1–29.
- [16] M. F. Betta, A. Ferone, G. Paderni, Estimates for solutions to nonlinear degenerate elliptic equations with lower order terms, *Ricerche di Matematica*, 69 (2020), 367–384.
- [17] L. Boccardo, Some elliptic problems with degenerate coercivity, *Advanced Nonlinear Studies*, 6 (2006), 1–12.
- [18] F. Deringoz, V.S. Guliyev, M.N. Omarova, and M.A Ragusa, Calderón-Zygmund operators and their commutators on generalized weighted Orlicz-Morrey spaces, *Bulletin of Mathematical Sciences*, (2022), doi: 10.1142/S1664360722500047.
- [19] G. Dong, X.C. Fang, Barrier solutions of elliptic differential equations in Musielak-Orlicz-Sobolev spaces, *Journal of Function Spaces*, (2021).
- [20] P. Drabek, F. Nicolosi, Existence of bounded solutions for some degenerated quasilinear elliptic equations, *Potential Theory and Degenerate Partial Differential Operators*, (1995), 475–493.
- [21] A. Elmahi, D. Meskine, Existence of solutions for elliptic equations having natural growth terms in Orlicz spaces, *Abstract and Applied Analysis*, 2004 (2004), 1031–1045.
- [22] A. Elmahi, D. Meskine, Non-linear elliptic problems having natural growth and  $L^1$  data in Orlicz spaces, *Annali di Matematica Pura ed Applicata*, 184,(2005), 161–184.
- [23] J.P. Gossez, V. Mustonen, Variational inequalities in Orlicz-Sobolev spaces, *Nonlinear Analysis: Theory, Methods & Applications*, 11 (1987), 379–492.
- [24] J.P. Gossez, Orlicz- Sobolev spaces and nonlinear elliptic boundary value problems, *Nonlinear Analysis, Nonlinear analysis, function spaces and applications*, BSB BG Teubner Verlagsgesellschaft, (1979), 59–94.

- [25] S. Huang, Q. Tian, J. Wang, J. Mu, Stability for noncoercive elliptic equations, *Electronic Journal of Differential Equations*, 242 (2016), 1–11.
- [26] J. Kačur, Solution of nonlinear degenerate elliptic-parabolic systems in Orlicz-Sobolev spaces, *BSB BG Teubner Verlagsgesellschaft*, (1990), 175–179.
- [27] M. Metwali, Nonlinear quadratic Volterra-Urysohn functional-integral equations in Orlicz spaces, *Filomat*, 35 (2021), 2963–2972.
- [28] G. Talenti, Nonlinear elliptic equations, rearrangements of functions and Orlicz spaces, *Annali di Matematica Pura ed Applicata*, 120 (1979), 159–184.
- [29] W. Zou, F. Li, Existence of solutions for degenerate quasilinear elliptic equations, *Nonlinear Analysis: Theory, Methods & Applications*, 73 (2010), 3069–3082.