



# A parametric unified Apostol-type Bernoulli, Euler, Genocchi, Fubini polynomials and numbers

Burak Kurt<sup>a</sup>

<sup>a</sup>Akdeniz University, Faculty of Educations Department of Mathematics, Antalya, TR-07058, Turkey

**Abstract.** In recent years, mathematicians ([1], [3], [5], [22], [23]) introduced and investigated the Fubini Apostol-type numbers and polynomials. They gave some recurrence relations explicit properties and identities for these polynomials. In [12], author considered unified degenerate Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials and gave some relations and identities for these polynomials.

In this article, we consider a parametric unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials. By using the monomiality principle, we give some relations for the parametric unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials. Furthermore, we give summation formula for these polynomials.

## 1. Introduction, Definitions and Notations

Special polynomials and numbers have significant roles in various branches of mathematics and theoretical physics. The Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials and numbers investigated by Luo ([14]-[16]). Also, he established several elementary properties and gave some explicit relationships and derived various explicit series representations for these polynomials. Srivastava in ([28], [29], [31]) investigated Apostol-Bernoulli, Euler and Genocchi polynomials including explicit representations in terms of a certain generalized Hurwitz-Lerch zeta function. Also, Srivastava et. al. ([23], [30]) considered a parametric type of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials.

Furthermore, Srivastava et. al. in [30] introduced and proved some relations for the three-variable unified Apostol-type  $q$ -polynomials. Masjed-Jamai et. al. ([6], [17], [18]) introduced and investigated a parametric Bernoulli, Euler and Genocchi polynomials. Ozarslan [19] presented a unified study of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials and obtained the explicit relations. Dattali et al. [2] considered monomiality principle for the Appell polynomials. Using monomiality principle, Khan et al. in ([8]-[10]) presented certain result for the 2-variable Apostol-type and related polynomials. Acala [1] introduced a unification of the generalized multiparameter Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials of higher order. The authors [12] considered and investigated unified degenerate Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials.

---

2020 *Mathematics Subject Classification.* Primary 11B68; Secondary 11B83, 05A15, 33B10, 33D45.

*Keywords.* The Apostol-Bernoulli; Apostol-Euler; Apostol-Genocchi; Fubini polynomials; Generalized Stirling numbers of the second kind; Parametric unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials; The Three-variable unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials; Quasi-monomial principle.

Received: 05 May 2022; Accepted: 13 February 2023

Communicated by Hari M. Srivastava

Research supported by the Scientific Research Project Administration of Akdeniz University.

Email address: burakurt@akdeniz.edu.tr (Burak Kurt)

As usual notations,  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{R}$  and  $\mathbb{C}$  for the set of natural integers, integers, real numbers and complex numbers, respectively. Also, we let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ ,  $\mathbb{Z}^- = \{-1, -2, \dots\}$ . We begin by introducing the following definition and notation (see also [13]-[24]). The generalized Apostol-Bernoulli polynomials  $\mathcal{B}_n^{(\alpha)}(x; \lambda)$  of order  $\alpha \in \mathbb{N}_0$ , the generalized Apostol-Euler polynomials  $\mathcal{E}_n^{(\alpha)}(x; \lambda)$  of order  $\alpha \in \mathbb{N}_0$  and the generalized Apostol-Genocchi polynomials  $\mathcal{G}_n^{(\alpha)}(x; \lambda)$  of order  $\alpha \in \mathbb{N}_0$  are defined by the following generating function ([13]-[16]), respectively

$$\sum_{n=0}^{\infty} \mathcal{B}_n^{(\alpha)}(x; \lambda) \frac{t^n}{n!} = \left( \frac{t}{\lambda e^t - 1} \right)^\alpha e^{xt},$$

( $|t| < 2\pi$  when  $\lambda = 1$ ;  $|t| < |\ln \lambda|$  when  $\lambda \neq 1$ ), (1)

$$\sum_{n=0}^{\infty} \mathcal{E}_n^{(\alpha)}(x; \lambda) \frac{t^n}{n!} = \left( \frac{2}{\lambda e^t + 1} \right)^\alpha e^{xt},$$

( $|t| < \pi$  when  $\lambda = 1$ ;  $|t| < |\ln(-\lambda)|$  when  $\lambda \neq 1$ ) (2)

and

$$\sum_{n=0}^{\infty} \mathcal{G}_n^{(\alpha)}(x; \lambda) \frac{t^n}{n!} = \left( \frac{2t}{\lambda e^t + 1} \right)^\alpha e^{xt},$$

( $|t| < \pi$  when  $\lambda = 1$ ;  $|t| < |\ln(-\lambda)|$  when  $\lambda \neq 1$ ). (3)

When  $x = 0$  in (1), (2) and (3), we have the Apostol-Bernoulli numbers  $\mathcal{B}_n^{(\alpha)}(0; \lambda) := \mathcal{B}_n^{(\alpha)}(\lambda)$ , the Apostol-Euler numbers  $\mathcal{E}_n^{(\alpha)}(0; \lambda) := \mathcal{E}_n^{(\alpha)}(\lambda)$  and the Apostol-Genocchi numbers  $\mathcal{G}_n^{(\alpha)}(0; \lambda) := \mathcal{G}_n^{(\alpha)}(\lambda)$ , respectively. Two variable Fubini polynomials of order  $\alpha$  are defined in ([7], [12], [23]) as follows

$$\frac{e^{xt}}{[1 - y(e^t - 1)]^\alpha} = \sum_{n=0}^{\infty} \mathcal{F}_n^{(\alpha)}(x, y) \frac{t^n}{n!}. \tag{4}$$

When  $\alpha = 1$ ,  $\mathcal{F}_n^{(1)}(x, y) = F_n(x, y)$  the two-variable Fubini polynomials. Moreover setting  $x = 0$  and  $y = 1$  in (4), we obtain the Fubini numbers  $\mathcal{F}_n^{(\alpha)}(1) := \mathcal{F}_n^{(\alpha)}$  of order  $\alpha \in \mathbb{N}$ .

Acala [1] defined the generalized multiparameter Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials as follows

$$\sum_{n=0}^{\infty} \mathcal{F}_{n,k}^{(\alpha)}(x, y; a, b, c, \lambda) \frac{t^n}{n!} = \left( \frac{a^{-t} t^k}{1 - y \left( \lambda \left( \frac{b}{a} \right)^t - 1 \right)} \right)^\alpha c^{xt}$$

$\left( \left| t \log \left( \frac{b}{a} \right) + \ln \left( \frac{\lambda y}{y + 1} \right) \right| < 2\pi; \alpha \in \mathbb{C}; a, b, c \in \mathbb{R}^+, x, y \in \mathbb{R}, k \in \mathbb{N}_0, 1^\alpha := 1 \right).$

The Stirling numbers of the second kind were defined by Srivastava et al. [25] as follows

$$\sum_{n=0}^{\infty} S(n, \nu, a, b, \beta) \frac{t^n}{n!} = \frac{(\beta^b e^t - a^b)^\nu}{\nu!} \tag{6}$$

where  $\nu \in \mathbb{N}$ ,  $a, b, \beta \in \mathbb{R}$ ,  $a \neq b$ .

Author [12] defined the higher-order unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials  $\mathcal{F}_n^{(\alpha)}(x, y; a, b, \beta, k)$  of order  $\alpha \in \mathbb{N}$ , by means of the following generating function

$$\sum_{n=0}^{\infty} \mathcal{F}_n^{(\alpha)}(x, y; a, b, \beta, k) \frac{t^n}{n!} = \left( \frac{t^k}{1 - y(\beta^b e^t - a^b)} \right)^\alpha e^{xt} \tag{7}$$

$$\left( \left| t + \ln \left( \frac{y\beta^b}{1 + ya^b} \right) \right| < 2\pi; a, b \in \mathbb{R}^+, x, y \in \mathbb{R}, k, \alpha \in \mathbb{N}_0 \right).$$

For  $x, y \in \mathbb{R}$ , the Taylor-Maclaurin expansions of the two functions  $e^{xt} \cos(yt)$  and  $e^{xt} \sin(yt)$  are given, respectively, by (see [24], [30])

$$e^{xt} \cos(yt) = \sum_{n=0}^{\infty} C_n(x, y) \frac{t^n}{n!} \tag{8}$$

and

$$e^{xt} \sin(yt) = \sum_{n=0}^{\infty} S_n(x, y) \frac{t^n}{n!} \tag{9}$$

where the functions  $C_n(x, y)$  and  $S_n(x, y)$  are defined as

$$C_n(x, y) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n}{2k} x^{n-2k} y^{2k} \tag{10}$$

and

$$S_n(x, y) = \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^k \binom{n}{2k+1} x^{n-2k-1} y^{2k+1}. \tag{11}$$

In [24], Srivastava H. M., Masjed-Jamei M. and Beyki M. R. introduced two parameters the Apostol-Bernoulli polynomials  $\mathcal{B}_n^{(\alpha)}(x; \lambda)$  of order  $\alpha \in \mathbb{N}_0$ , the Apostol-Euler polynomials  $\mathcal{E}_n^{(\alpha)}(x; \lambda)$  of order  $\alpha \in \mathbb{N}_0$  and the Apostol-Genocchi polynomials  $\mathcal{G}_n^{(\alpha)}(x; \lambda)$  of order  $\alpha \in \mathbb{N}_0$  in [23] Srivastava and Kizilates defined the parametric kinds of above functions as

$$\left( \frac{t}{\lambda e^t - 1} \right)^\alpha e^{xt} \cos(yt) = \sum_{n=0}^{\infty} \mathcal{B}_n^{(c,\alpha)}(x, y; \lambda) \frac{t^n}{n!} \tag{12}$$

and

$$\left( \frac{t}{\lambda e^t - 1} \right)^\alpha e^{xt} \sin(yt) = \sum_{n=0}^{\infty} \mathcal{B}_n^{(s,\alpha)}(x, y; \lambda) \frac{t^n}{n!}; \tag{13}$$

$$\left( \frac{2}{\lambda e^t + 1} \right)^\alpha e^{xt} \cos(yt) = \sum_{n=0}^{\infty} \mathcal{E}_n^{(c,\alpha)}(x, y; \lambda) \frac{t^n}{n!} \tag{14}$$

and

$$\left( \frac{2}{\lambda e^t + 1} \right)^\alpha e^{xt} \sin(yt) = \sum_{n=0}^{\infty} \mathcal{E}_n^{(s,\alpha)}(x, y; \lambda) \frac{t^n}{n!}; \tag{15}$$

$$\left( \frac{2t}{\lambda e^t + 1} \right)^\alpha e^{xt} \cos(yt) = \sum_{n=0}^{\infty} \mathcal{G}_n^{(c,\alpha)}(x, y; \lambda) \frac{t^n}{n!} \tag{16}$$

and

$$\left( \frac{2t}{\lambda e^t + 1} \right)^\alpha e^{xt} \sin(yt) = \sum_{n=0}^{\infty} \mathcal{G}_n^{(s,\alpha)}(x, y; \lambda) \frac{t^n}{n!}, \tag{17}$$

respectively.

From the equations (12)-(17), we define two parametric kinds of the unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials as

$$\left(\frac{t^k}{1 - y(\beta^b e^t - a^b)}\right)^\alpha e^{xt} \cos(ut) = \sum_{n=0}^\infty \mathcal{F}_n^{(c,\alpha)}(x, y, u; a, b, \beta, k) \frac{t^n}{n!} \tag{18}$$

and

$$\left(\frac{t^k}{1 - y(\beta^b e^t - a^b)}\right)^\alpha e^{xt} \sin(ut) = \sum_{n=0}^\infty \mathcal{F}_n^{(s,\alpha)}(x, y, u; a, b, \beta, k) \frac{t^n}{n!}. \tag{19}$$

Some general polynomials have been described with the help of monomiality principle Khan et al. ([8], [9]) defined the 2-variable general polynomials (2VGP)  $G_n(x, y)$  are defined by

$$e^{xt} \varphi(y, t) = \sum_{n=0}^\infty G_n(x, y) \frac{t^n}{n!}, G_0(x, y) \neq 1. \tag{20}$$

The 2VGP  $G_n(x, y)$  are quasi-monimal ([8], [9]) with respect to the following multiplicative and derivative operators;

$$\widehat{M} = x + \frac{\varphi'(y, D_x)}{\varphi(y, D_x)}, \left(D_x := \frac{d}{dx}; \varphi'(y, t) = \frac{d}{dt} \varphi(y, t)\right) \tag{21}$$

and

$$\widehat{P}_G = D_x. \tag{22}$$

According to monomiality principle, the polynomials  $G_n(x, y)$  satisfy the following relations

$$\widehat{M}_G \{G_n(x, y)\} = G_{n+1}(x, y), \tag{23}$$

$$\widehat{P}_G \{G_n(x, y)\} = nG_{n-1}(x, y), \tag{24}$$

$$\widehat{M}_G \widehat{P}_G \{G_n(x, y)\} = nG_n(x, y) \tag{25}$$

and

$$\exp(t \widehat{M}_G) \{1\} = \sum_{n=0}^\infty G_n(x, y) \frac{t^n}{n!}, |t| < \infty. \tag{26}$$

### 2. The 3-Variable Unified Apostol-type Bernoulli, Euler, Genocchi and Fubini Polynomials

In this section, we consider the 3-variable unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials (3VUIABEGFP). Using the quasi-monomiality principle, we give some identities and related relations for these polynomials.

We define the 3-variable generalized unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials  $\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k)$  of order  $\alpha \in \mathbb{N}_0$  as

$$\sum_{n=0}^\infty {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} = \left(\frac{t^k}{1 - y(\beta^b e^t - a^b)}\right)^\alpha e^{xt} \varphi(z, t) \tag{27}$$

$$\left(\left|t + \ln\left(\frac{y\beta^b}{1 + ya^b}\right)\right| < 2\pi; a, b \in \mathbb{R}^+, x, y \in \mathbb{R}, k, \alpha \in \mathbb{N}, \beta \in \mathbb{R}\right).$$

For  $\alpha = 1$ , we simply denote  ${}_G\mathcal{F}_n^{(1)}(x, y, z; a, b, \beta, k)$  by  ${}_G\mathcal{F}_n(x, y, z; a, b, \beta, k)$ .

**Remark 2.1.** Setting  $k = \alpha = 1, y = -2, a = b = 1$  and  $\beta = \frac{1}{2}$  in (27), we get

$${}_G\mathcal{F}_n^{(1)}\left(x, -2, z; 1, 1, \frac{\lambda}{2}, 1\right) = \mathcal{B}_n(x, z; \lambda).$$

**Remark 2.2.** Choosing  $k = 0, \alpha = 1, y = -\frac{1}{2}, a = b = 1$  and  $\beta = \lambda$  in (27), we get

$${}_G\mathcal{F}_n^{(1)}\left(x, -\frac{1}{2}, z; 1, 1, \lambda, 0\right) = \mathcal{E}_n(x, z; \lambda).$$

**Remark 2.3.** Letting  $k = 1, \alpha = 1, y = -\frac{1}{2}, a = b = 1$  and  $\beta = \lambda$  in (27), we get

$${}_G\mathcal{F}_n^{(1)}\left(x, -\frac{1}{2}, z; 1, 1, \lambda, 1\right) = \mathcal{G}_n(x, z; \lambda).$$

**Remark 2.4.** Setting  $k = 0, \alpha = 1, a = b = 1$  and  $\beta = \lambda$  in (27), we get

$${}_G\mathcal{F}_n^{(1)}(x, y, z; 1, 1, \lambda, 0) = \mathcal{F}_n(x, y, z; \lambda).$$

**Theorem 2.5.** The 3-variable generalized unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials are quasi-monomial with respect to the following multiplicative and derivative operators

$$\widehat{M}_{G\mathcal{F}_n^{(\alpha)}} = x + \alpha \left( \frac{k}{D_x} - \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right) + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} \tag{28}$$

and

$$\widehat{P}_{G\mathcal{F}_n^{(\alpha)}} = D_x. \tag{29}$$

*Proof.* Differentiating Eq. (27) partially with respect to  $t$  gives

$$\begin{aligned} & \sum_{n=0}^{\infty} {}_G\mathcal{F}_{n+1}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left\{ x + \alpha \left( \frac{k}{t} + \frac{y\beta^b e^t}{1 - y(\beta^b e^t - a^b)} \right) + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} \right\} {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!}. \end{aligned} \tag{30}$$

If  $\varphi(z, t)$  is an invertible series and  $\frac{\varphi'(z,t)}{\varphi(z,t)}$  has Taylor series expansion in powers of  $t$ , we write as

$$D_x(e^{xt}\varphi(z, t)) = t(e^{xt}\varphi(z, t)). \tag{31}$$

From (28) and (29), we can write the equation (30) as

$$\begin{aligned} & \sum_{n=0}^{\infty} {}_G\mathcal{F}_{n+1}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left\{ x + \alpha \left( \frac{k}{D_x} + \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right) + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} \right\} {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!}. \end{aligned}$$

From the last equations, we have result.

By using (31), differentiating Eq. (27) partially with respect to  $x$  gives

$$\begin{aligned} & D_x \left\{ \sum_{n=0}^{\infty} {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \right\} \\ &= \sum_{n=1}^{\infty} {}_G\mathcal{F}_{n-1}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{(n-1)!}. \end{aligned} \tag{32}$$

Using (24) and (32), we get (29).  $\square$

**Theorem 2.6.** The (3VGUABEGFP)  ${}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k)$  satisfy the following differential equation

$$\left(x D_x + \alpha \left(\frac{k}{D_x} - \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)}\right) D_x + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} D_x - n\right) {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) = 0. \tag{33}$$

*Proof.* Using the operators (28) and (29) and (25), we get (33).  $\square$

**Theorem 2.7.** The following relation holds true:

$${}_G\mathcal{F}_n^{(\alpha)}(x + u, y, z; a, b, \beta, k) = \sum_{m=0}^n \binom{n}{m} {}_G\mathcal{F}_{n-m}^{(\alpha)}(x, y, z; a, b, \beta, k) u^m. \tag{34}$$

The proof of this theorem is easy prove from (27).

**Theorem 2.8.** There is the following relation between the (3VGUABEGFP) and the generalized Stirling numbers of the second kinds

$$\begin{aligned} & {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) \\ &= \sum_{r=0}^{n-k\alpha} \binom{n-k\alpha}{r} (n+1-k\alpha)(n+2-k\alpha)\cdots n \sum_{l=0}^{\infty} \binom{l+\alpha-1}{l} \\ & \quad \times l! y^l S(r, l, a, b, \beta) G_{n-k\alpha-r}(x, z), \end{aligned} \tag{35}$$

where  $n \geq k\alpha$ .

*Proof.* From (6), (20) and (27), we write as

$$\begin{aligned} & \sum_{n=0}^{\infty} {}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} = t^{\alpha k} [1 - y(\beta^b e^t - a^b)]^{(-\alpha)} e^{xt} \varphi(z, t) \\ &= t^{\alpha k} \sum_{l=0}^{\infty} \binom{l+\alpha-1}{l} l! y^l \sum_{r=0}^{\infty} S(r, l, a, b, \beta) \frac{t^r}{r!} \sum_{m=0}^{\infty} G_m(x, z) \frac{t^m}{m!} \\ &= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \binom{l+\alpha-1}{l} l! y^l \sum_{r=0}^n \binom{n}{r} S(r, l, a, b, \beta) G_{n-r}(x, z) \frac{t^{n+r}}{n!} \\ &= \sum_{n=k\alpha}^{\infty} \sum_{r=0}^{n-k\alpha} \binom{n-k\alpha}{r} (n+1-k\alpha)(n+2-k\alpha)\cdots n \sum_{l=0}^{\infty} \binom{l+\alpha-1}{l} \\ & \quad \times l! S(r, l, a, b, \beta) G_{n-k\alpha-r}(x, z) \frac{t^n}{n!}. \end{aligned} \tag{36}$$

Comparing the coefficients of the both sides of equation (36), we have (35).  $\square$

**Theorem 2.9.** The (3VGUABEGFP)  ${}_G\mathcal{F}_n^{(\alpha)}(x, y, z; a, b, \beta, k)$  satisfy the following summation formulas

$${}_G\mathcal{F}_{n+m}^{(\alpha)}(v, y, z; a, b, \beta, k) = \sum_{p=0}^n \sum_{q=0}^m \binom{n}{p} \binom{m}{q} (v-x)^{p+q} {}_G\mathcal{F}_{n+m-p-q}^{(\alpha)}(x, y, z; a, b, \beta, k). \tag{37}$$

*Proof.* We first replace  $t$  by  $t + u$  in (27). We then rewrite the generating function (27) as follows

$$\begin{aligned} & \left( \frac{(t + u)^k}{1 - y(\beta^b e^{t+u} - a^b)} \right)^\alpha e^{x(t+u)} \varphi(z, t + u) \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!}. \end{aligned}$$

Upon replacing  $x$  by  $v$  in the above equation, we get

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(v, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!} \\ &= e^{(v-x)(t+u)} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!}. \end{aligned}$$

Which, upon expanding the exponential equation, yields

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(v, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!} \\ &= \sum_{N=0}^{\infty} \frac{[(v-x)(t+u)]^N}{N!} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!}. \end{aligned} \tag{38}$$

Now, by applying the following known series identity ([26],p. 52, Eq. 1.6 (2))

$$\sum_{N=0}^{\infty} f(N) \frac{(x+y)^N}{N!} = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} f(n+m) \frac{x^n}{n!} \frac{y^m}{m!} \tag{39}$$

in the right hand side of (38), we get

$$\begin{aligned} & \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} (v-x)^{p+q} \frac{t^p}{p!} \frac{u^q}{q!} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(x, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!} \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} {}_G\mathcal{F}_{n+m}^{(\alpha)}(v, y, z; a, b, \beta, k) \frac{t^n}{n!} \frac{u^m}{m!}. \end{aligned} \tag{40}$$

Finally, upon first replacing  $n$  by  $n - p$  and  $m$  by  $m - q$  and then applying a known result ([26, p. 100, Lemma 1, Eq. 2.1 (1)]) by using Cauchy product in the left-hand side of the above equation (40) and comparing the coefficients of  $\frac{t^n}{n!}$  and  $\frac{u^m}{m!}$  on both sides of the resulting equation. We complete our demonstration of the assertion (37) Theorem 5.  $\square$

### 3. A Parametric Generalized Unified Apostol-type Bernoulli, Euler, Genocchi and Fubini Polynomials

In this section, by using monomiality principle and operational methods. We introduce and investigate the parametric generalized unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials. We give some identities and relations for these polynomials.

**Theorem 3.1.** *The generating functions for the parametric unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials  ${}_G\mathcal{F}_n^{(c,\alpha)}(x, y, z, u; a, b, \beta, k)$  and  ${}_G\mathcal{F}_n^{(s,\alpha)}(x, y, z, u; a, b, \beta, k)$  are given as follows*

$$\sum_{n=0}^{\infty} {}_G\mathcal{F}_n^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) \frac{t^n}{n!} = \left( \frac{t^k}{1 - y(\beta^b e^t - a^b)} \right)^\alpha e^{xt} \varphi(z, t) \cos(ut) \tag{41}$$

and

$$\sum_{n=0}^{\infty} {}_G\mathcal{F}_n^{(s,\alpha)}(x, y, z, u; a, b, \beta, k) \frac{t^n}{n!} = \left( \frac{t^k}{1 - y(\beta^b e^t - a^b)} \right)^\alpha e^{xt} \varphi(z, t) \sin(ut), \tag{42}$$

$$\left( \left| t + \ln \left( \frac{y\beta^b}{1 + ya^b} \right) \right| < 2\pi, a, b \in \mathbb{R}^+, x, y \in \mathbb{R}, k, \alpha \in \mathbb{N}, \beta \in \mathbb{R} \right)$$

respectively.

*Proof.* In Eq. (18) replacing  $x$  and  $u$  by the multiplicative operator  $\widehat{M}_G$  of the 2VGP  $G_n(x, y)$  and  $u$  respectively, gives

$$\sum_{n=0}^{\infty} \mathcal{F}_n^{(c,\nu)} \left( x + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)}, u \right) \frac{t^n}{n!} = \left( \frac{t^k}{1 - y(\beta^b e^t - a^b)} \right)^\alpha e^{xt} \varphi(z, t) \cos(ut).$$

Using the equation (41), we have

$${}_G\mathcal{F}_n^{(c,\nu)}(x, y, z, u; a, b, \beta, k) = \mathcal{F}_n^{(c,\nu)} \left( x + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)}, u \right). \tag{43}$$

Similarly, we can prove the assertion in Eq. (42).  $\square$

**Theorem 3.2.** The parametric unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials  ${}_G\mathcal{F}_n^{(c,\alpha)}(x, y, z, u; a, b, \beta, k)$  and  ${}_G\mathcal{F}_n^{(s,\alpha)}(x, y, z, u; a, b, \beta, k)$  are quasi-monomial with respect to the following multiplicative and derivate operators:

$$\widehat{M} {}_G\mathcal{F}_n^{(c,\alpha)} = x + \alpha \left[ \frac{k}{D_x} + \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right] + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} - u \tan(uD_x) \tag{44}$$

and

$$\widehat{P} {}_G\mathcal{F}_n^{(c)} = D_x; \tag{45}$$

$$\widehat{M} {}_G\mathcal{F}_n^{(s,\alpha)} = x + \alpha \left[ \frac{k}{D_x} + \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right] + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} + u \cot(uD_x) \tag{46}$$

and

$$\widehat{P} {}_G\mathcal{F}_n^{(s)} = D_x \tag{47}$$

respectively.

*Proof.* Differentiating Eq. (41) partially with respect to  $t$  gives to

$$\begin{aligned} & \sum_{n=0}^{\infty} {}_G\mathcal{F}_{n+1}^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left\{ x + \alpha \left( \frac{k}{t} + \frac{y\beta^b e^t}{1 - y(\beta^b e^t - a^b)} \right) + \frac{\varphi'(z, t)}{\varphi(z, t)} - u \tan(ut) \right\} \\ & \quad \times {}_G\mathcal{F}_n^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) \frac{t^n}{n!}. \end{aligned} \tag{48}$$

Applying the identity

$$D_x(e^{xt} \varphi(z, t)) = t(e^{xt} \varphi(z, t)) \tag{49}$$



in Eq. (48) and comparing the coefficients of  $\frac{t^n}{n!}$  in both sides, we get

$$\begin{aligned} & {}_G\mathcal{F}_{n+1}^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) \\ &= \left\{ x + \alpha \left( \frac{k}{D_x} + \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right) + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} - u \tan(uD_x) \right\} \\ & \times {}_G\mathcal{F}_n^{(c,\alpha)}(x, y, z, u; a, b, \beta, k). \end{aligned}$$

Which in view of the monomiality principle given in Eq. (23) mean the assertion in the equation (44). Differentiating on both sides of the equation (41) with respect to  $x$  gives

$$\begin{aligned} & D_x \left\{ \sum_{n=0}^{\infty} {}_G\mathcal{F}_n^{(c,\nu)}(x, y, z, u; a, b, \beta, k) \frac{t^n}{n!} \right\} \\ &= \sum_{n=1}^{\infty} {}_G\mathcal{F}_{n-1}^{(c,\nu)}(x, y, z, u; a, b, \beta, k) \frac{t^n}{(n-1)!}. \end{aligned} \tag{50}$$

When comparing the coefficients of  $t^n$  on both sides, we obtain the monomiality principle property given in Eq. (24) for the equation (45).

Also, using similar arguments as the proof of Eq. (44) and (45). We can prove equations (46) and (47).  $\square$

**Theorem 3.3.** *The parametric unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials  ${}_G\mathcal{F}_n^{(c,\alpha)}(x, y, z, u; a, b, \beta, k)$  and  ${}_G\mathcal{F}_n^{(s,\alpha)}(x, y, z, u; a, b, \beta, k)$  satisfy the following differential equations*

$$\begin{aligned} & \left\{ xD_x + \alpha \left( \frac{k}{D_x} + \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right) D_x + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} D_x - u \tan(uD_x) D_x - n \right\} \\ & \times {}_G\mathcal{F}_n^{(c,\nu)}(x, y, z, u; a, b, \beta, k) \\ &= 0 \end{aligned} \tag{51}$$

and

$$\begin{aligned} & \left\{ xD_x + \alpha \left( \frac{k}{D_x} + \frac{y\beta^b e^{D_x}}{1 - y(\beta^b e^{D_x} - a^b)} \right) D_x + \frac{\varphi'(z, D_x)}{\varphi(z, D_x)} D_x + u \cot(uD_x) D_x - n \right\} \\ & \times {}_G\mathcal{F}_n^{(s,\nu)}(x, y, z, u; a, b, \beta, k) \\ &= 0 \end{aligned} \tag{52}$$

respectively.

*Proof.* Using operators (44) and (45) and employing the monomiality principle in the equation (25).

We get (51), similarly, we can prove equation (52).  $\square$

**Theorem 3.4.** *The following relations hold true:*

$$\begin{aligned} & {}_G\mathcal{F}_n^{(c,\alpha)}(x + x_1, y, z, u + u_1; a, b, \beta, k) \\ &= \sum_{r=0}^n \binom{n}{r} \left\{ {}_G\mathcal{F}_{n-r}^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) C_r(x_1, u_1) \right. \\ & \left. - {}_G\mathcal{F}_{n-r}^{(s,\alpha)}(x_1, y, z, u_1; a, b, \beta, k) S_r(x, u) \right\} \end{aligned} \tag{53}$$

and

$${}_G\mathcal{F}_n^{(s,\alpha)}(x + x_1, y, z, u + u_1; a, b, \beta, k)$$

$$\begin{aligned}
 &= \sum_{r=0}^n \binom{n}{r} \left\{ {}_G\mathcal{F}_{n-r}^{(s,\alpha)}(x, y, z, u; a, b, \beta, k) C_r(x_1, u_1) \right. \\
 &\quad \left. + {}_G\mathcal{F}_{n-r}^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) S_r(x_1, u_1) \right\}. \tag{54}
 \end{aligned}$$

*Proof.* From (41), (8) and (9), we write as

$$\begin{aligned}
 &\sum_{n=0}^{\infty} {}_G\mathcal{F}_n^{(c,\alpha)}(x + x_1, y, z, u + u_1; a, b, \beta, k) \frac{t^n}{n!} \\
 &= \sum_{m=0}^{\infty} {}_G\mathcal{F}_m^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) \frac{t^m}{m!} \sum_{r=0}^{\infty} C_r(x_1, u_1) \frac{t^r}{r!} \\
 &\quad - \sum_{m=0}^{\infty} {}_G\mathcal{F}_m^{(s,\alpha)}(x_1, y, z, u_1; a, b, \beta, k) \frac{t^m}{m!} \sum_{r=0}^{\infty} S_r(x, u) \frac{t^r}{r!}.
 \end{aligned}$$

By using Cauchy product and comparing the coefficients of  $\frac{t^n}{n!}$ , we have (53).

Similarly, from (42), (8) and (9), we have (54).  $\square$

**Corollary 3.5.** For  $x = x_1$  and  $u = u_1$  in (53) and (54), we have

$$\begin{aligned}
 &{}_G\mathcal{F}_n^{(c,\alpha)}(2x, y, z, 2u; a, b, \beta, k) \\
 &= \sum_{r=0}^n \binom{n}{r} \left\{ {}_G\mathcal{F}_{n-r}^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) C_r(x, u) - {}_G\mathcal{F}_{n-r}^{(s,\alpha)}(x, y, z, u; a, b, \beta, k) S_r(x, u) \right\}
 \end{aligned}$$

and

$$\begin{aligned}
 &{}_G\mathcal{F}_n^{(s,\alpha)}(2x, y, z, 2u; a, b, \beta, k) \\
 &= \sum_{r=0}^n \binom{n}{r} \left\{ {}_G\mathcal{F}_{n-r}^{(s,\alpha)}(x, y, z, u; a, b, \beta, k) C_r(x, u) + {}_G\mathcal{F}_{n-r}^{(c,\alpha)}(x, y, z, u; a, b, \beta, k) S_r(x, u) \right\}.
 \end{aligned}$$

#### 4. Conclusion

We introduced the generalized unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials and gave some relations by using the monomiality principle. We proved summation formula for these polynomials. Also, we gave some recurrence relation for the parametric generalized unified Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials.

#### 5. Acknowledgments

The author would like to thank an anonymous referee for the helpful comments.

#### References

- [1] N. G. Acala, A unification of the generalized multiparameter Apostol-type Bernoulli, Euler, Fubini and Genocchi polynomials of higher order, *European J. of Pure Math.*, 13(3), (2020) 587-607.
- [2] G. Dattali, M. Migliorati and H. M. Srivastava, A class of Bessel summation formulas and associated operational methods, *Fract. Calc. Appl. Anal.*, 7(2), (2004), 169-176.
- [3] R. S. Goma and A. M. Magar, Generalized Fubini Apostol-type polynomials and probabilistic applications. *Int. J. Math. Math. Sci.*, (2022), Article ID 2853920, 1-12.
- [4] Y. He, S. Araci and H. M. Srivastava, Some new formulas for the products of the Apostol-type polynomials, *Adv. Differ. Equ.*, (2016), Article 287, 1-18.

- [5] S. Jin, M. C. Dağlı and F. Qi, Degenerate Fubini-type polynomials and numbers, degenerate Apostol-Bernoulli polynomials and numbers and degenerate Apostol-Euler polynomials and numbers, *Axioms*, 11(2022), Article ID 477, 1-10.
- [6] M. M-Jamai, Z. Moalemi, W. Koepf and H. M. Srivastava, An extension of the Taylor series expansion by using the Bell polynomials, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM*, (2019), 113, 1445-1461.
- [7] D. S. Kim, T. Kim, H.-I. Kwon and J.-W. Park, Two variable higher order Fubini polynomials, *J. Korean Math. Soc.*, (2018), 55(4), 975-986.
- [8] S. Kahn and N. Raza, General-Appell polynomials within the context of monomiality principle, *Int. J. Anal.*, (2013), Article ID 328032, 1-12.
- [9] S. Kahn, G. Yasmin and M. Riyasat, Certain result for the 2 -variable Apostol-type and related polynomials, *Comput. Math. Appl.*, (2015), 69, 1367-1382.
- [10] S. Khan, T. Nahid and M. Riyasat, On degenerate Apostol-type polynomials and applications, *Boletín de la Sociedad Matemática Mexicana*, (2019), 25, 509–528.
- [11] I. Kucukoglu, Y. Simsek and H. M. Srivastava, A new family of Lerch-type zeta functions interplating a certain class of higher order Apostol-type numbers and Apostol-type polynomials, *Questiones Math.*, (2019), 42(4), 465-478.
- [12] B. Kurt, Unified degenerate Apostol-type Bernoulli, Euler, Genocchi and Fubini polynomials, *J. Math. Comput. Sci.*, (2022), 25, 259-268.
- [13] D.-Q. Lu and H. M. Srivastava, Some series identities involving the generalized Apostol-type and related polynomials, *Comput. Math. Appl.*, (2011), 62, 3591-3602.
- [14] Q.-M. Luo, Apostol-Euler polynomials of higher order and Gaussian hypergeometric functions, *Taiwan. J. Math.*, (2006), 10, 917-925.
- [15] Q.-M. Luo and H. M. Srivastava, Some generalization of the Apostol-Bernoulli and Apostol-Euler polynomials, *J. Math. Anal. Appl.*, (2005), 308, 290-302.
- [16] Q.-M. Luo and H. M. Srivastava, Some generalizations of the Apostol-Genocchi polynomials and the Stirling numbers of the second kind, *Appl. Math. Comput.*, 217(2011), 5702-5728.
- [17] M. Masjed-Jamei, M. R. Beyki and W. Koepf, A new type of Euler polynomials and numbers, *Mediterr. J. Math.*, (2018), 15, Article ID 138, 1-17.
- [18] M. Masjed-Jamei and W. Koepf, Symbolic computation of some power-trigonometric series, *J. Symb. Comput.*, (2017), 80, 273-284.
- [19] M. A. Ozarslan, Unified Apostol-Bernoulli, Euler and Genocchi polynomials, *Comput. Math. Appl.*, (2011), 62, 2452-2462.
- [20] S. K. Sharma, W. A. Khan and C. S. Ryoo, A parametric kind of the degenerate Fubini numbers and polynomials, *Mathematics*, 8(2020), Article ID 405, 1-13.
- [21] H. M. Srivastava and J. Choi, *Zeta and  $q$ -zeta functions and associated series and integrals*, Elsevier, Amsterdam, 2012.
- [22] H. M. Srivastava, R. Srivastava, A. Muhyi, G. Yasmin, H. Islahi and S. Araci, Construction of a ne family of Fubini-type polynomials and its applications, *Adv. Differ. Equ.*, (2021), Article number 36(2021), 1-25.
- [23] H. M. Srivastava and C. Kizilates, A parametric kind of the Fubini-type polynomials, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM*, (2019), 113, 3253-3267.
- [24] H. M. Srivastava, Masjed-Jamei M. and M. R. Beyki, A parametric type of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials, *Appl. Math. Inform. Sci.*, (2018), 12, 907-917.
- [25] H. M. Srivastava, B. Kurt and V. Kurt, Identities and relation involving the modified degenerate Hermite-based Apostol-Bernoulli and Apostol-Euler polynomials, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM*, (2019), 113(2), 1299-1313.
- [26] H. M. Srivastava and H. L. Manocha, *A treatise an generating functions*, Halsted Press (Ellis Horwood Limited, Chichester), Willes, New york 1984.
- [27] H. M. Srivastava, M. Masjed-Jamei and M. R. Beyki, Some new generalizations and applications of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials, *Rocky Mountain J. Math.*, (2019), 49, 681-697.
- [28] H. M. Srivastava, Some generalizations and basic (or  $q$ -) extensions of the Bernoulli, Euler and Genocchi polynomials, *Appl. Math. Inform. Sci.*, (2011), 5(5), 390-444.
- [29] H. M. Srivastava, Some families of generating functions associated with orthogonal polynomials and other higher Transcendental functions, *Mathematics*, 10(2022), Article ID 3730, 1-23.
- [30] H. M. Srivastava, S. Khan, S. Araci, M. Acikgoz and M. Riyasat, A general class of the three-variable unified Apostol-type  $q$ -polynomials and multiple power  $q$ -sums, *Bull. Iran. Math. Soc.*, 46 (2020), 519–542.
- [31] H. M. Srivastava, A survey of some recent developments of higher transcendental functions of analytic number theory and applied mathematics, *Symmetry*, 13(2021), Article ID 2294, 1-22.
- [32] D.-D. Su and Y. He, Some identities for the two variable Fubini polynomials, *Mathematics*, 7(2019), Article ID 115 1-11, .
- [33] W. Wang, C. Jia and T. Wang, Some results on the Apostol-Bernoulli and Apostol-Euler polynomials, *Comput. Math. Appl.*, (2008), 55, 1322-1332.