



Canonical F -planar mappings of spaces with affine connection onto 3-symmetric spaces

Patrik Peška^a, Vladimir Berezovski^b, Yevhen Cherevko^c, Miloš Z. Petrović^d

^aFaculty of Science, Palacký University in Olomouc, 17. listopada 12, 77146 Olomouc, Czech Republic

^bUman National University of Horticulture, Institutskaya 1, 20300, Uman, Ukraine

^cOdesa National University of Technology, Kanatnaya Str. 112, 65039, Odesa, Ukraine

^dUniversity of Niš, Faculty of Agriculture in Kruševac, Kosančićeva 4, 37000 Kruševac, Serbia

Abstract. We consider canonical F -planar mappings of spaces with affine connection onto 3-symmetric spaces. The main equations for the mappings have been obtained as a closed system of PDEs of Cauchy type in covariant derivatives. We have found the maximum numbers of essential parameters which the general solution of the system depends on.

1. Introduction.

The paper is devoted to further study of the theory of almost geodesic mappings of affinely connected spaces. The theory goes back to the paper by T. Levi-Civita [20], in which the problem on the search for Riemannian spaces with common geodesics was stated and solved in a special coordinate system. We note a remarkable fact that this problem is related to the study of equations of dynamics of mechanical systems.

The theory of geodesic mappings has been developed by T. Thomas, H. Weyl, P.A. Shirokov, A.S. Solodovnikov, N.S. Sinyukov, A.V. Aminova, J. Mikeš, and others, see [23, 31].

Issues arisen by the exploration were studied by V.F. Kagan, G. Vrančeanu, Ya.L. Shapiro, D.V. Vedenyapin. The authors discover special classes of $(n - 2)$ -spaces. In [24] A.Z. Petrov introduced the notion of quasi-geodesic mappings. In particular, holomorphically projective mappings of Kählerian spaces are special quasi-geodesic mappings; they were examined by T. Otsuki, Y. Tashiro, M. Prvanović, J. Mikeš, and others, see [23, 31].

A natural generalization of these classes of mappings is the class of almost geodesic mappings introduced by Sinyukov (see [30–32]). He also specified three types of almost geodesic mappings π_1, π_2, π_3 . The theory of almost geodesic mappings was developed by V.S. Sobchuk [33, 34], N.Y. Yablonskaya [40], V.E. Berezovski, J. Mikeš [2–15, 39], Lj.S. Velimirović, N. Vesić, M.S. Stanković, [25, 27, 28, 35–38] et al.

2020 *Mathematics Subject Classification.* Primary 53B05; Secondary 53B35.

Keywords. F -planar mapping; Space with affine connection; Symmetric space; 2-symmetric space; 3-symmetric space.

Received: 05 January 2023; Accepted: 23 February 2023

Communicated by Mića Stanković

The research of the first author was supported by the project IGA PrF 2023010 (Palacký University Olomouc) and the research of the fourth author was financed by the Ministry of Science, Technological Development and Innovation, Republic of Serbia (Contract reg. no. 451-03-47/2023-01/200383).

Email addresses: Patrik.Peska@upl.cz (Patrik Peška), berez.volod@gmail.com (Vladimir Berezovski), cherevko@usa.com (Yevhen Cherevko), petrovic.milos@ni.ac.rs (Miloš Z. Petrović)

F -planar mappings were introduced by N.S. Sinyukov and J. Mikeš (see [21]) as a further generalization of almost geodesic and holomorphically projective ones. In the same manner, F -planar mappings were studied between manifolds endowed with non-symmetric linear connection [26].

Definition and theorems of the F -planar mappings theory which are given below are presented in monographs [22, 23] and in an expository article [19].

Our study is restricted to the local theory, and we assume that all geometric objects we consider not only continuous but also sufficiently smooth.

2. Basic definitions of the F -planar mappings theory of spaces with affine connection.

Let us consider an n -dimensional space A_n with an affine torsion-free connection ∇ and affinator structure $F (\neq Id)$. The space is referred to a local coordinate system $x = (x^1, x^2, \dots, x^n)$, $\Gamma_{ij}^h(x)$ and $F_i^h(x) (\neq a \cdot \delta_i^h)$ are components of ∇ and F , where δ_i^h is the Kronecker delta and a is some function.

A curve ℓ given by the equation $x^h = x^h(t)$ is called F -planar if its tangent vector $\lambda^h(t) \equiv dx^h(t)/dt$ subjected by the parallel translation along the curve belongs to the distribution generated by the vector fields λ and $F\lambda$.

According to this definition, a curve ℓ is F -planar if and only if the following condition holds

$$\lambda^h_{,\alpha} \lambda^\alpha = \rho_1(x) \lambda^h + \rho_2(x) \lambda^\alpha F_{\alpha}^h,$$

where $\rho_1(t)$ and $\rho_2(t)$ are some functions of the parameter t , we denote by comma “,” the covariant derivative with respect to the connection ∇ of the space A_n .

The set of F -planar curves in a space A_n is a very wide one. It includes geodesics, planar curves, quasi-geodesic curves, and others.

Suppose two affinely connected spaces A_n and \bar{A}_n are given and there are defined structural affinars F and \bar{F} respectively. J. Mikeš and N.S. Sinyukov [21] introduced the following terms:

A mapping $\pi : A_n \rightarrow \bar{A}_n$ is an F -planar mapping if any F -planar curve of A_n is mapped under f onto an \bar{F} -planar curve in \bar{A}_n .

Let us consider the spaces A_n and \bar{A}_n which are referred to a common coordinate system x^1, x^2, \dots, x^n with respect to the F -planar mapping. The tensor

$$P_{ij}^h(x) = \bar{\Gamma}_{ij}^h(x) - \Gamma_{ij}^h(x), \tag{1}$$

is called a *deformation tensor* [1, 31]. Here $\Gamma_{ij}^h(x)$ and $\bar{\Gamma}_{ij}^h(x)$ are components of the affine connection in A_n and \bar{A}_n .

From [19] it follows that a mapping $\pi : A_n \rightarrow \bar{A}_n$ ($n > 2$) is F -planar if and only if the deformation tensor $P_{ij}^h(x)$ of the mapping π in a common coordinate system x^1, x^2, \dots, x^n satisfies the condition

$$P_{ij}^h = \delta_{(i}^h \psi_{j)} + F_{(i}^h \varphi_{j)},$$

where $\psi_i(x)$ and $\varphi_i(x)$ are some covectors, we denote by the round parentheses an operation called *symmetrization* without division with respect to the indices i and j .

An F -planar mapping π for which $\psi_i \equiv 0$ is called *canonical*. It is known that any F -planar mapping can be written as the composition of a canonical F -planar mapping and a geodesic mapping. The latter may be referred to as a trivial F -planar mapping.

Hence, canonical F -planar mappings in a common coordinate system x^1, x^2, \dots, x^n with respect to the mapping are determined by the equations

$$P_{ij}^h = F_{(i}^h \varphi_{j)}. \tag{2}$$

Suppose that the affinator F_i^h defined in the space A_n satisfies the condition

$$F_{\alpha}^h F_i^{\alpha} = e \delta_i^h, \tag{3}$$

where $e = \mp 1$. Such F -planar mappings we denote by $\pi(e)$, where $e = \mp 1$.

3. Canonical F -planar mappings $\pi(e)$ ($e = \mp 1$) of spaces with affine connections onto 3-symmetric spaces.

A space \bar{A}_n with an affine connection is called (locally) symmetric if its Riemann tensor field is parallel. Symmetric spaces were introduced by P.A. Shirokov [29] and É. Cartan [17], see also S. Helgason [18]. Precisely, symmetric spaces satisfy the conditions $\bar{R}_{ijk;m}^h = 0$, where \bar{R}_{ijk}^h is the Riemann tensor of the space \bar{A}_n , the symbol “;” denotes a covariant derivative with respect to the connection $\bar{\nabla}$ of the space \bar{A}_n .

A space \bar{A}_n with an affine connection is called 2-symmetric and 3-symmetric if its Riemann tensor \bar{R}_{ijk}^h satisfies the conditions $\bar{R}_{ijk;m_1m_2}^h = 0$ and $\bar{R}_{ijk;m_1m_2m_3}^h = 0$.

Canonical F -planar mappings $\pi(e)$ ($e = \mp 1$) of spaces with affine connection onto 2-symmetric spaces were studied in [16].

Obviously, symmetric spaces are 2-symmetric spaces, and 2-symmetric spaces are 3-symmetric ones.

Let us consider canonical F -planar mappings $\pi(e)$ ($e = \mp 1$) of spaces with affine connection onto 3-symmetric spaces \bar{A}_n . The mapping are characterized by the equations (2). Assume that the spaces A_n and \bar{A}_n are referred to a common coordinate system x^1, x^2, \dots, x^n and the affinor F_i^h defined in the space A_n satisfies the condition (3).

Since $\bar{R}_{ijk;m}^h = \frac{\partial \bar{R}_{ijk}^h}{\partial x^m} + \bar{\Gamma}_{m\alpha}^h \bar{R}_{ijk}^\alpha - \bar{\Gamma}_{mi}^\alpha \bar{R}_{\alpha jk}^h - \bar{\Gamma}_{mj}^\alpha \bar{R}_{iak}^h - \bar{\Gamma}_{mk}^\alpha \bar{R}_{ij\alpha}^h$, then taking account of (1) we can obtain

$$\bar{R}_{ijk;m}^h = \bar{R}_{ijk;m}^h + P_{m\alpha}^h \bar{R}_{ijk}^\alpha - P_{mi}^\alpha \bar{R}_{\alpha jk}^h - P_{mj}^\alpha \bar{R}_{iak}^h - P_{mk}^\alpha \bar{R}_{ij\alpha}^h \tag{4}$$

According to the definition of covariant derivative

$$\left(\bar{R}_{ijk;m}^h\right)_{,\rho_1} = \frac{\partial \bar{R}_{ijk;m}^h}{\partial x^{\rho_1}} + \Gamma_{\alpha\rho_1}^h \bar{R}_{ijk;m}^\alpha - \Gamma_{i\rho_1}^\alpha \bar{R}_{\alpha jk;m}^h - \Gamma_{j\rho_1}^\alpha \bar{R}_{iak;m}^h - \Gamma_{k\rho_1}^\alpha \bar{R}_{ij\alpha;m}^h - \Gamma_{m\rho_1}^\alpha \bar{R}_{ijk;\alpha}^h$$

and taking account of (1), we have

$$\left(\bar{R}_{ijk;m}^h\right)_{,\rho_1} = \bar{R}_{ijk;m\rho_1}^h - P_{\alpha\rho_1}^h \bar{R}_{ijk;m}^\alpha + P_{i\rho_1}^\alpha \bar{R}_{\alpha jk;m}^h + P_{j\rho_1}^\alpha \bar{R}_{iak;m}^h + P_{k\rho_1}^\alpha \bar{R}_{ij\alpha;m}^h + P_{m\rho_1}^\alpha \bar{R}_{ijk;\alpha}^h \tag{5}$$

Differentiating (4) with respect to x^{ρ_1} in the space A_n , we get

$$\left(\bar{R}_{ijk;m}^h\right)_{,\rho_1} = \bar{R}_{ijk;m\rho_1}^h + P_{m\alpha,\rho_1}^h \bar{R}_{ijk}^\alpha + P_{m\alpha}^h \bar{R}_{ijk,\rho_1}^\alpha - P_{mi,\rho_1}^\alpha \bar{R}_{\alpha jk}^h - P_{mi}^\alpha \bar{R}_{\alpha jk,\rho_1}^h - P_{mj,\rho_1}^\alpha \bar{R}_{iak}^h - P_{mj}^\alpha \bar{R}_{iak,\rho_1}^h - P_{mk,\rho_1}^\alpha \bar{R}_{ij\alpha}^h - P_{mk}^\alpha \bar{R}_{ij\alpha,\rho_1}^h \tag{6}$$

Substituting in (5) from (6), we have

$$\bar{R}_{ijk;m\rho_1}^h = \bar{R}_{ijk;m\rho_1}^h - P_{\alpha\rho_1}^h \bar{R}_{ijk;m}^\alpha + P_{i\rho_1}^\alpha \bar{R}_{\alpha jk;m}^h + P_{j\rho_1}^\alpha \bar{R}_{iak;m}^h + P_{k\rho_1}^\alpha \bar{R}_{ij\alpha;m}^h + P_{m\rho_1}^\alpha \bar{R}_{ijk;\alpha}^h - P_{m\alpha,\rho_1}^h \bar{R}_{ijk}^\alpha - P_{m\alpha}^h \bar{R}_{ijk,\rho_1}^\alpha + P_{mi,\rho_1}^\alpha \bar{R}_{\alpha jk}^h + P_{mi}^\alpha \bar{R}_{\alpha jk,\rho_1}^h + P_{mj,\rho_1}^\alpha \bar{R}_{iak}^h + P_{mj}^\alpha \bar{R}_{iak,\rho_1}^h + P_{mk,\rho_1}^\alpha \bar{R}_{ij\alpha}^h + P_{mk}^\alpha \bar{R}_{ij\alpha,\rho_1}^h \tag{7}$$

Taking account of (2) and (4) the equations (7) can be written as

$$\bar{R}_{ijk;m\rho_1}^h = \bar{R}_{ijk;m\rho_1}^h + \Theta_{ijkm\rho_1}^h \tag{8}$$

where

$$\Theta_{ijkm\rho_1}^h = -F_{(\alpha\varphi\rho_1)}^h \left(\bar{R}_{ijk;m}^\alpha + \Theta_{ijkm}^\alpha\right) + F_{(i\varphi\rho_1)}^\alpha \left(\bar{R}_{\alpha jk;m}^h + \Theta_{\alpha jkm}^h\right) + F_{(j\varphi\rho_1)}^\alpha \left(\bar{R}_{iak;m}^h + \Theta_{iakm}^h\right) + F_{(k\varphi\rho_1)}^\alpha \left(\bar{R}_{ij\alpha;m}^h + \Theta_{ij\alpha m}^h\right) + F_{(m\varphi\rho_1)}^\alpha \left(\bar{R}_{ijk;\alpha}^h + \Theta_{ijk\alpha}^h\right) - F_{(m\varphi\alpha),\rho_1}^h \bar{R}_{ijk}^\alpha - F_{(m\varphi\alpha)}^h \bar{R}_{ijk,\rho_1}^\alpha + F_{(m\varphi i),\rho_1}^\alpha \bar{R}_{\alpha jk}^h + F_{(m\varphi i)}^\alpha \bar{R}_{\alpha jk,\rho_1}^h + F_{(m\varphi j),\rho_1}^\alpha \bar{R}_{iak}^h + F_{(m\varphi j)}^\alpha \bar{R}_{iak,\rho_1}^h + F_{(m\varphi k),\rho_1}^\alpha \bar{R}_{ij\alpha}^h + F_{(m\varphi k)}^\alpha \bar{R}_{ij\alpha,\rho_1}^h \tag{9}$$

$$\Theta_{ijkm}^h = F_{(m\varphi\alpha)}^h \bar{R}_{ijk}^\alpha - F_{(m\varphi i)}^\alpha \bar{R}_{\alpha jk}^h - F_{(m\varphi j)}^\alpha \bar{R}_{iak}^h - F_{(m\varphi k)}^\alpha \bar{R}_{ij\alpha}^h \tag{10}$$

Taking account of structure of the tensor Θ_{ijkm}^h defined by the formulas (10), one could see that the tensor $\Theta_{ijkm\rho_1}^h$ defined by the formulas (9), depends on the tensors $F_k^h, \bar{R}_{ijk}^h, \varphi_k$ and their covariant derivatives with respect to the connection of the space A_n . Also, we assume that the tensor F_k^h called affinator is given, and the tensor satisfies the conditions (3).

According to the definition of covariant derivative

$$\left(\bar{R}_{ijk;m\rho_1}^h\right)_{,\rho_2} = \frac{\partial \bar{R}_{ijk;m\rho_1}^h}{\partial x^{\rho_2}} + \Gamma_{\alpha\rho_2}^h \bar{R}_{ijk;m\rho_1}^\alpha - \Gamma_{i\rho_2}^\alpha \bar{R}_{\alpha jk;m\rho_1}^h - \Gamma_{j\rho_2}^\alpha \bar{R}_{i\alpha k;m\rho_1}^h - \Gamma_{k\rho_2}^\alpha \bar{R}_{ij\alpha;m\rho_1}^h - \Gamma_{m\rho_2}^\alpha \bar{R}_{ijk;\alpha\rho_1}^h - \Gamma_{\rho_1\rho_2}^\alpha \bar{R}_{ijk;m\alpha}^h$$

and taking account of (1), we get

$$\left(\bar{R}_{ijk;m\rho_1}^h\right)_{,\rho_2} = \bar{R}_{ijk;m\rho_1\rho_2}^h - P_{\alpha\rho_2}^h \bar{R}_{ijk;m\rho_1}^\alpha + P_{i\rho_2}^\alpha \bar{R}_{\alpha jk;m\rho_1}^h + P_{j\rho_2}^\alpha \bar{R}_{i\alpha k;m\rho_1}^h + P_{k\rho_2}^\alpha \bar{R}_{ij\alpha;m\rho_1}^h + P_{m\rho_2}^\alpha \bar{R}_{ijk;\alpha\rho_1}^h + P_{\rho_1\rho_2}^\alpha \bar{R}_{ijk;m\alpha}^h. \tag{11}$$

Because of (2) and (8) it follows from (11) that

$$\begin{aligned} \left(\bar{R}_{ijk;m\rho_1}^h\right)_{,\rho_2} &= \bar{R}_{ijk;m\rho_1\rho_2}^h - F_{(\alpha}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;m\rho_1}^\alpha - \Theta_{ijkm\rho_1}^\alpha\right) + F_{(i}^h \varphi_{\rho_2)} \left(\bar{R}_{\alpha jk;m\rho_1}^h - \Theta_{\alpha jkm\rho_1}^h\right) \\ &+ F_{(j}^h \varphi_{\rho_2)} \left(\bar{R}_{i\alpha k;m\rho_1}^h - \Theta_{i\alpha km\rho_1}^h\right) + F_{(k}^h \varphi_{\rho_2)} \left(\bar{R}_{ij\alpha;m\rho_1}^h - \Theta_{ij\alpha m\rho_1}^h\right) + F_{(m}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;\alpha\rho_1}^h - \Theta_{ijk\alpha\rho_1}^h\right) + F_{(\rho_1}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;m\alpha}^h - \Theta_{ijkm\alpha}^h\right). \end{aligned} \tag{12}$$

Let us differentiate the equations (8) covariantly with respect to x^{ρ_2} in the space A_n . Taking account of (12), we get

$$\begin{aligned} \bar{R}_{ijk;m\rho_1\rho_2}^h &= \bar{R}_{ijk;m\rho_1\rho_2}^h - F_{(\alpha}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;m\rho_1}^\alpha - \Theta_{ijkm\rho_1}^\alpha\right) + F_{(i}^h \varphi_{\rho_2)} \left(\bar{R}_{\alpha jk;m\rho_1}^h - \Theta_{\alpha jkm\rho_1}^h\right) \\ &+ F_{(j}^h \varphi_{\rho_2)} \left(\bar{R}_{i\alpha k;m\rho_1}^h - \Theta_{i\alpha km\rho_1}^h\right) + F_{(k}^h \varphi_{\rho_2)} \left(\bar{R}_{ij\alpha;m\rho_1}^h - \Theta_{ij\alpha m\rho_1}^h\right) \\ &+ F_{(m}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;\alpha\rho_1}^h - \Theta_{ijk\alpha\rho_1}^h\right) + F_{(\rho_1}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;m\alpha}^h - \Theta_{ijkm\alpha}^h\right) + \Theta_{ijkm\rho_1\rho_2}^h. \end{aligned} \tag{13}$$

Suppose that the space with affine connection \bar{A}_n is a 3-symmetric space, i.e. $\bar{R}_{ijk;m_1m_2m_3}^h = 0$, and from (13) we get

$$\begin{aligned} \bar{R}_{ijk;m\rho_1\rho_2}^h &= -F_{(\alpha}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;m\rho_1}^\alpha - \Theta_{ijkm\rho_1}^\alpha\right) + F_{(i}^h \varphi_{\rho_2)} \left(\bar{R}_{\alpha jk;m\rho_1}^h - \Theta_{\alpha jkm\rho_1}^h\right) \\ &+ F_{(j}^h \varphi_{\rho_2)} \left(\bar{R}_{i\alpha k;m\rho_1}^h - \Theta_{i\alpha km\rho_1}^h\right) + F_{(k}^h \varphi_{\rho_2)} \left(\bar{R}_{ij\alpha;m\rho_1}^h - \Theta_{ij\alpha m\rho_1}^h\right) \\ &+ F_{(m}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;\alpha\rho_1}^h - \Theta_{ijk\alpha\rho_1}^h\right) + F_{(\rho_1}^h \varphi_{\rho_2)} \left(\bar{R}_{ijk;m\alpha}^h - \Theta_{ijkm\alpha}^h\right) + \Theta_{ijkm\rho_1\rho_2}^h \end{aligned} \tag{14}$$

where the tensor $\Theta_{ijkm\rho_1}^h$ is defined by formulas (9) and (10).

It is easy to see, that because of (9) and (10) the right hand side of the equations (14) depends on tensors $F_k^h, \bar{R}_{ijk}^h, \varphi_k$ and on their first and second covariant derivatives with respect to the connection of the space A_n . Also, we assume, that the affinator F_i^h and its covariant derivatives $F_{i,j}^h, F_{i,jk}^h$ are known.

It is known [31] that the Riemann tensors of the spaces with affine connection A_n and \bar{A}_n are related to each other by the equations

$$\bar{R}_{ijk}^h = R_{ijk}^h + P_{ik,j}^h - P_{ij,k}^h + P_{ik}^\alpha P_{\alpha j}^h - P_{ij}^\alpha P_{\alpha k}^h. \tag{15}$$

Taking account of structure of the deformation tensor (2), transforming the equations (15), we get

$$\varphi_{i,j} F_k^h + \varphi_{k,j} F_i^h - \varphi_{i,k} F_j^h - \varphi_{j,k} F_i^h = B_{ijk}^h, \tag{16}$$

where

$$B_{ijk}^h = \bar{R}_{ijk}^h - R_{ijk}^h - \varphi_i \left(F_{k,j}^h - F_{j,k}^h + e\delta_k^h \varphi_j + \varphi_\alpha F_k^\alpha F_j^h - e\delta_j^h \varphi_k - \varphi_\alpha F_j^\alpha F_k^h\right) - \varphi_k \left(F_{i,j}^h + \varphi_\alpha F_i^\alpha F_j^h\right) + \varphi_j \left(F_{i,k}^h + \varphi_\alpha F_i^\alpha F_k^h\right).$$

Contracting (16) with the affnor F_ρ^m for indices ρ and h , we have

$$\delta_k^m \varphi_{i,j} + \delta_i^m \varphi_{k,j} - \delta_j^m \varphi_{i,k} - \delta_i^m \varphi_{j,k} = e B_{ijk}^\alpha F_\alpha^m. \tag{17}$$

Contracting (17) for m indices k , we get

$$\varphi_{k,j} - \varphi_{j,k} = \frac{e}{n+1} B_{\beta jk}^\alpha F_\alpha^\beta. \tag{18}$$

Finally, let us contract (17) with respect to m and k . We obtain

$$n\varphi_{i,j} - \varphi_{j,i} = e B_{ij\beta}^\alpha F_\alpha^\beta. \tag{19}$$

Taking account of (18), the equations (19) could be written in the form

$$\varphi_{i,j} = \frac{e}{n-1} \left(B_{ij\beta}^\alpha - \frac{1}{n+1} B_{\beta ji}^\alpha \right) F_\alpha^\beta. \tag{20}$$

It should be remarked that the formulas (20) are obtained for the general case of canonical F -planar mappings $\pi(e)$ ($e = \mp 1$) of spaces with affine connection.

Let us introduce the tensors \bar{R}_{ijkm}^h and $\bar{R}_{ijkm\rho_1}^h$ defined by

$$\bar{R}_{ijk,m}^h = \bar{R}_{ijkm}^h, \tag{21}$$

$$\bar{R}_{ijkm,\rho_1}^h = \bar{R}_{ijkm\rho_1}^h. \tag{22}$$

Taking account of (21) and (22), we could write the equations (14) in the form

$$\begin{aligned} \bar{R}_{ijkm\rho_1,\rho_2}^h = & -F_{(\alpha}^h \varphi_{\rho_2)} (\bar{R}_{ijkm\rho_1}^\alpha - \Theta_{ijkm\rho_1}^\alpha) + F_{(i}^h \varphi_{\rho_2)} (\bar{R}_{\alpha jkm\rho_1}^h - \Theta_{\alpha jkm\rho_1}^h) + F_{(j}^h \varphi_{\rho_2)} (\bar{R}_{iakm\rho_1}^h - \Theta_{iakm\rho_1}^h) \\ & + F_{(k}^h \varphi_{\rho_2)} (\bar{R}_{ij\alpha m\rho_1}^h - \Theta_{ij\alpha m\rho_1}^h) + F_{(m}^h \varphi_{\rho_2)} (\bar{R}_{ijk\alpha\rho_1}^h - \Theta_{ijk\alpha\rho_1}^h) + F_{(\rho_1}^h \varphi_{\rho_2)} (\bar{R}_{ijkma}^h - \Theta_{ijkma}^h) + \Theta_{ijkm\rho_1,\rho_2}^h, \end{aligned} \tag{23}$$

where

$$\begin{aligned} \Theta_{ijkm\rho_1}^h = & -F_{(\alpha}^h \varphi_{\rho_1)} (\bar{R}_{ijkm}^\alpha + \Theta_{ijkm}^\alpha) + F_{(i}^h \varphi_{\rho_1)} (\bar{R}_{\alpha jkm}^h + \Theta_{\alpha jkm}^h) + F_{(j}^h \varphi_{\rho_1)} (\bar{R}_{iakm}^h + \Theta_{iakm}^h) + F_{(k}^h \varphi_{\rho_1)} (\bar{R}_{ij\alpha m}^h + \Theta_{ij\alpha m}^h) \\ & + F_{(m}^h \varphi_{\rho_1)} (\bar{R}_{ijk\alpha}^h + \Theta_{ijk\alpha}^h) - F_{(m}^h \varphi_{\alpha),\rho_1)} \bar{R}_{ijk}^\alpha - F_{(m}^h \varphi_{\alpha)} \bar{R}_{ijk\rho_1}^\alpha + F_{(m}^h \varphi_{\rho_1),\rho_1)} \bar{R}_{\alpha jk}^h + F_{(m}^h \varphi_{\rho_1)} \bar{R}_{\alpha jk\rho_1}^h + F_{(m}^h \varphi_{\rho_1),\rho_1)} \bar{R}_{iak}^h \\ & + F_{(m}^h \varphi_{\rho_1)} \bar{R}_{iak\rho_1}^h + F_{(m}^h \varphi_{\rho_1),\rho_1)} \bar{R}_{ij\alpha}^h + F_{(m}^h \varphi_{\rho_1)} \bar{R}_{ij\alpha\rho_1}^h, \end{aligned} \tag{24}$$

the tensor Θ_{ijkm}^h is defined by the formulas (10).

In the following we have assumed that in (24) covariant derivatives of the vector φ_i are expressed according to (20); in the tensor $\Theta_{ijkm\rho_1,\rho_2}^h$ we could see at (23), covariant derivatives of the vector φ_i are also expressed according to (20), and covariant derivatives of the tensor \bar{R}_{ijk}^h are expressed according to (21).

Obviously, in the space A_n the equations (20), (21), (22), and (23) form a system of PDEs of Cauchy type with respect to the functions $\varphi_i(x)$, $\bar{R}_{ijk}^h(x)$, $\bar{R}_{ijkm}^h(x)$, $\bar{R}_{ijkm\rho_1}^h(x)$.

The functions $\bar{R}_{ijk}^h(x)$, and also $\bar{R}_{ijkl}^h(x)$, $\bar{R}_{ijklm}^h(x)$, must satisfy the algebraic conditions

$$\bar{R}_{i(jk)}^h = 0, \quad \bar{R}_{(ijk)}^h = 0, \quad \bar{R}_{i(jkl)}^h = 0, \quad \bar{R}_{(ijk)l}^h = 0, \quad \bar{R}_{i(jk)lm}^h = 0, \quad \bar{R}_{(ijk)lm}^h = 0. \tag{25}$$

Hence we have proved the theorem.

Theorem 3.1. *In order that a space A_n with an affine connection admit a canonical F -planar mapping $\pi(e)$ ($e = \mp 1$) onto a 3-symmetric space \bar{A}_n , it is necessary and sufficient that the mixed system of differential equations of Cauchy type in covariant derivatives (20), (21), (22), (23), and (25) have a solution with respect to the unknown functions $\varphi_i(x)$, $\bar{R}_{ijk}^h(x)$, $\bar{R}_{ijkl}^h(x)$, $\bar{R}_{ijklm}^h(x)$.*

Obviously, the general solution of the closed mixed system of Cauchy type (20), (21), (22), (23), and (25) depends on no more than

$$1/3 n^2 (n^2 - 1) (1 + n + n^2) + n$$

essential parameters.

References

- [1] O.O. Belova, Connections in fiberings associated with the Grassman manifold and the space of centered planes. *Journal of Mathematical Sciences*. Springer New York. Vol. 162, Issue 5, 2009, 605-632.
- [2] V.E. Berezovski, J. Mikeš, On the classification of almost geodesic mappings of affine-connected spaces. In *Proceedings of the Differential Geometry and Applications Conference, Dubrovnik, Yugoslavia, 26 June–3 July 1988*, pp. 41–48.
- [3] V.E. Berezovski, J. Mikeš, On a classification of almost geodesic mappings of affine connection spaces. *Acta Univ. Palacki. Olomuc. Math.* 35 (1996) 21–24.
- [4] V.E. Berezovski, S. Bácsó, J. Mikeš, Almost geodesic mappings of affinely connected spaces that preserve the Riemannian curvature, *Ann. Math. Inf.* 45 (2015) 3–10.
- [5] V.E. Berezovskii, N.I. Guseva, J. Mikeš, On special first-type almost geodesic mappings of affine connection spaces preserving a certain tensor, *Math. Notes* 98 (2015) 515–518.
- [6] V.E. Berezovski, M. Jukl, L. Juklová, Almost geodesic mappings of the first type onto symmetric spaces, In *Proc. 16th Conf. Appl. Math. (APLIMAT 2017)*, Bratislava, Slovakia, 31 January–2 February 2017.
- [7] V.E. Berezovski, J. Mikeš, On almost geodesic mappings of the type π_1 of Riemannian spaces preserving a system n -orthogonal hypersurfaces, *Rend. Circ. Mat. Palermo II* 59 (1999) 103–108.
- [8] V.E. Berezovski, J. Mikeš, Almost geodesic mappings of type π_1 onto generalized Ricci-symmetric manifolds, *Uch. zap. Kazan. Univ. Ser. Fiz.-Math.* 151 (2009) 9–14.
- [9] V.E. Berezovski, J. Mikeš, On canonical almost geodesic mappings of the first type of affinely connected spaces, *Russ. Math.* 58 (2014) 1–5.
- [10] V.E. Berezovski, J. Mikeš, Almost geodesic mappings of spaces with affine connection, *J. Math. Sci.* 207 (2015) 389–409.
- [11] V.E. Berezovski, J. Mikeš, A. Vanžurová, Almost geodesic mappings onto generalized Ricci-symmetric manifolds, *Acta Math. Acad. Paedag. Nyiregyhaziensis* 26 (2010) 221–230.
- [12] V.E. Berezovski, J. Mikeš, A. Vanžurová, Fundamental PDE's of the canonical almost geodesic mappings of type π_1 , *Bull. Malays. Math. Sci. Soc.* 2 (2014) 647–659.
- [13] V.E. Berezovski, J. Mikeš, H. Chudá, O.Y. Chepurna, On canonical almost geodesic mappings which preserve the Weyl projective tensor, *Russ. Math.* 61 (2017) 1–5.
- [14] V.E. Berezovski, J. Mikeš, L. Rýparová, A. Sabykanov, On canonical almost geodesic mappings of type $\pi_2(e)$, *Mathematics* 8(1):54 (2020).
- [15] V.E. Berezovski, Y. Cherevko, I. Hinterleitner, P. Peška, Geodesic mappings of spaces with affine connections onto generalized symmetric and Ricci-symmetric spaces, *Mathematics* 8(9):1560 (2020).
- [16] V.E. Berezovski, I.A. Kuzmina, J. Mikeš, Canonical F -planar mappings of spaces with affine connection to two symmetric spaces, *Lobachevskii J. Math.* 43(3) (2022) 533–538.
- [17] É. Cartan, Sur une classe remarquable d'espaces de Riemann, I, II, *Bull. S.M.F.* 54 (1926) 214–264; 55 (1927) 114–134.
- [18] S. Helgason, *Differential geometry and symmetric spaces*, Acad. Press, New York, 1962.
- [19] I. Hinterleitner, J. Mikeš, On F -planar mappings of spaces with affine connections, *Note Mat.* 27:1 (2007) 111–118.
- [20] T. Levi-Civita, Sulle trasformazioni delle equazioni dinamiche, *Ann. Mat.* 24 (1896) 252–300.
- [21] J. Mikeš, N.S. Sinyukov, On quasiplanar mappings of spaces of affine connection, *Sov. Math.* 27:1 (1983) 63–70.
- [22] J. Mikeš et al., *Differential Geometry of Special Mappings*, Palacky Univ. Press, Olomouc, 2015.
- [23] J. Mikeš et al., *Differential Geometry of Special Mappings*, (2nd edition), Palacky Univ. Press, Olomouc, 2019.
- [24] A.Z. Petrov, Modeling of physical fields, *Gravitation Gen. Relat.* 4 (1968) 7–21.
- [25] M.Z. Petrović, M.S. Stanković, Special almost geodesic mappings of the first type of non-symmetric affine connection spaces, *Bull. Malays. Math. Sci. Soc.* 40 (2017) 1353–1362.
- [26] M.Z. Petrović, M.S. Stanković, *A note on F-planar mappings of manifolds with non-symmetric linear connection* *Int. J. Geom. Methods Mod. Phys.* Vol. 16, No. 05 (2019) 1950078
- [27] M.Z. Petrović, Canonical almost geodesic mappings of type $\theta\pi_2(0, F)$, $\theta \in \{1, 2\}$ between generalized parabolic Kähler manifolds, *Miskolc Math. Notes* 19 (2018) 469–482.
- [28] M.Z. Petrović, Special almost geodesic mappings of the second type between generalized Riemannian spaces, *Bull. Malays. Math. Sci. Soc.* 42 (2019) 707–727.
- [29] A.P. Shirokov, Structures on differentiable manifolds. *Progress Math.* 9 (1971) 137–207.
- [30] N.S. Sinyukov, Almost geodesic mappings of affinely connected and Riemannian spaces, *Sov. Math.* 4 (1963) 1086–1088.
- [31] N.S. Sinyukov, *Geodesic mappings of Riemannian spaces*, Nauka, Moscow, Russia, 1979.
- [32] N.S. Sinyukov, Almost-geodesic mappings of affinely connected and Riemann spaces, *J. Sov. Math.* 25 (1984) 1235–1249.
- [33] V.S. Sobchuk, Almost geodesic mappings of Riemannian spaces onto symmetric Riemannian spaces, *Mat. Zametki* 17 (1975) 757–763.
- [34] V.S. Sobchuk, J. Mikeš, O. Pokorná, On almost geodesic mappings π_2 between semisymmetric Riemannian spaces, *Novi Sad J. Math.* 9 (1999) 309–312.
- [35] M.S. Stanković, On canonically almost geodesic mappings of the second type of affine spaces, *Filomat* 13 (1999) 105–144.
- [36] M.S. Stanković, M.L. Zlatanović, N.O. Vesić, Basic equations of G -almost geodesic mappings of the second type, which have the property of reciprocity, *Czech. Math. J.* 65 (2015) 787–799.
- [37] N.O. Vesić, M.S. Stanković, Invariants of special second-type almost geodesic mappings of generalized Riemannian space, *Mediterr. J. Math.* 15:2, 60 (2018) 1–12.
- [38] N.O. Vesić, L.S. Velimirović, M.S. Stanković, Some invariants of equitorus third type almost geodesic mappings, *Mediterr. J. Math.* 13 (2016) 4581–4590.

- [39] H. Vavříková, J. Mikeš, O. Pokorná, G. Starko, On fundamental equations of almost geodesic mappings $\pi_2(e)$, *Russ. Math.* 1 (2007) 8–12.
- [40] N.V. Yablonskaya, Special groups of almost geodesic transformations of spaces with affine connection, *Sov. Math.* 30 (1986) 105–108.