



Approximation results for Beta Jakimovski-Leviatan type operators via q -analogue

Md. Nasiruzzaman^a, Mohammed A.O. Tom^a, Stefano Serra-Capizzano^{b,c}, Nadeem Rao^d, Mohammad Ayman-Mursaleen^{e,*}

^aDepartment of Mathematics, Faculty of Science,

University of Tabuk, PO Box 4279, Tabuk-71491, Saudi Arabia

^bDepartment of Science & High Technology, University of Insubria, Via Valleggio 11, 22100 Como, Italy

^cDepartment of Information Technology, Division of Scientific Computing, Uppsala University,
L Gerhyddsv. 2, Hus 2, P.O. Box 337, SE-751 05 Uppsala, Sweden

^dDepartment of Mathematics, University Center for Research & Developement, Chandigarh University, Mohali-140413, Punjab, India

^eSchool of Information & Physical Sciences, The University of Newcastle, University Drive, Callaghan, NSW 2308, Australia

Abstract. We construct a new version of q -Jakimovski-Leviatan type integral operators and show that set of all continuous functions f defined on $[0, \infty)$ are uniformly approximated by our new operators. Finally we construct the Stancu type operators and obtain approximation properties in weighted spaces. Moreover, with the aid of modulus of continuity we discuss the rate of convergence, Lipschitz type maximal approximation and some direct theorems.

1. Preliminaries and introduction

In 1880, Appell investigated a class of polynomials which were named in his honor as Appell polynomials (see [11]). Much later, Jakimovski and Leviatan introduced and modified the Appell polynomials [15] in 1969. The tool for defining the new class is defined below:

$$R(u)e^{uy} = \sum_{k=0}^{\infty} \beta_k(y)u^k, \quad (1)$$

where $\beta_k(y) = \sum_{i=0}^s \alpha_i \frac{y^{s-i}}{(s-i)!}$ ($s \in \mathbb{N}$, the set of positive numbers) and $R(u) = \sum_{k=0}^{\infty} \alpha_k u^k$, $R(1) \neq 0$.

We recall some basic notations regarding the q -calculus (see [19–21]). For each non-negative integer s , the q -integer is defined as

$$[s]_q = \begin{cases} \frac{1-q^s}{1-q}, & q \neq 1 \\ s, & q = 1 \end{cases} \quad \text{for } s \in \mathbb{N} \text{ and } [0]_q = 0.$$

2020 Mathematics Subject Classification. Primary 41A25, 41A36; Secondary 33C45.

Keywords. Appell polynomials; q -Appell polynomials; Jakimovski-Leviatan operators; Korovkin's theorem; modulus of continuity.

Received: 06 December 2022; Revised: 21 February 2023; Accepted: 11 April 2023

Communicated by Miodrag Spalević

* Corresponding author: Mohammad Ayman-Mursaleen

Email addresses: nasir3489@gmail.com, mfarooq@ut.edu.sa (Md. Nasiruzzaman), mohammedkawal@gmail.com (Mohammed A.O. Tom), stefano.serrac@uninsubria.it, stefano.serra@it.uu.se (Stefano Serra-Capizzano), nadeemrao1990@gmail.com, nadeem.e14515@cumail.in (Nadeem Rao), mohdaymanm@gmail.com, mohammad.mursaleen@uon.edu.au (Mohammad Ayman-Mursaleen)

For $|q| < 1$, the q -factorial $[s]_q!$ is defined by

$$[s]_q! = \begin{cases} 1 & (s = 0) \\ \prod_{k=1}^s [k]_q & (s \in \mathbb{N}). \end{cases} \quad (2)$$

In the standard approach the exponential functions for q -calculus:

$$e_q(y) = \sum_{k=0}^{\infty} \frac{y^k}{[k]_q!}. \quad (3)$$

The improper integral of function f is formally defined by as

$$\int_0^{\infty/A} f(y) d_q y = (1-q) \sum_{s \in \mathbb{N}} f\left(\frac{q^s}{A}\right) \frac{q^s}{A}, \quad A \in \mathbb{R} - \{0\}. \quad (4)$$

Al-Salam (see [6, 25]) introduced the family of q -Appell polynomials through the generating functions $R_q(t) = \sum_{s=0}^{\infty} R_{s,q} \frac{t^s}{[s]_q!}$, $R_q(1) \neq 0$. We have

$$R_{r,q}(y) = \sum_{k=0}^r \left[\begin{array}{c} r \\ k \end{array} \right]_{q,y} A_{r-k,q} y^k, \quad (r \in \mathbb{N})$$

and q -differential, $D_{q,y}(R_{r,q}(y)) = [r]R_{r-1,q}(y)$, $r = 1, 2, \dots$. We $D_{q,y}(R_{1,q}(y)) = [1]_q R_{0,q}(y) = K$, where $R_{0,q}(y)$ is a nonzero constant. Moreover, we define $R_q(t)e_q(ty) = \sum_{s=0}^{\infty} R_{r,q}(y) \frac{t^r}{[r]_q!}$, $0 < q < 1$.

In the recent years the Jakimovski-Leviatan type operators have attracted attention in the mathematical community and several papers have been published (for example, see [1–4, 7, 8, 10, 14, 22–24, 29, 31, 36–38, 40] and further recent works such as [5, 12, 13, 28, 30, 32–35, 43–45] on related topics).

Our aim is to construct the extended form in terms of the parametric Stancu variant of recent q -Jakimovski-Leviatan-Beta type integral operators and show that our newly positive linear operators are convergent to the identity acting on the space of continuous functions. Further, we obtain Korovkin type results, estimates on the rate of convergence, as well as direct theorems.

For all $y \in [0, \infty)$, $R_{r,q}(y) \geq 0$ and $R_q(1) \neq 0$, the Jakimovski-Leviatan type operators suppose $\mathcal{T}_{s,q}$ are given by (see [9])

$$\mathcal{T}_{s,q}(f(t); y) = \frac{e_q(-[s]y)}{R_q(1)} \sum_{r=0}^{\infty} \frac{R_{r,q}([s]y)}{[r]_q!} \frac{\mathcal{K}(A, r+1)}{B_q(r+1, s)} \int_0^{\infty/A} \frac{t^r}{(1+t)_q^{r+s+1}} f(q^r t) d_q t, \quad (5)$$

where $s \in \mathbb{N}$, $0 < q < 1$ and $f \in C[0, \infty)$, the latter being the set of all continuous functions on $[0, \infty)$.

Lemma 1.1. [9]

$$\begin{aligned} \sum_{r=0}^{\infty} \frac{R_{r,q}([s]y)}{[r]_q!} &= R_q(1)e_q([s]y), \\ \sum_{r=0}^{\infty} r \frac{R_{r,q}([s]y)}{[r]_q!} &= [[s]R_q(1)y + R'_q(1)]e_q([s]y), \\ \sum_{r=0}^{\infty} r^2 \frac{R_{r,q}([s]y)}{[r]_q!} &= [[s]^2 R_q(1)y^2 + 2[s]R'_q(1)y + R''_q(1)]e_q([s]y), \end{aligned}$$

$$\sum_{r=0}^{\infty} r^3 \frac{R_{r,q}([s]y)}{[r]!} = \left[[s]^3 R_q(1)y^3 + 3[s]^2 R'_q(1)y^2 + 3[s] R''_q(1)y + R'''_q(1) \right] e_q([s]y),$$

$$\sum_{r=0}^{\infty} r^4 \frac{R_{r,q}([s]y)}{[r]!} = \left[[s]^4 R_q(1)y^4 + 4[s]^3 R'_q(1)y^3 + 6[s]^2 R''_q(1)y^2 + 4[s] R'''_q(1)y + R^{(4)}_q(1) \right] e_q([s]y).$$

Lemma 1.2. [9] For the test function $f(t) = 1, t, t^2, t^3, t^4$ the following identities hold true for s large enough, where in the expression of $\mathcal{T}_{s,q}(f(t); y)$ the term t is the dummy variable and y the global one.

$$\begin{aligned}
(1) \quad \mathcal{T}_{s,q}(1; y) &= 1; \\
(2) \quad \mathcal{T}_{s,q}(t; y) &= \frac{1}{q[s-1]} + \frac{1}{[s-1]} \left([s]_q y + \frac{R'_q(1)}{R_q(1)} \right); \\
(3) \quad \mathcal{T}_{s,q}(t^2; y) &= \frac{(1+q)}{q^3[s-1][s-2]} + \frac{(1+2q)}{q^2[s-1][s-2]} \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \\
&\quad + \frac{1}{[s-1][s-2]} \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right); \\
(4) \quad \mathcal{T}_{s,q}(t^3; y) &= \frac{1+2q+2q^2+q^3}{q^4[s-1][s-2][s-3]} + \frac{(1+3q+4q^2+3q^3)}{q^3[s-1][s-2][s-3]} \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \\
&\quad + \frac{(1+2q+3q^2)}{q[s-1][s-2][s-3]} \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \\
&\quad + \frac{q^2}{[s-1][s-2][s-3]} \left([s]^3 y^3 + 3[s]^2 \frac{R'_q(1)}{R_q(1)} y^2 + 3[s] \frac{R''_q(1)}{R_q(1)} y + \frac{R'''_q(1)}{R_q(1)} \right); \\
(5) \quad \mathcal{T}_{s,q}(t^4; y) &= \frac{1+3q+5q^2+6q^3+5q^4+3q^5+q^6}{q^5[s-1][s-2][s-3][s-4]} \\
&\quad + \frac{(1+5q+10q^2+13q^3+12q^4+7q^5+2q^6)}{q^3[s-1][s-2][s-3][s-4]} \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \\
&\quad + \frac{(1+3q+7q^2+9q^3+9q^4+6q^5)}{q[s-1][s-2][s-3][s-4]} \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \\
&\quad + \frac{(q^2+2q^3+2q^4+2q^5+q^6+2q^7)}{[s-1][s-2][s-3][s-4]} \\
&\quad \times \left([s]^3 y^3 + 3[s]^2 \frac{R'_q(1)}{R_q(1)} y^2 + 3[s] \frac{R''_q(1)}{R_q(1)} y + \frac{R'''_q(1)}{R_q(1)} \right) \\
&\quad + \frac{q^6}{[s-1][s-2][s-3][s-4]} \\
&\quad \times \left([s]^4 y^4 + 4[s]^3 \frac{R'_q(1)}{R_q(1)} y^3 + 6[s]^2 \frac{R''_q(1)}{R_q(1)} y^2 + 4[s] \frac{R'''_q(1)}{R_q(1)} y + \frac{R^{(4)}_q(1)}{R_q(1)} \right).
\end{aligned}$$

2. Operators and its associated Moments

We suppose $y \in [0, \infty)$, $R_{r,q}(y) \geq 0$, $R_q(1) \neq 0$ and $f \in C_\lambda[0, \infty) = \{f \in C[0, \infty) : f(t) = O(t^\lambda), \text{ as } t \rightarrow \infty\}$. Then for any $0 \leq \nu \leq \mu$ and $0 < q < 1$, we construct

$$\mathcal{W}_{s,q}(f; y) = \frac{e_q(-[s]y)}{R_q(1)} \sum_{r=0}^{\infty} \frac{R_{r,q}([s]y)}{[r]!} \frac{\mathcal{K}(A, r+1)}{B_q(r+1, s)} \int_0^{\infty/A} \frac{t^r}{(1+t)^{q^{r+s+1}}} f\left(\frac{q^r([s]t+\nu)}{[s]+\mu}\right) d_q t, \quad (6)$$

where $\lambda > s, s \in \mathbb{N}$, $[s] = [s]_q$, $[r] = [r]_q$ and

$$B_q(r, s) = \mathcal{K}(A, r) \int_0^{\infty/A} \frac{y^{r-1}}{(1+y)_q^{r+s}} d_q y = \frac{[r-1]}{[s]} B_q(r-1, s+1), \quad r > 1, s \geq 0,$$

with

$$\mathcal{K}(A, r+1) = q^r \mathcal{K}(A, r),$$

$$\mathcal{K}(A, r) = q^{\frac{r(r-1)}{2}}, \quad \mathcal{K}(A, 0) = 1.$$

Lemma 2.1. For all $y \geq 0$, $\mathcal{W}_{s,q}(1; y) = 1$ and the following identities hold true.

- (1) $\mathcal{W}_{s,q}(t; y) = \left(\frac{[s]}{[s]+\mu} \right) \mathcal{T}_{s,q}(t; y) + \left(\frac{\nu}{[s]+\mu} \right) \mathcal{T}_{s,q}(1; y),$
- (2) $\mathcal{W}_{s,q}(t^2; y) = \left(\frac{[s]}{[s]+\mu} \right)^2 \mathcal{T}_{s,q}(t^2; y) + \frac{2\nu[s]}{([s]+\mu)^2} \mathcal{T}_{s,q}(t; y) + \left(\frac{\nu}{[s]+\mu} \right)^2 \mathcal{T}_{s,q}(1; y),$
- (3) $\mathcal{W}_{s,q}(t^3; y) = \left(\frac{[s]}{[s]+\mu} \right)^3 \mathcal{T}_{s,q}(t^3; y) + \frac{3\nu[s]^2}{([s]+\mu)^3} \mathcal{T}_{s,q}(t^2; y)$
 $+ \frac{3\nu^2[s]}{([s]+\mu)^3} \mathcal{T}_{s,q}(t; y) + \left(\frac{\nu}{[s]+\mu} \right)^3 \mathcal{T}_{s,q}(1; y),$
- (4) $\mathcal{W}_{s,q}(t^4; y) = \left(\frac{[s]}{[s]+\mu} \right)^4 \mathcal{T}_{s,q}(t^4; y) + \frac{4\nu[s]^3}{([s]+\mu)^4} \mathcal{T}_{s,q}(t^3; y) + \frac{6\nu^2[s]^2}{([s]+\mu)^4} \mathcal{T}_{s,q}(t^2; y)$
 $+ \frac{4\nu^3[s]}{([s]+\mu)^4} \mathcal{T}_{s,q}(t; y) + \left(\frac{\nu}{[s]+\mu} \right)^4 \mathcal{T}_{s,q}(1; y).$

Lemma 2.2. Take the test functions $f(t) = 1, t, t^2, t^3, t^4$. For any $s \in \mathbb{N} - \{1, 2, 3, 4\}$, the operators in (6) satisfy the following identities.

- 1° $\mathcal{W}_{s,q}(1; y) = 1;$
- 2° $\mathcal{W}_{s,q}(t; y) = \frac{[s]}{q([s]+\mu)[s-1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s]+\mu};$
- 3° $\mathcal{W}_{s,q}(t^2; y) = \left(\frac{[s]}{[s]+\mu} \right)^2 \frac{1}{q^3[s-1][s-2]} \left\{ 1 + q + (q + 2q^2) \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \right.$
 $\left. + q^3 \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \right\}$
 $- \frac{2\nu[s]}{q[s-1]([s]+\mu)^2} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \left(\frac{\nu}{[s]+\mu} \right)^2;$
- 4° $\mathcal{W}_{s,q}(t^3; y) = \left(\frac{[s]}{[s]+\mu} \right)^3 \frac{1}{q^4[s-1][s-2][s-3]} \left\{ (1 + 2q + 2q^2 + q^3) \right.$
 $\left. + (q + 3q^2 + 4q^3 + 3q^4) \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \right.$
 $\left. + (q^3 + 2q^4 + 3q^5) \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \right\}$

$$\begin{aligned}
& +q^6 \left([s]^3 y^3 + 3[s]^2 \frac{R'_q(1)}{R_q(1)} y^2 + 3[s] \frac{R''_q(1)}{R_q(1)} y + \frac{R'''_q(1)}{R_q(1)} \right) \} \\
& + \frac{3\nu[s]^2}{([s]+\mu)^3} \frac{1}{q^3[s-1][s-2]} \left\{ \left(1+q+(q+2q^2) \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \right. \right. \\
& \left. \left. + q^3 \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \right) \right. \\
& \left. - \frac{2\nu[s]}{q[s-1]([s]+\mu)^2} \left(1+q[s]y + \frac{qP'_q(1)}{R_q(1)} \right) + \left(\frac{\nu}{[s]+\mu} \right)^2 \right\} \\
& + \frac{3\nu^2[s]}{([s]+\mu)^3} \frac{1}{q[s-1]} \left\{ \left(1+q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s]+\mu} \right\} + \left(\frac{\nu}{[s]+\mu} \right)^3 ; \\
5^\circ \quad \mathcal{W}_{s,q}(t^4; y) &= \left(\frac{[s]}{[s]+\mu} \right)^4 \frac{1}{q^5[s-1][s-2][s-3][s-4]} \\
& \times \left\{ (1+3q+5q^2+6q^3+5q^4+3q^5+q^6) \right. \\
& + \left(q^2+5q^3+10q^4+13q^5+12q^6+7q^7+2q^8 \right) \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \\
& + \left(q^4+3q^4+7q^6+9q^7+9q^8+6q^9 \right) \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \\
& + \left(q^7+2q^8+2q^9+2q^{10}+q^{11}+2q^{12} \right) \\
& \times \left([s]^3 y^3 + 3[s]^2 \frac{R'_q(1)}{R_q(1)} y^2 + 3[s] \frac{R''_q(1)}{R_q(1)} y + \frac{R'''_q(1)}{R_q(1)} \right) \\
& + q^{11} \left([s]^4 y^4 + 4[s]^3 \frac{R'_q(1)}{R_q(1)} y^3 + 6[s]^2 \frac{R''_q(1)}{R_q(1)} y^2 + 4[s] \frac{R'''_q(1)}{R_q(1)} y + \frac{R^{(4)}_q(1)}{R_q(1)} \right) \} \\
& + \frac{4\nu[s]^3}{([s]+\mu)^4} \frac{1}{q^4[s-1][s-2][s-3]} \left\{ (1+2q+2q^2+q^3) \right. \\
& + \left(q+3q^2+4q^3+3q^4 \right) \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \\
& + \left(q^3+2q^4+3q^5 \right) \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \\
& + q^6 \left([s]^3 y^3 + 3[s]^2 \frac{R'_q(1)}{R_q(1)} y^2 + 3[s] \frac{R''_q(1)}{R_q(1)} y + \frac{R'''_q(1)}{R_q(1)} \right) \} \\
& + \frac{3\nu[s]^2}{([s]+\mu)^3} \frac{1}{q^3[s-1][s-2]} \left\{ \left(1+q+(q+2q^2) \left([s]y + \frac{R'_q(1)}{R_q(1)} \right) \right. \right. \\
& \left. \left. + q^3 \left([s]^2 y^2 + \frac{2[s]R'_q(1)}{R_q(1)} y + \frac{R''_q(1)}{R_q(1)} \right) \right) \right. \\
& \left. - \frac{2\nu[s]}{q[s-1]([s]+\mu)^2} \left(1+q[s]y + \frac{qP'_q(1)}{R_q(1)} \right) + \left(\frac{\nu}{[s]+\mu} \right)^2 \right\} \\
& + \frac{3\nu^2[s]}{([s]+\mu)^3} \frac{1}{q[s-1]} \left\{ \left(1+q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s]+\mu} \right\} + \left(\frac{\nu}{[s]+\mu} \right)^3 \}
\end{aligned}$$

$$\begin{aligned}
& + \frac{6\nu^2[s]^2}{([s] + \mu)^4} \frac{1}{q^3[s - 1][s - 2]} \left\{ \left(1 + q + (q + 2q^2) \left(([s]y + \frac{R'_q(1)}{R_q(1)}) \right. \right. \right. \\
& \left. \left. \left. + q^3 \left([s]^2y^2 + \frac{2[s]R'_q(1)}{R_q(1)}y + \frac{R''_q(1)}{R_q(1)} \right) \right) \right. \right. \\
& \left. \left. - \frac{2\nu[s]}{q[s - 1]([s] + \mu)^2} \left(1 + q[s]y + \frac{qP'_q(1)}{R_q(1)} \right) + \left(\frac{\nu}{[s] + \mu} \right)^2 \right\} \right. \\
& \left. + \frac{4\nu^3[s]}{([s] + \mu)^4} \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} + \left(\frac{\nu}{[s] + \mu} \right)^4. \right.
\end{aligned}$$

Remark 2.3. For the choices for $\nu = \mu = 0$ then our new operators defined by (6) simply reduces to those studied in the recent published article [9] so that $\mathcal{W}_{s,q}(t^i; y)$ is the same as $\mathcal{T}_{s,q}(t^i; y)$ for all $i = 0, 1, 2, 3, 4$.

Lemma 2.4. Take $\gamma_i(y) = (t - y)^i$ for $i = 1, 2$. Then for every $y \in [0, \infty)$, $0 < q < 1$, $R_{r,q}(y) \geq 0$ with $R_q(1) \neq 0$ if we put

$$(\delta_{s,q})^i = \mathcal{W}_{s,q}(\gamma_i(y); y) \text{ for } i = 1, 2, s > 1, 2. \quad (7)$$

Furthermore the following equalities hold.

$$\begin{aligned}
& (\delta_{s,q})^i = \begin{cases} \left(\frac{[s]}{[s] + \mu} \right)^2 \frac{1}{q^3[s - 1][s - 2]} \left\{ 1 + q + (q + 2q^2) \left(([s]y + \frac{R'_q(1)}{R_q(1)}) \right. \right. \\ \left. \left. + q^3 \left([s]^2y^2 + \frac{2[s]R'_q(1)}{R_q(1)}y + \frac{R''_q(1)}{R_q(1)} \right) \right) \right. \\ \left. + \frac{2\nu[s]}{q[s - 1]([s] + \mu)^2} \left(1 + q[s]y + \frac{qP'_q(1)}{R_q(1)} \right) + \left(\frac{\nu}{[s] + \mu} \right)^2 \right. \\ \left. - \frac{2[s]y}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) - \frac{2\nu y}{[s] + \mu} + y^2, \right. \\ \text{for } i = 2, s > 2, \\ \left. \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - y, \right. \\ \text{for } i = 1, s > 1. \end{cases}
\end{aligned}$$

3. Approximation in Weighted Space

Suppose that $C_b[0, \infty)$ be the set of all bounded and continuous functions on $[0, \infty)$ with the supremum norm:

$$\|f\|_{C_b} = \sup_{y \in [0, \infty)} |f(y)|.$$

Let

$$E := \left\{ f : y \in [0, \infty), \frac{f(y)}{1 + y^2} \text{ is convergent as } y \rightarrow \infty \right\}.$$

We choose $q = q_s$ where $0 < q_s < 1$ such that

$$\lim_s q_s \rightarrow 1, \quad \lim_\tau q_s^s \rightarrow \alpha \quad (8)$$

Theorem 3.1. For any function $f \in C[0, \infty) \cap E$, we have

$$\lim_{s \rightarrow \infty} \mathcal{W}_{s,q}(f; y) \rightarrow f(y)$$

uniformly convergent on each compact subset of $[0, \infty)$.

Proof. We take in account the Korovkin theorem [26]. Indeed, as $s \rightarrow \infty$, the operators $\mathcal{W}_{s,q}$ converges to the identity operator on the set of all continuous functions defined on $[0, \infty)$, because

$$\lim_{s \rightarrow \infty} \mathcal{W}_{s,q}(t^j; y) = y^j, \quad j = 0, 1, 2.$$

When $s \rightarrow \infty$, from (8), it is immediate to verify that $\frac{1}{[s]_{qs}} \rightarrow 0$ and $\frac{[s]_{ds}}{[s-1]_{qs}} \rightarrow 1$. Consequently

$$\lim_{s \rightarrow \infty} \mathcal{W}_{s,q}(1; y) = 1, \quad \lim_{s \rightarrow \infty} \mathcal{W}_{s,q}(t; y) = y, \quad \lim_{s \rightarrow \infty} \mathcal{W}_{s,q}^*(t^2; y) = y^2,$$

which completes the proof. \square

Suppose that $\phi(y)$ is a continuous and strictly increasing function on $[0, \infty)$ and take $\varrho(y) = 1 + \phi^2(y)$, $\lim_{y \rightarrow \infty} \varrho(y) \rightarrow \infty$ such that $\phi(y) \rightarrow y$. Let $B_\varrho[0, \infty)$ be a set of functions defined on $[0, \infty)$, such that $|f(y)| \leq M_f \varrho(y)$, where M_f is any positive constant. Furthermore, we denote the set all continuous functions on $[0, \infty)$ by $C[0, \infty)$ and its subsets $C_\varrho[0, \infty)$ with $C_\varrho[0, \infty) = B_\varrho[0, \infty) \cap C[0, \infty)$. It is well known for the sequence of linear positive operators $\{L_s\}_{s \geq 1}$ (see [17]) maps $C_\varrho[0, \infty) \rightarrow B_\varrho[0, \infty)$ if and only if

$$|L_s(\varrho; y)| \leq C\varrho(y),$$

where C is a positive constant. For $m \in \mathbb{N}$, let us denote

$$C_\varrho^m[0, \infty) = \left\{ f \in C_\varrho[0, \infty) : \lim_{y \rightarrow \infty} \frac{f(y)}{\varrho(y)} = c \in \mathbb{R} \right\}. \quad (9)$$

Theorem 3.2. [17, 18] Suppose $\{L_s\}_{s \geq 1}$ the positive linear operators which acting from $C_\varrho[0, \infty) \rightarrow B_\varrho[0, \infty)$ for $j = 0, 1, 2$ such that $\lim_{s \rightarrow \infty} \|L_s(\phi_j) - y^j\|_{\varrho(y)} = 0$. Then, for every $f \in C_\varrho[0, \infty)$, it follows that

$$\lim_{s \rightarrow \infty} \|L_s(f) - f\|_{\varrho(y)} = 0.$$

Theorem 3.3. Take $\varphi \in C_\varrho[0, \infty)$ which acting from $C_\varrho[0, \infty) \rightarrow B_\varrho[0, \infty)$ for $j = 0, 1, 2$ then for all $y \in [0, \infty)$ and $\varphi \in C_\varrho[0, \infty)$ with $R_{r,q}(y) \geq 0$, $R_q(1) \neq 0$ it follows that

$$\lim_{s \rightarrow \infty} \|\mathcal{W}_{s,q}(\varphi) - \varphi\|_{\varrho(y)} = 0.$$

Proof. Supposing $\lambda^j = t^j$ for $j = 0, 1, 2$, then by the use of the Korovkin theorem (see [26]), we directly conclude that

$$\lim_{s \rightarrow \infty} \|\mathcal{W}_{s,q}(\varphi^j) - \varphi^j\|_{\varrho(y)} = 0, \quad j = 0, 1, 2.$$

By considering the Lemma 2.1, we obtain

$$\|\mathcal{W}_{s,q}(\lambda^0) - 1\|_{\varrho(y)} = \sup_{y \in [0, \infty)} \frac{|\mathcal{W}_{s,q}(1; y) - 1|}{\varrho(y)} = 0.$$

For $j = 1$, we can write

$$\begin{aligned} \|\mathcal{W}_{s,q}(\lambda^1) - y\|_{\varrho(y)} &= \sup_{y \in [0, \infty)} \frac{|\mathcal{W}_{s,q}(\lambda^1; y) - y|}{\varrho(y)} \\ &= \sup_{y \in [0, \infty)} \frac{y}{\varrho(y)} \left| \left(\frac{[s]^2}{([s] + \mu)[s - 1]} - 1 \right) \right| \\ &\quad + \sup_{y \in [0, \infty)} \frac{1}{\varrho(y)} \left| \frac{[s]}{([s] + \mu)[s - 1]} \frac{R'_q(1)}{R_q(1)} + \frac{\nu}{[s] + \mu} \right|. \end{aligned}$$

If $s \rightarrow \infty$, then easily we get $\|\mathcal{W}_{s,q}(\lambda^1) - y\|_{\varrho(y)} \rightarrow 0$.

For $j = 2$, we get

$$\begin{aligned} \|\mathcal{W}_{s,q}(\lambda^2) - y^2\|_{\varrho(y)} &= \sup_{y \in [0, \infty)} \frac{|\mathcal{W}_{s,q}(\lambda^2; y) - y^2|}{\varrho(y)} \\ &= \sup_{y \in [0, \infty)} \frac{y^2}{\varrho(y)} \left| \left(\frac{[s]^4}{([s] + \mu)^2[s - 1][s - 2]} - 1 \right) \right| \\ &\quad + \sup_{y \in [0, \infty)} \frac{y}{\varrho(y)} \left| \frac{[s]^2}{q^3([s] + \mu)^2[s - 1]} \left\{ \frac{1 + q + (q + 2q^2)[s]}{[s - 2]} \right. \right. \\ &\quad \left. \left. + \frac{[s]}{[s - 2]} \frac{2q^3R'_q(1)}{R_q(1)} - 2\nu q^2 \right\} \right| \\ &\quad + \sup_{y \in [0, \infty)} \frac{1}{\varrho(y)} \left| \frac{[s]}{q^3([s] + \mu)^2[s - 1]} \left\{ \frac{(1 + q)[s]}{[s - 2]} \right. \right. \\ &\quad \left. \left. + \frac{q(1 + 2q)[s]}{[s - 2]} \frac{R'_q(1)}{R_q(1)} + \frac{[s]}{[s - 2]} \frac{q^3R''_q(1)}{R_q(1)} - 2\nu q^2 \left(1 + \frac{qR'_q(1)}{R_q(1)} \right) \right\} \right| \end{aligned}$$

Thus we easily infer $\|\mathcal{W}_{s,q}(\lambda^2) - y^2\|_{\varrho(y)} \rightarrow 0$, as $s \rightarrow \infty$. \square

Theorem 3.4. For all $\varphi \in C_\varrho^m[0, \infty)$, $m \in \mathbb{N}$ and any $\xi \in [0, \infty)$, the operators $\mathcal{W}_{s,q}$ are such that the following limit relationship is satisfied:

$$\lim_{s \rightarrow \infty} \sup_{y \in [0, \infty)} \frac{|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)|}{(\varrho(y))^{\xi+1}} = 0.$$

Proof. By the virtue of $|\varphi(y)| \leq \|(1 + y^2)\varphi\|_{\varrho(y)}$ and for any positive number $y_0 \in \mathbb{R}$ (the real number), we easily obtain that

$$\begin{aligned} \lim_{s \rightarrow \infty} \sup_{y \in [0, \infty)} \frac{|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)|}{(\varrho(y))^{\xi+1}} &\leq \sup_{y \leq y_0} \frac{|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)|}{(\varrho(y))^{\xi+1}} + \sup_{y \geq y_0} \frac{|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)|}{(\varrho(y))^{\xi+1}} \\ &\leq \|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)\|_{C[0, y_0]} \\ &\quad + \|\varphi\|_{\varrho(y)} \sup_{y \geq y_0} \frac{|\mathcal{W}_{s,q}(1 + t^2; y) - \varphi(y)|}{(\varrho(y))^{\xi+1}} + \sup_{y \geq y_0} \frac{|\varphi(y)|}{(\varrho(y))^{\xi+1}} \\ &= \mathcal{M}_1 + \mathcal{M}_2 + \mathcal{M}_3, (\text{suppose}). \end{aligned}$$

Thus

$$\mathcal{M}_3 = \sup_{y \geq y_0} \frac{|\varphi(y)|}{(\varrho(y))^{\xi+1}} \leq \sup_{y \geq y_0} \frac{\|\varphi\|_{\varrho(y)}(1+y^2)}{(\varrho(y))^{\xi+1}} \leq \frac{\|\varphi\|_{\varrho(y)}}{(1+y_0^2)^\xi}. \quad (10)$$

In view of Lemma 2.2, we deduce that

$$\limsup_{s \rightarrow \infty} \sup_{y \geq y_0} \frac{\mathcal{W}_{s,q}(1+t^2; y)}{\varrho(y)} = 1,$$

while, for any $\epsilon > 0$, there exists $s_1 \in \mathbb{N}$ such that $s \geq s_1$ satisfying

$$\sup_{y \geq y_0} \frac{\mathcal{W}_{s,q}(1+t^2; y)}{\varrho(y)} \leq \frac{(1+y_0^2)^\xi}{\|\varphi\|_{\varrho(y)}} \frac{\epsilon}{3} + 1.$$

Thus, for any $s \geq s_1$, we have

$$\mathcal{M}_2 = \|\varphi\|_{\varrho(y)} \sup_{y \geq y_0} \frac{\mathcal{W}_{s,q}(1+t^2; y)}{(\varrho(y))^{\xi+1}} \leq \frac{\|\varphi\|_{\varrho(y)}}{(1+y_0^2)^\xi} + \frac{\epsilon}{3}. \quad (11)$$

Taking into account the relations in (10) and (11), we infer

$$\mathcal{M}_2 + \mathcal{M}_3 \leq 2 \frac{\|\varphi\|_{\varrho(y)}}{(1+y_0^2)^\xi} + \frac{\epsilon}{3}.$$

Choosing y_0 such that $\frac{\|\varphi\|_{\varrho(y)}}{(1+y_0^2)^\xi} \leq \frac{\epsilon}{6}$, we find

$$\mathcal{M}_2 + \mathcal{M}_3 \leq \frac{2\epsilon}{3}, \quad \text{for all } s \geq s_1, \quad (12)$$

and similarly, for $s_2 \geq s$ such that

$$\mathcal{M}_1 = \|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)\|_{C[0, y_0]} \leq \frac{\epsilon}{3}. \quad (13)$$

Next, choose $s_3 = \max(s_1, s_2)$. From (12) and (13), it is plain to deduce

$$\sup_{y \in [0, \infty)} \frac{|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)|}{(\varrho(y))^{\xi+1}} < \epsilon,$$

which completes the proof of Theorem 3.4. \square

4. Order of Approximation

In the present section we study the order of convergence in terms of modulus of continuity functions in terms of Peetres K -functional and Lipschitz type maximal function. Suppose any $\delta > 0$, then for all $f \in C[a, b]$ (spaces of all continuous functions on $[a, b]$) the classical modulus of continuity is defined as

$$\omega(f; \delta) = \sup_{(y_1, y_2) \in [a, b] \times [a, b], |y_1 - y_2| \leq \delta} |f(y_1) - f(y_2)|.$$

Theorem 4.1. [46] For any sequence of positive linear operators suppose $\{L\}_{s \geq 1} : [y_1, y_2] \rightarrow C[u, v]$ and $[u, v] \subseteq [y_1, y_2]$, we obtain

1. if $f \in C[y_1, y_2]$ and $y \in [u, v]$, then

$$\begin{aligned} |L_s(f; y) - f(y)| &\leq |f(y)||L_s(1; y) - 1| \\ &+ \left\{ L_s(1; y) + \frac{1}{\delta} \sqrt{L_s((t-y)^2; y)} \sqrt{L_s(1; y)} \right\} \omega(f; \delta). \end{aligned}$$

2. Furthermore, for any $\varphi' \in C[y_1, y_2]$, for all $y \in [u, v]$, one has

$$\begin{aligned} |L_s(\varphi; y) - \varphi(y)| &\leq |\varphi(y)||L_s(1; y) - 1| + |\varphi'(y)||L_s(t-y; y)| \\ &+ L_s((t-y)^2; y) \left\{ \sqrt{L_s(1; y)} + \frac{1}{\delta} \sqrt{L_s((t-y)^2; y)} \right\} \omega(\varphi'; \delta). \end{aligned}$$

Theorem 4.2. Suppose $f \in C_\varrho[0, \infty)$ and $y \in [0, \infty)$. Then the inequality

$$|\mathcal{W}_{s,q}(f; y) - f(y)| \leq 2\omega\left(f; \sqrt{\delta_{s,q}^*(y)}\right)$$

holds, with $\delta = \sqrt{\delta_{s,q}^*(y)} = \sqrt{\mathcal{W}_{s,q}((t-y)^2; y)}$.

Proof. By considering Lemma 2.2 and Theorem 4.1, we directly obtain

$$\begin{aligned} |\mathcal{W}_{s,q}(f; y) - f(y)| &\leq |f(y)||\mathcal{W}_{s,q}(1; y) - 1| + \left\{ \mathcal{W}_{s,q}(1; y) \right. \\ &+ \left. \frac{1}{\delta} \sqrt{\mathcal{W}_{s,q}((t-y)^2; y)} \sqrt{\mathcal{W}_{s,q}(1; y)} \right\} \omega(f; \delta), \end{aligned}$$

on choosing $\delta = \sqrt{\delta_{s,q}^*(y)} = \sqrt{\mathcal{W}_{s,q}((t-y)^2; y)}$. The desired results follows now from a denumerability argument. \square

Theorem 4.3. For $y \in [0, \infty)$, for all $\varphi' \in C_\varrho[0, \infty)$, the inequality

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &\leq \left| \left(\frac{[s]^2}{([s] + \mu)[s-1]} - 1 \right) y + \frac{[s]}{([s] + \mu)[s-1]} \frac{R'_q(1)}{R_q(1)} + \frac{\nu}{[s] + \mu} \right| |\varphi'(y)| \\ &+ 2\delta_{s,q}^*(y)\omega\left(\varphi'; \sqrt{\delta_{s,q}^*(y)}\right) \end{aligned}$$

holds, where $\delta_{s,q}^*(y) = \mathcal{W}_{s,q}((t-y)^2; y)$.

Proof. By taking in account Lemma 2.2, Lemma 2.4 and Theorem 4.1, we deduce that

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &\leq |\mathcal{W}_{s,q}(1; y) - 1||\varphi(y)| + |\varphi'(y)||\mathcal{W}_{s,q}(t-y; y)| \\ &+ \mathcal{W}_{s,q}((t-y)^2; y) \left\{ \sqrt{\mathcal{W}_{s,q}(1; y)} + \frac{1}{\delta} \sqrt{\mathcal{W}_{s,q}((t-y)^2; y)} \right\} \omega(\varphi'; \delta). \end{aligned}$$

In the light of Theorem 4.2, by using $\delta = \sqrt{\delta_{s,q}^*(y)} = \sqrt{\mathcal{W}_{s,q}((t-y)^2; y)}$, the claimed result follows. \square

5. Approximation on Lipschitz space and Peetre's K -functional

In the present section we provide Lipschitz type maximal approximation results and a local direct theorem for our new operators (6). To this end, we recall the following facts regarding Lipschitz type maximal functions. Indeed, for any choice of real parameters $\mu_1, \mu_2 > 0$ and $\chi \in (0, 1]$, we can define (see [39])

$$Lip_{\mathcal{L}}^\chi = \left\{ \varphi \in C_b[0, \infty) : |\varphi(t) - \varphi(y)| \leq \mathcal{L} \frac{|t-y|^\chi}{(\mu_1 y^2 + \mu_2 y + t)^{\frac{\chi}{2}}}; y, t \in [0, \infty) \right\},$$

where \mathcal{L} is denoted for Lipschitz constant.

Theorem 5.1. For any $\varphi \in Lip_{\mathcal{L}}^{\chi}$ the operators $\mathcal{W}_{s,q}$ verify the inequality

$$|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| \leq \mathcal{L} \left(\frac{\delta_{s,q}^*(y)}{(\mu_1 y^2 + \mu_2 y)} \right)^{\frac{\chi}{2}},$$

where $\delta_{s,q}^*(y)$ is given by Theorem 4.2.

Proof. Take $\varphi \in Lip_{\mathcal{L}}^{\chi}$ for $0 < \chi \leq 1$. First we show that the result is true for $\chi = 1$. For any $\mu_1, \mu_2 \geq 0$ we know the inequality $(\mu_1 y^2 + \mu_2 y + t)^{-1/2} \leq (\mu_1 y^2 + \mu_2 y)^{-1/2}$, thus by use of Cauchy-Schwarz inequality it can be written as:

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &\leq |\mathcal{W}_{s,q}(|\varphi(t) - \varphi(y)|; y)| + |\varphi(y)| |(1; y) - 1| \\ &\leq \mathcal{W}_{s,q} \left(\frac{|t - y|}{(\mu_1 y^2 + \mu_2 y + t)^{\frac{1}{2}}}; y \right) \\ &\leq \mathcal{L}(\mu_1 y^2 + \mu_2 y)^{-1/2} \mathcal{W}_{s,q}(|t - y|; y) \\ &\leq \mathcal{L}(\mu_1 y^2 + \mu_2 y)^{-1/2} |\mathcal{W}_{s,q}((t - y)^2; y)|^{1/2}. \end{aligned}$$

It can be concluded that the result is valid for $\chi = 1$. Now we verify the results for $0 < \chi < 1$. More precisely, we simply use the monotonicity property of the operators $\mathcal{W}_{s,q}$ and we use the Hölder's inequality. In this manner we have

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &\leq \mathcal{W}_{s,q}(|\varphi(t) - \varphi(y)|; y) \\ &\leq \left(\mathcal{W}_{s,q} \left(\left| \varphi(t) - \varphi(y) \right|^{\frac{2}{\chi}}; y \right) \right)^{\frac{\chi}{2}} \left(\mathcal{W}_{s,q}(1; y) \right)^{\frac{2-\chi}{2}} \\ &\leq \mathcal{L} \left\{ \frac{\mathcal{W}_{s,q}((t - y)^2; y)}{\mu_1 y^2 + \mu_2 y + t} \right\}^{\frac{\chi}{2}} \\ &\leq \mathcal{L}(\mu_1 y^2 + \mu_2 y)^{-\chi/2} \left\{ \mathcal{W}_{s,q}((t - y)^2; y) \right\}^{\frac{\chi}{2}} \\ &\leq \mathcal{L}(\mu_1 y^2 + \mu_2 y)^{-\chi/2} \left(\mathcal{W}_{s,q}(t - y)^2; y \right)^{\frac{\chi}{2}} \\ &= \mathcal{L} \left(\frac{\delta_{s,q}^*(y)}{(\mu_1 y^2 + \mu_2 y)} \right)^{\frac{\chi}{2}}, \end{aligned}$$

which gives the desired proof. \square

Here we also compute few other local approximation of $\mathcal{W}_{s,q}$ in terms of modulus of continuity of order one, by making use of Lipschitz maximal spaces. For any $\varphi \in C_b[0, \infty)$, $0 < \chi \leq 1$ and $t, y \in [0, \infty)$, we recall (see [27])

$$\omega_{\chi}(\varphi; y) = \sup_{t \neq y, t \in [0, \infty)} \frac{|\varphi(t) - \varphi(y)|}{|t - y|^{\chi}}, \quad (14)$$

where ω_{χ} denotes the modulus of continuity order one.

Theorem 5.2. For any $\varphi \in C_b[0, \infty)$, we have

$$|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| \leq \left(\delta_{s,q}^*(y) \right)^{\frac{\chi}{2}} \omega_{\chi}(\varphi; y),$$

where $\omega_{\chi}(f; y)$ is defined by (14) and $\delta_{s,q}^*(y)$ is given in Theorem 4.2.

Proof. By exploiting the Hölder inequality, it is easy to deduce the following chain of relationships

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &\leq \mathcal{W}_{s,q}(|\varphi(t) - \varphi(y)|; y) \\ &\leq \omega_\lambda(\varphi; y) \mathcal{W}_{s,q}(|t - y|^\lambda; y) \\ &\leq \omega_\lambda(\varphi; y) (\mathcal{W}_{s,q}(1; y))^{\frac{2-\lambda}{2}} (\mathcal{W}_{s,q}(|t - y|^2; y))^{\frac{\lambda}{2}} \\ &= \omega_\lambda(\varphi; y) (\mathcal{W}_{s,q}((t - y)^2; y))^{\frac{\lambda}{2}}, \end{aligned}$$

so that the proof is concluded. \square

Next, denoting by $C_b[0, \infty)$ the class of all continuous and bounded functions on the semi-axis $[0, \infty)$, we define

$$C_b^2[0, \infty) = \{\varphi \in C_b[0, \infty) : \varphi', \varphi'' \in C_b[0, \infty)\}, \quad (15)$$

$$\|\varphi\|_{C_b^2[0, \infty)} = \|\varphi\|_{C_b[0, \infty)} + \|\varphi'\|_{C_b[0, \infty)} + \|\varphi''\|_{C_b[0, \infty)}, \quad (16)$$

$$\|\varphi\|_{C_b[0, \infty)} = \sup_{y \in [0, \infty)} |\varphi(y)|. \quad (17)$$

Theorem 5.3. For all $\varphi \in C_b^2[0, \infty)$ the operators $\mathcal{W}_{s,q}(\cdot; \cdot)$ are such that

$$|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| \leq (2\Omega_{s,q}^*(y) + \delta_{s,q}^*(y)) \frac{\|\varphi\|_{C_b^2[0, \infty)}}{2},$$

where $\delta_{s,q}^*(y)$ is given in Theorem 4.2. Furthermore, we have

$$\Omega_{s,q}^*(y) = \mathcal{W}_{s,q}(t - y; y) = \left\{ \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - y \right\}.$$

Proof. We take $\varphi \in C_b^2[0, \infty)$. Then for any $\psi \in (y, t)$ the Taylor series expansions lead to

$$\varphi(t) = \varphi(y) + \varphi'(y)(t - y) + \varphi''(\psi) \frac{(t - y)^2}{2}.$$

Therefore

$$\mathcal{W}_{s,q}(\varphi; y) - \varphi(y) = \varphi'(y) \mathcal{W}_{s,q}(t - y; y) + \frac{\varphi''(\psi)}{2} \mathcal{W}_{s,q}((t - y)^2; y),$$

and hence

$$|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| \leq \left\{ \Omega_{s,q}^*(y) \right\} \|\varphi'\|_{C_b[0, \infty)} + \left\{ \delta_{s,q}^*(y) \right\} \frac{\|\varphi''\|_{C_b[0, \infty)}}{2}.$$

Furthermore, by invoking Lemma 2.4, we deduce

$$\Omega_{s,q}^*(y) = \mathcal{W}_{s,q}(t - y; y) = \left\{ \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - y \right\}$$

and, by Theorem 4.2, the required value of $\delta_{s,q}^*(y)$ is obtained.

Because of (16), it is easy to conclude that $\|\varphi'\|_{C_b[0, \infty)} \leq \|\varphi\|_{C_b^2[0, \infty)}$ and $\|\varphi''\|_{C_b[0, \infty)} \leq \|\varphi\|_{C_b^2[0, \infty)}$. As a consequence we find

$$|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| \leq \left\{ \Omega_{s,q}^*(y) \right\} \|\varphi\|_{C_b^2[0, \infty)} + \left\{ \delta_{s,q}^*(y) \right\} \frac{\|\varphi\|_{C_b^2[0, \infty)}}{2},$$

where the latter inequality completes the desired proof. \square

In this section, we directly provide estimates in the form of direct theorems. For the considered type of approximation results we use Peetre's K -functional introduced by J. Peetre [42] in 1968 and defined as follows

$$K_\varphi(\varphi; \delta) = \inf \left\{ \left(\| \varphi - \phi \|_{C_b[0, \infty)} + \delta \| \phi \|_{C_b^2[0, \infty)} \right) : \phi \in C_b^2[0, \infty) \right\}. \quad (18)$$

Furthermore we consider Peetre's K -functional in relation with the modulus of continuity of order two ω_2 . More precisely there exists a positive real number M for which

$$K_\varphi(\varphi; \delta) \leq M\omega_2(\varphi; \delta^{\frac{1}{2}}), \quad \delta > 0, \quad (19)$$

where

$$\omega_2(\varphi; \delta) = \sup_{0 < \lambda < \delta} \sup_{y \in [0, \infty)} |(\varphi(y + 2\lambda) - 2\varphi(y + \lambda) + \varphi(y))|. \quad (20)$$

Moreover, from [16], we recall that there exists an absolute constant $C > 0$ satisfying

$$K_\varphi(\varphi; \delta) \leq C\{\omega_2(\varphi; \sqrt{\delta}) + \min(1, \delta) \| \varphi \| \}. \quad (21)$$

Theorem 5.4. Let $q = q_s$ be the sequences of positive numbers such that $0 < q_s < 1$. Then for any $\varphi \in C_b[0, \infty)$, there exists a positive constant C such that

$$\begin{aligned} |W_{s,q}(\varphi; y) - \varphi(y)| &\leq 2C \left\{ \varphi; \sqrt{\frac{2\Omega_{s,q}^*(y) + \delta_{s,q}^*(y)}{2}} \right. \\ &\quad \left. + \min \left(1, \frac{2\Omega_{s,q}^*(y) + \delta_{s,q}^*(y)}{2} \right) \| \varphi \|_{C_b[0, \infty)} \right\}. \end{aligned}$$

Proof. We use Theorem 5.3 by considering any $\phi \in C_b[0, \infty)$. In this way we $|W_{s,q}(\varphi - \phi; y)| \leq \| \varphi - \phi \|_{C_b}$. Therefore, it is obvious that

$$\begin{aligned} |W_{s,q}(\varphi; y) - \varphi(y)| &\leq |W_{s,q}(\varphi - \phi; y)| + |W_{s,q}(\phi; y) - \phi(y)| + |\varphi(y) - \phi(y)| \\ &\leq 2 \| \varphi - \phi \|_{C_b[0, \infty)} + (2\Omega_{s,q}^*(y) + \delta_{s,q}^*(y)) \| \phi \|_{C_b^2[0, \infty)} \\ &= 2 \left(\| \varphi - \phi \|_{C_b[0, \infty)} + \frac{2\Omega_{s,q}^*(y) + \delta_{s,q}^*(y)}{2} \| \phi \|_{C_b^2[0, \infty)} \right). \end{aligned}$$

By applying the infimum over all $\phi \in C_b^2[0, \infty)$ and by exploiting relations (18) and (19), we deduce

$$|W_{s,q}(\varphi; y) - \varphi(y)| \leq 2K_\varphi \left(\varphi; \frac{2\Omega_{s,q}^*(y) + \delta_{s,q}^*(y)}{2} \right).$$

Hence, by virtue of (21), our desired results follow and the proof is complete. \square

Theorem 5.5. For any $\phi \in C_b^2[0, \infty)$, we define an auxiliary operator $X_{s,q}$ such that

$$X_{s,q}(\phi; y) = W_{s,q}(\phi; y) + \phi(y) - \phi \left\{ W_{s,q}(t; y) \right\}. \quad (22)$$

Then for all $\psi \in C_b^2[0, \infty)$ we get the inequality

$$|X_{s,q}(\psi; y) - \psi(y)| \leq \left\{ \delta_{s,q}^*(y) + \left(W_{s,q}(t; y) - y \right)^2 \right\} \| \psi'' \|,$$

where $W_{s,q}(t; y) = \frac{|s|}{q(|s|+\mu)(|s|-1)} \left(1 + q[s]y + \frac{qR_q'(1)}{R_q(1)} \right) + \frac{\nu}{|s|+\mu}$ defined by Lemma 2.2 and $\delta_{s,q}^*(y)$ is defined by Theorem 4.2.

Proof. For all $\psi \in C_b^2[0, \infty)$, it is obvious that $X_{s,q}(1; y) = 1$ and

$$X_{s,q}(t; y) = \mathcal{W}_{s,q}(t; y) + y - \left\{ \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} \right\} = y.$$

For any $\psi \in C_b^2[0, \infty)$ the Taylor series expansions lead to

$$\psi(t) = \psi(y) + (t - y)\psi'(y) + \int_y^t (t - \theta)\psi''(\theta)d\theta.$$

Thus

$$\begin{aligned} X_{s,q}(\psi; y) - \psi(y) &= \psi'(y)X_{s,q}(t - y; y) + X_{s,q}\left(\int_y^t (t - \theta)\psi''(\theta)d\theta; y\right) \\ &= X_{s,q}\left(\int_y^t (t - \theta)\psi''(\theta)d\theta; y\right) \\ &= \mathcal{W}_{s,q}\left(\int_y^t (t - \theta)\psi''(\theta)d\theta; y\right) + \int_y^y (y - \theta)\psi''(\theta)d\theta; y \\ &\quad - \int_y^{\mathcal{W}_{s,q}(t; y)} \left\{ \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - \theta \right\} \psi''(\theta)d\theta; \\ |X_{s,q}(\psi; y) - \psi(y)| &\leq \left| \mathcal{W}_{s,q}\left(\int_y^t (t - \theta)\psi''(\theta)d\theta; y\right) \right| \\ &\quad + \left| \int_y^{\mathcal{W}_{s,q}(t; y)} \psi''(\theta)d\theta \right|. \end{aligned}$$

In view of the known inequalities

$$\left| \int_y^t (t - \theta)\psi''(\theta)d\theta \right| \leq (t - y)^2 \|\psi''\|$$

and

$$\begin{aligned} &\left| \int_y^{\mathcal{W}_{s,q}(t; y)} \left\{ \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - \theta \right\} \psi''(\theta)d\theta \right| \\ &\leq \left\{ \frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - y \right\}^2 \|\psi''\|, \end{aligned}$$

we deduce

$$\begin{aligned} |X_{s,q}(\psi; y) - \psi(y)| &\leq \left\{ \mathcal{W}_{s,q}\left((t - y)^2; x\right) \right. \\ &\quad \left. + \left(\frac{[s]}{q([s] + \mu)[s - 1]} \left(1 + q[s]y + \frac{qR'_q(1)}{R_q(1)} \right) + \frac{\nu}{[s] + \mu} - y \right)^2 \right\} \|\psi''\|. \end{aligned}$$

As a consequence the proof is complete. \square

Theorem 5.6. Let $\phi \in C_b^2[0, \infty)$. Then for every $\varphi \in C_b[0, \infty)$, the operators $\mathcal{W}_{s,q}$ satisfy the relation

$$|\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| \leq C \left[\omega_2 \left\{ \varphi; \frac{1}{2} \sqrt{\Theta_{s,q}^*(y)} \right\} + \min \left\{ 1; \frac{1}{4} \left(\Theta_{s,q}^*(y) \right) \right\} \|\varphi\|_{C_b[0, \infty)} \right]$$

$$+ \omega_x(\varphi; |\Delta_{s,q}^*(y)|),$$

where $\delta_{s,q}^*(y)$ is obtained in Theorem 4.2 and $\Theta_{s,q}^*(y) = \delta_{s,q}^*(y) + \left(\frac{[s]}{q([s]+\mu)[s-1]} \left(1 + q[s]y + \frac{qR_q'(1)}{R_q(1)} \right) + \frac{\nu}{[s]+\mu} \right)^2$ and $\Delta_{s,q}^*(y) = \frac{[s]}{q([s]+\mu)[s-1]} \left(1 + q[s]y + \frac{qR_q'(1)}{R_q(1)} \right) + \frac{\nu}{[s]+\mu} - y$.

Proof. In order to prove the claimed thesis, we use the results from Theorem 5.5. Thus, take $\phi \in C_b^2[0, \infty)$. Hence for all $\varphi \in C_b[0, \infty)$, we deduce that

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &= |X_{s,q}(\varphi; y) - \varphi(y) + \varphi(\mathcal{W}_{s,q}(t; y)) - \varphi(y)| \\ &\leq |X_{s,q}(\varphi - \phi; y)| + |X_{s,q}(\phi; y) - \phi(y)| \\ &+ |\phi(y) - \varphi(y)| + |\varphi(\mathcal{W}_{s,q}(t; y)) - \varphi(y)| \\ &\leq 4\|\varphi - \phi\| + \omega_x(\varphi; |\mathcal{W}_{s,q}(t; y) - y|) \\ &+ \left\{ \delta_{s,q}^*(y) + (\mathcal{W}_{s,q}(t; y))^2 \right\} \|\phi''\|. \end{aligned}$$

Applying the infimum for all $\phi \in C_b^2[0, \infty)$, from equality (18), we conclude that

$$\begin{aligned} |\mathcal{W}_{s,q}(\varphi; y) - \varphi(y)| &\leq 4K_\varphi \left[\varphi; \frac{1}{4} \left\{ \delta_{s,q}^*(y) + (\mathcal{W}_{s,q}(t; y))^2 \right\} \right] \\ &+ \omega_x(\varphi; |\mathcal{W}_{s,q}(t; y) - y|) \\ &\leq C \left[\omega_2 \left\{ \varphi; \frac{1}{2} \sqrt{\delta_{s,q}^*(y) + (\mathcal{W}_{s,q}(t; y))^2} \right\} \right. \\ &+ \min \left\{ 1; \frac{1}{4} \left(\delta_{s,q}^*(y) + (\mathcal{W}_{s,q}(t; y))^2 \right) \right\} \|\varphi\|_{C_b[0,\infty)} \\ &+ \left. \omega_x(\varphi; |\mathcal{W}_{s,q}(t; y) - y|) \right]. \end{aligned}$$

In this way the theorem is proven. \square

References

- [1] T. Acar, A. M. Acu, N. Manav, *Approximation of functions by genuine Bernstein-Durrmeyer type operators*, J. Math. Inequalities, 12 (4) (2018), 975–987.
- [2] T. Acar, M. Mursaleen, S.A. Mohiuddine, *Stancu type (p, q) -Szász-Mirakyan-Baskakov operators*, Communications Faculty of Sciences University of Ankara Series A1: Mathematics and Statistics, 67 (1) (2018), 116–128.
- [3] T. Acar, A. Aral, *Pointwise Convergence of q -Bernstein Operators and Their q -derivatives*, Numerical Functional Analysis and Optimization, 36 (3) (2015), 287–304.
- [4] T. Acar, *(p, q) -Generalization of Szasz-Mirakyan operators*, Mathematical Methods in the Applied Sciences, 39 (10), (2016), 2685–2695.
- [5] A.A.H.A. Al-Abeid, M. Ayman-Mursaleen, M. Mursaleen: *Szász type operators involving Charlier polynomials and approximation properties*, FILOMAT, 35(15), 2021, 5149–5159.
- [6] W. A. Al-Salam, *q -Appell polynomials*, Ann. Mat. Pura Appl., 4 (1967), 31–45.
- [7] A. Alotaibi, *On the Approximation by bivariate Szász-Jakimovski-Leviatan-type operators of unbounded sequences of positive numbers*, Mathematics 11(4) (2023), 1009.
- [8] A. Alotaibi, *Approximation of GBS type q -Jakimovski-Leviatan-Beta integral operators in Bögel space*, Mathematics 10(5) (2022), 675.

- [9] A. Alotaibi, M. Mursaleen, *Approximation of Jakimovski-Leviatan-Beta type integral operators via q-calculus*, AIMS Mathematics **5**(4) (2020), 3019–3034.
- [10] K. J. Ansari, F. Özger, Z. Ödemiş Özger: *Numerical and theoretical approximation results for Schurer-Stancu operators with shape parameter λ* , Computational and Applied Mathematics, **41**, Article number: 181 (2022).
- [11] P. Appell, *Une classe de polynômes*, Ann. Sci. École Norm. Sup., **9** (1880), 119–144.
- [12] Reşat Aslan: *Approximation properties of univariate and bivariate new class λ -Bernstein-Kantorovich operators and its associated GBS operators*, Computational and Applied Mathematics volume 42, Article number: 34, (2023).
- [13] R. Aslan, M. Mursaleen: *Some approximation results on a class of new type λ -Bernstein polynomials*, Journal of Mathematical Inequalities **16**(2) (2022), 445–462.
- [14] M. Ayman-Mursaleen, S. Serra-Capizzano: *Statistical Convergence via q-Calculus and a Korovkin's Type Approximation Theorem*, AXIOMS **11**(2) 2022, 70.
- [15] İ. Büyükyazıcı, H. Tanberkan, S. Serenbay, C. Atakut, *Approximation by Chlodowsky type Jakimovski-Leviatan operators*, J. Comput. Appl. Math., **259** (2014), 153–163.
- [16] A. Ciupa, *A class of integral Favard-Szász type operators*, Stud. Univ. Babeş-Bolyai, Math., **40** (1995), 39–47.
- [17] A.D. Gadziev: *Theorems of the type of P.P. Korovkin's theorems*. Mat. Zametki **20**(5), 781–786 (in Russian) (1976), Math. Notes. **20**(5–6), 995–998 (Engl. Trans.) (1976)
- [18] A.D. Gadjev: *The convergence problem for a sequence of positive linear operators on bounded sets and theorems analogous to that of P. P. Korovkin*. Dokl. Akad. Nauk SSSR **218**(5), (1974), Transl. Soviet Math. Dokl. **15**(5), 1433–1436 (1974)
- [19] F. H. Jackson, *On q-definite integrals*, Quart. J. Pure Appl. Math., **41** (1910), 193–203.
- [20] V. Kac, P. Cheung, *Quantum Calculus*, Universitext, Springer-Verlag, New York, 2002.
- [21] V. Kac, A. De Sole, *On integral representations of q-gamma and q-beta functions*, Mathematica, **9** (2005), 11–29.
- [22] A. Kajla, S. A. Mohiuddine, A. Alotaibi, *Blending-type approximation by Lupaş-Durrmeyer type operators involving Pólya distribution*, Math. Meth. Appl. Sci. **44** (2021) 9407–9418.
- [23] A. Kajla, S. A. Mohiuddine, A. Alotaibi, *Durrmeyer-type generalization of μ -Bernstein operators*, Filomat **36**(1) (2022) 349–360.
- [24] A. Kajla, T. Acar, *Modified α -Bernstein operators with better approximation properties*, Annals of Functional Analysis **10** (4), 2019, 570–582.
- [25] M. E. Keleşteri, N. I. Mahmudov, *A study on q-Appell polynomials from determinantal point of view*, Appl. Math. Comp., **260** (2015), 351–369.
- [26] P. P. Korovkin, *Linear Operators and Approximation Theory*, Hindustan Publ. Co., Delhi 1960.
- [27] B. Lenze: *On Lipschitz type maximal functions and their smoothness spaces*. Nederl. Akad. Indag. Math. **50**, (1988) 53–63.
- [28] V. N. Mishra, P. Patel, *On generalized integral Bernstein operators based on q-integers*, Appl. Math. Comput., **242** (2014), 931–944.
- [29] M. Mursaleen, K. J. Ansari, M. Nasiruzzaman, *Approximation by q-analogue of Jakimovski-Leviatan operators involving q-Appell polynomials*, Iranian J. Sci. Tech. A, **41** (2017), 891–900.
- [30] M. Mursaleen, Md. Nasiruzzaman, *Approximation of modified Jakimovski-Leviatan-Beta type operators*, Constr. Math. Anal., **1** (2) (2018), 88–98.
- [31] M. Mursaleen, T. Khan, *On approximation by Stancu type Jakimovski-Leviatan-Durrmeyer operators*, Azerbaijan J. Math., **7** (2017), 16–26.
- [32] S.A. Mohiuddine, T. Acar, A. Alotaibi, *Construction of a new family of Bernstein-Kantorovich operators*. Math. Meth. Appl. Sci. **40** (2017), 7749–7759.
- [33] S.A. Mohiuddine, B. Hazarika, M.A. Alghamdi, *Ideal relatively uniform convergence with Korovkin and Voronovskaya types approximation theorems*, Filomat. **33**(14) (2019), 4549–4560.
- [34] S.A. Mohiuddine, F. Özger, *Approximation of functions by Stancu variant of Bernstein-Kantorovich operators based on shape parameter α* , Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales, Serie A. RACSAM, **114** (2020), Article id: 70.
- [35] S. A. Mohiuddine, K. K. Singh, A. Alotaibi, *On the order of approximation by modified summation-integral-type operators based on two parameters*, Demonstr. Math. **56** (2023), 20220182.
- [36] M. Nasiruzzaman, A. F. Aljohani, *Approximation by Szász-Jakimovski-Leviatan type operators via aid of Appell polynomials*, J. Funct. Spaces **2020** (2020), Article ID 9657489.
- [37] M. Nasiruzzaman, A. F. Aljohani, *Approximation by parametric extension of Szász-Mirakjan-Kantorovich operators involving the Appell polynomials*, J. Funct. Spaces **2020** (2020), Article ID 9657489.
- [38] M. Nasiruzzaman, M. Mursaleen, *Approximation by Jakimovski-Leviatan-Beta operators in weighted space*, Adv. Difference Equ. **2020** (2020):393.
- [39] M.A. Ozarslan, H. Aktuğlu: *Local approximation for certain King type operators*, Filomat, **27**, 173–181 (2013)
- [40] F. Özger, E. Aljimi, M. Temizer Ersoy: *Rate of Weighted Statistical Convergence for Generalized Blending-Type Bernstein-Kantorovich Operators*, Mathematics **2022**, **10**(12), 2027.
- [41] F. Özger, H. M. Srivastava, S. A. Mohiuddine, *Approximation of functions by a new class of generalized Bernstein-Schurer operators*, Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat. RACSAM (2020) 114:173.
- [42] J. Peetre, *Noteas de mathematica 39, Rio de Janeiro, Instituto de Mathematica Pura e Applicada, Conselho Nacional de Pesquisas*, 1968.
- [43] M. Qasim, A. Khan, Z. Abbas, F. Özger P. Raina: *Approximation by generalized Baskakov-beta operators and its convergence properties*, <https://doi.org/10.1007/s41478-022-00528-5>.
- [44] M. Raiz, A. Kumar, V. N. Mishra, N. Rao: *Dunkl Analogue of Szász Schurer Beta Operators and their Approximation Behavior*, Mathematical Foundation of Computing, **5**(4), 2022, 315–330.
- [45] N. Rao, A. Wafi, A. M. Acu, *q-Szász-Durrmeyer type operators based on Dunkl analogue*, Complex Anal. Oper. Theory, **13** (2019), 915–934.
- [46] O. Shisha, B. Bond: *The degree of convergence of sequences of linear positive operators*, Proc. Nat. Acad. Sci. USA., **60**, 1196–1200 (1968)