



On the Roter type of generalised Wintgen ideal Legendrian submanifolds

Aleksandar Šebeković^a

^aDepartment of Sciences and Mathematics, State University of Novi Pazar, Vuka Karadžica 9, 36300 Novi Pazar, Serbia,
<http://orcid.org/0000-0002-3185-5197>

Abstract. I. Mihai obtained an inequality relating intrinsic normalised scalar curvature and extrinsic squared mean curvature and normalised normal curvature of Legendrian submanifolds M^n in Sasakian space forms $\widetilde{M}^{2n+1}(c)$. In this paper, for the class of generalised Wintgen ideal Legendrian submanifolds M^n of Sasakian space form $\widetilde{M}^{2n+1}(c)$, we study relationship between some properties concerning their Deszcz symmetry and their Roter type.

1. Preliminaries ([2], [3], [16], [17])

Let M^n be an n -dimensional Riemannian manifold with metric tensor g (g is positive definite $(0, 2)$ -tensor). With R we denote $(0, 4)$ -Riemann–Christoffel curvature tensor and with S the $(0, 2)$ -Ricci tensor on M^n . The Ricci tensor S is symmetric and all its eigenvalues are real and S determines an orthogonal set of eigendirections on M^n , which are the intrinsic (Ricci) principal directions on M^n .

For two $(0, 2)$ tensors t and r , we denote with \wedge Nomizu–Kulkarni product defined by

$$(t \wedge r)(X, Y, Z, W) = t(X, W)r(Y, Z) + t(Y, Z)r(X, W) - t(X, Z)r(Y, W) - t(Y, W)r(X, Z),$$

whereby X, Y, Z, W are tangent vector fields on M^n . Now, for plane π spanned by tangent vector fields X and Y , the sectional curvature $K(\pi)$ is given by

$$K(\pi) = \frac{R(X, Y, Y, X)}{\frac{1}{2}(g \wedge g)(X, Y, Y, X)}.$$

Riemannian manifold with constant sectional curvature c , $K = c$, is called a real space form of curvature c , denoted with $M^n(c)$.

If Ricci tensor S is proportional to metric tensor g on manifold M^n ($S = \lambda g$, λ is some function), we say that M^n is an Einstein space and every 3-dimensional Einstein space has constant sectional curvature. If the Ricci tensor S has an eigenvalue of multiplicity $\geq n - 1$ on M^n , we say that M^n is quasi-Einstein.

2020 Mathematics Subject Classification. 53B20, 53B25, 53A07, 53C42.

Keywords. Generalised Wintgen ideal Legendrian submanifolds; Deszcz symmetry; Roter space.

Received: 12 December 2022; Accepted: 14 February 2023

Communicated by Zoran Rakić and Mića Stanković

Email address: asebekovic@np.ac.rs (Aleksandar Šebeković)

The Weyl conformal curvature tensor C is defined by

$$C = R - \left(\frac{1}{n-2} g \wedge S + \frac{\tau}{(n-1)(n-2)} g \wedge g \right).$$

For manifold M^n , $n = 3$, it is known that $C \equiv 0$. If $C \equiv 0$ for $n \geq 4$, we say that M^n is conformally flat.

A Riemannian manifold M^n ($n \geq 3$) is called a Roter space when its Riemann–Christofel curvature tensor R satisfies the equality

$$R = \tilde{\lambda} (g \wedge g) + \tilde{\mu} (g \wedge S) + \tilde{\nu} (S \wedge S), \tag{1}$$

for some functions $\tilde{\lambda}, \tilde{\mu}, \tilde{\nu} : M^n \rightarrow \mathbb{R}$, [9]. The Roter spaces, from an algebraic point of view, may be considered as the simplest Riemannian manifolds, which are the next to the real space forms [5].

It is obvious that the real space forms $M^n(c)$ are Roter spaces for which $\tilde{\lambda} = \frac{c}{2}$ and $\tilde{\mu} = \tilde{\nu} = 0$. Einstein Roter spaces are real space forms. Also, all 3-dimensional Riemannian manifolds and all conformally flat Riemannian manifolds M^n ($n \geq 4$) are Roter spaces for which

$$\tilde{\lambda} = \frac{\tau}{2(n-1)(n-2)}, \quad \tilde{\mu} = \frac{1}{n-2}, \quad \tilde{\nu} = 0.$$

The Deszcz symmetric spaces, from a geometric point of view, may be considered to be the simplest Riemannian manifolds next to the real space form [5]. The Riemannian spaces M^n ($n \geq 3$) are Deszcz symmetric if $(0, 6)$ -tensors $R \circ R$ and $Q(g, R)$ are proportional, i.e.,

$$R \circ R = L Q(g, R), \tag{2}$$

for some function $L : M^n \rightarrow \mathbb{R}$, [8]. $(0, 6)$ -tensor $R \circ R$ is defined by

$$\begin{aligned} (R \circ R)(X_1, X_2, X_3, X_4; X, Y) &= (R(X, Y) \circ R)(X_1, X_2, X_3, X_4) \\ &= -R(R(X, Y)X_1, X_2, X_3, X_4) - R(X_1, R(X, Y)X_2, X_3, X_4) \\ &\quad - R(X_1, X_2, R(X, Y)X_3, X_4) - R(X_1, X_2, X_3, R(X, Y)X_4), \end{aligned}$$

and Tachibana $(0, 6)$ -tensor $Q(g, R)$ is given by

$$\begin{aligned} Q(g, R)(X_1, X_2, X_3, X_4; X, Y) &= ((X \wedge_g Y) \circ R)(X_1, X_2, X_3, X_4) \\ &= -R((X \wedge_g Y)X_1, X_2, X_3, X_4) - R(X_1, (X \wedge_g Y)X_2, X_3, X_4) \\ &\quad - R(X_1, X_2, (X \wedge_g Y)X_3, X_4) - R(X_1, X_2, X_3, (X \wedge_g Y)X_4), \end{aligned}$$

where \wedge_g is metric endomorphisam defined by

$$(X \wedge_g Y)Z = g(Y, Z)X - g(X, Z)Y.$$

For Riemannian manifold M^n ($n \geq 3$) we say that it is Ricci pseudo-symmetric if

$$(R \circ S)(p) = L_s(p) Q(g, S)(p), \quad \forall p \in M^n,$$

where $R \circ S$ and $Q(g, S)$ are $(0, 4)$ -tensors given by

$$\begin{aligned} (R \circ S)(X, Y, Z, W) &= -S(R(Z, W)X, Y) - S(X, R(Z, W)Y), \\ Q(g, S) &= -S((Z \wedge_g W)X, Y) - S(X, (Z \wedge_g W)Y), \end{aligned}$$

for $X, Y, Z, W \in TM^n$.

Similarly, Riemannian manifold M^n ($n \geq 4$) has pseudo-symmetric Weyl tensor C if

$$C \circ C = L_C Q(g, C),$$

for some functions $L_C : M^n \rightarrow \mathbb{R}$ (on the open part of M^n where $Q(g, C) \neq 0$) for $(0, 6)$ -tensors $C \circ C$ and $Q(g, C)$.

It is known ([6], [11], [10], [13], [14]) that:

- a) the open submanifold \mathcal{U} of a Riemannian manifold M^n of Roter type is Deszcz symmetric and has pseudo-symmetric Weyl conformal tensor C ;
- b) the open submanifold \mathcal{U} of a Deszcz symmetric space with pseudo-symmetric Weyl tensor C is a space of Roter type.

Let M^n be n -dimensional Riemannian submanifold in $(m + n)$ -dimensional real space form $\widetilde{M}^{n+m}(c)$, and let g, ∇ and $\widetilde{g}, \widetilde{\nabla}$ be the metric (Riemannian) and the corresponding Levi-Civita connection on M^n and $\widetilde{M}^{n+m}(c)$, respectively. Let X, Y, Z, \dots be the tangent vector fields on M^n and ξ, η, \dots be the normal vector fields on $\widetilde{M}^{n+m}(c)$. Then we have the formula of Gauss ($\widetilde{\nabla}_X Y = \nabla_X Y + h(X, Y)$) and Weingarten ($\widetilde{\nabla}_X \xi = -A_\xi(X) + \nabla_X^\perp \xi$) which decompose the vector fields $\widetilde{\nabla}_X Y$ and $\widetilde{\nabla}_X \xi$ on their tangential ($\nabla_X Y$ and $A_\xi(X)$) and normal ($h(X, Y)$ and $\nabla_X^\perp \xi$) components along M^n in $\widetilde{M}^{n+m}(c)$, respectively. With h and A_ξ we denote the second fundamental form and the shape operator of M^n with respect to ξ (normal vector field), such that

$$g(h(X, Y), \xi) = g(A_\xi(X), Y).$$

∇^\perp denote the connection in the normal bundle.

Let $\{E_1, E_2, \dots, E_n, \xi_1, \dots, \xi_m\}$ be any local orthonormal frame field on M^n in $\widetilde{M}^{n+m}(c)$. Then the mean curvature vector field of M^n in $\widetilde{M}^{n+m}(c)$ is defined by

$$\vec{H} = \frac{1}{n} \operatorname{tr} h = \frac{1}{n} \sum_{i=1}^n h(E_i, E_i) = \frac{1}{n} \sum_{\alpha=1}^m (\operatorname{tr} A_\alpha) \xi_\alpha.$$

For submanifold M^n in $\widetilde{M}^{n+m}(c)$ we say that it is:

- (i) totally geodesic when $h = 0$,
- (ii) totally umbilical when $h = g\vec{H}$,
- (iii) minimal when $\vec{H} = 0$,
- (iv) pseudo-umbilical when $A_{\vec{H}} = \lambda I_d$ (where I_d denote identity operator on TM and λ is some real function on M^n).

The normalised scalar curvature of M^n is given by

$$\rho = \frac{2}{n(n-1)} \sum_{i < j}^n R(E_i, E_j, E_j, E_i),$$

where

$$R(X, Y, Z, W) = \widetilde{g}(h(Y, Z), h(X, W)) - \widetilde{g}(h(X, Z), h(Y, W)) + c(g(Y, Z)g(X, W) - g(X, Z)g(Y, W))$$

is Riemann-Christoffel curvature tensor of M^n in $\widetilde{M}^{n+m}(c)$.

The normalised normal scalar curvature function of M^n at a point p is given by

$$\rho^\perp(p) = \frac{2}{n(n-1)} \sqrt{\sum_{i < j}^n \sum_{\alpha < \beta}^m R^\perp(E_i, E_j, \xi_\alpha, \xi_\beta)^2},$$

where R^\perp is the curvature tensor of normal space and $\{\xi_1, \dots, \xi_m\}$ is an orthonormal frame field of that space, $R^\perp(X, Y; \xi, \eta) = g([A_\xi, A_\eta]X, Y)$, whereby $[A_\xi, A_\eta] = A_\xi A_\eta - A_\eta A_\xi$.

We further will be concerned with Wintgen ideal submanifolds. The original inequality is obtained for surfaces M^2 in \mathbb{E}^4 by Wintgen in 1979. He proved that, intrinsic invariant of M^2 , Gauss curvature K and extrinsic invariants, the squared of mean curvature H^2 and normal curvature K^\perp , satisfy the inequality $K \leq H^2 - K^\perp$, and also characterised equality case [21]. After that, Rouxel [19], Rodriguez-Guadalupe [15],

De Smet, Dillen, Verstraelen and Vrancken [7] gave some generalizations of this results. And, finally, Choi and Lu [4] and Ge–Tang [12] proved that for submanifolds M^n in a real space form $\widetilde{M}^{n+m}(c)$ holds inequality

$$\rho \leq H^2 - \rho^\perp + c. \tag{3}$$

They also proved that equality in inequality (3) holds if the shape operators of the submanifold take the special forms for suitable adapted orthonormal frame $\{E_1, \dots, E_n, \xi_1, \dots, \xi_m\}$ on M^n in $\widetilde{M}^{n+m}(c)$.

The submanifolds M^n in $\widetilde{M}^{n+m}(c)$ for which hold equality in inequality (3) are called Wintgen ideal submanifolds. In [5] the authors studied Wintgen ideal submanifolds M^n ($n \geq 4$) in real space forms $\widetilde{M}^{n+m}(c)$ which are Roter spaces and proved that such submanifold is Deszcz symmetric if and only if it is Roter space.

2. Generalised Wintgen inequality for Legendrian submanifolds

A $(2m + 1)$ -dimensional Riemannian manifold $(\widetilde{M}^{2m+1}(c), g)$ is Sasakian manifold if the triple (ϕ, ξ, η) (ϕ is an endomorphism of tangent bundle of $T\widetilde{M}^{2m+1}(c)$; η is 1-form and ξ is vector field called characteristic vector field) satisfy:

$$\begin{aligned} \phi^2 &= -I_d + \eta \otimes \xi, \quad \eta(\xi) = -1, \quad \phi \xi = 0, \quad \eta \circ \phi = 0, \\ g(\phi X, \phi Y) &= g(X, Y) - \eta(X)\eta(Y), \quad \eta(X) = g(X, \xi), \\ (\widetilde{\nabla}_X \phi)Y &= -g(X, Y)\xi + \eta(Y)X, \quad \widetilde{\nabla}_X \xi = \phi X, \end{aligned}$$

where X and Y are vector fields on $\widetilde{M}^{2m+1}(c)$ and $\widetilde{\nabla}$ denotes the Riemannian connection with respect to g . If a plane π is spanned by X and ϕX , then a plane section π in $T_p \widetilde{M}^{2m+1}$ is called a ϕ -section, where X is a unit tangent vector which is orthogonal to ξ . The sectional curvature of a ϕ -section is called ϕ -sectional curvature and a Sasakian manifold with constant ϕ -sectional curvature c is called a Sasakian space form $\widetilde{M}^{2m+1}(c)$. On a Sasakian space form $\widetilde{M}^{2m+1}(c)$ the curvature tensor \widetilde{R} is given by, [1]

$$\begin{aligned} \widetilde{R}(X, Y)Z &= \frac{c+3}{4}\{g(Y, Z)X - g(X, Z)Y\} + \frac{c-1}{4}\{\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + \\ &+ g(X, Z)\eta(Y)\xi - g(Y, Z)\eta(X)\xi + g(\phi Y, Z)\phi X - g(\phi X, Z)\phi Y - 2g(\phi X, Y)\phi Z\}, \end{aligned}$$

for the tangent vector fields X, Y, Z on $\widetilde{M}^{2m+1}(c)$.

Let M^n be an n -dimensional submanifold of a Sasakian space form $\widetilde{M}^{2m+1}(c)$. Then the Gauss equation is given by

$$\widetilde{R}(X, Y, Z, W) = R(X, Y, Z, W) + g(h(X, W), h(Y, Z)) - g(h(X, Z), h(Y, W)),$$

whereby R and h are the Riemann curvature tensor and second fundamental form, respectively, of M^n , and X, Y, Z, W are vectors tangent to M^n . For every $p \in M^n$, the mean curvature is given by

$$H(p) = \frac{1}{n} \sum_{i=1}^n h(E_i, E_i),$$

where $\{E_1, E_2, \dots, E_n, \dots, E_{2m+1}\}$ is an orthonormal basis of $T_p \widetilde{M}^{2m+1}$.

C -totally real submanifold is a submanifold M^n normal to ξ in a Sasakian manifold, i.e. $\phi(T_p M^n) \subset T_p^\perp M^n$, for every $p \in M^n$. If $m \equiv n$, then M^n is called Legendrian submanifold.

Let M^n be an n -dimensional Legendrian submanifold of a Sasakian space form $\widetilde{M}^{2n+1}(c)$ and $\{E_1, \dots, E_n\}$ an orthonormal frame on M^n and $\{E_{n+1}, \dots, E_{2n}, E_{2n+1} = \xi\}$ an orthonormal frame in the normal bundle $T^\perp M^n$. The Gauss equation is given by

$$\begin{aligned} R(X, Y, Z, W) &= \frac{c+3}{4}\{g(X, Z)g(Y, W) - g(Y, Z)g(X, W)\} + \\ &+ g(h(X, Z), h(Y, W)) - g(h(X, W), h(Y, Z)). \end{aligned}$$

I. Mihai in [18] established a generalised Wintgen inequality for Legendrian submanifolds in Sasakian space forms.

Theorem 2.1 ([18]). *Let M^n be an n -dimensional Legendrian submanifold of a Sasakian space form $\widetilde{M}^{2n+1}(c)$. Then*

$$(\rho^\perp)^2 \leq \left(\|H\|^2 - \rho + \frac{c+3}{4} \right)^2 + \frac{4}{n(n-1)} \left(\rho - \frac{c+3}{4} \right) \frac{c-1}{4} + \frac{(c-1)^2}{8n(n-1)}, \tag{4}$$

and equality holds if and only if with respect to suitable orthonormal frames $\{E_1, \dots, E_n\}$ and $\{E_{n+1}, \dots, E_{2n}, E_{2n+1} = \xi\}$, the shape operators of M^n in $\widetilde{M}^{2n+1}(c)$ are given by:

$$A_{E_{n+1}} = \begin{bmatrix} \lambda_1 & \mu & 0 & \cdots & 0 \\ \mu & \lambda_1 & 0 & \cdots & 0 \\ 0 & 0 & \lambda_1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_1 \end{bmatrix}, \quad A_{E_{n+2}} = \begin{bmatrix} \lambda_2 + \mu & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 - \mu & 0 & \cdots & 0 \\ 0 & 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_2 \end{bmatrix},$$

$$A_{E_{n+3}} = \begin{bmatrix} \lambda_3 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_3 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_3 \end{bmatrix}, \quad A_{E_{n+4}} = \cdots = A_{E_{2n}} = A_{E_{2n+1}} = 0,$$

whereby $\lambda_1, \lambda_2, \lambda_3$ and μ are real functions on M^n .

Legendrian submanifolds M^n in a Sasakian space forms $\widetilde{M}^{2n+1}(c)$ satisfying equality in generalised Wintgen inequality (4) are called generalized Wintgen ideal Legendrian submanifolds. A frame $\{E_1, E_2, \dots, E_n, E_{n+1}, \dots, E_{2n+1}\}$ from Theorem 2.1 is called Choi–Lu frame on such M^n in $\widetilde{M}^{2n+1}(c)$.

3. Main result

From Theorem 2.1, using Gauss equation, we obtain, [20], that all components of $(0, 4)$ curvature tensor R of generalised Wintgen ideal Legendrian submanifold M^n of a Sasakian space form $\widetilde{M}^{2n+1}(c)$ are zero, except these:

$$R_{1221} = 2\mu^2 - c_1, \quad R_{1kk1} = -\lambda_2\mu - c_1, \quad k \geq 3, \quad R_{2kk2} = \lambda_2\mu - c_1, \quad k \geq 3,$$

$$R_{1kk2} = -\lambda_1\mu, \quad k \geq 3, \quad R_{kllk} = -c_1, \quad k \neq l, \quad k, l \geq 3,$$

whereby $c_1 = \frac{c+3}{4} + \lambda_1^2 + \lambda_2^2 + \lambda_3^2$.

The nontrivial components of $(0, 2)$ -Ricci tensor S of such submanifold are, [20]:

$$S_{11} = 2\mu^2 - (n-1)c_1 - (n-2)\lambda_2\mu,$$

$$S_{22} = 2\mu^2 - (n-1)c_1 + (n-2)\lambda_2\mu,$$

$$S_{12} = -(n-2)\lambda_1\mu,$$

$$S_{kk} = -(n-1)c_1, \quad k \geq 3.$$

The equation (1) in local components looks like:

$$R_{ijkl} = \widetilde{\lambda}(g_{il}g_{jk} - g_{ik}g_{jl}) + \widetilde{\mu}(g_{il}S_{jk} + g_{jk}S_{il} - g_{ik}S_{jl} - g_{jl}S_{ik}) + \widetilde{\nu}(S_{il}S_{jk} - S_{ik}S_{jl}). \tag{5}$$

The condition (5) for M^n to be a Roter space is equivalent to the following system of linear equations:

$$\left. \begin{aligned} 2\mu^2 - c_1 &= \tilde{\lambda} - 2(n-1)c_1\tilde{\mu} + ((2\mu^2 - (n-1)c_1)^2 - (n-2)^2\mu^2(\lambda_1^2 + \lambda_2^2))\tilde{v}, \\ -\lambda_2\mu - c_1 &= \tilde{\lambda} + (2\mu^2 - 2(n-1)c_1 - (n-2)\lambda_2\mu)\tilde{\mu} + \\ &\quad + (n-1)c_1(-2\mu^2 + (n-1)c_1 + (n-2)\lambda_2\mu)\tilde{v}, \\ \lambda_1\mu &= -(n-2)\lambda_1\mu\tilde{\mu} + (n-1)(n-2)\lambda_1\mu c_1\tilde{v}, \\ \lambda_2\mu - c_1 &= \tilde{\lambda} + (2\mu^2 - 2(n-1)c_1 + (n-2)\lambda_2\mu)\tilde{\mu} + \\ &\quad + (n-1)c_1(-2\mu^2 + (n-1)c_1 - (n-2)\lambda_2\mu)\tilde{v}, \\ -c_1 &= \tilde{\lambda} - (n-1)c_1\tilde{\mu} + (n-1)^2c_1^2\tilde{v}. \end{aligned} \right\} \quad (6)$$

For the Deszcz symmetric generalised Wintgen ideal Legendrian submanifold M^n in a Sasakian space form $\tilde{M}^{2n+1}(c)$ the system (6) of linear equations is valid if and only if

- (i) $\mu = 0$ or
- (ii) $\mu \neq 0$ and $\lambda_1 = \lambda_2 = 0$.

In case (i), we have that M^n is itself a space form and hence a Roter space. In case (ii) from system (6), we obtain

$$\tilde{\lambda} = \left(\frac{c+3}{4} + \lambda_3^2\right)\left(2\mu^2 - (n-1)^2\left(\frac{c+3}{4} + \lambda_3^2\right)\right), \quad \tilde{\mu} = \frac{(n-1)\left(\frac{c+3}{4} + \lambda_3^2\right)}{2\mu^2}, \quad \tilde{v} = -\frac{1}{4\mu^2},$$

as its unique solution. We thus obtained the following result:

Theorem 3.1. *Let M^n be a generalised Wintgen ideal Legendrian submanifold of a Sasakian space form $\tilde{M}^{2n+1}(c)$, $n \geq 4$. Then M^n is Deszcz symmetric if and only if it is a Roter space.*

References

- [1] D. Blair, Riemannian Geometry of Contact and Symplectic Manifolds, Birkhäuser, Boston, 2002.
- [2] B. Y. Chen, Geometry of Submanifolds, Marcel Dekker, New York, 1973.
- [3] B. Y. Chen, Pseudo-Riemannian Geometry, δ -invariants and Applications, World Scientific, Publ. Co, Singapore, 2011.
- [4] T. Choi, Z. Lu, On the DDVV conjecture and the comass in calibrated geometry I, Math. Z. 260 (2008) 409–429.
- [5] S. Decu, M. Petrović-Torgašev, A. Šebeković, L. Verstraelen, On the Roter Type of Wintgen Ideal Submanifolds, Rev. Roumanie Math. Pures. Appl. 57 (2012) 75–90.
- [6] F. Defever, R. Deszcz, On Semi-Riemannian Manifolds Satisfying the Condition $R \circ R = Q(S, R)$ (in Geometry and Topology of Submanifolds, Vol. III, L. Verstraelen, A. West), World Scientific Publ. Co., Singapore, pp. 108–130, 1991.
- [7] P. J. De Smet, F. Dillen, L. Verstraelen, L. Vrancken, A Pointwise Inequality in Submanifold Theory, Arc. Math. (Brno) 35 (1999) 115–128.
- [8] R. Deszcz, On Pseudosymmetric Spaces, Bull. Soc. Math. Belg. Serie A 44 (1992), 1–34.
- [9] R. Deszcz, On Some Akivis–Goldberg Type Metrics, Publ. Inst. Math. (Beograd) (N. S.) 74(88) (2003), 71–83.
- [10] R. Deszcz, M. Hotloš, On Hypersurfaces with Type Number Two in Spaces of Constant Curvature, Ann. Univ. Budap. Rolando Eötvös, Sect. Math. 46 (2003) 19–34.
- [11] R. Deszcz, Ş Yaprak, Curvature Properties of Certain Pseudosymmetric Manifolds, Publ. Math. Debrecin 45 (1994) 333–345.
- [12] J. Ge, Z. Tang, A Proof of the DDVV Conjecture and its Equality Case, Pacific J. Math 237 (2008) 87–95.
- [13] M. Glogowska, Curvature Conditions on Hypersurfaces with Two Distinct Principal Curvatures (in PDEs, Submanifolds and Affine Differential Geometry), Banach Center Publications, Polish Academy of Sciences, Institute of Mathematics, Vol. 69, Warszawa, eds. B. Opozda, U. Simon, M. Wiehe, pp. 133–143, 2005.
- [14] M. Glogowska, On Roter Type Manifolds (in Pure and Applied Differential Geometry, F. Dillen, I. Van de Woestijne), Shaker Verlag, Aachen, pp. 114–122, 2007.
- [15] I. V. Guadalupe, L. Rodriguez, Normal Curvature of Surfaces in Space Forms, Pacific J. Math. 106 (1983) 95–103.
- [16] S. Kobayashi, K. Nomizu, Foundations of Differential Geometry I, John Wiley and Sons, New York, 1963.
- [17] S. Kobayashi, K. Nomizu, Foundations of Differential Geometry II, John Wiley and Sons, New York, 1969.
- [18] I. Mihai, On the Generalized Wintgen Inequality for Legendrian Submanifolds in Sasakian Space Forms, Tahoku Math. J. 69 (2017) 45–53.
- [19] B. Rouxel, Sur quelques propriétés anallagmatiques de l’espaces euclidien \mathbb{E}^4 , Mémoire couronné par l’Académie Royale de Belgique, 1982.
- [20] A. Šebeković, M. Petrović-Torgašev, A. Pantić, Pseudosymmetry Properties of Generalised Wintgen Ideal Legendrian Submanifolds, Filomat 33/4 (2019) 1209–1215.
- [21] P. Wintgen, Sur l’inégalité de Chen–Willmore, C. R. Bend. Sci 288 (1979) 993–995.