



2-normal composition operators with linear fractional symbols on H^2

Rong Yang^a, Songxiao Li^{b,*}, Lian Hu^a

^a*Institute of Fundamental and Frontier Sciences, University of Electronic Science and Technology of China,
610054, Chengdu, Sichuan, P.R. China*

^b*Department of Mathematics, Shantou University, 515063, Shantou, Guangdong, P.R. China*

Abstract. In this paper, some sufficient and necessary conditions for the composition operator C_φ to be 2-normal are investigated when the symbol φ is a linear fractional self-map of \mathbb{D} .

1. Introduction

Let H be a complex Hilbert space, $B(H)$ be the space of all bounded linear operators defined in H . An operator $T \in B(H)$ is called normal if it satisfies the condition $[T, T^*] = 0$, where $[T, T^*] = TT^* - T^*T$. An operator $T \in B(H)$ is subnormal if there is a Hilbert space K containing H and a normal operator M on K such that $MH \subset H$ and $T = M|_H$. An operator T is called quasinormal if $[T, T^*T] = 0$. An operator $T \in B(H)$ is called p -hyponormal if $(T^*T)^p \geq (TT^*)^p$, where $0 < p \leq 1$. If $p = 1$, T is said to be hyponormal. An operator $T \in B(H)$ is called binormal when $[T^*T, TT^*] = 0$. An operator T is said to belong to Θ class if $[T^*T, T + T^*] = 0$. From [1, 9], we see that

quasinormal \subset binormal

normal \subset quasinormal \subset subnormal \subset hyponormal.

The operator T is said to be n -normal if T^* commutes with T^n , that is $[T^*, T^n] = 0$. When $n = 2$, the operator T is called 2-normal, that is, $[T^*, T^2] = 0$. It is clear that a normal operator is a 2-normal operator, but the converse is not true.

Let \mathbb{D} denote the open unit disk in the complex plane \mathbb{C} . Let $H(\mathbb{D})$ be the space of those analytic functions on \mathbb{D} . The Hardy space $H^2(\mathbb{D})$ is the space of all $f \in H(\mathbb{D})$ for which

$$\|f\|_{H^2(\mathbb{D})}^2 = \sum_{n=0}^{\infty} |a_n|^2 < \infty,$$

where $\{a_n\}$ is the sequence of Maclaurin coefficients for f . The space H^2 is a reproducing kernel Hilbert space. In other word, for any $w \in \mathbb{D}$ and $f \in H^2$, there exists a unique function $K_w \in H^2$ such that

$$f(w) = \langle f, K_w \rangle.$$

2020 *Mathematics Subject Classification.* Primary 30H10; Secondary 47B38.

Keywords. Hardy space; Composition operator; Normal.

Received: 10 April 2023; Accepted: 06 June 2023

Communicated by Dragan S. Djordjević

Research supported by Guangdong Basic and Applied Basic Research Foundation (no. 2022A1515010317, no. 2023A1515010614).

* Corresponding author: Songxiao Li

Email addresses: yangrong071428@163.com (Rong Yang), jyu1sx@163.com (Songxiao Li), hl152808@163.com (Lian Hu)

It is well known that $K_w(z) = \frac{1}{1-\bar{w}z}$.

Let φ be an analytic self-map of \mathbb{D} . The composition operator C_φ with symbol φ is defined by

$$C_\varphi f = f \circ \varphi.$$

It is easy to see that $C_\varphi^* K_\alpha(z) = K_{\varphi(\alpha)}(z)$ for any $\alpha \in \mathbb{D}$. For $f \in L^\infty(\partial\mathbb{D})$ and $g \in H^2$, the Toeplitz operator T_f on H^2 is defined by $T_f(g) = P(fg)$, where P denotes the orthogonal projection of L^2 onto H^2 . It is easy to check that

$$T_f^* K_\alpha = \overline{f(\alpha)} K_\alpha$$

for any $\alpha \in \mathbb{D}$ and $f \in H^\infty(\mathbb{D})$, the bounded analytic function space in \mathbb{D} .

H. Schwarz [11] showed that C_φ is normal if and only if $\varphi(z) = az$ with $|a| \leq 1$. S. Jung, Y. Kim and E. Ko [10] proved that C_φ is quasinormal if and only if C_φ is normal, where $\varphi(z) = \frac{az+b}{cz+d}$ is a linear fractional self-map of \mathbb{D} with $\varphi(0) = 0$. Also they proved that, when $\varphi(z) = \frac{z}{uz+v}$ with $u \neq 0$ and $|v| \geq 1 + |u|$, C_φ is binormal if and only if C_φ is hyponormal, or C_φ is subnormal. Fatehi, Shaabani and Thompson studied hyponormal and quasinormal weighted composition operators on H^2 and the weighted Bergman space A_α^2 in [8]. For more study on composition operators on H^2 , see [2–11].

In this paper, we discuss 2-normal composition operators with linear fractional symbols on H^2 . The necessary and sufficient conditions for C_φ to be 2-normal are given when $\varphi(z) = \frac{az+b}{cz+d}$ is a linear fractional self-map of \mathbb{D} . In particular, when $\varphi(z) = \frac{az+b}{cz+d}$ is a linear fractional self-map of \mathbb{D} , we prove that C_φ is 2-normal if and only if C_φ is normal when $b = 0$ or $c = 0$. We also give an example of a linear fractional self-map φ which induces a 2-normal operator C_φ but not a normal operator.

2. Auxiliary results

In this section, we state some lemmas which will be used in this paper.

Lemma 1. [11] Let φ be an analytic self-map of \mathbb{D} . Then C_φ is normal if and only if $\varphi(z) = az$ with $|a| \leq 1$.

Lemma 2. [3, Theorem 2] Let $\varphi(z) = \frac{az+b}{cz+d}$ be a linear fractional transformation mapping \mathbb{D} into itself, where $ad - bc \neq 0$. Then $\sigma(z) = \frac{\bar{a}z - \bar{c}}{-bz + d}$ maps \mathbb{D} into itself, $g(z) = \frac{1}{-bz + d}$ and $h(z) = cz + d$ are in H^∞ , and

$$C_\varphi^* = T_g C_\sigma T_h^*.$$

The map σ is called the Krein adjoint of φ . g and h are called the Cowen auxiliary functions.

Lemma 3. Let $\varphi(z) = \frac{az+b}{cz+d}$ be a linear fractional self-map of \mathbb{D} . Then

$$\begin{aligned} C_\varphi^* C_\varphi C_\varphi K_\alpha(z) &= \frac{c(a+d)}{c(a+d) - (a^2 + bc)\bar{\alpha}} K_{\varphi(0)}(z) \\ &\quad + \left(\frac{bc + d^2}{bc + d^2 - b(a+d)\bar{\alpha}} - \frac{c(a+d)}{c(a+d) - (a^2 + bc)\bar{\alpha}} \right) K_{\varphi(\sigma_2(\alpha))}(z) \end{aligned}$$

for any $\alpha \in \mathbb{D}$ with $c(a+d) \neq (a^2 + bc)\bar{\alpha}$, where $\sigma(z) = \frac{\bar{a}z - \bar{c}}{-bz + d}$.

Proof. Let $\alpha \in \mathbb{D}$ with $c(a+d) \neq (a^2 + bc)\bar{\alpha}$. Then

$$\begin{aligned} C_\varphi^* C_\varphi C_\varphi K_\alpha(z) &= C_\varphi^* K_\alpha(\varphi_2(z)) = C_\varphi^* \frac{1}{1 - \bar{\alpha}\varphi_2(z)} \\ &= C_\varphi^* \frac{1}{1 - \bar{\alpha} \frac{(a^2 + bc)z + ab + bd}{(ac + cd)z + bc + d^2}} = C_\varphi^* \frac{Az + B}{Cz + D}, \end{aligned}$$

where

$$\begin{aligned} A &= ac + cd, & B &= bc + d^2, \\ C &= c(a + d) - \bar{\alpha}(a^2 + bc), & D &= (bc + d^2) - b(a + d)\bar{\alpha}. \end{aligned}$$

From [10, Lemma 2.3] and $C \neq 0$, we have

$$C_\varphi^* \frac{Az + B}{Cz + D} = \frac{A}{C} K_{\varphi(0)}(z) + \left(\frac{B}{D} - \frac{A}{C} \right) K_{\varphi(-\frac{\bar{C}}{D})}(z).$$

Since $\frac{A}{C} = \frac{c(a+d)}{c(a+d) - \bar{\alpha}(a^2+bc)}$,

$$\frac{B}{D} - \frac{A}{C} = \frac{bc + d^2}{bc + d^2 - b(a + d)\bar{\alpha}} - \frac{c(a + d)}{c(a + d) - (a^2 + bc)\bar{\alpha}},$$

and

$$-\frac{\bar{C}}{D} = \frac{\alpha(\bar{a}^2 + \bar{b}c) - (\bar{a}c + \bar{c}d)}{(\bar{b}c + \bar{d}^2) - \alpha(\bar{a}b + \bar{b}d)} = \sigma_2(\alpha),$$

we obtain

$$\begin{aligned} C_\varphi^* C_\varphi C_\varphi K_\alpha(z) &= \frac{c(a + d)}{c(a + d) - (a^2 + bc)\bar{\alpha}} K_{\varphi(0)}(z) \\ &+ \left(\frac{bc + d^2}{bc + d^2 - b(a + d)\bar{\alpha}} - \frac{c(a + d)}{c(a + d) - (a^2 + bc)\bar{\alpha}} \right) K_{\varphi(\sigma_2(\alpha))}(z). \end{aligned}$$

□

Lemma 4. Let $\varphi(z) = \frac{az+b}{cz+d}$ be a linear fractional self-map of \mathbb{D} . Then

$$\begin{aligned} C_\varphi C_\varphi C_\varphi^* K_\alpha(z) &= \frac{c(a + d)}{c(a + d) - (a^2 + bc)\overline{\varphi(\alpha)}} \\ &+ \left(\frac{bc + d^2}{bc + d^2 - b(a + d)\overline{\varphi(\alpha)}} - \frac{c(a + d)}{c(a + d) - (a^2 + bc)\overline{\varphi(\alpha)}} \right) K_{\sigma_2(\varphi(\alpha))}(z) \\ &= \frac{(\bar{c}\bar{\alpha} + \bar{d})[(ac + cd)z + bc + d^2]}{(\bar{c}\bar{\alpha} + \bar{d})[(ac + cd)z + bc + d^2] - (\bar{a}\bar{\alpha} + \bar{b})[(a^2 + bc)z + ab + bd]} \end{aligned}$$

for any $\alpha \in \mathbb{D}$ with $c(a + d) \neq (a^2 + bc)\overline{\varphi(\alpha)}$, where $\sigma(z) = \frac{\bar{a}z - \bar{c}}{-bz + \bar{d}}$.

Proof. For any $\alpha \in \mathbb{D}$ with $c(a + d) \neq (a^2 + bc)\overline{\varphi(\alpha)}$, according to the proof of [10, Lemma 2.3], we have that

$$\begin{aligned} C_\varphi C_\varphi C_\varphi^* K_\alpha(z) &= \frac{c(a + d)}{c(a + d) - (a^2 + bc)\overline{\varphi(\alpha)}} \\ &+ \left(\frac{bc + d^2}{bc + d^2 - b(a + d)\overline{\varphi(\alpha)}} - \frac{c(a + d)}{c(a + d) - (a^2 + bc)\overline{\varphi(\alpha)}} \right) K_{\sigma_2(\varphi(\alpha))}(z). \end{aligned}$$

On the other hand,

$$\begin{aligned} C_\varphi C_\varphi C_\varphi^* K_\alpha(z) &= C_\varphi C_\varphi K_{\varphi(\alpha)}(z) = K_{\varphi(\alpha)}(\varphi_2(z)) = \frac{1}{1 - \overline{\varphi(\alpha)}\varphi_2(z)} \\ &= \frac{1}{1 - \frac{\bar{a}\bar{\alpha} + \bar{b}}{\bar{c}\bar{\alpha} + \bar{d}} \frac{(a^2 + bc)z + ab + bd}{(ac + cd)z + bc + d^2}} \\ &= \frac{(\bar{c}\bar{\alpha} + \bar{d})[(ac + cd)z + bc + d^2]}{(\bar{c}\bar{\alpha} + \bar{d})[(ac + cd)z + bc + d^2] - (\bar{a}\bar{\alpha} + \bar{b})[(a^2 + bc)z + ab + bd]}. \end{aligned}$$

□

Lemma 5. Let φ be an analytic self-map of \mathbb{D} . If C_φ is 2-normal, then $\overline{\varphi(0)}\varphi_2(0) = 0$.

Proof. Note that

$$\langle C_\varphi^* C_\varphi C_\varphi K_0, K_0 \rangle = \langle K_0 \circ \varphi_2, K_0 \rangle = K_0(\varphi_2(0)) = \frac{1}{1 - \overline{0}\varphi_2(0)} = 1$$

and

$$\langle C_\varphi C_\varphi C_\varphi^* K_0, K_0 \rangle = \langle K_{\varphi(0)} \circ \varphi_2, K_0 \rangle = \frac{1}{1 - \overline{\varphi(0)}\varphi_2(0)}.$$

Since C_φ is 2-normal, we get the desired result. \square

As an application of Lemma 5, we get the following simple example.

Example 1. If $\varphi(z) = \frac{1}{2}iz + \frac{1}{4}$, then $\varphi_2(z) = -\frac{1}{4}z + \frac{1}{8}i + \frac{1}{4}$. Since $\overline{\varphi(0)}\varphi_2(0) \neq 0$, C_φ is not 2-normal by Lemma 5.

3. Main results and proofs

3.1. Automorphism

Theorem 1. Let φ be an automorphism of \mathbb{D} . Then the following statements are equivalent.

- (i) C_φ is 2-normal;
- (ii) $\varphi(z) = -\lambda z, |\lambda| = 1$ or $\varphi(z) = \frac{z-a}{\bar{a}z-1}$ for $a \in \mathbb{D}$.

Proof. (ii) \Rightarrow (i). If $\varphi(z) = -\lambda z, |\lambda| = 1$, then C_φ is normal by Lemma 1 and hence C_φ is 2-normal. If $\varphi(z) = \frac{z-a}{\bar{a}z-1}$ for $a \in \mathbb{D}$, we note that $(\varphi \circ \varphi)(z) = z$, which implies that

$$C_\varphi^* C_\varphi C_\varphi = C_\varphi C_\varphi C_\varphi^*.$$

Thus, C_φ is 2-normal.

(i) \Rightarrow (ii). Assume that C_φ is 2-normal and $\varphi(z) = \frac{\lambda(z-a)}{\bar{a}z-1}$, where $a \in \mathbb{D}$ and $|\lambda| = 1$. We note that

$$\overline{\varphi(0)}\varphi_2(0) = \frac{|a|^2(\lambda - 1)}{\lambda|a|^2 - 1} = 0$$

from Lemma 5. Then $|a|^2(\lambda - 1) = 0$. Hence $|a| = 0$ or $\lambda = 1$.

If $a = 0$, then $\varphi(z) = -\lambda z, |\lambda| = 1$. If $\lambda = 1$, then $\varphi(z) = \frac{z-a}{\bar{a}z-1}$, as desired. The proof is complete. \square

3.2. Linear fractional self-maps with $\varphi(0) = 0$

Theorem 2. Let $\varphi(z) = \frac{az+b}{cz+d}$ be a linear fractional self-map of \mathbb{D} and $\varphi(0) = 0$. Then C_φ is 2-normal if and only if C_φ is normal.

Proof. Sufficiency. It is obvious.

Necessity. Assume that C_φ is 2-normal. Since $\varphi(0) = 0, a \neq 0$, and $\varphi(\mathbb{D}) \subset \mathbb{D}$, we can set

$$\varphi(z) = \frac{z}{mz + n},$$

where $m = \frac{c}{a}, n = \frac{d}{a}$ and $|n| \geq 1 + |m|$. If $m = 0$, then $\varphi(z) = \frac{z}{n}$. So C_φ is normal.

Now we assume that $m \neq 0$. Then $|n| > 1$. For Lemma 3 we obtain that

$$C_\varphi^* C_\varphi C_\varphi K_\alpha(z) = \frac{m(1+n)}{m(1+n) - \bar{\alpha}} - \frac{\bar{\alpha}}{m(1+n) - \bar{\alpha}} \frac{1}{1 - \frac{\bar{\alpha}-m(1+n)}{m(\bar{\alpha}-m(1+n))+|n|^2} z} \tag{1}$$

for $\bar{\alpha} \neq m(1+n)$. From Lemma 4 we get that

$$C_\varphi C_\varphi C_\varphi^* K_\alpha(z) = \frac{m(1+n)(\overline{m\alpha} + \bar{n})}{m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}} - \frac{\bar{\alpha}}{m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}} \frac{1}{1 - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{(\overline{m\alpha} + \bar{n})n^2} z} \tag{2}$$

for $\bar{\alpha} \neq m(1+n)(\overline{m\alpha} + \bar{n})$. Since C_φ is 2-normal, by (1) and (2) we get that

$$\begin{aligned} & \frac{m(1+n)}{m(1+n) - \bar{\alpha}} - \frac{\bar{\alpha}}{m(1+n) - \bar{\alpha}} \frac{1}{1 - \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n} z}} \\ &= \frac{m(1+n)(\overline{m\alpha} + \bar{n})}{m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}} - \frac{\bar{\alpha}}{m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}} \frac{1}{1 - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{(\overline{m\alpha} + \bar{n})n^2} z} \end{aligned}$$

for any $\alpha \in \mathbb{D}$ with $\bar{\alpha} \neq m(1+n)$ and $\bar{\alpha} \neq m(1+n)(\overline{m\alpha} + \bar{n})$. That is,

$$\begin{aligned} 0 &= \frac{\bar{\alpha}m(1+n)(\overline{m\alpha} + \bar{n} - 1)}{[m(1+n) - \bar{\alpha}][m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}]} - \frac{\bar{\alpha}}{m(1+n) - \bar{\alpha}} \frac{1}{1 - \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n} z}} \\ &+ \frac{\bar{\alpha}}{m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}} \frac{1}{1 - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{(\overline{m\alpha} + \bar{n})n^2} z}, \end{aligned}$$

which gives that

$$\begin{aligned} 0 &= \bar{\alpha}m(1+n)(\overline{m\alpha} + \bar{n} - 1) \left(1 - \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n} z} \right) \left(1 - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{(\overline{m\alpha} + \bar{n})n^2} z \right) \\ &- \bar{\alpha}[m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}] \left(1 - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{(\overline{m\alpha} + \bar{n})n^2} z \right) \\ &+ \bar{\alpha}[m(1+n) - \bar{\alpha}] \left(1 - \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n} z} \right) \end{aligned}$$

for any $\alpha \in \mathbb{D}$ with $\bar{\alpha} \neq m(1+n)$ and $\bar{\alpha} \neq m(1+n)(\overline{m\alpha} + \bar{n})$. Multiply this by $(\overline{m\alpha} + \bar{n})$, we get

$$\begin{aligned} 0 &= \bar{\alpha}m(1+n)(\overline{m\alpha} + \bar{n} - 1) \left(1 - \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n} z} \right) \\ &\cdot \left(\overline{m\alpha} + \bar{n} - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{n^2} z \right) \\ &- \bar{\alpha}[m(1+n)(\overline{m\alpha} + \bar{n}) - \bar{\alpha}] \left(\overline{m\alpha} + \bar{n} - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{n^2} z \right) \\ &+ \bar{\alpha}[m(1+n) - \bar{\alpha}] \left(1 - \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n} z} \right) (\overline{m\alpha} + \bar{n}) \end{aligned} \tag{3}$$

for any $\alpha \in \mathbb{D}$ with $\bar{\alpha} \neq m(1+n)$ and $\bar{\alpha} \neq m(1+n)(\overline{m\alpha} + \bar{n})$. Since (3) holds for any $z \in \mathbb{D}$, the coefficient of z^2 in (3) must be zero. This implies that

$$\bar{\alpha}m(1+n)(\overline{m\alpha} + \bar{n} - 1) \frac{\bar{\alpha} - m(1+n)}{\overline{m(\bar{\alpha} - m(1+n)) + |n|^2 n}} - \frac{\bar{\alpha} - m(1+n)(\overline{m\alpha} + \bar{n})}{n^2} = 0.$$

Since $m \neq 0$, from the last equality we obtain that

$$\bar{\alpha}(1+n)(\overline{m\alpha} + \bar{n} - 1) = 0$$

for any $\alpha \in \mathbb{D}$ with $\bar{\alpha} \neq m(1+n)$ and $\bar{\alpha} \neq m(1+n)(\overline{m\alpha} + \bar{n})$. After a calculation, we get

$$\overline{m}(1+n)\bar{\alpha}^2 + (1+n)(\bar{n} - 1)\bar{\alpha} = 0$$

for any $\alpha \in \mathbb{D}$ with $\bar{\alpha} \neq m(1+n)$ and $\bar{\alpha} \neq m(1+n)(\overline{m\alpha} + \bar{n})$, which is a contradiction. So $m = 0$ and $\varphi(z) = \frac{z}{n}$, $|n| \geq 1$. Therefore C_φ is normal. The proof is complete. \square

3.3. Linear fractional self-maps with $c = 0$

Lemma 6. [9] If $\varphi(z) = \frac{az+b}{cz+d}$ is a linear fractional self-map into \mathbb{D} and $c = 0$, then

$$C_\varphi = C_{\tilde{\sigma}}^* T_{\tilde{g}}^*$$

where $\tilde{\sigma}(z) = \frac{\bar{a}z}{-bz+d}$ and $\tilde{g}(z) = \frac{\bar{d}}{-bz+d}$.

Theorem 3. Let $\varphi(z) = \frac{az+b}{cz+d}$ be a linear fractional self-map into \mathbb{D} with $c = 0$. Then C_φ is 2-normal if and only if C_φ is normal.

Proof. Sufficiency. It is obvious.

Necessity. Suppose that C_φ is 2-normal, that is,

$$C_\varphi^* C_\varphi C_\varphi K_\alpha(z) = C_\varphi C_\varphi C_\varphi^* K_\alpha(z)$$

for any $\alpha, z \in \mathbb{D}$. Since $c = 0$, we set $\varphi(z) = sz + t$, where $s = \frac{a}{d}$, $t = \frac{b}{d}$ and $|s| + |t| \leq 1$. Put

$$\sigma(z) = \frac{\bar{s}z}{1-\bar{t}z}, g(z) = \frac{1}{1-\bar{t}z}.$$

According to the proof of [9, Theorem 2.4], by Lemma 6 we obtain that

$$\begin{aligned} C_\varphi^* C_\varphi C_\varphi K_\alpha(z) &= C_\varphi^* C_\sigma^* T_g^* C_\sigma^* T_g K_\alpha(z) = \overline{g(\alpha)g(\sigma(\alpha))} K_{\varphi(\sigma_2(\alpha))}(z) \\ &= \frac{1}{1-t(s+1)\bar{\alpha} - [\bar{t} + (|s|^2s - |t|^2s - |t|^2)\bar{\alpha}]z}. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} C_\varphi C_\varphi C_\varphi^* K_\alpha(z) &= C_\sigma^* T_g^* C_\sigma^* T_g K_{\varphi(\alpha)}(z) = \overline{g(\varphi(\alpha))g(\sigma(\varphi(\alpha)))} K_{\sigma_2(\varphi(\alpha))}(z) \\ &= \frac{1}{1-t\overline{\varphi(\alpha)}} \frac{1}{1-t\overline{\sigma(\varphi(\alpha))}} \frac{1}{1-\overline{\sigma_2(\varphi(\alpha))}z} \\ &= \frac{1}{1-t\overline{\varphi(\alpha)}} \frac{1}{1-t\overline{\sigma(\varphi(\alpha))}} \frac{1}{1-\frac{s\overline{\sigma(\varphi(\alpha))}}{1-t\overline{\sigma(\varphi(\alpha))}}z} \\ &= \frac{1}{1-t\overline{\varphi(\alpha)}} \frac{1}{1-t\overline{\sigma(\varphi(\alpha))} - s\overline{\sigma(\varphi(\alpha))}z} \\ &= \frac{1}{1-t\overline{\varphi(\alpha)} - st\overline{\varphi(\alpha)} - s^2\overline{\varphi(\alpha)}z} \\ &= \frac{1}{1-t\bar{s}\bar{\alpha} - |t|^2 - t|s|^2\bar{\alpha} - s|t|^2 - s|s|^2\bar{\alpha}z - s^2\bar{t}z}. \end{aligned}$$

Since C_φ is 2-normal, we get

$$\begin{aligned} &\frac{1}{1-t(s+1)\bar{\alpha} - [\bar{t} + (|s|^2s - |t|^2s - |t|^2)\bar{\alpha}]z} \\ &= \frac{1}{1-t\bar{s}\bar{\alpha} - |t|^2 - t|s|^2\bar{\alpha} - s|t|^2 - s|s|^2\bar{\alpha}z - s^2\bar{t}z} \end{aligned} \tag{4}$$

for any $\alpha, z \in \mathbb{D}$. Taking $\alpha = 0$ in (4), we obtain

$$\frac{1}{1-|t|^2 - s|t|^2 - s^2\bar{t}z} = \frac{1}{1-\bar{t}z} \tag{5}$$

for any $z \in \mathbb{D}$. So we have $t = 0$ or $s = -1$. If $s = -1$, we know that $t = 0$ since $|s| + |t| \leq 1$. Therefore, by Lemma 1 we get the desired result. The proof is complete. \square

3.4. Linear fractional self-maps with $\varphi(0) \neq 0$ and $a = 0$

Lemma 7. Let $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ be a constant function. Then C_φ is 2-normal if and only if φ is zero on \mathbb{D} .

Proof. Let $\varphi(z) \equiv c$ for some $c \in \mathbb{D}$. Then

$$C_\varphi^* C_\varphi C_\varphi K_\alpha(z) = C_\varphi^* C_\varphi K_\alpha(\varphi(z)) = \frac{1}{1 - \bar{\alpha}c} C_\varphi^* C_\varphi K_0(z) = \frac{1}{1 - \bar{\alpha}c} K_{\varphi(0)}(z) = \frac{1}{1 - \bar{\alpha}c} \frac{1}{1 - \bar{c}z},$$

and

$$C_\varphi C_\varphi C_\varphi^* K_\alpha(z) = K_{\varphi(\alpha)}(\varphi_2(z)) = \frac{1}{1 - \overline{\varphi(\alpha)}\varphi_2(z)} = \frac{1}{1 - |c|^2}$$

for any $\alpha, z \in \mathbb{D}$. Hence C_φ is 2-normal if and only if $c = 0$. \square

Theorem 4. Let $\varphi(z) = \frac{az+b}{cz+d}$ be a linear fractional self-map of \mathbb{D} with $\varphi(0) \neq 0$ and $a = 0$. Then C_φ is not 2-normal.

Proof. Since $\varphi(0) \neq 0$ and $a = 0$, we can set $\varphi(z) = \frac{1}{uz+v}$ where $u = \frac{c}{b}$ and $v = \frac{d}{b}$. If $u = 0$, then $\varphi(z) = \frac{1}{v} \neq 0$ and so C_φ is not 2-normal from Lemma 7. Suppose C_φ is 2-normal and $u \neq 0$ and $v \neq 0$. From Lemma 3, we get

$$C_\varphi^* C_\varphi C_\varphi K_\alpha(z) = \frac{uv}{uv - u\bar{\alpha}} K_{\varphi(0)}(z) + \left(\frac{u + v^2}{u + v^2 - v\bar{\alpha}} - \frac{uv}{uv - u\bar{\alpha}} \right) K_{\varphi(\sigma_2(\alpha))}(z) \tag{6}$$

for any $z \in \mathbb{D}$ and $\alpha \in \mathbb{D}$ with $uv \neq u\bar{\alpha}$. From Lemma 4, we have

$$C_\varphi C_\varphi C_\varphi^* K_\alpha(z) = \frac{(\bar{u}\bar{\alpha} + \bar{v})(uvz + u + v^2)}{(\bar{u}\bar{\alpha} + \bar{v})(uvz + u + v^2) - (uz + v)} \tag{7}$$

for any $z \in \mathbb{D}$ and $\alpha \in \mathbb{D}$. In particular, taking $\alpha = 0$ in (6) and (7), we get

$$C_\varphi^* C_\varphi C_\varphi K_0(z) = \frac{\bar{v}}{\bar{v} - z} \tag{8}$$

and

$$C_\varphi C_\varphi C_\varphi^* K_0(z) = \frac{\bar{v}(uvz + u + v^2)}{\bar{v}(uvz + u + v^2) - (uz + v)}. \tag{9}$$

Since C_φ is 2-normal, (8) and (9) are equal, that is,

$$\frac{\bar{v}}{\bar{v} - z} = \frac{\bar{v}(uvz + u + v^2)}{\bar{v}(uvz + u + v^2) - (uz + v)} \tag{10}$$

for any $z \in \mathbb{D}$. After a calculation, we obtain that

$$u|v|^2 z^2 + |v|^2 vz - |v|^2 = 0$$

for any $z \in \mathbb{D}$. Since $v \neq 0$, dividing both sides by $|v|^2$, we get

$$uz^2 + vz - 1 = 0 \tag{11}$$

for any $z \in \mathbb{D}$, which is a contradiction. Hence C_φ is not 2-normal. \square

3.5. Linear fractional self-maps with $\varphi(0) \neq 0$, $a \neq 0$ and $c \neq 0$

Lemma 8. Let $\varphi(z) = \frac{az+b}{cz+1}$ be a linear fractional self-map of \mathbb{D} with $\varphi(0) \neq 0$, $a \neq 0$ and $c \neq 0$. If C_φ is 2-normal, then

$$(a^2 + bc)\overline{\varphi(0)} \neq c(a + 1).$$

Proof. If $(a^2 + bc)\overline{\varphi(0)} = c(a + 1)$, then we obtain that $a^2\bar{b} = (a + 1 - |b|^2)c$. Since $a \neq 0$ and $\varphi(0) = b \neq 0$, it must be hold that $a + 1 - |b|^2 \neq 0$, and so $c = \frac{a^2\bar{b}}{a+1-|b|^2}$. Therefore,

$$\varphi(z) = \frac{az + b}{\frac{a^2\bar{b}}{a+1-|b|^2}z + 1}.$$

After a calculation,

$$\overline{\varphi(0)}\varphi_2(0) = \frac{(a + 1 - |b|^2)|b|^2}{1 + (a - 1)|b|^2}.$$

Since $a + 1 - |b|^2 \neq 0$ and $b \neq 0$, $\overline{\varphi(0)}\varphi_2(0) \neq 0$. Thus C_φ is not 2-normal by Lemma 5. \square

Theorem 5. Let $\varphi(z) = \frac{az+b}{cz+1}$ be a linear fractional self-map with $\varphi(0) \neq 0$, $a \neq -1$ and $c \neq 0$, then C_φ is not 2-normal.

Proof. Assume that C_φ is 2-normal. By Lemma 3, we get

$$\begin{aligned} C_\varphi^*C_\varphi C_\varphi K_\alpha(z) &= \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\bar{\alpha}} K_{\varphi(0)}(z) \\ &+ \left(\frac{bc + 1}{bc + 1 - b(a + 1)\bar{\alpha}} - \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\bar{\alpha}} \right) K_{\varphi(\sigma_2(a))}(z) \end{aligned} \tag{12}$$

for any $\alpha \in \mathbb{D}$ with $c(a + 1) \neq (a^2 + bc)\bar{\alpha}$.

From Lemma 4, we see

$$\begin{aligned} C_\varphi C_\varphi C_\varphi^* K_\alpha(z) &= \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\overline{\varphi(\alpha)}} \\ &+ \left(\frac{bc + 1}{bc + 1 - b(a + 1)\overline{\varphi(\alpha)}} - \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\overline{\varphi(\alpha)}} \right) K_{\sigma_2(\varphi(\alpha))}(z) \end{aligned} \tag{13}$$

for any $\alpha \in \mathbb{D}$ with $c(a + 1) \neq (a^2 + bc)\overline{\varphi(\alpha)}$. Since $c(a + 1) \neq 0$ and $(a^2 + bc)\overline{\varphi(0)} \neq c(a + 1)$ from Lemma 8, we can take $\alpha = 0$ in (12) and (13). Then

$$C_\varphi^*C_\varphi C_\varphi K_0(z) = \frac{1}{1 - \bar{b}z} \tag{14}$$

and

$$\begin{aligned} &C_\varphi C_\varphi C_\varphi^* K_0(z) \\ &= \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\bar{b}} + \left(\frac{bc + 1}{bc + 1 - b(a + 1)\bar{b}} - \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\bar{b}} \right) K_{\sigma_2(\varphi(0))}(z) \\ &= \frac{c(a + 1)}{c(a + 1) - (a^2 + bc)\bar{b}} - \frac{\bar{b}[(bc + 1)(a^2 + bc) - bc(a + 1)^2]}{[bc + 1 - |b|^2(a + 1)][c(a + 1) - (a^2 + bc)\bar{b}]} \frac{1}{1 - \frac{(a^2+bc)\bar{b}-(a+1)c}{-|b|^2(a+1)+bc+1}z}. \end{aligned} \tag{15}$$

Let

$$A = bc + 1 - |b|^2(a + 1), \quad B = c(a + 1) - (a^2 + bc)\bar{b}.$$

Since C_φ is 2-normal, (14) and (15) are equal, i.e.,

$$\frac{1}{1 - \bar{b}z} = \frac{c(a+1)}{B} - \frac{\bar{b}[(bc+1)(a^2+bc) - bc(a+1)^2]}{AB} \frac{A}{A+Bz}$$

for any $z \in \mathbb{D}$. Hence,

$$\frac{\bar{b}[(bc+1)(a^2+bc) - bc(a+1)^2]A}{AB(A+Bz)} = \frac{c(a+1)(1 - \bar{b}z) - B}{B(1 - \bar{b}z)}$$

for any $z \in \mathbb{D}$. Calculating the above equation and noting that the coefficients of z^2 must be zero, we obtain that $AB^2c(a+1)\bar{b} = 0$. However, $b, c(a+1), A$ and B are all nonzero, which gives a contradiction. Hence C_φ is not 2-normal. \square

In the end of this paper, we give an example of a linear fractional self-map φ which induces a 2-normal operator C_φ but not a normal operator.

Example 2. Let $\varphi(z) = \frac{-z+b}{cz+1}$ be a linear fractional self-map of \mathbb{D} with $\varphi(0) \neq 0$ and $c \neq 0$. Since

$$\varphi_2(z) = \varphi(\varphi(z)) = z, \quad C_\varphi^2 = C_{\varphi_2} = I$$

and so

$$C_\varphi^* C_\varphi C_\varphi = C_\varphi C_\varphi C_\varphi^*.$$

Hence C_φ is a 2-normal operator. However, according to Lemma 1 we see that C_φ is not a normal operator.

Data Availability No data were used to support this study.

Conflicts of Interest The authors declare that they have no conflicts of interest.

References

- [1] J. Conway, *The Theory of Subnormal Operators*, Mathematical Surveys and Monographs, **36**, American Mathematical Society, Providence, RI, 1991. xvi+436 pp.
- [2] C. Cowen, Composition operators on H^2 , *J. Operator Theory* **9** (1983), 77–106.
- [3] C. Cowen, Linear fractional composition operators on H^2 , *Integral Equations Operator Theory* **11**(2) (1988), 151–160.
- [4] C. Cowen, E. Ko, Hermitian weighted composition operators on H^2 , *Trans. Amer. Math. Soc.* **362** (2010), 5771–5801.
- [5] C. Cowen, B. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics. CRC Press, Boca Raton, 1995.
- [6] M. Fatehi, M. Shaabani, Normal, cohyponormal and normaloid weighted composition operators on the Hardy and weighted Bergman spaces, *J. Korean Math. Soc.* **54** (2017), 599–612.
- [7] M. Fatehi, M. Shaabani, Norms of hyponormal weighted composition operators on the Hardy and weighted Bergman spaces, *Oper. Matrices* **12** (2018), 997–1007.
- [8] M. Fatehi, M. Shaabani and D. Thompson, Quasinormal and hyponormal weighted composition operators on H^2 and A_α^2 with linear fractional compositional symbol, *Complex Anal. Oper. Theory* **12** (2018), 1767–1778.
- [9] S. Jung, Y. Kim and E. Ko, Composition operators for which $C_\varphi C_\varphi^*$ and $C_\varphi + C_\varphi^*$ commute, *Complex Var. Elliptic Equ.* **59** (2014), 1608–1625.
- [10] S. Jung, Y. Kim and E. Ko, Characterizations of binormal composition operators with linear fractional symbols on H^2 , *Appl. Math. Comput.* **261** (2015), 252–263.
- [11] H. Schwartz, *Composition operators on H^p* , Thesis (Ph.D.) The University of Toledo, 1969, 84 pp.