

A NUMERICAL PROCEDURE FOR COEFFICIENTS IN GENERALIZED GAUSS-TURÁN QUADRATURES

Gradimir V. Milovanović* and Miodrag M. Spalević

ABSTRACT. A numerical procedure for the coefficients in the generalized Gauss-Turán quadrature formulas is presented. The corresponding nodes as the zeros of s -orthogonal polynomials can be determined by a stable algorithm given in [10]. A numerical example is included.

1. Introduction

We consider the generalized Gauss-Turán quadrature formula (see [17])

$$(1.1) \quad \int_{\mathbb{R}} f(t) d\lambda(t) = \sum_{i=0}^{2s} \sum_{\nu=1}^n A_{i,\nu} f^{(i)}(\tau_{\nu}) + R_n(f),$$

where $d\lambda(t)$ is a nonnegative measure on the real line \mathbb{R} , with compact or infinite support, for which all moments

$$\mu_k = \int_{\mathbb{R}} t^k d\lambda(t), \quad k = 0, 1, \dots,$$

exist and are finite, and $\mu_0 > 0$. The formula (1.1) is exact for all polynomials of degree at most $2(s+1)n - 1$, i.e.,

$$R_n(f) = 0 \quad \text{for } f \in \mathcal{P}_{2(s+1)n-1}.$$

The knots τ_{ν} ($\nu = 1, \dots, n$) in (1.1) are the zeros of the monic polynomial $\pi_n^s(t)$, which minimizes the following integral

$$\int_{\mathbb{R}} \pi_n(t)^{2s+2} d\lambda(t),$$

Received 11.11.1993; Revised February 15.02.1994

1991 Mathematics Subject Classification: 03F65, 13A99

* Supported by Grant 0401A of RFNS through Math. Inst. SANU

where $\pi_n(t) = t^n + a_{n-1}t^{n-1} + \dots + a_1t + a_0$. This polynomial π_n^s is known as s -orthogonal (or s -self associated) polynomial with respect to the measure $d\lambda(t)$ (for some details see [2-7], [11-13]). For $s = 0$, we have the standard case of orthogonal polynomials, and (1.1) then becomes well-known Gauss-Christoffel formula.

In [10] one of us gave a stable method for numerical constructing s -orthogonal polynomials and obtaining the nodes of the generalized Gauss-Turán quadrature formula (1.1). It was an iterative method with quadratic convergence based on a discretized Stieltjes procedure and the Newton-Kantorovič method.

In this paper, in Section 2, we give a numerical procedure for finding the coefficients $A_{i,\nu}$ in (1.1). An alternative method was given by Stroud and Stancu [16] (see also [15]). A numerical example is given in Section 3.

2. The Coefficients in the Generalized Gauss-Turán Quadrature

Let $\tau_\nu = \tau_\nu(s, n)$, $\nu = 1, \dots, n$, be the zeros of the s -orthogonal polynomial $\pi_n(t) (\equiv p_n^s(t))$. If we define ω_ν by

$$\omega_\nu(t) = \left(\frac{\pi_n(t)}{t - \tau_\nu} \right)^{2s+1}, \quad \nu = 1, \dots, n,$$

then the coefficients $A_{i,\nu}$ in the generalized Gauss-Turán quadrature (1.1) can be expressed in the form (see [15])

$$A_{i,\nu} = \frac{1}{i!(2s-i)!} \left[D^{2s-i} \frac{1}{\omega_\nu(t)} \int_{\mathbb{R}} \frac{\pi_n(x)^{2s+1} - \pi_n(t)^{2s+1}}{x-t} d\lambda(x) \right]_{t=\tau_\nu},$$

where D is the standard differentiation operator. Especially, for $i = 2s$, we have

$$A_{2s,\nu} = \frac{1}{(2s)!(\pi_n'(\tau_\nu))^{2s+1}} \int_{\mathbb{R}} \frac{\pi_n(x)^{2s+1}}{t - \tau_\nu} d\lambda(x),$$

i.e.,

$$A_{2s,\nu} = \frac{B_\nu^{(s)}}{(2s)!(\pi_n'(\tau_\nu))^{2s}}, \quad \nu = 1, \dots, n,$$

where $B_\nu^{(s)}$ are the Christoffel numbers of the following Gaussian quadrature (with respect to the measure $d\mu(t) = \pi_n^{2s}(t)d\lambda(t)$),

$$\int_{\mathbb{R}} g(t) d\mu(t) = \sum_{\nu=1}^n B_\nu^{(s)} g(\tau_\nu) + R_n(g), \quad R_n(\mathcal{P}_{2n-1}) = 0.$$

Since $B_\nu^{(s)} > 0$, we conclude that $A_{2s,\nu} > 0$. The expressions for the other coefficients ($i < 2s$) become very complicated. For the numerical calculation we can use a triangular system of linear equations obtained from the formula (1.1) by replacing f with the Newton polynomials: $1, t - \tau_1, \dots, (t - \tau_1)^{2s+1}, (t - \tau_1)^{2s+1}(t - \tau_2), \dots, (t - \tau_1)^{2s+1}(t - \tau_2)^{2s+1} \dots (t - \tau_n)^{2s}$.

Here, we give a method for the numerical calculation of coefficients of the generalized Gauss-Turán quadrature formula (1.1), starting from the Hermite interpolation problem

$$H_m^{(i)}(\tau_\nu) = f^{(i)}(\tau_\nu),$$

where $\nu = 1, \dots, n$; $i = 0, 1, \dots, \alpha_\nu - 1$, $\alpha_1 + \dots + \alpha_n = m + 1$.

Taking $\alpha_i = 2s + 1$, $i = 1, \dots, n$ and integrating $f(t) - H_m(t) = r(f; t)$, we obtain

$$(2.1) \quad \int_{\mathbb{R}} f(t) d\lambda(t) = \sum_{i=0}^{2s} \sum_{\nu=1}^n f^{(i)}(\tau_\nu) \int_{\mathbb{R}} l_{\nu,i}(t) d\lambda(t) + R_n(f),$$

where

$$l_{\nu,i}(t) = \frac{1}{i!} \sum_{k=0}^{2s-i} \frac{1}{k!} \left[\frac{(t - \tau_\nu)^{2s+1}}{\Omega(t)} \right]_{t=\tau_\nu}^{(k)} \frac{\Omega(t)}{(t - \tau_\nu)^{2s-i-k+1}},$$

$$\Omega(t) = [(t - \tau_1)(t - \tau_2) \dots (t - \tau_n)]^{2s+1},$$

and $R_n(f) = \int_{\mathbb{R}} r(f; t) d\lambda(t)$ is the corresponding remainder term.

Hence, (2.1) becomes the generalized Gauss-Turán quadrature formula (1.1), where $A_{i,\nu}$ are the Cotes numbers of higher order and given by

$$A_{i,\nu} = \int_{\mathbb{R}} l_{\nu,i}(t) d\lambda(t),$$

i.e.,

$$A_{i,\nu} = \frac{1}{i!} \sum_{k=0}^{2s-i} \frac{1}{k!} \left[\frac{(t - \tau_\nu)^{2s+1}}{\Omega(t)} \right]_{t=\tau_\nu}^{(k)} \int_{\mathbb{R}} \Omega_{\nu,i+k}(t) d\lambda(t),$$

where $i = 0, 1, \dots, 2s$; $\nu = 1, \dots, n$ and

$$(2.2) \quad \Omega_{\nu,j}(t) = \frac{\Omega(t)}{(t - \tau_\nu)^{2s-j+1}} = (t - \tau_\nu)^j \prod_{j \neq \nu} (t - \tau_j)^{2s+1}.$$

For $i + k \leq 2s$, we can see that $\Omega_{\nu, i+k}(t)$ is a polynomial of degree at most

$$(n-1)(2s+1) + 2s = (2s+1)n - 1 \leq 2(s+1)n - 1 = 2N - 1,$$

where $N = (s+1)n$.

Hence, the problem of determining of the coefficients of the Gauss-Turán quadrature formula (1.1) is reduced to determination of integrals in (2.2). All of the above integrals in can be found exactly, except for rounding errors, by using a Gauss-Christoffel quadrature formula with respect to the measure $d\lambda(t)$,

$$(2.3) \quad \int_{\mathbb{R}} g(t) d\lambda(t) = \sum_{k=1}^N A_k^{(N)} g(\tau_k^{(N)}) + R_N(g),$$

taking $N = (s+1)n$ knots. This formula is exact for all polynomials of degree at most $2N - 1 = 2(s+1)n - 1$.

In order to calculate the derivatives

$$(2.4) \quad \left[\frac{(t - \tau_\nu)^{2s+1}}{\Omega(t)} \right]_{t=\tau_\nu}^{(k)} \quad (k = 0, 1, \dots, 2s; \nu = 1, \dots, n),$$

we need the following auxiliary result:

Lemma 2.1. *If $g \in C^{(m)}(E)$, $m \in \mathbb{N}_0$, $E \subset \mathbb{R}$, then*

$$(e^g)^{(0)} = e^g, \quad (e^g)^{(p)} = \sum_{l=1}^p \binom{p-1}{l-1} g^{(l)} (e^g)^{(p-l)}, \quad p = 1, \dots, m.$$

Proof. Since $(e^g)' = g'e^g$, applying the Leibnitz's formula for the derivative of the product of functions, we have

$$\begin{aligned} (e^g)^{(p)} &= (g'e^g)^{(p-1)} = \sum_{j=0}^{p-1} \binom{p-1}{j} (g')^{(j)} (e^g)^{(p-j-1)} \\ &= \sum_{l=1}^p \binom{p-1}{l-1} g^{(l)} (e^g)^{(p-l)}, \end{aligned}$$

where $p = 1, \dots, m$. \square

Let $a = \tau_0 < \tau_1 < \dots < \tau_n < \tau_{n+1} = b$, where $[a, b]$ is the smallest closed interval containing $\text{supp}(d\lambda)$, or $a = -\infty$, $b = +\infty$. For $t \in (\tau_{\nu-1}, \tau_{\nu+1})$ we define u_ν by

$$u_\nu(t) = \prod_{i \neq \nu} (t - \tau_i)^{-(2s+1)} = (-1)^{n-\nu} \exp \left[-(2s+1) \sum_{i \neq \nu} \log |t - \tau_i| \right],$$

i.e., $u_\nu(t) = (-1)^{n-\nu} e^{h_\nu(t)}$, where

$$h_\nu(t) = -(2s+1) \sum_{i \neq \nu} g_i(t), \quad g_i(t) = \log |t - \tau_i|.$$

Since $g_i^{(j)}(\tau_\nu) = (-1)^{j-1} (j-1)! (\tau_\nu - \tau_i)^{-j}$, $j \geq 1$, we have

$$h_\nu^{(j)}(\tau_\nu) = -(2s+1) (-1)^{j-1} (j-1)! \sum_{i \neq \nu} (\tau_\nu - \tau_i)^{-j}.$$

It is clear that the derivatives in (2.4) are exactly the derivatives of $u_\nu(t)$ in the point $t = \tau_\nu$. Thus, using Lemma 2.1, we can express them in terms of $h_\nu^{(j)}(\tau_\nu)$.

This numerical method for calculating the coefficients $A_{i,\nu}$ can be summarized in the following form:

Proposition 2.2. *Let τ_ν , $\nu = 1, \dots, n$, be zeros of the s -orthogonal polynomial $\pi_n^s(t)$, with respect to the measure $d\lambda(t)$ on \mathbb{R} . Then, coefficients of the generalized Gauss-Turán quadrature formula,*

$$\int_{\mathbb{R}} f(t) d\lambda(t) = \sum_{i=0}^{2s} \sum_{\nu=1}^n A_{i,\nu} f^{(i)}(\tau_\nu) + R(f),$$

can be expressed in the form

$$A_{i,\nu} = \frac{1}{i!} (-1)^{n-\nu} \sum_{k=0}^{2s-i} \frac{1}{k!} [e^{h_\nu(t)}]_{t=\tau_\nu}^{(k)} \sum_{j=1}^N A_j^{(N)} \Omega_{\nu, i+k}(\tau_j^{(N)}),$$

where $A_j^{(N)}$ and $\tau_j^{(N)}$ are weights and nodes of the Gauss-Christoffel quadrature formula (2.3) in $N = (s+1)n$ points, the polynomial $\Omega_{\nu,j}(t)$ is given by (2.2), and $[e^{h_\nu(t)}]_{t=\tau_\nu}^{(k)}$ is determined by Lemma 2.1.

To conclude this section we mention a particularly interesting case of the Chebyshev measure $d\lambda(t) = (1-t^2)^{-1/2} dt$. In 1930, S. Bernstein [1] showed

that the monic Chebyshev polynomial $\hat{T}_n(t) = T_n(t)/2^{n-1}$ minimizes all integrals of the form

$$\int_{-1}^1 \frac{|\pi_n(t)|^{k+1}}{\sqrt{1-t^2}} dt, \quad k \geq 0.$$

Thus, the Chebyshev-Turán formula

$$(2.5) \quad \int_{-1}^1 \frac{f(t)}{\sqrt{1-t^2}} dt = \sum_{i=0}^{2s} \sum_{\nu=1}^n A_{i,\nu} f^{(i)}(\tau_\nu) + R_n(f),$$

with $\tau_\nu = \cos \frac{(2\nu-1)\pi}{2n}$, $\nu = 1, \dots, n$ is exact for all polynomials of degree at most $2(s+1)n - 1$. Turán has stated a problem of explicit determination of $A_{i,\nu}$ and its behavior as $n \rightarrow +\infty$ (see Problem XXVI in [18]). Some characterizations and solution for $s = 2$ were obtained by Micchelli and Rivlin [9], Riess [14], and Varma [19]. One simple answer to Turán question was given by Kis [9].

3. Numerical Example

In this section we give an example when is preferable to use a formula of Turán type instead of the standard Gaussian formula

$$(3.1) \quad \int_{\mathbb{R}} f(t) d\lambda(t) = \sum_{\nu=1}^n A_\nu f(t_\nu) + R_n(f),$$

for which $R_n(\mathcal{R}_{2n-1}) = 0$. All computations were done on the MICROVAX 3400 computer in Q-arithmetic (machine precision $\approx 1.93 \times 10^{-34}$).

Consider the following simple numerical example

$$I = \int_{-1}^1 e^t \sqrt{1-t^2} dt = 1.7754996892121809468785765372 \dots$$

Here we have $f(t) = e^t$ and $d\lambda(t) = \sqrt{1-t^2} dt$ on $[-1, 1]$ (the Chebyshev measure of the second kind). Notice that $f^{(i)}(t) = f(t)$ for every $i \geq 0$.

The Gaussian formula (3.1) and the corresponding Gauss-Turán formula (1.1) give

$$(3.2) \quad I \approx I_n^G = \sum_{\nu=1}^n A_\nu e^{t_\nu}$$

and

$$(3.3) \quad I \approx I_{n,s}^T = \sum_{\nu=1}^n C_{\nu}^{(s)} e^{\tau_{\nu}},$$

respectively, where $C_{\nu}^{(s)} = \sum_{i=0}^{2s} A_{i,\nu}$.

Table 3.1 shows the relative errors $|(I_{n,s}^T - I)/I_{n,s}^T|$ for $n = 1(1)5$ and $s = 0(1)5$. (Numbers in parentheses indicate decimal exponents and m.p. is the machine precision.)

TABLE 3.1
Relative errors in quadrature sums $I_{n,s}^T$

n	$s = 0$	$s = 1$	$s = 2$	$s = 3$	$s = 4$	$s = 5$
1	1.15(-1)	4.71(-3)	9.72(-5)	1.21(-6)	1.01(-8)	5.98(-11)
2	2.38(-3)	2.05(-7)	3.06(-12)	1.36(-17)	2.40(-23)	1.88(-29)
3	1.97(-5)	1.15(-12)	4.02(-21)	9.26(-31)	m.p.	m.p.
4	8.76(-8)	1.71(-18)	4.68(-31)	m.p.	m.p.	m.p.
5	2.43(-10)	9.40(-25)	m.p.	m.p.	m.p.	m.p.

For $s = 0$ the quadrature formula (3.3) reduces to (3.2), i.e., $I_{n,0}^T \equiv I_n^G$. Notice that Turán formula (3.3) with n nodes has the same degree of exactness as Gaussian formula with $(s + 1)n$ nodes.

REFERENCES

- [1] S. BERNSTEIN, *Sur les polinomes orthogonaux relatifs à un segment fini*, J. Math. Pures Appl. **9**(9) (1930), 127-177.
- [2] H. ENGELS, *Numerical Quadrature and Cubature*, Academic Press, London, 1980.
- [3] W. GAUTSCHI, *A survey of Gauss-Christoffel quadrature formulae*, In: E. B. Christoffel (P. L. Butzer and F. Fehér, eds.), Birkhäuser, Basel, 1981, pp. 72-147.
- [4] W. GAUTSCHI, *On generating orthogonal polynomials*, SIAM J. Sci. Statist. Comput. **3** (1982), 289-317.
- [5] A. GHIZZETTI and A. OSSICINI, *Su un nuovo tipo di sviluppo di una funzione in serie di polinomi*, Rend. Accad. Naz. Lincei **8** (43) (1967), 21-29.
- [6] S. GUERRA, *Polinomi generati da successioni peso e teoremi di rappresentazione di una funzione in serie di tali polinomi*, Rend. Ist. Mat. Univ. Trieste **8** (1976), 172-194.
- [7] S. GUERRA, *Su un determinate collegato ad un sistema di polinomi ortogonali*, Rend. Ist. Mat. Univ. Trieste **10** (1978), 66-79.
- [8] O. KIS, *Remark on mechanical quadrature*, Acta Math. Acad. Sci. Hungar. **8** (1957), 473-476 (Russian).

- [9] C. A. MICCHELLI and T. J. RIVLIN, *Turán formulae highest precision quadrature rules for Chebyshev coefficients*, IBM J. Res. Develop. **16** (1972), 372–379.
- [10] G. V. MILOVANOVIĆ, *Construction of s -orthogonal polynomials and Turán quadrature formulae*, In: Numerical Methods and Approximation Theory III (Niš, 1987) (G. V. Milovanović, ed.), Univ. Niš, Niš, 1988, pp. 311–328.
- [11] A. OSSICINI, *Costruzione di formule di quadratura di tipo Gaussiano*, Ann. Mat. Pura Appl. (4) **72** (1966), 213–238.
- [12] A. OSSICINI and F. ROSATI, *Funzioni caratteristiche nelle formule di quadratura gaussiane con nodi multipli*, Bull. Un. Mat. Ital. (4) **11** (1975), 224–237.
- [13] A. OSSICINI and F. ROSATI, *Sulla convergenza dei funzionali ipergaussiani*, Rend. Mat. (6) bf11 (1978), 97–108.
- [14] R. D. RIESS, *Gauss-Turán quadratures of Chebyshev type and error formulae*, Computing **15** (1975), 173–179.
- [15] D. D. STANCU, *Asupra unor formule generale de integrate numerica*, Acad. R. P. Romine. Stud. Cere. Mat. **9** (1958), 209–216.
- [16] A. H. STROUD and D. D. STANCU, *Quadrature formulas with multiple Gaussian nodes*, J. SIAM Numer. Anal. Ser. B **2** (1965), 129–143.
- [17] P. TURÁN, *On the theory of mechanical quadrature*, Acta Sci. Math. Szeged **12** (1950), 30–37.
- [18] P. TURÁN, *On some open problems of approximation theory*, J. Approx. Theory **29** (1980), 23–85.
- [19] A. K. VARMA, *On optimal quadrature formulae*, Studia Sci. Math. Hungar. **19** (1984), 437–446.

UNIVERSITY OF NIŠ, FACULTY OF ELECTRONIC ENGINEERING, DEPARTEMENT OF MATHEMATICS, P. O. Box 73, 18000 NIŠ, YUGOSLAVIA.

UNIVERSITY "SVETOZAR MARKOVIĆ", FACULTY OF NATURAL SCIENCES AND MATHEMATICS, DEPARTMENT OF MATHEMATICS, R. DOMANOVIĆA 12, 34000 KRAGUJEVAC, YUGOSLAVIA.