

SOME CLASSES OF SIMULTANEOUS COSPECTRAL GRAPHS FOR ADJACENCY, LAPLACIAN AND NORMALIZED LAPLACIAN MATRICES

A. DAS¹ AND P. PANIGRAHI²

ABSTRACT. In this paper we construct several classes of non-regular graphs which are co-spectral with respect to all the three matrices, namely, adjacency, Laplacian and normalized Laplacian, and hence we answer a question asked by S. Butler. We make these constructions starting with two pairs (G_1, H_1) and (G_2, H_2) of A -cospectral regular graphs, taking their R -graph $\mathcal{R}(G_i)$, $\mathcal{R}(H_i)$, $i = 1, 2$, and finally making some kind of partial joins between $\mathcal{R}(G_1)$ and $\mathcal{R}(G_2)$; and $\mathcal{R}(H_1)$ and $\mathcal{R}(H_2)$. Moreover, we determine the number of spanning trees and the Kirchhoff index of the newly constructed graphs.

1. INTRODUCTION

In recent years, construction of cospectral graphs for different matrices is one of the interesting research problem in the area of spectral graph theory. Here we construct some graphs which give an answer to the question “Is there an example of two non-regular graphs which are cospectral with respect to the adjacency, combinatorial Laplacian and normalized Laplacian at the same time?” asked by Butler [2]. To present the results of the paper we need some definitions and terminology as follow. All graphs considered in the paper are simple and undirected. For any graph G , we take $V(G)$ and $E(G)$ as the vertex set and edge set of G , respectively. The *adjacency matrix* of graph G , denoted by $A(G)$, is a square matrix whose rows and columns are indexed by vertices of graph G , and the $(u, v)^{\text{th}}$ entry is 1 if and only if vertex u is adjacent to vertex v and 0 otherwise. If $D(G)$ is the diagonal matrix of vertex degrees in G , then

Key words and phrases. Adjacency matrix, Laplacian matrix, normalized Laplacian matrix, R -vertex-vertex join, R -edge-edge join, R -edge-vertex join.

2010 *Mathematics Subject Classification.* Primary: 05C50. Secondary: 47A75.

<https://doi.org/10.46793/KgJMat2608.1243D>

Received: December 07, 2023.

Accepted: November 19, 2024.

the *Laplacian matrix* $L(G)$ is defined as $L(G) = D(G) - A(G)$ and the *normalized Laplacian matrix* $\mathcal{L}(G)$ of G is defined as $\mathcal{L}(G) = I - D(G)^{-1/2}A(G)D(G)^{-1/2}$ with the convention that $D(G)^{-1}(u, u) = 0$ if degree of u is zero. For a given square matrix M of size n , we denote the characteristic polynomial $\det(xI_n - M)$ by $f_M(x)$. The eigenvalues of $A(G)$, $L(G)$ and $\mathcal{L}(G)$ are denoted by $\lambda_1(G) \geq \lambda_2(G) \geq \dots \geq \lambda_n(G)$, $0 = \mu_1(G) \leq \mu_2(G) \leq \dots \leq \mu_n(G)$, and $0 = \delta_1(G) \leq \delta_2(G) \leq \dots \leq \delta_n(G) \leq 2$, respectively, where n is the number of vertices of G . The multiset of eigenvalues of $A(G)$ (respectively, $L(G)$, $\mathcal{L}(G)$) is called the *adjacency* (respectively, *Laplacian*, *normalized Laplacian*) *spectrum* of G , and denoted by A -spectrum (respectively, L -spectrum, $\mathcal{L}$ -spectrum). Two graphs are said to be A -cospectral (respectively, L -cospectral, $\mathcal{L}$ -cospectral) if they have the same A -spectrum (respectively, L -spectrum, $\mathcal{L}$ -spectrum).

The R -graph $\mathcal{R}(G)$ [6] of a graph G is the graph obtained from G by introducing a new vertex u_e for each $e \in E(G)$ and making u_e adjacent to both the end vertices of e . The set of such new vertices is denoted by $I(G)$, i.e., $I(G) = V(\mathcal{R}(G)) \setminus V(G)$. The partial joins of R -graphs which are considered in the paper are given in the definition below.

Definition 1.1. Let G_1 and G_2 be two vertex-disjoint graphs with number of vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then the following hold.




- (i) The R -vertex-vertex join of G_1 and G_2 , denoted by $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$, is the graph obtained from $\mathcal{R}(G_1)$ and $\mathcal{R}(G_2)$ by joining each vertex of $V(G_1)$ with every vertex of $V(G_2)$. The graph $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ has $n_1 + n_2 + m_1 + m_2$ vertices and $3m_1 + n_1n_2 + 3m_2$ edges.
- (ii) The R -edge-edge join of G_1 and G_2 , denoted by $\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)$, is the graph obtained from $\mathcal{R}(G_1)$ and $\mathcal{R}(G_2)$ by joining each vertex of $I(G_1)$ with every vertex of $I(G_2)$. The graph $\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)$ has $n_1 + n_2 + m_1 + m_2$ vertices and $m_1(3 + m_2) + 3m_2$ edges.
- (iii) The R -edge-vertex join of G_1 and G_2 , denoted by $\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)$, is the graph obtained from $\mathcal{R}(G_1)$ and $\mathcal{R}(G_2)$ by joining each vertex of $I(G_1)$ with every vertex of $V(G_2)$. The graph $\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)$ has $n_1 + n_2 + m_1 + m_2$ vertices and $m_1(3 + n_2) + 3m_2$ edges. (We note that R -vertex-edge join of G_1 and G_2 is isomorphic to the R -edge-vertex join of G_2 and G_1 .)

Example 1.1. Let us consider two graphs $G_1 = P_3$ and $G_2 = P_4$ (see Figures 1, 2 and 3). The set of dark vertices of G_1 and G_2 are $I(G_1)$ and $I(G_2)$, respectively.

In the following lemma we find the degrees of vertices in the above constructed graphs.

Lemma 1.1. (i) The degree of any vertex v in $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ is given by

$$d_{\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)}(v) = \begin{cases} n_2 + 2d_{G_1}(v), & \text{if } v \in V(G_1), \\ 2, & \text{if } v \in I(G_1) \cup I(G_2), \\ n_1 + 2d_{G_2}(v), & \text{if } v \in V(G_2). \end{cases}$$

FIGURE 1. R -vertex-vertex join of P_3 and P_4 FIGURE 2. R -edge-edge join of P_3 and P_4 FIGURE 3. R -edge-vertex join of P_3 and P_4

(ii) The degree of any vertex v in $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ is given by

$$d_{\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)}(v) = \begin{cases} 2d_{G_1}(v), & \text{if } v \in V(G_1), \\ 2 + m_2, & \text{if } v \in I(G_1), \\ 2d_{G_2}(v), & \text{if } v \in V(G_2), \\ 2 + m_1, & \text{if } v \in I(G_2). \end{cases}$$

(iii) The degree of any vertex v in $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ is given by

$$d_{\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)}(v) = \begin{cases} 2d_{G_1}(v), & \text{if } v \in V(G_1), \\ 2 + n_2, & \text{if } v \in I(G_1), \\ 2d_{G_2}(v) + m_1 & \text{if } v \in V(G_2), \\ 2, & \text{if } v \in I(G_2). \end{cases}$$

For two matrices A and B , of same size $m \times n$, the *Hadamard product* $A \bullet B$ of A and B is a matrix of the same size $m \times n$ with entries given by $(A \bullet B)_{ij} = (A)_{ij} \cdot (B)_{ij}$ (that is entrywise multiplication). Hadamard product is commutative, that is $A \bullet B = B \bullet A$.

Notation. Throughout the paper, for any positive integers k , n_1 and n_2 , I_k denotes the identity matrix of size k , $J_{n_1 \times n_2}$ denotes $n_1 \times n_2$ matrix whose all entries are 1, $\mathbf{1}_n$ stands for the column vector of size n with all entries equal to 1, $K_{n \times n}$ denotes an $n \times n$ matrix whose all entries are the same. In other words, $K_{n \times n} = \alpha J_{n \times n}$, for a real number α . For any positive integers s and t , $O_{s \times t}$ denotes the zero matrix of size $s \times t$.

To prove our results we need some basics as given below.

Lemma 1.2 (Schur complement [7]). *Suppose that the order of all four matrices M , N , P and Q satisfy the rules of operations on matrices. Then we have*

$$\begin{aligned} \begin{vmatrix} M & N \\ P & Q \end{vmatrix} &= |Q| \cdot |M - NQ^{-1}P| \quad (\text{if } Q \text{ is a non-singular square matrix}) \\ &= |M| \cdot |Q - PM^{-1}N| \quad (\text{if } M \text{ is a non-singular square matrix}). \end{aligned}$$

Lemma 1.3 ([7]). *For a square matrix A of size n and a scalar α , $\det(A + \alpha J_{n \times n}) = \det(A) + \alpha \mathbf{1}_n^T \text{adj}(A) \mathbf{1}_n$, where $\text{adj}(A)$ is the adjugate matrix of A .*

Lemma 1.4. *For any real numbers $c, d > 0$, we have*

$$(cI_n - dJ_{n \times n})^{-1} = \frac{1}{c}I_n + \frac{d}{c(c - nd)}J_{n \times n}.$$

Proof. We have

$$\begin{aligned} (cI_n - dJ_{n \times n})^{-1} &= \frac{\text{adj}(cI_n - dJ_{n \times n})}{\det(cI_n - dJ_{n \times n})} = \frac{c^{n-2}(c - nd)I_n + c^{n-2}dJ_{n \times n}}{c^{n-1}(c - nd)} \\ &= \frac{1}{c}I_n + \frac{d}{c(c - nd)}J_{n \times n}. \quad \square \end{aligned}$$

For a graph G on n vertices and m edges, the *vertex-edge incidence matrix* $R(G)$ of G is a matrix of size $n \times m$, with entry $r_{ij} = 1$ if the i^{th} vertex is incident to the j^{th} edge, and 0 otherwise. In particular, if G is an r -regular graph then $R(G)R(G)^T = A(G) + rI_n = 2rI_n - L(G) = r(2I_n - \mathcal{L}(G))$.

Notation. The M -coronal of an $n \times n$ matrix M , denoted by $\Gamma_M(x)$, is defined [5,10] as the sum of the entries of the matrix $(xI_n - M)^{-1}$, that is, $\Gamma_M(x) = \mathbf{1}_n^T(xI_n - M)^{-1}\mathbf{1}_n$.

Lemma 1.5 ([5]). *If M is an $n \times n$ matrix with each row sum equal to a constant t , then $\Gamma_M(x) = \frac{n}{x-t}$.*

Butler [2] constructed non-regular bipartite graphs which are cospectral with respect to both the adjacency and normalized Laplacian matrices, and then asked for existence of non-regular graphs which are cospectral with respect to all the three matrices, namely, adjacency, Laplacian and normalized Laplacian. In this paper we construct several classes of such graphs taking help of the operations R -vertex-vertex join, R -edge-edge join, and R -edge-vertex join. We also find the number of spanning trees and Kirchhoff index for all the partial join of R -graphs constructed here.

2. ADJACENCY, LAPLACIAN AND NORMALIZED LAPLACIAN SPECTRA OF THE GRAPHS

In this section we consider regular graphs G_i on n_i vertices, m_i edges, and with degree of regularity r_i , $i = 1, 2$. To obtain the required matrices we label the vertices of the graphs in the following way. Let $V(G_1) = \{v_1, \dots, v_{n_1}\}$, $I(G_1) = \{e_1, \dots, e_{m_1}\}$, $V(G_2) = \{u_1, \dots, u_{n_2}\}$, $I(G_2) = \{f_1, \dots, f_{m_2}\}$. Then, $V(G_1) \cup I(G_1) \cup V(G_2) \cup I(G_2)$ is a partition for all $V(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2))$, $V(\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2))$ and $V(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2))$.

Lemma 2.1. For $i = 1, 2$, let G_i be a graph with n_i vertices and m_i edges. Then, we have the following:

$$\begin{aligned}
 & \text{(i)} \\
 & A(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} A(G_1) & R(G_1) & J_{n_1 \times n_2} & O_{n_1 \times m_2} \\ R(G_1)^T & O_{m_1} & O_{m_1 \times n_2} & O_{m_1 \times m_2} \\ J_{n_2 \times n_1} & O_{n_2 \times m_1} & A(G_2) & R(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & R(G_2)^T & O_{m_2} \end{pmatrix}; \\
 & \text{(ii)} \\
 & A(\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} A(G_1) & R(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ R(G_1)^T & O_{m_1} & O_{m_1 \times n_2} & J_{m_1 \times m_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & A(G_2) & R(G_2) \\ O_{m_2 \times n_1} & J_{m_2 \times m_1} & R(G_2)^T & O_{m_2} \end{pmatrix}; \\
 & \text{(iii)} \\
 & A(\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} A(G_1) & R(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ R(G_1)^T & O_{m_1} & J_{m_1 \times n_2} & O_{m_1 \times m_2} \\ O_{n_2 \times n_1} & J_{n_2 \times m_1} & A(G_2) & R(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & R(G_2)^T & O_{m_2} \end{pmatrix}.
 \end{aligned}$$

Theorem 2.1. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the adjacency spectrum of $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ consists of:

- (i) two roots of the equation $x^2 - \lambda_i(G_1)x - r_1 - \lambda_i(G_1)$, for every eigenvalue $\lambda_i(G_1)$, $i = 2, 3, \dots, n_1$, of $A(G_1)$;
- (ii) two roots of the equation $x^2 - \lambda_j(G_2)x - r_2 - \lambda_j(G_2)$, for every eigenvalue $\lambda_j(G_2)$, $j = 2, 3, \dots, n_2$, of $A(G_2)$,
- (iii) the eigenvalue 0 with multiplicity $m_1 + m_2 - n_1 - n_2$;
- (iv) four roots of the equation $x^4 - (r_1 + r_2)x^3 - (2r_1 + n_1n_2 + 2r_2 - r_1r_2)x^2 + 4r_1r_2x + 4r_1r_2 = 0$.

Proof. The adjacency characteristic polynomial of $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ is

$$\begin{aligned}
 f_{A(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2))}(x) &= \det \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & -J_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -R(G_1)^T & xI_{m_1} & O_{m_1 \times n_2} & O_{m_1 \times m_2} \\ -J_{n_2 \times n_1} & O_{n_2 \times m_1} & xI_{n_2} - A(G_2) & -R(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & -R(G_2)^T & xI_{m_2} \end{pmatrix} \\
 &= x^{m_2} \det(S),
 \end{aligned}$$

where

$$\begin{aligned}
 S &= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & -J_{n_1 \times n_2} \\ -R(G_1)^T & xI_{m_1} & O_{m_1 \times n_2} \\ -J_{n_2 \times n_1} & O_{n_2 \times m_1} & xI_{n_2} - A(G_2) \end{pmatrix} \\
 &\quad - \begin{pmatrix} O_{n_1 \times m_2} \\ O_{m_1 \times m_2} \\ -R(G_2) \end{pmatrix} \frac{1}{x} \begin{pmatrix} O_{m_2 \times n_1} & O_{m_2 \times m_1} & -R(G_2)^T \end{pmatrix}
 \end{aligned}$$

$$= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & -J_{n_1 \times n_2} \\ -R(G_1)^T & xI_{m_1} & O_{m_1 \times n_2} \\ -J_{n_2 \times n_1} & O_{n_2 \times m_1} & xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T \end{pmatrix}.$$

Hence,

$$\begin{aligned} \det(S) &= \det\left(xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T\right) \det(W) \\ &= \prod_{j=1}^{n_2} \left\{ x - \lambda_j(G_2) - \frac{r_2}{x} - \frac{\lambda_j(G_2)}{x} \right\} \det(W), \end{aligned}$$

where

$$\begin{aligned} W &= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) \\ -R(G_1)^T & xI_{m_1} \end{pmatrix} \\ &\quad - \begin{pmatrix} -J_{n_1 \times n_2} \\ O_{m_1 \times n_2} \end{pmatrix} \left(xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T \right)^{-1} \begin{pmatrix} -J_{n_2 \times n_1} & O_{n_2 \times m_1} \end{pmatrix} \\ &= \begin{pmatrix} xI_{n_1} - A(G_1) - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x)J_{n_1 \times n_1} & -R(G_1) \\ -R(G_1)^T & xI_{m_1} \end{pmatrix}. \end{aligned}$$

Then,

$$\begin{aligned} \det(W) &= x^{m_1} \det \left(xI_{n_1} - A(G_1) - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x)J_{n_1 \times n_1} - \frac{1}{x}R(G_1)R(G_1)^T \right) \\ &= x^{m_1} \left[\det \left(xI_{n_1} - A(G_1) - \frac{1}{x}R(G_1)R(G_1)^T \right) \right. \\ &\quad \left. - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \mathbf{1}_{n_1}^T \operatorname{adj} \left(xI_{n_1} - A(G_1) - \frac{1}{x}R(G_1)R(G_1)^T \right) \mathbf{1}_{n_1} \right] \\ &= x^{m_1} \det \left(xI_{n_1} - A(G_1) - \frac{1}{x}R(G_1)R(G_1)^T \right) \\ &\quad \times \left[1 - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \mathbf{1}_{n_1}^T \left(xI_{n_1} - A(G_1) - \frac{1}{x}R(G_1)R(G_1)^T \right)^{-1} \mathbf{1}_{n_1} \right] \\ &= x^{m_1} \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{r_1}{x} - \frac{\lambda_i(G_1)}{x} \right\} \\ &\quad \times \left[1 - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \Gamma_{A(G_1) + \frac{1}{x}R(G_1)R(G_1)^T}(x) \right] \\ &= x^{m_1} \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{r_1}{x} - \frac{\lambda_i(G_1)}{x} \right\} \left[1 - \frac{n_2}{x - r_2 - \frac{2r_2}{x}} \cdot \frac{n_1}{x - r_1 - \frac{2r_1}{x}} \right]. \end{aligned}$$

Therefore,

$$f_{A(\mathcal{R}(G_1))\check{\vee}\mathcal{R}(G_2)}(x) = x^{m_1} x^{m_2} \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{r_1}{x} - \frac{\lambda_i(G_1)}{x} \right\}$$

$$\begin{aligned}
& \times \prod_{j=1}^{n_2} \left\{ x - \lambda_j(G_2) - \frac{r_2}{x} - \frac{\lambda_j(G_2)}{x} \right\} \\
& \times \left[1 - \frac{n_2}{x - r_2 - \frac{2r_2}{x}} \cdot \frac{n_1}{x - r_1 - \frac{2r_1}{x}} \right] \\
& = x^{m_1 - n_1} x^{m_2 - n_2} \prod_{i=2}^{n_1} \left\{ x^2 - \lambda_i(G_1)x - r_1 - \lambda_i(G_1) \right\} \\
& \times \prod_{j=2}^{n_2} \left\{ x^2 - \lambda_j(G_2)x - r_2 - \lambda_j(G_2) \right\} \left[x^4 - (r_1 + r_2)x^3 \right. \\
& \quad \left. - (2r_1 + n_1n_2 + 2r_2 - r_1r_2)x^2 + 4r_1r_2x + 4r_1r_2 \right],
\end{aligned}$$

and the result follows immediately. $\square$

Theorem 2.2. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the adjacency spectrum of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ consists of:

- (i) two roots of the equation $x^2 - \lambda_i(G_1)x - r_1 - \lambda_i(G_1)$, for every eigenvalue $\lambda_i(G_1)$, $i = 2, 3, \dots, n_1$, of $A(G_1)$;
- (ii) two roots of the equation $x^2 - \lambda_j(G_2)x - r_2 - \lambda_j(G_2)$, for every eigenvalue $\lambda_j(G_2)$, $j = 2, 3, \dots, n_2$, of $A(G_2)$;
- (iii) the eigenvalue 0 with multiplicity $m_1 + m_2 - n_1 - n_2$;
- (iv) four roots of the equation $x^4 - (r_1 + r_2)x^3 - (2r_1 + m_1m_2 + 2r_2 - r_1r_2)x^2 + (4r_1r_2 + m_1m_2r_1 + m_1m_2r_2)x + 4r_1r_2 - m_1m_2r_1r_2 = 0$.

Proof. The adjacency characteristic polynomial of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ is

$$\begin{aligned}
f_{A(\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2))}(x) &= \det \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -R(G_1)^T & xI_{m_1} & O_{m_1 \times n_2} & -J_{m_1 \times m_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & xI_{n_2} - A(G_2) & -R(G_2) \\ O_{m_2 \times n_1} & -J_{m_2 \times m_1} & -R(G_2)^T & xI_{m_2} \end{pmatrix} \\
&= x^{m_2} \det(S),
\end{aligned}$$

where

$$\begin{aligned}
S &= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & O_{n_1 \times n_2} \\ -R(G_1)^T & xI_{m_1} & O_{m_1 \times n_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & xI_{n_2} - A(G_2) \end{pmatrix} \\
&\quad - \begin{pmatrix} O_{n_1 \times m_1} \\ -J_{m_1 \times m_2} \\ -R(G_2) \end{pmatrix} \frac{1}{x} \begin{pmatrix} O_{m_2 \times n_1} & -J_{m_2 \times m_1} & -R(G_2)^T \end{pmatrix} \\
&= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & O_{n_1 \times n_2} \\ -R(G_1)^T & xI_{m_1} - \frac{m_2}{x} J_{m_1 \times m_1} & -\frac{1}{x} J_{m_1 \times m_2} R(G_2)^T \\ O_{n_2 \times n_1} & -\frac{1}{x} R(G_2) J_{m_2 \times m_1} & xI_{n_2} - A(G_2) - \frac{1}{x} R(G_2) R(G_2)^T \end{pmatrix}.
\end{aligned}$$

Hence,

$$\begin{aligned}\det(S) &= \det \left(xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T \right) \det(W) \\ &= \prod_{j=1}^{n_2} \left\{ x - \lambda_j(G_2) - \frac{r_2}{x} - \frac{\lambda_j(G_2)}{x} \right\} \det(W),\end{aligned}$$

where

$$\begin{aligned}W &= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) \\ -R(G_1)^T & xI_{m_1} - \frac{m_2}{x}J_{m_1 \times m_1} \end{pmatrix} - \begin{pmatrix} O_{n_1 \times n_2} \\ -\frac{1}{x}J_{m_1 \times m_2}R(G_2)^T \end{pmatrix} \\ &\quad \times \left(xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T \right)^{-1} \begin{pmatrix} O_{n_2 \times n_1} & -\frac{1}{x}R(G_2)J_{m_2 \times m_1} \end{pmatrix} \\ &= \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) \\ -R(G_1)^T & xI_{m_1} - \frac{m_2}{x}J_{m_1 \times m_1} - \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)}J_{m_1 \times m_1} \end{pmatrix}.\end{aligned}$$

Then,

$$\begin{aligned}&\det(W) \\ &= \det \left(xI_{m_1} - \frac{m_2}{x}J_{m_1 \times m_1} - \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)}J_{m_1 \times m_1} \right) \\ &\quad \times \det \left(xI_{n_1} - A(G_1) - R(G_1) \right. \\ &\quad \times \left. \left(xI_{m_1} - \frac{m_2}{x}J_{m_1 \times m_1} - \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)}J_{m_1 \times m_1} \right)^{-1} R(G_1)^T \right) \\ &= x^{m_1} \left\{ 1 - \left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) \frac{m_1}{x} \right\} \det \left[xI_{n_1} - A(G_1) \right. \\ &\quad \left. - R(G_1) \left\{ \frac{1}{x}I_{m_1} + \frac{\left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) J_{m_1 \times m_1}}{x \left(x - m_1 \left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) \right)} \right\} R(G_1)^T \right] \\ &= x^{m_1} \left\{ 1 - \left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) \frac{m_1}{x} \right\} \\ &\quad \times \det \left(xI_{n_1} - A(G_1) - \frac{1}{x}R(G_1)R(G_1)^T \right. \\ &\quad \left. - \frac{\left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) R(G_1)J_{m_1 \times m_1}R(G_1)^T}{x \left(x - m_1 \left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) \right)} \right) \\ &= x^{m_1} \left\{ 1 - \left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) \frac{m_1}{x} \right\}\end{aligned}$$

$$\begin{aligned}
& \times \det \left(xI_{n_1} - A(G_1) - \frac{1}{x}R(G_1)R(G_1)^T \right. \\
& \quad \left. - r_1^2 \frac{(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)})}{x(x - m_1(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x))))} J_{n_1 \times n_1} \right) \\
& = x^{m_1} \left\{ 1 - \left(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)} \right) \frac{m_1}{x} \right\} \\
& \quad \times \det \left(xI_{n_1} - A(G_1) - \frac{1}{x}(r_1I_{n_1} + A(G_1)) \right) \\
& \quad \times \left[1 - \frac{r_1^2(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x)})\Gamma_{A(G_1)+\frac{1}{x}R(G_1)R(G_1)^T(x)}}{x(x - m_1(\frac{m_2}{x} + \frac{r_2^2}{x^2}\Gamma_{A(G_2)+\frac{1}{x}R(G_2)R(G_2)^T(x))))} \right] \\
& = x^{m_1} \left\{ 1 - \left(\frac{m_2}{x} + \frac{r_2^2 n_2}{x^2(x - r_2 - \frac{2r_2}{x})} \right) \frac{m_1}{x} \right\} \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{1}{x}(r_1 + \lambda_i(G_1)) \right\} \\
& \quad \times \left[1 - \frac{r_1^2 \left(\frac{m_2}{x} + \frac{r_2^2 n_2}{x^2(x - r_2 - \frac{2r_2}{x})} \right) n_1}{x \left\{ x - m_1 \left(\frac{m_2}{x} + \frac{r_2^2 n_2}{x^2(x - r_2 - \frac{2r_2}{x})} \right) \right\} (x - r_1 - \frac{2r_1}{x})} \right].
\end{aligned}$$

Therefore,

$$\begin{aligned}
f_{A(\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2))}(x) & = x^{m_1} x^{m_2} \left\{ 1 - \left(\frac{m_2}{x} + \frac{r_2^2 n_2}{x^2(x - r_2 - \frac{2r_2}{x})} \right) \frac{m_1}{x} \right\} \\
& \quad \times \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{r_1}{x} - \frac{\lambda_i(G_1)}{x} \right\} \\
& \quad \times \prod_{j=1}^{n_2} \left\{ x - \lambda_j(G_2) - \frac{r_2}{x} - \frac{\lambda_j(G_2)}{x} \right\} \\
& \quad \times \left[1 - \frac{r_1^2 \left(\frac{m_2}{x} + \frac{r_2^2 n_2}{x^2(x - r_2 - \frac{2r_2}{x})} \right) n_1}{x \left\{ x - m_1 \left(\frac{m_2}{x} + \frac{r_2^2 n_2}{x^2(x - r_2 - \frac{2r_2}{x})} \right) \right\} (x - r_1 - \frac{2r_1}{x})} \right] \\
& = x^{m_1-n_1} x^{m_2-n_2} \prod_{i=2}^{n_1} \{x^2 - \lambda_i(G_1)x - r_1 - \lambda_i(G_1)\} \\
& \quad \times \prod_{j=2}^{n_2} \{x^2 - \lambda_j(G_2)x - r_2 - \lambda_j(G_2)\} \\
& \quad \times [x^4 - (r_1 + r_2)x^3 - (2r_1 + m_1m_2 + 2r_2 - r_1r_2)x^2 \\
& \quad + (4r_1r_2 + m_1m_2r_1 + m_1m_2r_2)x + 4r_1r_2 - m_1m_2r_1r_2],
\end{aligned}$$

and hence the result follows. $\square$

Theorem 2.3. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the adjacency spectrum of $\mathcal{R}(G_1) \bar{\vee} \mathcal{R}(G_2)$ consists of:

- (i) two roots of the equation $x^2 - \lambda_i(G_1)x - r_1 - \lambda_i(G_1)$, for every eigenvalue $\lambda_i(G_1)$, $i = 2, 3, \dots, n_1$, of $A(G_1)$;
- (ii) two roots of the equation $x^2 - \lambda_j(G_2)x - r_2 - \lambda_j(G_2)$, for every eigenvalue $\lambda_j(G_2)$, $j = 2, 3, \dots, n_2$, of $A(G_2)$;
- (iii) the eigenvalue 0 with multiplicity $m_1 + m_2 - n_1 - n_2$;
- (iv) four roots of the equation $x^4 - (r_1 + r_2)x^3 - (2r_1 + m_1n_2 + 2r_2 - r_1r_2)x^2 + (4r_1r_2 + r_1m_1n_2)x + 4r_1r_2 = 0$.

Proof. The adjacency characteristic polynomial of $\mathcal{R}(G_1) \bar{\vee} \mathcal{R}(G_2)$ is

$$f_{A(\mathcal{R}(G_1) \bar{\vee} \mathcal{R}(G_2))}(x) = \det \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -R(G_1)^T & xI_{m_1} & -J_{m_1 \times n_2} & O_{m_1 \times m_2} \\ O_{n_2 \times n_1} & -J_{n_2 \times m_1} & xI_{n_2} - A(G_2) & -R(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & -R(G_2)^T & xI_{m_2} \end{pmatrix} \\ = x^{m_2} \det(S),$$

where

$$S = \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & O_{n_1 \times n_2} \\ -R(G_1)^T & xI_{m_1} & -J_{m_1 \times n_2} \\ O_{n_2 \times n_1} & -J_{n_2 \times m_1} & xI_{n_2} - A(G_2) \end{pmatrix} \\ - \begin{pmatrix} O_{n_1 \times m_2} \\ O_{m_1 \times m_2} \\ -R(G_2) \end{pmatrix} \frac{1}{x} \begin{pmatrix} O_{m_2 \times n_1} & O_{m_2 \times m_1} & -R(G_2)^T \end{pmatrix} \\ = \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) & O_{n_1 \times n_2} \\ -R(G_1)^T & xI_{m_1} & -J_{m_1 \times n_2} \\ O_{n_2 \times n_1} & -J_{n_2 \times m_1} & xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T \end{pmatrix}.$$

Hence,

$$\det(S) = \det\left(xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T\right) \det(W) \\ = \prod_{j=1}^{n_2} \left\{x - \lambda_j(G_2) - \frac{r_2}{x} - \frac{\lambda_j(G_2)}{x}\right\} \det(W),$$

where

$$W = \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) \\ -R(G_1)^T & xI_{m_1} \end{pmatrix} \\ - \begin{pmatrix} O_{n_1 \times n_2} \\ -J_{m_1 \times n_2} \end{pmatrix} \left(xI_{n_2} - A(G_2) - \frac{1}{x}R(G_2)R(G_2)^T\right)^{-1} \begin{pmatrix} O_{n_2 \times n_1} & -J_{n_2 \times m_1} \end{pmatrix} \\ = \begin{pmatrix} xI_{n_1} - A(G_1) & -R(G_1) \\ -R(G_1)^T & xI_{m_1} - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T} J_{m_1 \times m_1} \end{pmatrix}.$$

Then,

$$\begin{aligned}
 \det(W) &= \det \left(xI_{m_1} - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T} J_{m_1 \times m_1} \right) \det \left(xI_{n_1} - A(G_1) \right. \\
 &\quad \left. - R(G_1)(xI_{m_1} - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T} J_{m_1 \times m_1})^{-1} R(G_1)^T \right) \\
 &= x^{m_1} \left(1 - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \frac{m_1}{x} \right) \det \left[xI_{n_1} - A(G_1) \right. \\
 &\quad \left. - R(G_1) \left\{ \frac{1}{x} I_{m_1} + \frac{\Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x)}{x(x - m_1 \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x))} J_{m_1 \times m_1} \right\} R(G_1)^T \right] \\
 &= x^{m_1} \left(1 - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \frac{m_1}{x} \right) \det \left(xI_{n_1} - A(G_1) - \frac{1}{x} R(G_1)R(G_1)^T \right. \\
 &\quad \left. - \frac{\Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x)}{x(x - m_1 \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x))} R(G_1)J_{m_1 \times m_1} R(G_1)^T \right) \\
 &= x^{m_1} \left(1 - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \frac{m_1}{x} \right) \det \left(xI_{n_1} - A(G_1) \right. \\
 &\quad \left. - \frac{1}{x} R(G_1)R(G_1)^T - r_1^2 \frac{\Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x)}{x(x - m_1 \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x))} J_{n_1 \times n_1} \right) \\
 &= x^{m_1} \left(1 - \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \frac{m_1}{x} \right) \\
 &\quad \det \left(xI_{n_1} - A(G_1) - \frac{1}{x} (r_1 I_{n_1} + A(G_1)) \right) \\
 &\quad \times \left[1 - \frac{r_1^2 \Gamma_{A(G_1) + \frac{1}{x}R(G_1)R(G_1)^T}(x) \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x)}{x \left(x - m_1 \Gamma_{A(G_2) + \frac{1}{x}R(G_2)R(G_2)^T}(x) \right)} \right] \\
 &= x^{m_1} \left(1 - \frac{m_1 n_2}{x(x - r_2 - \frac{2r_2}{x})} \right) \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{1}{x} (r_1 + \lambda_i(G_1)) \right\} \\
 &\quad \times \left[1 - \frac{r_1^2 n_1 n_2}{x(x - r_1 - \frac{2r_1}{x}) \left(x - \frac{m_1 n_2}{x - r_2 - \frac{2r_2}{x}} \right) (x - r_2 - \frac{2r_2}{x})} \right].
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 f_{A(\mathcal{R}(G_1) \bar{\vee} \mathcal{R}(G_2))}(x) &= x^{m_1} x^{m_2} \left(1 - \frac{m_1 n_2}{x \left(x - r_2 - \frac{2r_2}{x} \right)} \right) \\
 &\quad \times \prod_{i=1}^{n_1} \left\{ x - \lambda_i(G_1) - \frac{r_1}{x} - \frac{\lambda_i(G_1)}{x} \right\} \\
 &\quad \times \prod_{j=1}^{n_2} \left\{ x - \lambda_j(G_2) - \frac{r_2}{x} - \frac{\lambda_j(G_2)}{x} \right\}
 \end{aligned}$$

$$\begin{aligned}
& \times \left[1 - \frac{r_1^2 n_1 n_2}{x \left(x - r_1 - \frac{2r_1}{x} \right) \left(x - \frac{m_1 n_2}{x - r_2 - \frac{2r_2}{x}} \right) \left(x - r_2 - \frac{2r_2}{x} \right)} \right] \\
& = x^{m_1 - n_1} x^{m_2 - n_2} \prod_{i=2}^{n_1} \{x^2 - \lambda_i(G_1)x - r_1 - \lambda_i(G_1)\} \\
& \quad \times \prod_{j=2}^{n_2} \{x^2 - \lambda_j(G_2)x - r_2 - \lambda_j(G_2)\} \left[x^4 - (r_1 + r_2)x^3 \right. \\
& \quad \left. - (2r_1 + m_1 n_2 + 2r_2 - r_1 r_2)x^2 + (4r_1 r_2 + r_1 m_1 n_2)x + 4r_1 r_2 \right],
\end{aligned}$$

and the result follows. $\square$

In the similar way we can obtain Laplacian and normalized Laplacian spectra of the partial join of R -graphs, which are given below.

Lemma 2.2. *We have the following Laplacian matrices:*

$$\begin{aligned}
& \text{(i)} \\
& L(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} (r_1 + n_2)I_{n_1} + L(G_1) & -R(G_1) & -J_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -R(G_1)^T & 2I_{m_1} & O_{m_1 \times n_2} & O_{m_1 \times m_2} \\ -J_{n_2 \times n_1} & O_{n_2 \times m_1} & (r_2 + n_1)I_{n_2} + L(G_2) & -R(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & -R(G_2)^T & 2I_{m_2} \end{pmatrix}; \\
& \text{(ii)} \\
& L(\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} r_1 I_{n_1} + L(G_1) & -R(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -R(G_1)^T & (2 + m_2)I_{m_1} & O_{m_1 \times n_2} & -J_{m_1 \times m_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & r_2 I_{n_2} + L(G_2) & -R(G_2) \\ O_{m_2 \times n_1} & -J_{m_2 \times m_1} & -R(G_2)^T & (2 + m_1)I_{m_2} \end{pmatrix}; \\
& \text{(iii)} \\
& L(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} r_1 I_{n_1} + L(G_1) & -R(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -R(G_1)^T & (2 + n_2)I_{m_1} & -J_{m_1 \times n_2} & O_{m_1 \times m_2} \\ O_{n_2 \times n_1} & -J_{n_2 \times m_1} & (r_2 + m_1)I_{n_2} + L(G_2) & -R(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & -R(G_2)^T & 2I_{m_2} \end{pmatrix}.
\end{aligned}$$

Theorem 2.4. *For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the Laplacian spectrum of $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ consists of:*

- (i) roots of the equation $x^2 - (2 + r_1 + n_2 + \mu_i(G_1))x + 2n_2 + 3\mu_i(G_1) = 0$, for every eigenvalue $\mu_i(G_1)$, $i = 2, 3, \dots, n_1$, of $L(G_1)$;
- (ii) roots of the equation $x^2 - (2 + r_2 + n_1 + \mu_j(G_2))x + 2n_1 + 3\mu_j(G_2) = 0$, for every eigenvalue $\mu_j(G_2)$, $j = 2, 3, \dots, n_2$, of $L(G_2)$;
- (iii) the eigenvalue 2 with multiplicity $m_1 + m_2 - n_1 - n_2$;
- (iv) four roots of the equation $x^4 - (4 + r_1 + r_2 + n_1 + n_2)x^3 + (4 + 4n_1 + 4n_2 + 2r_1 + 2r_2 + r_1 r_2 + r_1 n_1 + r_2 n_2)x^2 - 2(2n_1 + 2n_2 + r_1 n_1 + r_2 n_2)x = 0$.

Theorem 2.5. *For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the Laplacian spectrum of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ consists of:*

- (i) roots of the equation $x^2 - (2 + r_1 + m_2 + \mu_i(G_1))x + r_1m_2 + 3\mu_i(G_1) + m_2\mu_i(G_1) = 0$, for every eigenvalue $\mu_i(G_1)$, $i = 2, 3, \dots, n_1$, of $L(G_1)$;
- (ii) roots of the equation $x^2 - (2 + r_2 + m_1 + \mu_j(G_2))x + r_2m_1 + 3\mu_j(G_2) + m_1\mu_j(G_2) = 0$, for every eigenvalue $\mu_j(G_2)$, $j = 2, 3, \dots, n_2$, of $L(G_2)$;
- (iii) the eigenvalue $2 + m_2$ with multiplicity $m_1 - n_1$;
- (iv) the eigenvalue $2 + m_1$ with multiplicity $m_2 - n_2$;
- (v) four roots of the equation $x^4 - (4 + r_1 + r_2 + m_1 + m_2)x^3 + (4 + 2r_1 + 2r_2 + r_1r_2 + r_1m_1 + r_2m_2 + 2m_1 + 2m_2 + r_1m_2 + r_2m_1)x^2 - (2r_1m_2 + 2r_2m_1 + r_1r_2m_1 + r_1r_2m_2)x = 0$.

Theorem 2.6. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the Laplacian spectrum of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ consists of:

- (i) roots of the equation $x^2 - (2 + r_1 + n_2 + \mu_i(G_1))x + r_1n_2 + 3\mu_i(G_1) + n_2\mu_i(G_1) = 0$, for every eigenvalue $\mu_i(G_1)$, $i = 2, 3, \dots, n_1$, of $L(G_1)$;
- (ii) roots of the equation $x^2 - (2 + r_2 + m_1 + \mu_j(G_2))x + 2m_1 + 3\mu_j(G_2) = 0$, for every eigenvalue $\mu_j(G_2)$, $j = 2, 3, \dots, n_2$, of $L(G_2)$;
- (iii) the eigenvalue $2 + n_2$ with multiplicity $m_1 - n_1$;
- (iv) the eigenvalue 2 with multiplicity $m_2 - n_2$;
- (v) four roots of the equation $x^4 - (4 + r_1 + r_2 + m_1 + n_2)x^3 + (4 + 2r_1 + 2r_2 + 4m_1 + 2n_2 + r_1r_2 + r_1m_1 + r_1n_2 + r_2n_2)x^2 - (4m_1 + 2r_1m_1 + 2r_1n_2 + r_1r_2n_2)x = 0$.

Lemma 2.3. We have the following normalized Laplacian matrices:

(i)

$$\mathcal{L}(\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} \mathcal{L}(G_1) \bullet B(G_1) & -cR(G_1) & -K_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -cR(G_1)^T & I_{m_1} & O_{m_1 \times n_2} & O_{m_1 \times m_2} \\ -K_{n_2 \times n_1} & O_{n_2 \times m_1} & \mathcal{L}(G_2) \bullet B(G_2) & -dR(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & -dR(G_2)^T & I_{m_2} \end{pmatrix},$$

where $K_{n_1 \times n_2}$ is the matrix of size $n_1 \times n_2$ with all entries equal to $\frac{1}{\sqrt{(2r_1+n_2)(2r_2+n_1)}}$, $B(G_1)$ is the $n_1 \times n_1$ matrix whose all diagonal entries are 1 and off-diagonal entries are $\frac{r_1}{2r_1+n_2}$, $B(G_2)$ is the $n_2 \times n_2$ matrix whose all diagonal entries are 1 and off-diagonal entries are $\frac{r_2}{2r_2+n_1}$, c is the constant whose value is $\frac{1}{\sqrt{2(2r_1+n_2)}}$, d is the constant whose value is $\frac{1}{\sqrt{2(2r_2+n_1)}}$;

(ii)

$$\mathcal{L}(\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} \mathcal{L}(G_1) \bullet B(G_1) & -cR(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -cR(G_1)^T & I_{m_1} & O_{m_1 \times n_2} & -K_{m_1 \times m_2} \\ O_{n_2 \times n_1} & O_{n_2 \times m_1} & \mathcal{L}(G_2) \bullet B(G_2) & -dR(G_2) \\ O_{m_2 \times n_1} & -K_{m_2 \times m_1} & -dR(G_2)^T & I_{m_2} \end{pmatrix},$$

where $K_{m_1 \times m_2}$ is the matrix of size $m_1 \times m_2$ with all entries equal to $\frac{1}{\sqrt{(2+m_2)(2+m_1)}}$, $B(G_1)$ is the $n_1 \times n_1$ matrix whose all diagonal entries are 1 and off-diagonal entries are $\frac{r_1}{2r_1}$, $B(G_2)$ is the $n_2 \times n_2$ matrix whose all diagonal entries are 1 and off-diagonal

entries are $\frac{r_2}{2r_2}$, c is the constant whose value is $\frac{1}{\sqrt{2r_1(2+m_2)}}$, d is the constant whose value is $\frac{1}{\sqrt{2r_2(2+m_1)}}$;

(iii)

$$\mathcal{L}(\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)) = \begin{pmatrix} \mathcal{L}(G_1) \bullet B(G_1) & -cR(G_1) & O_{n_1 \times n_2} & O_{n_1 \times m_2} \\ -cR(G_1)^T & I_{m_1} & -K_{m_1 \times n_2} & O_{m_1 \times m_2} \\ O_{n_2 \times n_1} & -K_{n_2 \times m_1} & \mathcal{L}(G_2) \bullet B(G_2) & -dR(G_2) \\ O_{m_2 \times n_1} & O_{m_2 \times m_1} & -dR(G_2)^T & I_{m_2} \end{pmatrix},$$

where $K_{m_1 \times n_2}$ is the matrix of size $m_1 \times n_2$ with all entries equal to $\frac{1}{\sqrt{(2+n_2)(2r_2+m_1)}}$, $B(G_1)$ is the $n_1 \times n_1$ matrix whose all diagonal entries are 1 and off-diagonal entries are $\frac{r_1}{2r_1}$, $B(G_2)$ is the $n_2 \times n_2$ matrix whose all diagonal entries are 1 and off-diagonal entries are $\frac{r_2}{2r_2}$, c is the constant whose value is $\frac{1}{\sqrt{2r_1(2+n_2)}}$, d is the constant whose value is $\frac{1}{\sqrt{2(2r_2+m_1)}}$.

Theorem 2.7. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the normalized Laplacian spectrum of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ consists of:

- (i) roots of the equation $2(2r_1+n_2)x^2 - 2(3r_1+2n_2+r_2\delta_i(G_1))x + 2n_2+3r_1\delta_i(G_1) = 0$, for every eigenvalue $\delta_i(G_1)$, $i = 2, 3, \dots, n_1$, of $\mathcal{L}(G_1)$;
- (ii) roots of the equation $2(2r_2+n_1)x^2 - 2(3r_2+2n_1+r_2\delta_j(G_2))x + 2n_1+3r_2\delta_j(G_2) = 0$, for every eigenvalue $\delta_j(G_2)$, $j = 2, 3, \dots, n_2$, of $\mathcal{L}(G_2)$;
- (iii) the eigenvalue 1 with multiplicity $m_1 + m_2 - n_1 - n_2$;
- (iv) four roots of the equation $(4r_1r_2 + 2r_1n_1 + 2r_2n_2 + n_1n_2)x^4 - (12r_1r_2 + 7r_1n_1 + 7r_2n_2 + 4n_1n_2)x^3 + (9r_1r_2 + 8r_1n_1 + 8r_2n_2 + 5n_1n_2)x^2 - (3r_1n_1 + 3r_2n_2 + 2n_1n_2)x = 0$.

Theorem 2.8. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the normalized Laplacian spectrum of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ consists of:

- (i) roots of the equation $2(2+m_2)x^2 - (6+3m_2+2\delta_i(G_1)+m_2\delta_i(G_1))x + m_2+3\delta_i(G_1)+m_2\delta_i(G_1) = 0$, for every eigenvalue $\delta_i(G_1)$, $i = 2, 3, \dots, n_1$, of $\mathcal{L}(G_1)$;
- (ii) roots of the equation $2(2+m_1)x^2 - (6+3m_1+2\delta_j(G_2)+m_1\delta_j(G_2))x + m_1+3\delta_j(G_2)+m_1\delta_j(G_2) = 0$, for every eigenvalue $\delta_j(G_2)$, $j = 2, 3, \dots, n_2$, of $\mathcal{L}(G_2)$;
- (iii) the eigenvalue 1 with multiplicity $m_1 + m_2 - n_1 - n_2$;
- (iv) four roots of the equation $4(4+2m_1+2m_2+m_1m_2)x^4 - 12(4+2m_1+2m_2+m_1m_2)x^3 + (36+22m_1+22m_2+9m_1m_2)x^2 - 2(3m_1+3m_2+m_1m_2)x = 0$.

Theorem 2.9. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then, the normalized Laplacian spectrum of $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$ consists of:

- (i) roots of the equation $2(2+n_2)x^2 - (6+3n_2+2\delta_i(G_1)+n_2\delta_i(G_1))x + n_2+3\delta_i(G_1)+n_2\delta_i(G_1) = 0$, for every eigenvalue $\delta_i(G_1)$, $i = 2, 3, \dots, n_1$, of $\mathcal{L}(G_1)$;
- (ii) roots of the equation $2(2r_2+m_1)x^2 - 2(3r_2+2m_1+r_2\delta_j(G_2))x + 2m_1+3r_2\delta_j(G_2) = 0$, for every eigenvalue $\delta_j(G_2)$, $j = 2, 3, \dots, n_2$, of $\mathcal{L}(G_2)$;
- (iii) the eigenvalue 1 with multiplicity $m_1 + m_2 - n_1 - n_2$;

- (iv) *four roots of the equation $(8r_2 + 4r_2n_2 + 4m_1 + 2m_1n_2)x^4 - (24r_2 + 12r_2n_2 + 14m_1 + 7m_1n_2)x^3 + (18r_2 + 11r_2n_2 + 16m_1 + 7m_1n_2)x^2 - (3r_2n_2 + 6m_1 + 2m_1n_2)x = 0$.*

3. SIMULTANEOUS COSPECTRAL GRAPHS

In this section we present the main result of the paper. We construct several classes of non-regular graphs which are cospectral with respect to all the three matrices, namely, adjacency, Laplacian and normalized Laplacian. For the construction of these graphs we consider two pairs of A -cospectral regular graphs, which are readily available in the literature, for example see [11]. Then we take partial join of R -graphs belong to different pairs.

The following lemma is immediate from the definition of Laplacian and normalized Laplacian matrices.

Lemma 3.1. (i) *If G is an r -regular graph, then $L(G) = rI_n - A(G)$ and $\mathcal{L}(G) = I_n - \frac{1}{r}A(G)$.*

(ii) *If G_1 and G_2 are A -cospectral regular graphs, then they are also cospectral with respect to the Laplacian and normalized Laplacian matrices.*

Observation. From all the theorems given in the previous section we observe that the adjacency, Laplacian and normalized Laplacian spectra of all the partial join graphs $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$, $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$, and $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$, depend only on the number of vertices, number of edges, degree of regularities, and the corresponding spectrum of G_1 and G_2 . Furthermore, we note that, although G_1 and G_2 are regular graphs, $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$, $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$, and $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ are non-regular graphs.

The following theorem is the main result of the paper.

Theorem 3.1. *Let $G_i, H_i, i = 1, 2$, be regular graphs, where G_1 need not be different from H_1 . If G_1 and H_1 are A -cospectral, and G_2 and H_2 are A -cospectral then $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$ (respectively, $\mathcal{R}(G_1)\bar{\vee}\mathcal{R}(G_2)$, $\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)$) and $\mathcal{R}(H_1)\check{\vee}\mathcal{R}(H_2)$ (respectively, $\mathcal{R}(H_1)\bar{\vee}\mathcal{R}(H_2)$, $\mathcal{R}(H_1)\check{\vee}\mathcal{R}(H_2)$) are simultaneously A -cospectral, L -cospectral and $\mathcal{L}$ -cospectral.*

Proof. Follows from Lemma 3.1 and the above observation. $\square$

4. SPANNING TREES AND KIRCHHOFF INDICES

Applying the results on Laplacian spectra, we find the number of spanning trees and Kirchhoff index of all the partial join graphs constructed in the paper.

Let $t(G)$ denote the number of *spanning trees* of G . It is well known [6] that if G is a connected graph on n vertices with Laplacian spectrum $0 = \mu_1(G) \leq \mu_2(G) \leq \dots \leq \mu_n(G)$, then $t(G) = \frac{\mu_2(G) \cdots \mu_n(G)}{n}$.

The *Kirchhoff index* of a graph G , denoted by $Kf(G)$, is defined as the sum of resistances between all pairs of vertices $[1, 9]$ in G . For a connected graph G on n vertices, the Kirchhoff index [8] can be expressed as $Kf(G) = n \sum_{i=2}^n \frac{1}{\mu_i(G)}$.

Theorem 4.1. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then,

$$\begin{aligned}
 & \text{(i)} \\
 & t(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)) \\
 &= \frac{2^{m_1+m_2-n_1-n_2} \cdot 2(2n_1+2n_2+r_1n_1+r_2n_2) \cdot \prod_{i=2}^{n_1} (2n_2+3\mu_i(G_1)) \cdot \prod_{j=2}^{n_2} (2n_1+3\mu_j(G_2))}{n_1+n_2+m_1+m_2}; \\
 & \text{(ii)} \\
 & t(\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)) \\
 &= (2+m_2)^{m_1-n_1} \cdot (2+m_1)^{m_2-n_2} \\
 & \quad \times \frac{(2r_1m_2+2r_2m_1+r_1r_2m_1+r_1r_2m_2) \cdot \prod_{i=2}^{n_1} (r_1m_2+3\mu_i(G_1)+m_2\mu_i(G_1)) \cdot \prod_{j=2}^{n_2} (r_2m_1+3\mu_j(G_2)+m_1\mu_j(G_2))}{n_1+n_2+m_1+m_2}; \\
 & \text{(iii)} \\
 & t(\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)) \\
 &= (2+n_2)^{m_1-n_1} \cdot 2^{m_2-n_2} \cdot (4m_1+2r_1m_1+2r_1n_2+r_1r_2n_2) \\
 & \quad \times \frac{\prod_{i=2}^{n_1} (r_1n_2+3\mu_i(G_1)+n_2\mu_i(G_1)) \cdot \prod_{j=2}^{n_2} (2m_1+3\mu_j(G_2))}{n_1+n_2+m_1+m_2}.
 \end{aligned}$$

Theorem 4.2. For $i = 1, 2$, let G_i be an r_i -regular graph with n_i vertices and m_i edges. Then,

$$\begin{aligned}
 & \text{(i)} \\
 & Kf(\mathcal{R}(G_1)\check{\vee}\mathcal{R}(G_2)) \\
 &= (n_1+n_2+m_1+m_2) \left(\frac{m_1+m_2-n_1-n_2}{2} \right. \\
 & \quad + \frac{4+4n_1+4n_2+2r_1+2r_2+r_1r_2+r_1n_1+r_2n_2}{2(2n_1+2n_2+r_1n_1+r_2n_2)} + \sum_{i=2}^{n_1} \frac{2+r_1+n_2+\mu_i(G_1)}{2n_2+3\mu_i(G_1)} \\
 & \quad \left. + \sum_{j=2}^{n_2} \frac{2+r_2+n_1+\mu_j(G_2)}{2n_1+3\mu_j(G_2)} \right); \\
 & \text{(ii)} \\
 & Kf(\mathcal{R}(G_1)\overline{\vee}\mathcal{R}(G_2)) \\
 &= (n_1+n_2+m_1+m_2) \left(\frac{m_1-n_1}{2+m_2} + \frac{m_2-n_2}{2+m_1} + \sum_{i=2}^{n_1} \frac{2+r_1+m_2+\mu_i(G_1)}{r_1m_2+3\mu_i(G_1)+m_2\mu_i(G_1)} \right. \\
 & \quad \left. + \frac{4+2r_1+2r_2+r_1r_2+r_1m_1+r_2m_2+2m_1+2m_2+r_1m_2+r_2m_1}{2r_1m_2+2r_2m_1+r_1r_2m_1+r_1r_2m_2} \right)
 \end{aligned}$$

$$(iii) \quad + \sum_{j=2}^{n_2} \frac{2 + r_2 + m_1 + \mu_j(G_2)}{r_2 m_1 + 3\mu_j(G_2) + m_1 \mu_j(G_2)} \Bigg);$$

$$\begin{aligned} & Kf(\mathcal{R}(G_1) \bar{\vee} \mathcal{R}(G_2)) \\ &= (n_1 + n_2 + m_1 + m_2) \left(\frac{m_1 - n_1}{2 + n_2} + \frac{m_2 - n_2}{2} \right. \\ &\quad + \frac{4 + 2r_1 + 2r_2 + 4m_1 + 2n_2 + r_1 r_2 + r_1 m_1 + r_1 n_2 + r_2 n_2}{4m_1 + 2r_1 m_1 + 2r_1 n_2 + r_1 r_2 n_2} \\ &\quad \left. + \sum_{i=2}^{n_1} \frac{2 + r_1 + n_2 + \mu_i(G_1)}{r_1 n_2 + 3\mu_i(G_1) + n_2 \mu_i(G_1)} + \sum_{j=2}^{n_2} \frac{2 + r_2 + m_1 + \mu_j(G_2)}{2m_1 + 3\mu_j(G_2)} \right). \end{aligned}$$

5. CONCLUDING REMARKS

The main result of the paper is based on regular A -cospectral graphs and certain operations on a pair of these graphs so that the operated (or resultant) graphs are non-regular and their adjacency, Laplacian and normalized Laplacian spectra depend on only the order, size, degree of regularity and spectra of the original graphs. Thus following the technique of this paper one may construct some more graphs like here.

REFERENCES

- [1] D. Bonchev, A. T. Balaban, X. Liu and D. J. Klein, *Molecular cyclicity and centrality of polycyclic graphs. I. Cyclicity based on resistance distances or reciprocal distances*, Internat. J. Quant. Chem. **50** (1994), 1–20. <https://doi.org/10.1002/qua.560500102>
- [2] S. Butler, *A note about cospectral graphs for the adjacency and normalized Laplacian matrices*, Linear Multilinear Algebra **58** (2010), 387–390. <https://doi.org/10.1080/03081080902722741>
- [3] S. Butler and J. Grout, *A construction of cospectral graphs for the normalized Laplacian*, Electron. J. Combin. **18** (2011), 20 pages. <https://doi.org/10.37236/718>
- [4] F. R. K. Chung, *Spectral Graph Theory*, CBMS. Reg. Conf. Ser. Math. 92, AMS, Providence, RI, 1997.
- [5] S. Y. Cui and G. X. Tian, *The spectrum and the signless Laplacian spectrum of coronae*, Linear Algebra Appl. **437** (2012), 1692–1703. <https://doi.org/10.1016/j.laa.2012.05.019>
- [6] D. M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs-Theory and Applications*, Third edition, Johann Ambrosius Barth. Heidelberg, 1995.
- [7] D. Cvetković, P. Rowlinson and S. Simić, *An Introduction to the Theory of Graph Spectra*, Cambridge University Press, 2009.
- [8] I. Gutman and B. Mohar, *The quasi-Wiener and the Kirchhoff indices coincide*, J. Chem. Inf. Comput. Sci. **36** (1996), 982–985.
- [9] D. J. Klein and M. Randić, *Resistance distance*, J. Math. Chem. **12** (1993), 81–95. <https://doi.org/10.1007/BF01164627>
- [10] C. McLeman and E. McNicholas, *Spectra of coronae*, Linear Algebra Appl. **435** (2011), 998–1007. <https://doi.org/10.1016/j.laa.2011.02.007>
- [11] E. R. van Dam and W. H. Haemers, *Which graphs are determined by their spectrum?*, Linear Algebra Appl. **373** (2003), 241–272. [https://doi.org/10.1016/S0024-3795\(03\)00483-X](https://doi.org/10.1016/S0024-3795(03)00483-X)

¹DEPARTMENT OF MATHEMATICS,
BETHUADAHARI COLLEGE,
NADIA,INDIA-741126
Email address: arpita.das1201@gmail.com

²DEPARTMENT OF MATHEMATICS,
INDIAN INSTITUTE OF TECHNOLOGY KHARAGPUR,
KHARAGPUR,INDIA-721302
Email address: pratima@maths.iitkgp.ernet.in
ORCID id: <https://orcid.org/0000-0002-3460-9506>