

**STRONG CONVERGENCE OF AN IMPLICIT ITERATION
PROCESS FOR ASYMPTOTICALLY NONEXPANSIVE IN THE
INTERMEDIATE SENSE MAPPINGS IN BANACH SPACES**

G. S. SALUJA

ABSTRACT. The aim of this article is to study an implicit iteration process with errors for a finite family of non-Lipschitzian asymptotically nonexpansive in the intermediate sense mappings in Banach spaces. Also we have established some strong convergence theorems for said scheme to converge to common fixed point. The results presented in this paper extend and improve the corresponding results of [1], [3]-[11], [13], [14], [16] and many others.

1. INTRODUCTION AND PRELIMINARIES

Let C be a nonempty subset of a real Banach space E . Let $T: C \rightarrow C$ be a mapping. We use $F(T)$ to denote the set of fixed points of T , that is, $F(T) = \{x \in C : Tx = x\}$. Recall the following concepts.

(1) T is nonexpansive if

$$(1.1) \quad \|Tx - Ty\| \leq \|x - y\|,$$

for all $x, y \in C$.

(2) T is asymptotically nonexpansive if there exists a sequence $\{a_n\}$ in $[1, \infty)$ with $a_n \rightarrow 1$ as $n \rightarrow \infty$ such that

$$(1.2) \quad \|T^n x - T^n y\| \leq a_n \|x - y\|,$$

for all $x, y \in C$ and $n \geq 1$.

(3) T is uniformly L -Lipschitzian if there exists a constant $L > 0$ such that

$$(1.3) \quad \|T^n x - T^n y\| \leq L \|x - y\|,$$

Key words and phrases. Asymptotically nonexpansive in the intermediate sense mapping, common fixed point, implicit iteration process with errors, Banach space, strong convergence.

2010 Mathematics Subject Classification. Primary: 47H09, Secondary: 47H10.

Received: August 21, 2011.

for all $x, y \in C$ and $n \geq 1$.

It is clear that every nonexpansive mapping is asymptotically nonexpansive and every asymptotically nonexpansive is uniformly Lipschitzian.

The class of asymptotically nonexpansive mappings was introduced by Goebel and Kirk [4] as a generalization of the class of nonexpansive mappings. T is said to be asymptotically nonexpansive in the intermediate sense [2] if it is continuous and the following inequality holds:

$$(1.4) \quad \limsup_{n \rightarrow \infty} \sup_{x, y \in C} \left(\|T^n x - T^n y\| - \|x - y\| \right) \leq 0.$$

From the above definitions, it follows that asymptotically nonexpansive mappings must be asymptotically nonexpansive in the intermediate sense and asymptotically quasi-nonexpansive mapping. But the converse does not hold as the following example shows:

Example 1.1. Let $X = \mathbb{R}$ be a normed linear space and $K = [0, 1]$. For each $x \in K$, we define

$$T(x) = \begin{cases} kx, & \text{if } x \neq 0, \\ 0, & \text{if } x = 0, \end{cases}$$

where $0 < k < 1$. Then

$$|T^n x - T^n y| = k^n |x - y| \leq |x - y|$$

for all $x, y \in K$ and $n \in \mathbb{N}$.

Thus T is an asymptotically nonexpansive mapping with constant sequence $\{1\}$ and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \{|T^n x - T^n y| - |x - y|\} &= \limsup_{n \rightarrow \infty} \{k^n |x - y| - |x - y|\} \\ &\leq 0 \end{aligned}$$

because $\lim_{n \rightarrow \infty} k^n = 0$ as $0 < k < 1$ and for all $x, y \in K$, $n \in \mathbb{N}$. Hence T is an asymptotically nonexpansive in the intermediate sense mapping.

Example 1.2. Let $X = \mathbb{R}$, $K = [-\frac{1}{\pi}, \frac{1}{\pi}]$ and $|k| < 1$. For each $x \in K$, define

$$T(x) = \begin{cases} kx \sin(1/x), & \text{if } x \neq 0, \\ 0, & \text{if } x = 0. \end{cases}$$

Then T is an asymptotically nonexpansive in the intermediate sense mapping but it is not asymptotically nonexpansive mapping.

Let E be a Hilbert space, let K be a nonempty closed convex subset of E , and let $\{T_1, T_2, \dots, T_N\}: K \rightarrow K$ be N nonexpansive mappings. In 2001, Xu and Ori [17] introduced the following implicit iteration process $\{x_n\}$ defined by

$$(1.5) \quad x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{n(\text{mod } N)} x_n, \quad n \geq 1,$$

where $x_0 \in K$ is an initial point, $\{\alpha_n\}_{n \geq 1}$ is a real sequence in $(0, 1)$ and proved the weakly convergence of the sequence $\{x_n\}$ defined by (1.5) to a common fixed point $p \in F = \bigcap_{i=1}^N F(T_i)$.

In 2003, Sun [13] introduced the following implicit iterative sequence $\{x_n\}$

$$(1.6) \quad x_n = (1 - \alpha_n)x_{n-1} + \alpha_n T_{i(n)}^{k(n)} x_n, \quad n \geq 1,$$

for a finite family of asymptotically quasi-nonexpansive self-mappings on a bounded closed convex subset K of a Hilbert space E with $\{\alpha_n\}$ a sequence in $(0, 1)$ and an initial point $x_0 \in K$, where $n = (k(n) - 1)N + i(n)$, $i(n) \in \{1, 2, \dots, N\}$, and proved the strong convergence of the sequence $\{x_n\}$ defined by (1.6) to a common fixed point $p \in F = \bigcap_{i=1}^N F(T_i)$.

Recently, in 2006, Gu [6] introduced the following implicit iterative sequence $\{x_n\}$ with errors

$$(1.7) \quad \begin{aligned} x_n &= (1 - \alpha_n)x_{n-1} + \alpha_n T_{i(n)}^{k(n)} y_n + u_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n T_{i(n)}^{k(n)} x_n + v_n, \quad n \geq 1, \end{aligned}$$

for a finite family of asymptotically nonexpansive mappings on a closed convex subset K of a Banach space X with $K + K \subset K$, $\{\alpha_n\}$ and $\{\beta_n\}$ be two sequences in $[0, 1]$, $\{u_n\}$ and $\{v_n\}$ be two sequences in K , and an initial point $x_0 \in K$, where $n = (k(n) - 1)N + i(n)$, $i(n) \in \{1, 2, \dots, N\}$, and proved the weak and strong convergence of the sequence $\{x_n\}$ defined by (1.7) to a common fixed point $p \in F = \bigcap_{i=1}^N F(T_i)$.

It should be pointed out that the sequence defined by (1.6) is a special case of the sequence defined by (1.7) with $u_n = v_n = 0$, $\beta_n = 0$, for all $n \geq 1$.

Recently concerning the convergence problems of an implicit (or nonimplicit) iterative process to a common fixed point for a finite family of asymptotically nonexpansive mappings (or nonexpansive mappings) in Hilbert spaces or uniformly convex Banach spaces have been considered by several authors (see, e.g., Bauschke [1], Chang and Cho [3], Goebel and Kirk [4], Gornicki [5], Gu [6], Halpern [7], Lions [8], Osilike [9], Reich [11], Schu [12], Sun [13], Tan and Xu [14], Wittmann [16], Xu and Ori [17] and Zhou and Chang [18]).

Inspired and motivated by [6, 13, 17] and many others, we introduce the following implicit iterative sequence $\{x_n\}$ with errors defined by

$$(1.8) \quad \begin{aligned} x_n &= \alpha_n x_{n-1} + \beta_n T_{i(n)}^{k(n)} y_n + \gamma_n u_n, \\ y_n &= \hat{\alpha}_n x_{n-1} + \hat{\beta}_n T_{i(n)}^{k(n)} x_n + \hat{\gamma}_n v_n, \quad n \geq 1, \end{aligned}$$

where $n = (k(n) - 1)N + i(n)$, $i(n) \in \{1, 2, \dots, N\} = I$, is called the implicit iterative sequence for a finite family of asymptotically nonexpansive in the intermediate sense mappings $\{T_i\}_{i=1}^N$, where $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\hat{\alpha}_n\}$, $\{\hat{\beta}_n\}$ and $\{\hat{\gamma}_n\}$ are six sequences in $[0, 1]$ satisfying $\alpha_n + \beta_n + \gamma_n = \hat{\alpha}_n + \hat{\beta}_n + \hat{\gamma}_n = 1$ for all $n \geq 1$, x_0 is a given point in K , as well as $\{u_n\}$ and $\{v_n\}$ are two bounded sequences in K .

The purpose of this article is to study an implicit iterative sequence with errors defined by (1.8) for a finite family of asymptotically nonexpansive in the intermediate sense mappings in Banach spaces and establish the strong convergence theorems for said iteration scheme and mappings.

In what follows we need the following lemmas to prove our main results:

Lemma 1.1. (see [15]) *Let $\{a_n\}$ and $\{b_n\}$ be sequences of nonnegative real numbers satisfying the inequality*

$$a_{n+1} \leq a_n + b_n, \quad n \geq 1.$$

If $\sum_{n=1}^{\infty} b_n < \infty$, then $\lim_{n \rightarrow \infty} a_n$ exists. In particular, if $\{a_n\}$ has a subsequence converging to zero, then $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 1.2. (Schu [12]) *Let E be a uniformly convex Banach space and $0 < a \leq t_n \leq b < 1$ for all $n \geq 1$. Suppose that $\{x_n\}$ and $\{y_n\}$ are sequences in E satisfying $\limsup_{n \rightarrow \infty} \|x_n\| \leq r$, $\limsup_{n \rightarrow \infty} \|y_n\| \leq r$, $\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n)y_n\| = r$, for some $r \geq 0$. Then $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.*

Lemma 1.3. *Let E be a real Banach space and C be a nonempty closed convex subset of E . Let $\{T_i\}_{i=1}^N: C \rightarrow C$ be N asymptotically nonexpansive in the intermediate sense mappings with $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Put*

$$(1.9) \quad A_n = \max \left\{ 0, \sup_{x, y \in C, n \geq 1} \left(\|T_i^n x - T_i^n y\| - \|x - y\| \right) : i \in I \right\},$$

where $n = (k - 1)N + i$, $k = k(n)$, $i = i(n) \in \{1, 2, \dots, N\} = I$, such that $\sum_{n=1}^{\infty} A_n < \infty$. Let $\{u_n\}$ and $\{v_n\}$ be two bounded sequences in C , and let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\hat{\alpha}_n\}$, $\{\hat{\beta}_n\}$ and $\{\hat{\gamma}_n\}$ be six sequences in $[0, 1]$ satisfying the following conditions:

- (i) $\alpha_n + \beta_n + \gamma_n = \hat{\alpha}_n + \hat{\beta}_n + \hat{\gamma}_n = 1$;
- (ii) $\tau = \sup\{\beta_n : n \geq 1\} < 1$;
- (iii) $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \hat{\gamma}_n < \infty$.

If $\{x_n\}$ is an implicit iterative sequence defined by (1.8), then for each $p \in F = \bigcap_{i=1}^N F(T_i)$ the limit $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists.

Proof. Since $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$, for any given $p \in F$, it follows from (1.8) and (1.9) that

$$(1.10) \quad \begin{aligned} \|x_n - p\| &= \left\| (1 - \beta_n - \gamma_n)x_{n-1} + \beta_n T_n^k y_n + \gamma_n u_n - p \right\| \\ &\leq (1 - \beta_n - \gamma_n) \|x_{n-1} - p\| + \beta_n \|T_n^k y_n - p\| \\ &\quad + \gamma_n \|u_n - p\| \end{aligned}$$

where $n = (k - 1)N + i$, $k = k(n)$, $i = i(n) \in \{1, 2, \dots, N\} = I$ and $T_n = T_{i(\text{mod } N)} = T_i$. This implies that

$$\begin{aligned}
 \|x_n - p\| &\leq (1 - \beta_n - \gamma_n) \|x_{n-1} - p\| + \beta_n \|T_i^k y_n - p\| \\
 &\quad + \gamma_n \|u_n - p\| \\
 &= (1 - \beta_n - \gamma_n) \|x_{n-1} - p\| + \beta_n \|T_i^k y_n - T_i^k p\| \\
 &\quad + \gamma_n \|u_n - p\| \\
 &\leq (1 - \beta_n - \gamma_n) \|x_{n-1} - p\| + \beta_n [\|y_n - p\| + A_n] \\
 &\quad + \gamma_n \|u_n - p\| \\
 &\leq (1 - \beta_n) \|x_{n-1} - p\| + \beta_n \|y_n - p\| + A_n \\
 (1.11) \quad &\quad + \gamma_n \|u_n - p\|.
 \end{aligned}$$

Again it follows from (1.8) and (1.9) that

$$\begin{aligned}
 \|y_n - p\| &\leq (1 - \hat{\beta}_n - \hat{\gamma}_n) \|x_{n-1} - p\| + \hat{\beta}_n \|T_i^k x_n - p\| \\
 &\quad + \hat{\gamma}_n \|v_n - p\| \\
 &= (1 - \hat{\beta}_n - \hat{\gamma}_n) \|x_{n-1} - p\| + \hat{\beta}_n \|T_i^k x_n - T_i^k p\| \\
 &\quad + \hat{\gamma}_n \|v_n - p\| \\
 &\leq (1 - \hat{\beta}_n) \|x_{n-1} - p\| + \hat{\beta}_n [\|x_n - p\| + A_n] \\
 &\quad + \hat{\gamma}_n \|v_n - p\| \\
 &\leq (1 - \hat{\beta}_n) \|x_{n-1} - p\| + \hat{\beta}_n \|x_n - p\| + A_n \\
 (1.12) \quad &\quad + \hat{\gamma}_n \|v_n - p\|.
 \end{aligned}$$

Substituting (1.12) into (1.11), we obtain that

$$\begin{aligned}
 \|x_n - p\| &\leq (1 - \beta_n \hat{\beta}_n) \|x_{n-1} - p\| + \beta_n \hat{\beta}_n \|x_n - p\| + (\beta_n + 1)A_n \\
 &\quad + \beta_n \hat{\gamma}_n \|v_n - p\| + \gamma_n \|u_n - p\| \\
 &\leq (1 - \beta_n \hat{\beta}_n) \|x_{n-1} - p\| + \beta_n \hat{\beta}_n \|x_n - p\| + 2A_n \\
 &\quad + \beta_n \hat{\gamma}_n \|v_n - p\| + \gamma_n \|u_n - p\|
 \end{aligned}$$

which implies that

$$(1.13) \quad (1 - \beta_n \hat{\beta}_n) \|x_n - p\| \leq (1 - \beta_n \hat{\beta}_n) \|x_{n-1} - p\| + \mu_n,$$

where $\mu_n = 2A_n + \beta_n \hat{\gamma}_n \|v_n - p\| + \gamma_n \|u_n - p\|$. By assumption $\sum_{n=1}^{\infty} A_n < \infty$, condition (iii) and boundedness of the sequences $\{\beta_n\}$, $\{\|u_n - p\|\}$, and $\{\|v_n - p\|\}$, we have $\sum_{n=1}^{\infty} \mu_n < \infty$. From condition (ii) we know that

$$\beta_n \hat{\beta}_n \leq \beta_n \leq \tau < 1,$$

and so

$$(1.14) \quad 1 - \beta_n \hat{\beta}_n \geq 1 - \tau > 0,$$

hence from (1.13) we have

$$\begin{aligned}
 \|x_n - p\| &\leq \|x_{n-1} - p\| + \frac{\mu_n}{1 - \beta_n \hat{\beta}_n} \\
 &\leq \|x_{n-1} - p\| + \frac{\mu_n}{1 - \tau} \\
 (1.15) \qquad &= \|x_{n-1} - p\| + B_n,
 \end{aligned}$$

where

$$B_n = \frac{\mu_n}{1 - \tau}.$$

By assumption of the theorem and condition (iii) we have that

$$\sum_{n=1}^{\infty} B_n = \sum_{n=1}^{\infty} \frac{\mu_n}{1 - \tau} < \infty.$$

Taking $A_n = \|x_{n-1} - p\|$ in inequality (1.15), we have

$$A_{n+1} \leq A_n + B_n, \quad \forall n \geq 1,$$

and satisfied all conditions in Lemma 1.1. Therefore the limit $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. Without loss of generality we may assume that $\lim_{n \rightarrow \infty} \|x_n - p\| = d$, where $d \geq 0$ is some nonnegative number. This completes the proof of Lemma 1.3. \square

2. MAIN RESULTS

We are now in a position to prove our main results in this paper.

Theorem 2.1. *Let E be a real Banach space and C be a nonempty closed convex subset of E . Let $\{T_i\}_{i=1}^N : C \rightarrow C$ be N asymptotically nonexpansive in the intermediate sense mappings with $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$. Put*

$$A_n = \max \left\{ 0, \sup_{x, y \in C, n \geq 1} \left(\|T_i^n x - T_i^n y\| - \|x - y\| \right) : i \in I \right\},$$

where $n = (k - 1)N + i$, $k = k(n)$, $i = i(n) \in \{1, 2, \dots, N\} = I$, such that $\sum_{n=1}^{\infty} A_n < \infty$. Let $\{u_n\}$ and $\{v_n\}$ be two bounded sequences in C , and let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\hat{\alpha}_n\}$, $\{\hat{\beta}_n\}$ and $\{\hat{\gamma}_n\}$ be six sequences in $[0, 1]$ satisfying the following conditions:

- (i) $\alpha_n + \beta_n + \gamma_n = \hat{\alpha}_n + \hat{\beta}_n + \hat{\gamma}_n = 1$;
- (ii) $\tau = \sup\{\beta_n : n \geq 1\} < 1$;
- (iii) $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \hat{\gamma}_n < \infty$.

Then the implicit iterative sequence $\{x_n\}$ defined by (1.8) converges strongly to a common fixed point $p \in F = \bigcap_{i=1}^N F(T_i)$ if and only if

$$(2.1) \qquad \liminf_{n \rightarrow \infty} d(x_n, F) = 0.$$

Proof. The necessity of condition (2.1) is obvious.

Next we prove the sufficiency of Theorem 2.1. For any given $p \in F$, it follows from (1.15) in Lemma 1.3 that

$$(2.2) \quad \|x_n - p\| \leq \|x_{n-1} - p\| + B_n, \quad \forall n \geq 1,$$

where

$$B_n = \frac{\mu_n}{1 - \tau},$$

with $\sum_{n=1}^{\infty} B_n < \infty$. Hence, we have

$$(2.3) \quad d(x_n, F) \leq d(x_{n-1}, p) + B_n, \quad \forall n \geq 1.$$

It follows from (2.3) and Lemma 1.1 that the limit $\lim_{n \rightarrow \infty} d(x_n, F)$ exists. By the condition (2.1), we have

$$\lim_{n \rightarrow \infty} d(x_n, F) = 0.$$

Next, we prove that the sequence $\{x_n\}$ is a Cauchy sequence in C . For any integer $m \geq 1$, we have from (2.2) that

$$(2.4) \quad \begin{aligned} \|x_{n+m} - p\| &\leq \|x_{n+m-1} - p\| + B_{n+m-1} \\ &\leq \|x_{n+m-2} - p\| + B_{n+m-2} + B_{n+m-1} \\ &\leq \dots \\ &\leq \|x_n - p\| + \sum_{k=n}^{n+m-1} B_k. \end{aligned}$$

Since $\liminf_{n \rightarrow \infty} d(x_n, F) = 0$, without loss of generality, we may assume that a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ and a sequence $\{p_{n_k}\} \subset F$ such that $\|x_{n_k} - p_{n_k}\| \rightarrow 0$ as $k \rightarrow \infty$. Then for any $\varepsilon > 0$, there exists $k_\varepsilon > 0$ such that

$$(2.5) \quad \|x_{n_k} - p_{n_k}\| < \frac{\varepsilon}{4} \quad \text{and} \quad \sum_{k=n_{k_\varepsilon}}^{\infty} B_k < \frac{\varepsilon}{4},$$

for all $k \geq k_\varepsilon$.

For any $m \geq 1$ and for all $n \geq n_{k_\varepsilon}$, by (2.5), we have

$$(2.6) \quad \begin{aligned} \|x_{n+m} - x_n\| &\leq \|x_{n+m} - p_{n_k}\| + \|x_n - p_{n_k}\| \\ &\leq \|x_{n_k} - p_{n_k}\| + \sum_{k=n_{k_\varepsilon}}^{\infty} B_k \\ &\quad + \|x_{n_k} - p_{n_k}\| + \sum_{k=n_{k_\varepsilon}}^{\infty} B_k \\ &= 2\|x_{n_k} - p_{n_k}\| + 2\sum_{k=n_{k_\varepsilon}}^{\infty} B_k \\ &< 2 \cdot \frac{\varepsilon}{4} + 2 \cdot \frac{\varepsilon}{4} = \varepsilon. \end{aligned}$$

This implies that $\{x_n\}$ is a Cauchy sequence in C . By the completeness of C , we can assume that $\lim_{n \rightarrow \infty} x_n = p^*$. Since the set of fixed points of an asymptotically nonexpansive in the intermediate sense mapping is closed, hence F is closed. This implies that $p^* \in F$, and so p^* is a common fixed point of the mappings $\{T_i\}_{i=1}^N$. This completes the proof of Theorem 2.1. \square

Theorem 2.2. *Let E be a real uniformly convex Banach space, C be a nonempty closed convex subset of E . Let $\{T_i\}_{i=1}^N: C \rightarrow C$ be N asymptotically nonexpansive in the intermediate sense mappings with $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$ and there exists an T_l , $1 \leq l \leq N$, which is semi-compact (without loss of generality, we can assume that T_1 is semi-compact). Put*

$$A_n = \max \left\{ 0, \sup_{x, y \in C, n \geq 1} \left(\|T_i^n x - T_i^n y\| - \|x - y\| \right) : i \in I \right\},$$

where $n = (k - 1)N + i$, $k = k(n)$, $i = i(n) \in \{1, 2, \dots, N\} = I$, such that $\sum_{n=1}^{\infty} A_n < \infty$. Let $\{u_n\}$ and $\{v_n\}$ be two bounded sequences in C , and let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\hat{\alpha}_n\}$, $\{\hat{\beta}_n\}$ and $\{\hat{\gamma}_n\}$ be six sequences in $[0, 1]$ satisfying the following conditions:

- (i) $\alpha_n + \beta_n + \gamma_n = \hat{\alpha}_n + \hat{\beta}_n + \hat{\gamma}_n = 1$;
- (ii) $\tau = \sup\{\beta_n : n \geq 1\} < 1$;
- (iii) $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \hat{\gamma}_n < \infty$;
- (iv) $0 < \tau_1 = \inf\{\beta_n : n \geq 1\} \leq \sup\{\beta_n : n \geq 1\} = \tau_2 < 1$;
- (v) $\hat{\beta}_n \rightarrow 0$, $(n \rightarrow \infty)$;
- (vi) $0 \leq \delta = \sup\{\hat{\beta}_n : n \geq 1\} < 1$;
- (vii) there exists a constant $L > 0$ such that, for any $i \in \{1, 2, \dots, N\}$, we have

$$\|T_i^n x - T_i^n y\| \leq L \|x - y\|, \forall n \geq 1,$$

for all $x, y \in C$.

Then the implicit iterative sequence $\{x_n\}$ defined by (1.8) converges strongly to a common fixed point of the mappings $\{T_i\}_{i=1}^N$.

Proof. First, we prove that

$$(2.7) \quad \lim_{n \rightarrow \infty} \|x_n - T_l x_n\| = 0, \quad \forall l = 1, 2, \dots, N.$$

Let $p \in F$. Put $d = \|x_n - p\|$, where $d \geq 0$ is some nonnegative number. It follows from (1.8) that

$$(2.8) \quad \begin{aligned} \|x_n - p\| &= \|(1 - \beta_n)[x_{n-1} - p + \gamma_n(u_n - x_{n-1})] \\ &\quad + \beta_n[T_i^k y_n - p + \gamma_n(u_n - x_{n-1})]\| \\ &\rightarrow d, \text{ as } n \rightarrow \infty. \end{aligned}$$

Again since $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists, so $\{x_n\}$ is a bounded sequence in C . By virtue of condition (iii) and the boundedness of sequences $\{x_n\}$ and $\{u_n\}$ we have

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \|x_{n-1} - p + \gamma_n(u_n - x_{n-1})\| &\leq \limsup_{n \rightarrow \infty} \|x_{n-1} - p\| \\
 &\quad + \limsup_{n \rightarrow \infty} \gamma_n \|u_n - x_{n-1}\| \\
 (2.9) \qquad \qquad \qquad &= d, \quad p \in F.
 \end{aligned}$$

It follows from (1.12) and condition (v) that

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \|T_n^k y_n - p + \gamma_n(u_n - x_{n-1})\| &\leq \limsup_{n \rightarrow \infty} \|T_n^k y_n - p\| \\
 &\quad + \limsup_{n \rightarrow \infty} \gamma_n \|u_n - x_{n-1}\| \\
 &\leq \limsup_{n \rightarrow \infty} [\|y_n - p\| + A_n] \\
 &\leq \limsup_{n \rightarrow \infty} [(1 - \hat{\beta}_n) \|x_{n-1} - p\| \\
 &\quad + \hat{\beta}_n \|x_n - p\| + A_n \\
 &\quad + \hat{\gamma}_n \|v_n - p\| + A_n] \\
 &\leq \limsup_{n \rightarrow \infty} (1 - \hat{\beta}_n) \|x_{n-1} - p\| \\
 &\quad + \limsup_{n \rightarrow \infty} \hat{\beta}_n \|x_n - p\| + 2 \limsup_{n \rightarrow \infty} A_n \\
 &\quad + \limsup_{n \rightarrow \infty} \hat{\gamma}_n \|v_n - p\| \\
 (2.10) \qquad \qquad \qquad &= d, \quad p \in F,
 \end{aligned}$$

where $n = (k - 1)N + i$, $k = k(n)$, $i = i(n) \in \{1, 2, \dots, N\} = I$ and $T_n = T_{i(\text{mod } N)} = T_i$.

Therefore from condition (iv), (2.8) – (2.10), and Lemma 1.2 we know that

$$(2.11) \qquad \qquad \qquad \lim_{n \rightarrow \infty} \|T_n^k y_n - x_{n-1}\| = 0.$$

From (1.8), (2.11) and condition (iii), we have

$$\begin{aligned}
 \|x_n - x_{n-1}\| &= \|\beta_n [T_i^k y_n - x_{n-1}] + \gamma_n(u_n - x_{n-1})\| \\
 &\leq \beta_n \|T_i^k y_n - x_{n-1}\| + \gamma_n \|u_n - x_{n-1}\| \\
 (2.12) \qquad \qquad \qquad &\rightarrow 0, \quad \text{as } n \rightarrow \infty,
 \end{aligned}$$

which implies that

$$(2.13) \qquad \qquad \qquad \lim_{n \rightarrow \infty} \|x_n - x_{n-1}\| = 0,$$

and so

$$(2.14) \qquad \qquad \qquad \lim_{n \rightarrow \infty} \|x_n - x_{n+j}\| = 0, \quad \forall j = 1, 2, \dots, N.$$

On the other hand, we have

$$(2.15) \quad \begin{aligned} \|x_n - T_n^k x_n\| &\leq \|x_n - x_{n-1}\| + \|x_{n-1} - T_n^k y_n\| \\ &\quad + \|T_n^k y_n - T_n^k x_n\|, \end{aligned}$$

where $n = (k-1)N + i$, $k = k(n)$ and $i = i(n)$.

Now, we consider the third term of the right hand side of (2.15). From (1.8), (1.9) and the condition (vi) we have

$$(2.16) \quad \begin{aligned} \|T_n^k y_n - T_n^k x_n\| &\leq \|y_n - x_n\| + A_n \\ &\leq \|\hat{\alpha}_n x_{n-1} + \hat{\beta}_n T_n^k x_n + \hat{\gamma}_n v_n - x_n\| + A_n \\ &\leq \hat{\alpha}_n \|x_{n-1} - x_n\| + \hat{\beta}_n \|T_n^k x_n - x_n\| \\ &\quad + \hat{\gamma}_n \|v_n - x_n\| + A_n \\ &\leq \hat{\alpha}_n \|x_{n-1} - x_n\| + \delta \|T_n^k x_n - x_n\| \\ &\quad + \hat{\gamma}_n \|v_n - x_n\| + A_n. \end{aligned}$$

Substituting (2.16) into (2.15), we obtain that

$$(2.17) \quad \begin{aligned} (1 - \delta) \|x_n - T_n^k x_n\| &\leq (1 + \hat{\alpha}_n) \|x_n - x_{n-1}\| + \|x_{n-1} - T_n^k y_n\| \\ &\quad + \hat{\gamma}_n \|v_n - x_n\| + A_n. \end{aligned}$$

Hence, by virtue of the condition (iii), (2.11) and (2.13), we have

$$(2.18) \quad (1 - \delta) \limsup_{n \rightarrow \infty} \|x_n - T_n^k x_n\| \leq 0.$$

From the condition (vi), $0 \leq \delta < 1$, hence from (2.18) we have

$$(2.19) \quad \lim_{n \rightarrow \infty} \|x_n - T_n^k x_n\| = 0.$$

Now, we prove that (2.7) holds. In fact, since for each $n > N$, $n = (n - N)(\text{mod } N)$ and $n = (k(n) - 1)N + i(n)$, hence $n - N = ((k(n) - 1) - 1)N + i(n) = (k(n - N) - 1)N + i(n - N)$, that is,

$$k(n - N) = k(n) - 1 \quad \text{and} \quad i(n - N) = i(n).$$

From (2.14), (2.19) and condition (vii) that

$$\begin{aligned}
 \|x_n - T_n x_n\| &\leq \|x_n - T_n^k x_n\| + \|T_n^k x_n - T_n x_n\| \\
 &\leq \|x_n - T_n^k x_n\| + L \|T_n^{k-1} x_n - x_n\| \\
 &\leq \|x_n - T_n^k x_n\| + L \left\{ \|T_n^{k-1} x_n - T_{n-N}^{k-1} x_{n-N}\| \right. \\
 &\quad \left. + \|T_{n-N}^{k-1} x_{n-N} - x_{n-N}\| + \|x_{n-N} - x_n\| \right\} \\
 &\leq \|x_n - T_n^k x_n\| + L^2 \|x_n - x_{n-N}\| \\
 &\quad + L \|T_{n-N}^{k-1} x_{n-N} - x_{n-N}\| + L \|x_{n-N} - x_n\| \\
 &= \|x_n - T_n^k x_n\| + L(L+1) \|x_n - x_{n-N}\| \\
 &\quad + L \|T_{n-N}^{k-1} x_{n-N} - x_{n-N}\| \\
 &\rightarrow 0 \text{ as } n \rightarrow \infty,
 \end{aligned}$$

which implies that

$$(2.20) \quad \lim_{n \rightarrow \infty} \|x_n - T_n x_n\| = 0,$$

and so, from condition (vii), (2.13) and (2.20), it follows that, for any $j = 1, 2, \dots, N$,

$$\begin{aligned}
 \|x_n - T_{n+j} x_n\| &\leq \|x_n - x_{n+j}\| + \|x_{n+j} - T_{n+j} x_{n+j}\| \\
 &\quad + \|T_{n+j} x_{n+j} - T_{n+j} x_n\| \\
 &\leq (1+L) \|x_n - x_{n+j}\| + \|x_{n+j} - T_{n+j} x_{n+j}\| \\
 &\rightarrow 0 \text{ as } n \rightarrow \infty,
 \end{aligned}$$

which implies that

$$(2.21) \quad \lim_{n \rightarrow \infty} \|x_n - T_{n+j} x_n\| = 0,$$

for all $j = 1, 2, \dots, N$.

Without loss of generality, we can assume that $n_k = i(\text{mod } N)$ for all k and some $i \in \{1, 2, \dots, N\}$. For any fixed $l \in \{1, 2, \dots, N\}$, we can find a $j \in \{1, 2, \dots, N\}$, independent of k , such that $i + j = l(\text{mod } N)$, and so $n_k + j = l(\text{mod } N)$ for all k . Hence, from (2.21), we have

$$(2.22) \quad \lim_{n_k \rightarrow \infty} \|x_{n_k} - T_l x_{n_k}\| = 0,$$

for all $l = 1, 2, \dots, N$.

Thus

$$(2.23) \quad \lim_{n \rightarrow \infty} \|x_n - T_l x_n\| = 0,$$

for all $l = 1, 2, \dots, N$.

That is, (2.7) holds.

For any given $p \in F = \bigcap_{i=1}^N F(T_i)$, by the same method as given in proving Lemma 1.3 and (2.23), we can prove that

$$(2.24) \quad \lim_{n \rightarrow \infty} \|x_n - p\| = d,$$

where $d \geq 0$ is some nonnegative number, and

$$(2.25) \quad \lim_{n \rightarrow \infty} \|x_n - T_l x_n\| = 0,$$

for all $l = 1, 2, \dots, N$.

Especially, we have

$$(2.26) \quad \lim_{n \rightarrow \infty} \|x_n - T_1 x_n\| = 0.$$

By the assumption of the theorem, T_1 is semi-compact, therefore it follows from (2.26) that there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $x_{n_i} \rightarrow x^* \in C$. Hence from (2.25) we have that

$$\|x^* - T_l x^*\| = \lim_{n_i \rightarrow \infty} \|x_{n_i} - T_l x_{n_i}\| = 0,$$

for all $l = 1, 2, \dots, N$, which implies that

$$x^* \in F = \bigcap_{i=1}^N F(T_i).$$

Take $p = x^*$ in (2.24), similarly we can prove that

$$\lim_{n \rightarrow \infty} \|x_n - x^*\| = d_1,$$

where $d_1 \geq 0$ is some nonnegative number. From $x_{n_i} \rightarrow x^*$ we know that $d_1 = 0$, i.e., $x_n \rightarrow x^*$. Thus $\{x_n\}$ converges strongly to a common fixed point of the mappings $\{T_i\}_{i=1}^N$. This completes the proof of Theorem 2.2. \square

Remark 2.1. Our results improves and extends the corresponding results of Chang and Cho [3] to the case of more general class of asymptotically nonexpansive mapping considered in this paper.

Remark 2.2. Our results also improve and extend the corresponding results of [1, 4, 5, 7, 8, 9, 11, 13, 14, 16] to the case of more general class of mappings, spaces and iteration schemes considered in this paper.

Remark 2.3. Our results also extends the corresponding results of Gu [6] to the case of more general class of asymptotically nonexpansive mapping and implicit iteration process with bounded errors considered in this paper.

Remark 2.4. Theorem 2.2 extends and improves Theorem 3.3 of Plubtieng and Wangkeeree [10] to the case of implicit iteration process with bounded errors for a finite family of mappings considered in this paper.

REFERENCES

- [1] H.H. Bauschke, *The approximation of fixed points of compositions of nonexpansive mappings in Hilbert space*, J. Math. Anal. Appl. **202** (1996), 150–159.
- [2] R. Bruck, T. Kuczumow and S. Reich, *Convergence of iterates of asymptotically nonexpansive mappings in Banach spaces with the uniform Opial property*, Collo. Math. **65(2)** (1993), 169–179.
- [3] S.S. Chang and Y.J. Cho, *The implicit iterative processes for asymptotically nonexpansive mappings*, Nonlinear Anal. Appl. **1** (2003), 369–382.
- [4] K. Goebel and W.A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **35** (1972), 171–174.
- [5] J. Gornicki, *Weak convergence theorems for asymptotically nonexpansive mappings in uniformly convex Banach spaces*, Comment. Math. Univ. Carolin. **301** (1989), 249–252.
- [6] F. Gu, *Convergence of implicit iterative process with errors for a finite family of asymptotically nonexpansive mappings*, Acta Math. Sinica **26(A)(7)** (2006), 1131–1143.
- [7] B. Halpern, *Fixed points of nonexpansive maps*, Bull. Amer. Math. Soc. **73** (1967), 957–961.
- [8] P.L. Lions, *Approximation de points fixes de contractions*, C. R. Acad. Sci. Paris Ser. I Math. **284** (1977), 1357–1359.
- [9] M.O. Osilike, *Implicit iteration process for common fixed point of a finite family of strictly pseudocontractive maps*, J. Math. Anal. Appl. **294** (2004), 73–81.
- [10] S. Plubtieng and R. Wangkeeree, *Noor iterations with error for Non-Lipschitzian mappings in Banach spaces*, Kyungpook Math. J. **46** (2006), 201–209.
- [11] S. Reich, *Strong convergence theorems for resolvents of accretive operators in Banach spaces*, J. Math. Anal. Appl. **75** (1980), 287–292.
- [12] J. Schu, *Weak and strong convergence theorems to fixed points of asymptotically nonexpansive mappings*, Bull. Austral. Math. Soc. **43** (1991), 153–159.
- [13] Z. Sun, *Strong convergence of an implicit iteration process for a finite family of asymptotically quasi-nonexpansive mappings*, J. Math. Anal. Appl. **286(1)** (2003), 351–358.
- [14] K.K. Tan and H.K. Xu, *The nonlinear ergodic theorem for asymptotically nonexpansive mappings in Banach spaces*, Proc. Amer. Math. Soc. **114** (1992), 399–404.
- [15] K.K. Tan and H.K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl. **178** (1993), 301–308.
- [16] R. Wittmann, *Approximation of fixed points of nonexpansive mappings*, Arch. Math. **58** (1992), 486–491.
- [17] H.K. Xu and R.G. Ori, *An implicit iteration process for nonexpansive mappings*, Numer. Funct. Anal. Optim. **22** (2001), 767–773.
- [18] Y.Y. Zhou and S.S. Chang, *Convergence of implicit iterative process for a finite family of asymptotically nonexpansive mappings in Banach spaces*, Numer. Funct. Anal. Optim. **23** (2002), 911–921.

DEPARTMENT OF MATHEMATICS AND INFORMATION TECHNOLOGY,
GOVT. NAGARJUNA P.G. COLLEGE OF SCIENCE,
RAIPUR - 492010 (C.G.), INDIA.
E-mail address: saluja_1963@rediffmail.com, saluja1963@gmail.com