

## SEIDEL ENERGY OF ITERATED LINE GRAPHS OF REGULAR GRAPHS

HARISHCHANDRA S. RAMANE<sup>1</sup>, IVAN GUTMAN<sup>2</sup>, AND MAHADEVAPPA M. GUNDLOOR<sup>3</sup>

ABSTRACT. The Seidel matrix  $S(G)$  of a graph  $G$  is the square matrix whose  $(i, j)$ -entry is equal to  $-1$  or  $1$  if the  $i$ -th and  $j$ -th vertices of  $G$  are adjacent or non-adjacent, respectively, and is zero if  $i = j$ . The Seidel energy of  $G$  is the sum of the absolute values of the eigenvalues of  $S(G)$ . We show that if  $G$  is regular of order  $n$  and of degree  $r \geq 3$ , then for each  $k \geq 2$ , the Seidel energy of the  $k$ -th iterated line graph of  $G$  depends solely on  $n$  and  $r$ . This result enables the construction of pairs of non-cospectral, Seidel equienergetic graphs of the same order.

### 1. INTRODUCTION

Let  $G$  be a simple, undirected graph with  $n$  vertices and  $m$  edges. The vertices of  $G$  are labeled as  $v_1, v_2, \dots, v_n$ . The *adjacency matrix* of  $G$  is the square matrix  $A(G) = (a_{ij})$ , in which  $a_{ij} = 1$  if  $v_i$  is adjacent to  $v_j$  and  $a_{ij} = 0$  otherwise. The eigenvalues of  $A(G)$  will be referred to as the *adjacency eigenvalues* of  $G$ , and labeled as  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ . These form the *adjacency spectrum* of  $G$  [4, 5].

The *Seidel matrix* of the graph  $G$  is the  $n \times n$  real symmetric matrix  $S(G) = (s_{ij})$ , where  $s_{ij} = -1$  if the vertices  $v_i$  and  $v_j$  are adjacent,  $s_{ij} = 1$  if the vertices  $v_i$  and  $v_j$  are not adjacent, and  $s_{ij} = 0$  if  $i = j$ . The eigenvalues of the Seidel matrix, labeled as  $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n$ , are said to be the *Seidel eigenvalues* of  $G$  and their collection is the *Seidel spectrum* of  $G$  [1]. Two non-isomorphic graphs are said to be *Seidel cospectral* if their Seidel spectra coincide.

The *Seidel energy* of the graph  $G$  is defined as [8]

$$(1.1) \quad SE = SE(G) = \sum_{i=1}^n |\sigma_i| .$$

---

*Key words and phrases.* Seidel spectrum, Seidel energy, line graph, regular graph.  
2010 *Mathematics Subject Classification.* Primary: 05C50, 05C07.

*Received:* February 15, 2015

*Accepted:* March 4, 2015.

The Seidel energy is conceived in full analogy with the ordinary graph energy  $E(G)$ , which is defined as [7]

$$(1.2) \quad E(G) = \sum_{i=1}^n |\lambda_i|$$

and whose theory is nowadays extensively elaborated [10].

The study of the Seidel energy started only quite recently [8], and only some of its basic properties have been established so far [6, 8, 11].

Two graphs  $G_1$  and  $G_2$  are said to be *equienergetic* if  $E(G_1) = E(G_2)$ . In a trivial manner, cospectral graphs are equienergetic. Much work has been done on the study of non-cospectral equienergetic graphs; for details see [10].

Two graphs  $G_1$  and  $G_2$  will be said to be *Seidel equienergetic* if  $SE(G_1) = SE(G_2)$ . Of course, Seidel cospectral graphs are Seidel equienergetic. We are interested in Seidel non-cospectral, Seidel equienergetic graphs, having same number of vertices.

Denote by  $\overline{G}$  the complement of the graph  $G$ . Then, evidently,  $S(G) = A(\overline{G}) - A(G)$  implying  $S(\overline{G}) = -S(G)$ . Consequently, if the Seidel eigenvalues of a graph  $G$  are  $\sigma_i, i = 1, 2, \dots, n$ , then the Seidel eigenvalues of  $\overline{G}$  are  $-\sigma_i, i = 1, 2, \dots, n$ . Thus, by Eq. (1.1),  $SE(\overline{G}) = SE(G)$ , that is  $G$  and  $\overline{G}$  are Seidel equienergetic.

In what follows we search for less trivial pairs of Seidel equienergetic graphs.

In this paper, based on a result that relates the Seidel eigenvalues of regular graphs with its adjacency eigenvalues, we show that  $SE$  of second and higher line graphs of regular graphs is fully determined by the order  $n$  and regularity  $r$ . By this, infinitely many pairs of Seidel equienergetic graphs can be constructed.

## 2. ON SPECTRA OF REGULAR GRAPHS AND THEIR LINE GRAPHS

The line graph [9] of the graph  $G$  will be denoted by  $L(G)$ . For  $k = 1, 2, \dots$ , the  $k$ -th iterated line graph of  $G$ , denoted by  $L^k(G)$ , is defined recursively as  $L^k(G) = L(L^{k-1}(G))$ , with  $L^0(G) = G$  and  $L^1(G) = L(G)$ .

The line graph of a regular graph is a regular graph. In particular, the line graph of a regular graph  $G$  of order  $n_0$  and of degree  $r_0$  is a regular graph of order  $n_1 = (r_0 n_0)/2$  and of degree  $r_1 = 2r_0 - 2$ . Consequently, the order and degree of  $L^k(G)$  are [2, 3]:

$$n_k = \frac{1}{2} r_{k-1} n_{k-1} \quad \text{and} \quad r_k = 2r_{k-1} - 2$$

where  $n_i$  and  $r_i$  stand for the order and degree of  $L^i(G)$ ,  $i = 1, 2, \dots$ . Therefore,

$$(2.1) \quad r_k = 2^k r_0 - 2^{k+1} + 2$$

and

$$(2.2) \quad n_k = \frac{n_0}{2^k} \prod_{i=0}^{k-1} r_i = n_0 \prod_{i=0}^{k-1} (2^{i-1} r_0 - 2^i + 1).$$

The basic spectral properties of regular graphs are well known. Here we state some results needed for the present considerations.

**Theorem 2.1.** [4, 5] *If  $G$  is an  $r$ -regular graph, then its maximum adjacency eigenvalue is equal to  $r$ .*

**Theorem 2.2.** [4, 5] *If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the adjacency eigenvalues of a regular graph  $G$  of order  $n$  and of degree  $r$ , then the adjacency eigenvalues of  $L(G)$  are*

$$\begin{cases} \lambda_i + r - 2, & i = 1, 2, \dots, n, \\ -2, & \frac{n(r-2)}{2} \text{ times.} \end{cases}$$

**Theorem 2.3.** [12, 13] *If  $G$  is a regular graph of order  $n$  and of degree  $r \geq 3$ , then*

$$E(L^2(G)) = 2nr(r-2).$$

**Theorem 2.4.** [1] *Let  $G$  be an  $r$ -regular graph on  $n$  vertices. If  $r, \lambda_2, \dots, \lambda_n$  are the adjacency eigenvalues of  $G$ , then its Seidel eigenvalues are  $n-1-2r$  and  $-1-2\lambda_i$ ,  $i = 2, 3, \dots, n$ .*

### 3. SEIDEL ENERGY

We first state a result for Seidel energy that is formally similar to Theorem 2.3.

**Theorem 3.1.** *If  $G$  is a regular graph of order  $n$  and of degree  $r \geq 3$ , then*

$$SE(L^2(G)) = \begin{cases} 4nr(r-2) + nr - 16r + 22, & \text{if } n \geq \frac{16r-22}{r(r-1)}, \\ 3nr(r-2), & \text{if } n < \frac{16r-22}{r(r-1)}. \end{cases}$$

*Proof.* Let  $r, \lambda_2, \dots, \lambda_n$  be the adjacency eigenvalues of the regular graph  $G$ . Then from Theorem 2.2, the adjacency eigenvalues of  $L(G)$  are

$$(3.1) \quad \begin{cases} 2r-2, \\ \lambda_i + r - 2, & i = 2, 3, \dots, n, \\ -2, & \frac{n(r-2)}{2} \text{ times.} \end{cases}$$

In view of the fact that  $L(G)$  is also a regular graph on  $nr/2$  vertices and of degree  $2r-2$ , applying Theorem 2.2 to Eq. (3.1), the adjacency eigenvalues of  $L^2(G)$  are found to be

$$(3.2) \quad \begin{cases} 4r-6, \\ \lambda_i + 3r - 6, & i = 2, 3, \dots, n, \\ 2r-6, & \frac{n(r-2)}{2} \text{ times,} \\ -2, & \frac{nr(r-2)}{2} \text{ times.} \end{cases}$$

Because  $L^2(G)$  is a regular graph on  $nr(r-1)/2$  vertices and of degree  $4r-6$ , applying Theorem 2.4 to Eq. (3.2), the Seidel eigenvalues of  $L^2(G)$  are calculated as

$$(3.3) \quad \begin{cases} \frac{1}{2}nr(r-1) + 11 - 8r, \\ -2\lambda_i - 6r + 11, & i = 2, 3, \dots, n, \\ -4r + 11, & \frac{n(r-2)}{2} \text{ times,} \\ 3, & \frac{nr(r-2)}{2} \text{ times.} \end{cases}$$

By Theorem 2.1, all adjacency eigenvalues of a regular graph of degree  $r$  satisfy the condition  $-r \leq \lambda_i \leq r$ . If  $r \geq 3$ , then  $4r-11 \geq 0$ , and because of  $\lambda_i \geq -r$ , we have  $2\lambda_i + 6r - 11 \geq 0$ . Bearing this in mind, the Seidel energy of  $L^2(G)$  is computed from Eq. (3.3) as:

$$\begin{aligned} SE(L^2(G)) &= \left| \frac{nr(r-1)}{2} + 11 - 8r \right| + \sum_{i=2}^n (2\lambda_i + 6r - 11) \\ &\quad + \frac{n(r-2)}{2}(4r-11) + \frac{nr(r-2)}{2}(3) \\ &= \left| \frac{nr(r-1)}{2} + 11 - 8r \right| - 2r + (n-1)(6r-11) \\ &\quad + \frac{n(r-2)}{2}(4r-11) + \frac{3nr(r-2)}{2} \end{aligned}$$

since  $\sum_{i=2}^n \lambda_i = -r$ .

If  $\frac{nr(r-1)}{2} + 11 - 8r \geq 0$  i.e.,  $n \geq \frac{16r-22}{r(r-1)}$ , then the above equality yields

$$SE(L^2(G)) = 4nr(r-2) + nr - 16r + 22.$$

If  $\frac{nr(r-1)}{2} + 11 - 8r < 0$  i.e.,  $n < \frac{16r-22}{r(r-1)}$ , then we get  $SE(L^2(G)) = 3nr(r-2)$ .  $\square$

**Corollary 3.1.** *Let  $G$  be a regular graph of order  $n_0$  and of degree  $r_0 \geq 3$ . Let  $n_k$  and  $r_k$  be the order and degree, respectively, of the  $k$ -th iterated line graph  $L^k(G)$  of  $G$ ,  $k \geq 1$ . Then*

$$SE(L^{k+1}(G)) = \begin{cases} 4n_k(r_k - 2) + 2n_k - 8(r_k + 2) + 22, & \text{if } n_k \geq \frac{8r_k - 6}{r_k}, \\ 3n_k(r_k - 2), & \text{if } n_k < \frac{8r_k - 6}{r_k}. \end{cases}$$

From Corollary 3.1 we see that the Seidel energy of the  $k$ -th iterated line graph of a regular graph  $G$  is fully determined by the order  $n_0$  and degree  $r_0$  of  $G$ .

## 4. SEIDEL EQUIENERGETIC GRAPHS

**Theorem 4.1.** *Let  $G_1$  and  $G_2$  be two regular graphs of the same order and of the same degree. Then for any  $k \geq 1$  the following holds:*

- (i)  $L^k(G_1)$  and  $L^k(G_2)$  are of the same order, same degree and have the same number of edges.
- (ii)  $L^k(G_1)$  and  $L^k(G_2)$  are Seidel cospectral if and only if  $G_1$  and  $G_2$  are adjacency cospectral.

*Proof.* Statement (i) follows from Eqs. (2.1) and (2.2), and the fact that the number of edges of  $L^k(G)$  is equal to the number of vertices of  $L^{k+1}(G)$ . It was earlier stated in [12, 13]. Statement (ii) follows from Theorems 2.2 and 2.4.  $\square$

Theorem 4.1 and Corollary 3.1 infer the following.

**Theorem 4.2.** *Let  $G_1$  and  $G_2$  be two Seidel non-cospectral regular graphs of the same order and of the same degree  $r \geq 3$ . Then for any  $k \geq 2$ , the iterated line graphs  $L^k(G_1)$  and  $L^k(G_2)$  form a pair of Seidel non-cospectral, Seidel equienergetic graphs of equal order and with equal number of edges.*

One should recall that the following fully analogous result was earlier obtained for ordinary graph energy, based on adjacency eigenvalues, cf. Eq. (1.2).

**Theorem 4.3.** [13] *Let  $G_1$  and  $G_2$  be two non-cospectral regular graphs of the same order and of the same degree  $r \geq 3$ . Then for any  $k \geq 2$ , the iterated line graphs  $L^k(G_1)$  and  $L^k(G_2)$  form a pair of non-cospectral, equienergetic graphs of equal order and with equal number of edges.*

Comparing Theorems 4.2 and 4.3 a deep analogy between ordinary graph energy and Seidel energy can be envisaged.

From Theorem 4.2, it is now easy to generate large families of Seidel equienergetic graphs. For instance, there are 2, 5, 19, and 85 connected regular graphs of degree 3 and of order 6, 8, 10, and 12, respectively. No two of these are Seidel cospectral as they are adjacency non-cospectral (see [4], pp. 268–269). Their second and higher iterated line graphs form families consisting of size 2, 5, 19, 85, . . . , Seidel equienergetic graphs.

**Acknowledgment:** The author H. S. Ramane is thankful to the University Grants Commission (UGC), Govt. of India for support through research grant under UPE FAR-II grant No. F 14-3/2012 (NS/PE). M. M. Gundloor is thankful to the Karnatak University for support through UGC-UPE scholarship No. KU/Sch/UGC-UPE/2013-14/1136.

## REFERENCES

- [1] A. E. Brouwer and W. H. Haemers, *Spectra of Graphs*, Springer, Berlin, 2012.
- [2] F. Buckley, *Iterated line graphs*, Congr. Numer. **33** (1981), 390–394.
- [3] F. Buckley, *The size of iterated line graphs*, Graph Theory Notes New York **25** (1993), 33–36.

- [4] D. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs – Theory and Application*, Academic Press, New York, 1980.
- [5] D. Cvetković, P. Rowlinson and S. Simić, *An Introduction to the Theory of Graph Spectra*, Cambridge Univ. Press, Cambridge, 2010.
- [6] E. Ghorbani, *On eigenvalues of Seidel matrices and Haemers conjecture*, arXiv:1301.0075v1.
- [7] I. Gutman, *The energy of a graph*, Ber. Math. Stat. Sect. Forschungsz. Graz **103** (1978), 1–22.
- [8] W. H. Haemers, *Seidel switching and graph energy*, MATCH Commun. Math. Comput. Chem. **68** (2012), 653–659.
- [9] F. Harary, *Graph Theory*, Addison–Wesley, Reading, 1969.
- [10] X. Li, Y. Shi and I. Gutman, *Graph Energy*, Springer, New York, 2012.
- [11] P. Nageswari and P. B. Sarasija, *Seidel energy and its bounds*, Int. J. Math. Anal. **8** (2014), 2869–2871.
- [12] H. S. Ramane, H. B. Walikar, S. B. Rao, B. D. Acharya, P. R. Hampiholi, S. R. Jog and I. Gutman, *Equienergetic graphs*, Kragujevac J. Math. **26** (2004), 5–13.
- [13] H. S. Ramane, H. B. Walikar, S. B. Rao, B. D. Acharya, P. R. Hampiholi, S. R. Jog and I. Gutman, *Spectra and energies of iterated line graphs of regular graphs*, Appl. Math. Lett. **18** (2005), 679–682.

<sup>1</sup>DEPARTMENT OF MATHEMATICS,  
KARNATAK UNIVERSITY,  
DHARWAD – 580003, INDIA  
*E-mail address:* [hsramane@yahoo.com](mailto:hsramane@yahoo.com)

<sup>2</sup>FACULTY OF SCIENCE,  
UNIVERSITY OF KRAGUJEVAC,  
KRAGUJEVAC, SERBIA, AND  
STATE UNIVERSITY OF NOVI PAZAR,  
NOVI PAZAR, SERBIA  
*E-mail address:* [gutman@kg.ac.rs](mailto:gutman@kg.ac.rs)

<sup>3</sup>DEPARTMENT OF MATHEMATICS,  
KARNATAK UNIVERSITY,  
DHARWAD – 580003, INDIA  
*E-mail address:* [madeshgundloor@gmail.com](mailto:madeshgundloor@gmail.com)