

**LOWER EXTREMITIES FOR GENERALIZED NORMALIZED
 δ -CASORATI CURVATURES OF BI-SLANT SUBMANIFOLDS IN
GENERALIZED COMPLEX SPACE FORMS**

MOHD. AQUIB¹, MOHAMMAD HASAN SHAHID¹, AND MOHAMMED JAMALI²

ABSTRACT. In this paper, we obtain the inequalities for the generalized normalized δ -Casorati curvature and the normalized scalar curvature for different submanifolds in generalized complex space form, which is based on an optimization procedure involving a quadratic polynomial in the components of the second fundamental form and characterizes the submanifolds on which equalities hold. We also develop, the same inequalities for semi-slant, hemi-slant, CR, slant, invariant and anti-invariant submanifolds in the same target space and consider the equality case. Moreover, we obtain a geometric inequality involving Casorati curvature for warped product bi-slant submanifolds in same ambient and obtain an obstruction result.

1. INTRODUCTION

The theory of Chen invariants, which establish the simple relationships between the main intrinsic invariants and the main extrinsic invariants of the submanifolds is one of the most interesting research areas of differential geometry started by Chen [5] in 1993. In the initial papers, Chen established inequalities between the scalar curvature, the sectional curvature (intrinsic invariants) and the squared norm of the mean curvature (the main extrinsic invariant) of a submanifold in a real space form. The same author obtained the inequalities for submanifolds between the k -Ricci curvature, the squared mean curvature, and the shape operator in the real space form with arbitrary codimension [4]. Since then, different geometers proved the similar inequalities for different submanifolds and ambient spaces [13, 15].

Key words and phrases. Casorati curvature, generalized complex space form, scalar curvature, warped product.

2010 *Mathematics Subject Classification.* Primary: 53B05. Secondary: 53B20, 53C40.

Received: May 08, 2017.

Accepted: July 26, 2017.

The Casorati curvature(extrinsic invariant) of a submanifold of a Riemannian manifold, introduced by Casorati is defined as the normalized square length of the second fundamental form [3]. The concept of Casorati curvature extends the concept of the principal direction of a hypersurface of a Riemannian manifold [9]. The geometrical meaning and the importance of the Casorati curvature has been discussed by distinguished geometers [6, 12, 19]. Therefore, it attracts the attention of geometers to obtain the optimal inequalities for the Casorati curvatures of the submanifolds of different ambient spaces [1, 7, 14].

In this article, we study the inequalities which relates the generalized normalized δ -Casorati curvature and the normalized scalar curvature for different submanifolds in generalized complex space form and consider the equality case of the inequality. We also obtain, a geometric inequality involving Casorati curvature for warped product submanifolds in same ambient and obtain a non-existence result.

2. PRELIMINARIES

Let \bar{M} be an almost Hermitian manifold with an almost complex structure J and a Riemannian metric g . An almost Hermitian manifold is said to be a nearly Kaehler manifold if $(\bar{\nabla}_X J)X = 0$ and becomes a Kaehler manifold if $\bar{\nabla}J = 0$ for all $X \in T\bar{M}$, where $\bar{\nabla}$ is the Levi-Civita connection of the Riemannian metric g .

Tricerri and Vanhecke [16] introduced the concept of generalized complex space form as a generalization of the complex space form. An almost Hermitian manifold \bar{M} is called the generalized complex space form, denoted by $\bar{M}(f_1, f_2)$, if the Riemannian curvature tensor \bar{R} satisfies

$$(2.1) \quad \begin{aligned} \bar{R}(X, Y)Z = & f_1\{g(Y, Z)X - g(X, Z)Y\} + f_2\{g(X, JZ)JY \\ & - g(Y, JZ)JX + 2g(X, JY)JZ\}, \end{aligned}$$

for all $X, Y, Z \in T\bar{M}$, where f_1 and f_2 are smooth functions on $\bar{M}(f_1, f_2)$.

Let M be an n -dimensional submanifold of a generalized complex space form $\bar{M}(f_1, f_2)$ of complex dimension m . Let ∇ and $\bar{\nabla}$ be the Levi-Civita connection on M and $\bar{M}(f_1, f_2)$ respectively. The Gauss and Weingarten equations are defined as

$$\begin{aligned} \bar{\nabla}_X Y &= \nabla_X Y + h(X, Y), \\ \bar{\nabla}_X \xi &= -S_\xi X + \nabla_X^\perp Y, \end{aligned}$$

for vector fields $X, Y \in TM$ and $\xi \in T^\perp M$, where h , S and ∇^\perp are the second fundamental form, the shape operator and the normal connection respectively. The second fundamental form and the shape operator are related by the following equation

$$g(h(X, Y), \xi) = g(S_\xi X, Y),$$

for vector fields $X, Y \in TM$ and $\xi \in T^\perp M$.

The equation of Gauss is given by

$$(2.2) \quad \bar{R}(X, Y, Z, W) = R(X, Y, Z, W) + g(h(X, Z), h(Y, W)) - g(h(X, W), h(Y, Z)),$$

TABLE 1. Definition

S.N.	\overline{M}	M	D_1	D_2	θ_1	θ_2
(1)	\overline{M}	bi-slant	slant	slant	slant angle	slant angle
(2)	\overline{M}	semi-slant	invariant	slant	0	slant angle
(3)	\overline{M}	hemi-slant	slant	anti-invariant	slant angle	$\frac{\pi}{2}$
(4)	\overline{M}	CR	invariant	anti-invariant	0	$\frac{\pi}{2}$
(5)	\overline{M}	slant	either $D_1 = 0$ or $D_2 = 0$		either $\theta_1 = \theta_2 = \theta$ or $\theta_1 = \theta_2 \neq \theta$	

for $X, Y, Z, W \in TM$, where \overline{R} and R represent the curvature tensor of $\overline{M}(f_1, f_2)$ and M respectively.

Let M be an n -dimensional submanifold of a generalized complex space form $\overline{M}(f_1, f_2)$ of complex dimension m . For any tangent vector field $X \in TM$, we can write $JX = PX + QX$, where P and Q are the tangential and normal components of JX respectively. If $P = 0$, the submanifold is said to be an anti-invariant submanifold and if $Q = 0$, the submanifold is said to be an invariant submanifold. The squared norm of P at $p \in M$ is defined as

$$(2.3) \quad \|P\|^2 = \sum_{i,j=1}^n g^2(Je_i, e_j),$$

where $\{e_1, \dots, e_n\}$ is any orthonormal basis of the tangent space T_pM .

A submanifold M of an almost Hermitian manifold \overline{M} is said to be a slant submanifold if for any $p \in M$ and a non zero vector $X \in T_pM$, the angle between JX and T_pM is constant, i.e., the angle does not depend on the choice of $p \in M$ and $X \in T_pM$. The angle $\theta \in [0, \frac{\pi}{2}]$ is called the slant angle of M in \overline{M} .

A submanifold M of an almost Hermitian manifold \overline{M} is said to be a bi-slant submanifold, if there exist two orthogonal distributions D_1 and D_2 , such that

- (i) TM admits the orthogonal direct decomposition, i.e., $TM = D_1 + D_2$;
- (ii) for $i = 1, 2$, D_i is the slant distribution with slant angle θ_i .

In fact, semi-slant submanifolds, hemi-slant submanifolds, CR-submanifolds, slant submanifolds can be obtained from bi-slant submanifolds in particular. We can see the case in the following table: Invariant and anti-invariant submanifolds are the slant submanifolds with slant angle $\theta = 0$ and $\theta = \frac{\pi}{2}$ respectively. If $0 < \theta < \frac{\pi}{2}$, then slant submanifold is called proper slant submanifold. It is said to be proper bi-slant if θ_i lies between 0 and $\frac{\pi}{2}$.

If M is a bi-slant submanifold in a generalized complex space form $\overline{M}(f_1, f_2)$, then one easily can see that

$$\|P\|^2 = \sum_{i,j=1}^n g^2(Pe_i, e_j) = 2(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2),$$

where $\dim D_1 = 2d_1$ and $\dim D_2 = 2d_2$.

Let M be a Riemannian manifold and $K(\pi)$ denotes the sectional curvature of M of the plane section $\pi \subset T_pM$ at a point $p \in M$. If $\{e_1, \dots, e_n\}$ and $\{e_{n+1}, \dots, e_{2m}\}$ be the orthonormal basis of T_pM and $T_p^\perp M$ at any $p \in M$, then the scalar curvature τ at that point is given by

$$(2.4) \quad \tau(p) = \sum_{1 \leq i < j \leq n} K(e_i \wedge e_j)$$

and the normalized scalar curvature ρ is defined as

$$\rho = \frac{2\tau}{n(n-1)}.$$

The mean curvature vector denoted by H is defined as

$$H = \frac{1}{n} \sum_{i=1}^n h(e_i, e_i).$$

We also put

$$h_{ij}^\gamma = g(h(e_i, e_j), e_\gamma), \quad i, j \in 1, 2, \dots, n, \gamma \in \{n+1, n+2, \dots, 2m\}.$$

The squared norm of the mean curvature of the submanifold is defined by

$$\|H\|^2 = \frac{1}{n^2} \sum_{\gamma=n+1}^{2m} \left(\sum_{i=1}^n h_{ii}^\gamma \right)^2,$$

and the squared norm of second fundamental form h is denoted by \mathcal{C} defined as

$$(2.5) \quad \mathcal{C} = \frac{1}{n} \sum_{\gamma=n+1}^{2m} \sum_{i,j=1}^n (h_{ij}^\gamma)^2$$

known as Casorati curvature of the submanifold.

If we suppose that L is an r -dimensional subspace of TM , $r \geq 2$, and $\{e_1, e_2, \dots, e_r\}$ is an orthonormal basis of L , then the scalar curvature of the r -plane section L is given as

$$\tau(L) = \sum_{1 \leq \gamma < \beta \leq r} K(e_\gamma \wedge e_\beta)$$

and the Casorati curvature \mathcal{C} of the subspace L is as follows

$$(2.6) \quad \mathcal{C}(L) = \frac{1}{r} \sum_{\gamma=n+1}^{2m} \sum_{i,j=1}^r (h_{ij}^\gamma)^2.$$

A point $p \in M$ is said to be an invariantly quasi-umbilical point if there exist $2m - n$ mutually orthogonal unit normal vectors $\xi_{n+1}, \dots, \xi_{2m}$ such that the shape operators with respect to all directions ξ_γ have an eigenvalue of multiplicity $n - 1$ and that for each ξ_γ the distinguished eigen direction is the same. The submanifold is said to be an invariantly quasi-umbilical submanifold if each of its points is an invariantly quasi-umbilical point.

The normalized δ -Casorati curvature $\delta_c(n - 1)$ and $\widehat{\delta}_c(n - 1)$ are defined as

$$(2.7) \quad [\delta_c(n - 1)]_p = \frac{1}{2}\mathcal{C}_p + \frac{n + 1}{2n} \inf\{\mathcal{C}(L)|L : \text{a hyperplane of } T_pM\}$$

and

$$(2.8) \quad [\widehat{\delta}_c(n - 1)]_p = 2\mathcal{C}_p + \frac{2n - 1}{2n} \sup\{\mathcal{C}(L)|L : \text{a hyperplane of } T_pM\}.$$

For a positive real number $t \neq n(n - 1)$, put

$$(2.9) \quad a(t) = \frac{1}{nt}(n - 1)(n + t)(n^2 - n - t),$$

then the generalized normalized δ -Casorati curvatures $\delta_c(t; n - 1)$ and $\widehat{\delta}_c(t; n - 1)$ are given as

$$[\delta_c(t; n - 1)]_p = t\mathcal{C}_p + a(t) \inf\{\mathcal{C}(L)|L : \text{a hyperplane of } T_pM\},$$

if $0 < t < n^2 - n$, and

$$[\widehat{\delta}_c(t; n - 1)]_p = r\mathcal{C}_p + a(t) \sup\{\mathcal{C}(L)|L : \text{a hyperplane of } T_pM\},$$

if $t > n^2 - n$.

3. GENERALIZED NORMALIZED δ -CASORATI CURVATURE

In this section we obtain the generalized normalized δ -Casorati curvatures for different submanifolds in generalized complex space form.

Theorem 3.1. *Let M be a bi-slant submanifold of a generalized complex space form $\overline{M}(f_1, f_2)$. Then*

(i) *the generalized normalized δ -Casorati curvature $\delta_c(t; n - 1)$ satisfies*

$$(3.1) \quad \rho \leq \frac{\delta_c(t; n - 1)}{n(n - 1)} + f_1 + \frac{6f_2}{n(n - 1)}(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2),$$

for any real number t such that $0 < t < n(n - 1)$, and

(ii) *the generalized normalized δ -Casorati curvature $\widehat{\delta}_c(t; n - 1)$ satisfies*

$$(3.2) \quad \rho \leq \frac{\widehat{\delta}_c(t; n - 1)}{n(n - 1)} + f_1 + \frac{6f_2}{n(n - 1)}(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2),$$

for any real number $t > n(n - 1)$. Moreover, the equality holds in (3.1) and (3.2) if and only if M is an invariantly quasi-umbilical submanifold with trivial normal connection in $\overline{M}(f_1, f_2)$, such that with respect to suitable tangent orthonormal frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{2m}\}$, the shape operator $S_r \equiv$

S_{e_r} , $r \in \{n + 1, \dots, 2m\}$, takes the following form

$$(3.3) \quad S_{n+1} = \begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 \\ 0 & a & 0 & \dots & 0 & 0 \\ 0 & 0 & a & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a & 0 \\ 0 & 0 & 0 & \dots & 0 & \frac{n(n-1)}{t}a \end{pmatrix}, S_{n+2} = \dots = S_{2m} = 0.$$

Proof. Let $\{e_1, \dots, e_n\}$ and $\{e_{n+1}, \dots, e_{2m}\}$ be the orthonormal basis of T_pM and $T_p^\perp M$ respectively at any point $p \in M$. Putting $X = W = e_i$, $Y = Z = e_j$, $i \neq j$ from (2.1), we have

$$(3.4) \quad \begin{aligned} \bar{R}(e_i, e_j, e_j, e_i) = & f_1\{g(e_j, e_j)g(e_i, e_i) - g(e_i, e_j)g(e_j, e_i)\} \\ & + f_2\{g(e_i, Je_j)g(Je_j, e_i) - g(e_j, Je_j)g(Je_i, e_i) \\ & + 2g(e_i, Je_j)g(Je_j, e_i)\}. \end{aligned}$$

From Gauss equation and (3.4), we have

$$(3.5) \quad \begin{aligned} R(e_i, e_j, e_j, e_i) = & f_1\{g(e_j, e_j)g(e_i, e_i) - g(e_i, e_j)g(e_j, e_i)\} \\ & + 3f_2\{g(e_i, Je_j)g(Je_j, e_i)\} - g(h(e_i, e_j), h(e_j, e_i)) \\ & + g(h(e_i, e_i), h(e_j, e_j)). \end{aligned}$$

By taking summation $1 \leq i, j \leq n$ and using (2.3) and (3.5), we get

$$(3.6) \quad 2\tau = n^2\|H\|^2 - n\mathcal{C} + n(n - 1)f_1 + 6f_2(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2).$$

Define the following function, denoted by \mathcal{Q} , a quadratic polynomial in the components of the second fundamental form

$$(3.7) \quad \mathcal{Q} = n(n - 1)f_1 + 6f_2(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2) + t\mathcal{C} + a(t)\mathcal{C}(L) - 2\tau,$$

where L is the hyperplane of T_pM . Without loss of generality, we suppose that L is spanned by e_1, \dots, e_{n-1} , it follows from (3.6) and (3.7) that

$$\mathcal{Q} = \frac{n + t}{n} \sum_{\gamma=n+1}^{2m} \sum_{i,j=1}^n (h_{ij}^\gamma)^2 + \frac{a(t)}{n - 1} \sum_{\gamma=n+1}^{2m} \sum_{i,j=1}^{n-1} (h_{ij}^\gamma)^2 - \sum_{\gamma=n+1}^{2m} \left(\sum_{i=1}^n h_{ii}^\gamma \right)^2,$$

which can be easily written as

$$(3.8) \quad \begin{aligned} \mathcal{Q} = & \sum_{\gamma=n+1}^{2m} \sum_{i=1}^{n-1} \left[\left(\frac{n + t}{n} + \frac{a(t)}{n - 1} \right) (h_{ii}^\gamma)^2 + \frac{2(n + t)}{n} (h_{in}^\gamma)^2 \right] \\ & + \sum_{n+1}^{2m} \left[2 \left(\frac{n + t}{n} + \frac{a(t)}{n - 1} \right) \sum_{(i<j)=1}^n (h_{ij}^\gamma)^2 - 2 \sum_{(i<j)=1}^n h_{ii}^\gamma h_{jj}^\gamma + \frac{t}{n} (h_{nn}^\gamma)^2 \right]. \end{aligned}$$

From (3.8), we can see that the critical points

$$h^c = (h_{11}^{n+1}, h_{12}^{n+1}, \dots, h_{nn}^{n+1}, \dots, h_{11}^{2m}, \dots, h_{nn}^{2m})$$

of \mathcal{Q} are the solutions of the following system of homogenous equations:

$$(3.9) \quad \begin{cases} \frac{\partial \mathcal{Q}}{\partial h_{ii}^\gamma} = 2 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right) (h_{ii}^\gamma) - 2 \sum_{k=1}^n h_{kk}^\gamma = 0, \\ \frac{\partial \mathcal{Q}}{\partial h_{nn}^\gamma} = \frac{2t}{n} h_{nn}^\gamma - 2 \sum_{k=1}^{n-1} h_{kk}^\gamma = 0, \\ \frac{\partial \mathcal{Q}}{\partial h_{ij}^\gamma} = 4 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right) (h_{ij}^\gamma) = 0, \\ \frac{\partial \mathcal{Q}}{\partial h_{in}^\gamma} = 4 \binom{n+t}{n} (h_{in}^\gamma) = 0, \end{cases}$$

where $i, j = \{1, 2, \dots, n-1\}, i \neq j$, and $\gamma \in \{n+1, n+2, \dots, 2m\}$.

Hence, every solution h^c has $h_{ij}^\gamma = 0$ for $i \neq j$ and the corresponding determinant to the first two equations of the above system is zero. Moreover, the Hessian matrix of \mathcal{Q} is of the following form

$$\mathcal{H}(\mathcal{Q}) = \begin{pmatrix} H_1 & O & O \\ O & H_2 & O \\ O & O & H_3 \end{pmatrix},$$

where

$$H_1 = \begin{pmatrix} 2 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right) - 2 & -2 & \dots & -2 & -2 \\ -2 & 2 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right) - 2 & \dots & -2 & -2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -2 & -2 & \dots & 2 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right) - 2 & -2 \\ -2 & -2 & \dots & -2 & \frac{2t}{n} \end{pmatrix},$$

H_2 and H_3 are the diagonal matrices and O is the null matrix of the respective dimensions. H_2 and H_3 are respectively given as

$$H_2 = \text{diag} \left(4 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right), 4 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right), \dots, 4 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right) \right),$$

and

$$H_3 = \text{diag} \left(\frac{4(n+t)}{n}, \frac{4(n+t)}{n}, \dots, \frac{4(n+t)}{n} \right).$$

Hence, we find that $\mathcal{H}(\mathcal{Q})$ has the following eigenvalues

$$\lambda_{11} = 0, \lambda_{22} = 2 \left(\frac{2t}{n} + \frac{a(t)}{n-1} \right), \lambda_{33} = \dots = \lambda_{nn} = 2 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right),$$

$$\lambda_{ij} = 4 \left(\frac{n+t}{n} + \frac{a(t)}{n-1} \right), \lambda_{in} = \frac{4(n+t)}{n},$$

for all $i, j \in \{1, 2, \dots, n-1\}, i \neq j$. Thus, \mathcal{Q} is parabolic and reaches at minimum $\mathcal{Q}(h^c) = 0$ for the solution h^c of the system (3.9). Hence $\mathcal{Q} \geq 0$ and hence we have

$$2\tau \leq t\mathcal{C} + a(t)\mathcal{C}(L) + n(n-1)f_1 + 6f_2(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2),$$

whereby, we obtain

$$\rho \leq \frac{t}{n(n-1)}\mathcal{C} + \frac{a(t)}{n(n-1)}\mathcal{C}(L) + f_1 + \frac{6f_2}{n(n-1)}(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2),$$

for every tangent hyperplane L of M . If we take the infimum over all tangent hyperplane L , the result trivially follows. Moreover the equality sign holds if and only if

$$(3.10) \quad h_{ij}^\gamma = 0, \text{ for all } i, j \in \{1, \dots, n\}, i \neq j \text{ and } \gamma \in \{n+1, \dots, 2m\}$$

and

$$(3.11) \quad h_{nm}^\gamma = \frac{n(n-1)}{t}h_{11}^\gamma = \dots = \frac{n(n-1)}{t}h_{n-1n-1}^\gamma, \text{ for all } \gamma \in \{n+1, \dots, 2m\}.$$

From (3.10) and (3.11), we obtain that the equality holds if and only if the submanifold is invariantly quasi-umbilical with normal connections in \overline{M} , such that the shape operator takes the form (3.3) with respect to the orthonormal tangent and orthonormal normal frames.

In the same way, we can prove (ii). □

Theorem 3.1 yield the following result.

Corollary 3.1. *Let M be a bi-slant submanifold of a generalized complex space form $\overline{M}(f_1, f_2)$. Then*

- (i) *the normalized δ -Casorati curvature $\delta_c(n-1)$ satisfies*

$$\rho \leq \delta_c(n-1) + f_1 + \frac{6f_2}{n(n-1)}(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2).$$

Moreover, the equality sign holds if and only if M is an invariantly quasi-umbilical submanifold with trivial normal connection in $\overline{M}(f_1, f_2)$, such that with respect to suitable tangent orthonormal frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{2m}\}$, the shape operator $S_r \equiv S_{e_r}, r \in \{n+1, \dots, 2m\}$, takes the following form

$$S_{n+1} = \begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 \\ 0 & a & 0 & \dots & 0 & 0 \\ 0 & 0 & a & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a & 0 \\ 0 & 0 & 0 & \dots & 0 & 2a \end{pmatrix}, S_{n+2} = \dots = S_{2m} = 0, \text{ and}$$

- (ii) *the normalized δ -Casorati curvature $\widehat{\delta}_c(n-1)$ satisfies*

$$\rho \leq \widehat{\delta}_c(n-1) + f_1 + \frac{6f_2}{n(n-1)}(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2).$$

Moreover, the equality sign holds if and only if M is an invariantly quasi-umbilical submanifold with trivial normal connection in $\overline{M}(f_1, f_2)$, such that

with respect to suitable tangent orthonormal frame $\{e_1, \dots, e_n\}$ a and normal orthonormal frame $\{e_{n+1}, \dots, e_{2m}\}$, the shape operator $S_r \equiv S_{e_r}$, $r \in \{n + 1, \dots, 2m\}$, takes the following form

$$S_{n+1} = \begin{pmatrix} 2a & 0 & 0 & \dots & 0 & 0 \\ 0 & 2a & 0 & \dots & 0 & 0 \\ 0 & 0 & 2a & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 2a & 0 \\ 0 & 0 & 0 & \dots & 0 & a \end{pmatrix}, S_{n+2} = \dots = S_{2m} = 0.$$

Proof. (i) One can easily see that

$$(3.12) \quad \left[\delta_c \left(\frac{n(n-1)}{2} : n-1 \right) \right]_p = n(n-1) [\delta_c(n-1)]_p,$$

at any point $p \in M$. Therefore, putting $t = \frac{n(n-1)}{2}$ in (3.1) and taking into account (3.12) we have our assertion.

Similarly, we obtain (ii). □

Moreover, we have the following.

Theorem 3.2. *Let M be submanifolds of a generalized complex space form $\overline{M}(f_1, f_2)$. Then we have the following table for generalized normalized δ -Casorati curvatures:*

Table 2: Generalized Normalized δ -Casorati curvatures

S.N.	$\overline{M}(f_1, f_2)$	M	Inequality
(1)	$\overline{M}(f_1, f_2)$	semi-slant	(a) $\rho \leq \frac{\delta_c(t;n-1)}{n(n-1)} + f_1 + \frac{6f_2}{n(n-1)}(d_1 + d_2 \cos^2 \theta_2)$ (b) $\rho \leq \frac{\widehat{\delta}_c(t;n-1)}{n(n-1)} + f_1 + \frac{6f_2}{n(n-1)}(d_1 + d_2 \cos^2 \theta_2)$
(2)	$\overline{M}(f_1, f_2)$	hemi-slant	(a) $\rho \leq \frac{\delta_c(t;n-1)}{n(n-1)} + f_1 + \frac{6f_2}{n(n-1)}d_1 \cos^2 \theta_1$ (b) $\rho \leq \frac{\widehat{\delta}_c(t;n-1)}{n(n-1)} + f_1 + \frac{6f_2}{n(n-1)}d_1 \cos^2 \theta_1$
(3)	$\overline{M}(f_1, f_2)$	CR	(a) $\rho \leq \frac{\delta_c(t;n-1)}{n(n-1)} + f_1 + \frac{6f_2}{n(n-1)}d_1$ (b) $\rho \leq \frac{\widehat{\delta}_c(t;n-1)}{n(n-1)} + f_1 + \frac{6f_2}{n(n-1)}d_1$

S.N.	$\overline{M}(f_1, f_2)$	M	Inequality
(4)	$\overline{M}(f_1, f_2)$	slant	(a) $\rho \leq \frac{\delta_c(t;n-1)}{n(n-1)} + f_1 + \frac{3f_2}{(n-1)} \cos^2 \theta$ (b) $\rho \leq \frac{\widehat{\delta}_c(t;n-1)}{n(n-1)} + f_1 + \frac{3f_2}{(n-1)} \cos^2 \theta$
(5)	$\overline{M}(f_1, f_2)$	invariant	(a) $\rho \leq \frac{\delta_c(t;n-1)}{n(n-1)} + f_1 + \frac{3f_2}{(n-1)}$ (b) $\rho \leq \frac{\widehat{\delta}_c(t;n-1)}{n(n-1)} + f_1 + \frac{3f_2}{(n-1)}$
(6)	$\overline{M}(f_1, f_2)$	anti-invariant	(a) $\rho \leq \frac{\delta_c(t;n-1)}{n(n-1)} + f_1$ (b) $\rho \leq \frac{\widehat{\delta}_c(t;n-1)}{n(n-1)} + f_1$

where in each case $0 < t < n(n-1)$ and $t > n(n-1)$ for (1) and (2) respectively for any real number t . Moreover, the equality holds if and only if M is an invariantly quasi-umbilical submanifold with trivial normal connection in $\overline{M}(f_1, f_2)$, such that with respect to suitable tangent orthonormal frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{2m}\}$, the shape operator $S_r \equiv S_{e_r}$, $r \in \{n+1, \dots, 2m\}$, takes the following form

$$(3.13) \quad S_{n+1} = \begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 \\ 0 & a & 0 & \dots & 0 & 0 \\ 0 & 0 & a & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a & 0 \\ 0 & 0 & 0 & \dots & 0 & \frac{n(n-1)}{t}a \end{pmatrix}, S_{n+2} = \dots = S_{2m} = 0.$$

Proof. First four results of the Theorem 3.2 can be simply obtained with the help of Table 1 and the results in Theorem 3.1. And the next two results of the Theorem 3.2 can be seen by putting $\theta = 0$ and $\theta = \frac{\pi}{2}$ in case of invariant and anti-invariant submanifold respectively in the result of slant submanifold given in Theorem 3.2. \square

As the consequence of the last theorem, we note the following corollary.

Corollary 3.2. *Let M be submanifolds of a generalized complex space form $\overline{M}(f_1, f_2)$. Then for the normalized δ -Casorati we have the following table*

Table 3: Normalized δ -Casorati curvatures

S.N.	$\overline{M}(f_1, f_2)$	M	Inequality
(1)	$\overline{M}(f_1, f_2)$	semi-slant	(a) $\rho \leq \delta_c(n - 1) + f_1 + \frac{6f_2}{n(n-1)}(d_1 + d_2 \cos^2 \theta_2)$ (b) $\rho \leq \widehat{\delta}_c(n - 1) + f_1 + \frac{6f_2}{n(n-1)}(d_1 + d_2 \cos^2 \theta_2)$
(2)	$\overline{M}(f_1, f_2)$	hemi-slant	(a) $\rho \leq \delta_c(n - 1) + f_1 + \frac{6f_2}{n(n-1)}d_1 \cos^2 \theta_1$ (b) $\rho \leq \widehat{\delta}_c(n - 1) + f_1 + \frac{6f_2}{n(n-1)}d_1 \cos^2 \theta_1$
(3)	$\overline{M}(f_1, f_2)$	CR	(a) $\rho \leq \delta_c(n - 1) + f_1 + \frac{6f_2}{n(n-1)}d_1$ (b) $\rho \leq \widehat{\delta}_c(n - 1) + f_1 + \frac{6f_2}{n(n-1)}d_1$
(4)	$\overline{M}(f_1, f_2)$	slant	(a) $\rho \leq \delta_c(n - 1) + f_1 + \frac{3f_2}{(n-1)}\cos^2 \theta$ (b) $\rho \leq \widehat{\delta}_c(n - 1) + f_1 + \frac{3f_2}{(n-1)}\cos^2 \theta$
(5)	$\overline{M}(f_1, f_2)$	invariant	(a) $\rho \leq \delta_c(n - 1) + f_1 + \frac{3f_2}{(n-1)}$ (b) $\rho \leq \widehat{\delta}_c(n - 1) + f_1 + \frac{3f_2}{(n-1)}$
(6)	$\overline{M}(f_1, f_2)$	anti-invariant	(a) $\rho \leq \delta_c(n - 1) + f_1$ (b) $\rho \leq \widehat{\delta}_c(n - 1) + f_1$

Moreover, the equality sign for the inequalities δ_c in the above holds if and only if M is an invariantly quasi-umbilical submanifold with trivial normal connection in $\overline{M}(f_1, f_2)$, such that with respect to suitable tangent orthonormal frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{2m}\}$, the shape operator $S_r \equiv S_{e_r}$, $r \in \{n + 1, \dots, 2m\}$, takes the following form

$$S_{n+1} = \begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 \\ 0 & a & 0 & \dots & 0 & 0 \\ 0 & 0 & a & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a & 0 \\ 0 & 0 & 0 & \dots & 0 & 2a \end{pmatrix}, S_{n+2} = \dots = S_{2m} = 0.$$

and the equality sign for the inequalities $\widehat{\delta}_c$ in the above table holds if and only if M is an invariantly quasi-umbilical submanifold with trivial normal connection in $\overline{M}(f_1, f_2)$, such that with respect to suitable tangent orthonormal frame $\{e_1, \dots, e_n\}$ and normal orthonormal frame $\{e_{n+1}, \dots, e_{2m}\}$, the shape operator $S_r \equiv S_{e_r}$, $r \in \{n+1, \dots, 2m\}$, takes the following form

$$S_{n+1} = \begin{pmatrix} 2a & 0 & 0 & \dots & 0 & 0 \\ 0 & 2a & 0 & \dots & 0 & 0 \\ 0 & 0 & 2a & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 2a & 0 \\ 0 & 0 & 0 & \dots & 0 & a \end{pmatrix}, S_{n+2} = \dots = S_{2m} = 0.$$

4. INEQUALITY FOR DOUBLY WARPED PRODUCT BI-SLANT SUBMANIFOLD

Let M_1 and M_2 be Riemannian manifolds of dimension n_1 and n_2 , equipped with Riemannian metric g_1 and g_2 , respectively. Let σ_1 and σ_2 be positive differentiable functions on M_1 and M_2 , respectively. Thus, the doubly warped product $M = \sigma_2 M_1 \times_{\sigma_1} M_2$ [17] of dimension n is defined based on the product manifold $M_1 \times M_2$ endowed with a metric $g = \sigma_2^2 g_1 + \sigma_1^2 g_2$.

In this section, mainly we will prove the following.

Theorem 4.1. *Let $\phi : M = \sigma_2 M_1 \times_{\sigma_1} M_2 \rightarrow \overline{M}(f_1, f_2)$ be an isometric immersion of an n -dimensional doubly warped product bi-slant submanifold of a $2m$ -dimensional generalized complex space form. Then*

$$(4.1) \quad \begin{aligned} n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} &\leq \frac{n^2}{2} \|H\|^2 + f_1 n_1 n_2 \\ &+ \frac{3}{2} f_2 (n_1 \cos^2 \theta_1 + n_2 \cos^2 \theta_2) - \frac{n}{2} \mathcal{C} + n \mathcal{C}(L), \end{aligned}$$

where $n_i = \dim M_i$, $i = 1, 2$, $n = n_1 + n_2$ and Δ_i is Laplacian operator on M_i , $i = 1, 2$.

Proof. Let us assume that $\phi : M = \sigma_2 M_1 \times_{\sigma_1} M_2 \rightarrow \overline{M}(f_1, f_2)$ be an isometric immersion of a warped product $\sigma_2 M_1 \times_{\sigma_1} M_2$ into a generalized complex space form $\overline{M}(f_1, f_2)$. Suppose that n_1, n_2 and n are the dimensions of M_1, M_2 and M , respectively. Then for the unit vector fields X, Z tangent to M_1, M_2 , respectively, it is easily seen that [17]

$$K(X \wedge Z) = \frac{1}{\sigma_1} \{(\nabla_X^1 X)\sigma_1 - X^2 \sigma_1\} + \frac{1}{\sigma_2} \{(\nabla_Z^2 Z)\sigma_2 - Z^2 \sigma_2\}.$$

Let us assume a local orthonormal frame $\{e_1, e_2, \dots, e_n\}$ such that e_1, e_2, \dots, e_{n_1} are tangent to M_1 and e_{n_1+1}, \dots, e_n are tangent to M_2 . Then

$$(4.2) \quad n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} = \sum_{1 \leq i \leq n_1} \sum_{n_1+1 \leq j \leq n} K(e_i \wedge e_j).$$

Further, equations (2.4) and (4.2) imply

$$(4.3) \quad n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} = \tau - \sum_{1 \leq j < k \leq n_1} K(e_j \wedge e_k) - \sum_{n_1+1 \leq s < t \leq n} K(e_s \wedge e_t).$$

Moreover, (2.1), (2.2) and (4.3) yield the following

$$(4.4) \quad \begin{aligned} n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} = & \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 \\ & - 3f_2 \sum_{1 \leq j < k \leq n_1} g^2(e_j, J e_k) - 3f_2 \sum_{n_1+1 \leq s < t \leq n} g^2(e_s, J e_t) \\ & - \sum_{r=n+1}^{2m} \sum_{1 \leq j < k \leq n_1} (h_{jj}^r h_{kk}^r - (h_{jk}^r)^2) \\ & - \sum_{r=n+1}^{2m} \sum_{n_1+1 \leq s < t \leq n} (h_{ss}^r h_{tt}^r - (h_{st}^r)^2). \end{aligned}$$

Using (2.5), (2.6) and (4.4), we obtain

$$(4.5) \quad \begin{aligned} n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} = & \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 \\ & - \frac{3}{2} f_2 (n_1 \cos^2 \theta_1 + n_2 \cos^2 \theta_2) - \frac{1}{2} n_1^2 \|H_1\|^2 \\ & - \frac{1}{2} n_2^2 \|H_2\|^2 + \frac{1}{2} n_1 \mathcal{C}(L) + \frac{1}{2} n_2 \mathcal{C}(L) \\ \leq & \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 \\ & - \frac{3}{2} f_2 (n_1 \cos^2 \theta_1 + n_2 \cos^2 \theta_2) + \frac{1}{2} n \mathcal{C}(L), \end{aligned}$$

where H_1 and H_2 are partial mean curvature vectors of M_1 and M_2 respectively. \square

An immediate consequence of the Theorem 4.1 is the following obstruction result.

Corollary 4.1. *Let $\phi : M = {}_{\sigma_2}M_1 \times_{\sigma_1} M_2 \rightarrow \overline{M}(f_1, f_2)$ be an isometric immersion of an n -dimensional doubly warped product bi-slant submanifold M in $2m$ -dimensional generalized complex space form $\overline{M}(f_1, f_2)$ such that σ_1 and σ_2 are harmonic functions. Then immersion of M can not be flat in $\overline{M}(f_1, f_2)$ if*

$$f_1 \leq \frac{1}{2n_1 n_2 - n^2 + n} [f_2 (n_1 \cos^2 \theta_1 + n_2 \cos^2 \theta_2) - n \mathcal{C}(L)].$$

Next, we state and prove.

Theorem 4.2. *Let $\phi : M = {}_{\sigma_2}M_1 \times_{\sigma_1} M_2 \rightarrow \overline{M}(f_1, f_2)$ be an isometric immersion of an n -dimensional doubly warped product submanifolds of a $2m$ -dimensional generalized complex space form. Then we have following table:*

Table 4: Doubly warped product inequalities

S.N.	$\overline{M}(f_1, f_2)$	M	Inequality
(1)	$\overline{M}(f_1, f_2)$	semi-slant	$n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} \leq \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 - \frac{3}{2} f_2 (n_1 + n_2 \cos^2 \theta_2) + \frac{1}{2} n \mathcal{C}(L)$
(2)	$\overline{M}(f_1, f_2)$	hemi-slant	$n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} \leq \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 - \frac{3}{2} f_2 (n_1 \cos^2 \theta_1) + \frac{1}{2} n \mathcal{C}(L)$
(3)	$\overline{M}(f_1, f_2)$	CR	$n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} \leq \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 - \frac{3}{2} f_2 n_1 + \frac{1}{2} n \mathcal{C}(L)$
(4)	$\overline{M}(f_1, f_2)$	slant	$n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} \leq \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 - \frac{3}{2} f_2 n \cos^2 \theta + \frac{1}{2} n \mathcal{C}(L)$
(5)	$\overline{M}(f_1, f_2)$	invariant	$n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} \leq \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 - \frac{3}{2} f_2 n + \frac{1}{2} n \mathcal{C}(L)$
(6)	$\overline{M}(f_1, f_2)$	anti-invariant	$n_2 \frac{\Delta_1 \sigma_1}{\sigma_1} + n_1 \frac{\Delta_2 \sigma_2}{\sigma_2} \leq \tau - \frac{f_1}{2} \{n(n-1)\} + f_1 n_1 n_2 + \frac{1}{2} n \mathcal{C}(L)$

Proof. We obtain the first four results of the Theorem 4.2 directly by using Table 1 and the result of the Theorem 4.1 and last two results by putting $\theta = 0$ and $\theta = \frac{\pi}{2}$ respectively in result of slant submanifold. \square

REFERENCES

- [1] M. Aquib, *Bounds for generalized normalized δ -Casorati curvatures for bi-slant submanifolds in T -space forms*, Filomat (to appear).
- [2] D. Blair and A. Ledger, *Quasi-umbilical minimal submanifolds of Euclidean space*, Bull. Belg. Math. Soc. Simon Stevin **51** (1977), 3–22.
- [3] F. Casorati, *Mesure de la courbure des surface suivant l’idee commune. Ses rapports avec les mesures de coubure gaussienne et moyenne*, Acta Math. **14** (1999), 95–110.
- [4] B.Y. Chen, *Relationship between Ricci curvature and shape operator for submanifolds with arbitrary codimensions*, Glasg. Math. J. **41** (1999), 33–41.
- [5] B.Y. Chen, *Some pinching and classification theorems for minimal submanifolds*, Arch. Math. **60** (1993), 568–578.
- [6] S. Decu, S. Haesen and L. Verstralelen, *Optimal inequalities involving Casorati curvatures*, Bull. Transilv. Univ. Brasov Ser. B **49** (2007), 85–93.
- [7] V. Ghisoiu, *Inequalities for the Casorati curvatures of the slant submanifolds in complex space forms*, Riemannian geometry and applications. Proceedings RIGA 2011, ed. Univ. Bucuresti, Bucharest (2011), 145–150.
- [8] A. Gray, *Nearly Kaehler manifolds*, J. Differential Geom. **4** (1970), 283–309.
- [9] S. Haesen, D. Kowalczyk and L. Verstralelen, *On the extrinsic principal directions of Riemannian submanifolds*, Note Mat. **29** (2009), 41–53.
- [10] S. Hong, K. Matsumoto and M.M. Tripathi, *Certain basic inequalities for submanifolds of locally conformal kaehler space forms*, SUT J. Math. **41**(1) (2005), 75–94.

- [11] T. Kashiwada, *Some properties of locally conformal Kaehler manifolds*, Hokkaido Math. J. **8** (1979), 191–198.
- [12] D. Kowalczyk, *Casorati curvatures*, Bull. Transilv. Univ. Brasov Ser. III **50**(1) (2008), 2009–2013.
- [13] J.S. Kim, Y.M. Song and M.M. Tripathi, *B. Y. Chen inequalities for submanifolds in generalised complex space forms*, Bull. Korean Math. Soc. **40**(3) (2003), 411–423.
- [14] C.W. Lee, J.W. Lee, G.E. Vilcu and D.W. Yoon, *Optimal inequalities for the Casorati curvatures of the submanifolds of Generalized space form endowed with semi-symmetric metric connections*, Bull. Korean Math. Soc. **52** (2015), 1631–1647.
- [15] K. Matsumoto, I. Mihai and A. Oiaga, *Ricci curvature of submanifolds in complex space form*, Rev. Roumaine Math. Pures Appl. **46** (2001), 775–782.
- [16] F. Tricerri and L. Vanhecke, *Curvature tensors on almost Hermitian manifolds*, Trans. Amer. Math. Soc. **26** (2003), 85–94.
- [17] B. Unal, *Doubly warped products*, Thesis(Ph.D.), University of Missouri, Columbia, 2000.
- [18] L. Vanhecke, *Almost Hermitian manifolds with J-invariant Riemann curvature tensor*, Rend. Semin. Mat. Univ. Politec. Torino **34** (1975-76), 487–498.
- [19] L. Verstralelen, *Geometry of submanifolds I, The first Casorati curvature indicatrices*, Kragujevac J. Math. **37** (2013), 5–23.

¹DEPARTMENT OF MATHEMATICS,
FACULTY OF NATURAL SCIENCES,
JAMIA MILLIA ISLAMIA,
NEW DELHI-110025, INDIA
Email address: aquib80@gmail.com
Email address: hasan_jmi@yahoo.com

²DEPARTMENT OF MATHEMATICS,
AL-FALAH UNIVERSITY,
HARYANA-121004, INDIA
Email address: jamali_dbd@yahoo.co.in