

REMARKS ON THE DEGREE KIRCHHOFF INDEX

M. MATEJIĆ, I. MILOVANOVIĆ, AND E. MILOVANOVIĆ

ABSTRACT. Let G be a simple connected graph with n vertices and m edges, with normalized Laplacian eigenvalues $\rho_1 \geq \rho_2 \geq \dots \geq \rho_{n-1} > \rho_n = 0$. The degree Kirchhoff index $Kf^*(G)$ is defined as $Kf^*(G) = 2m \sum_{i=1}^{n-1} \frac{1}{\rho_i}$. In this paper we obtain lower and upper bounds for $Kf^*(G)$.

1. INTRODUCTION

Let $G = (V, E)$, $V = \{1, 2, \dots, n\}$, be a simple connected graph with n , $n \geq 3$, vertices and m edges. Denote with $\Delta = d_1 \geq d_2 \geq \dots \geq d_n = \delta > 0$, a sequence of vertex degrees of G , \mathbf{A} the adjacency matrix of graph, and by $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n)$ the diagonal matrix of its vertex degrees. Matrix $\mathbf{L} = \mathbf{D} - \mathbf{A}$ is the Laplacian matrix of G . Eigenvalues of \mathbf{L} , $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} > \mu_n = 0$, form the so-called Laplacian spectrum of graph G . Provided that the graph G has no isolated vertices, the normalized Laplacian matrix is defined as $\mathbf{L}^* = \mathbf{D}^{-1/2} \mathbf{L} \mathbf{D}^{-1/2} = \mathbf{I} - \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}$. Its eigenvalues, $\rho_1 \geq \rho_2 \geq \dots \geq \rho_{n-1} > \rho_n = 0$, represent normalized Laplacian eigenvalues of G . The following identities are valid for ρ_i , $i = 1, 2, \dots, n - 1$, (see [5])

$$\sum_{i=1}^{n-1} \rho_i = n \quad \text{and} \quad \sum_{i=1}^{n-1} \rho_i^2 = n + 2R_{-1},$$

where

$$R_{-1} = \sum_{i \sim j} \frac{1}{d_i d_j},$$

is the general Randić index (also called branching index) introduced in [19]. A symbol $i \sim j$ denotes that vertices i and j are adjacent.

Key words and phrases. Degree Kirchhoff index, Laplacian eigenvalues (of graph), vertex degree.
2010 *Mathematics Subject Classification.* Primary: 05C12. Secondary: 05C50.

Received: May 30, 2017.

Accepted: August 30, 2017.

In [9], Klein and Randić, introduced the notion of resistance distance, r_{ij} , defined as the resistance between the nodes i and j in an electrical network corresponding to the graph G in which all edges are replaced by unit resistors. The sum of resistance distances of all pairs of vertices of a graph G is named as the Kirchhoff index, i.e.,

$$Kf(G) = \sum_{i < j} r_{ij}.$$

Eventually, it was shown [7] (see also [8]) that the resistance distance can be expressed in terms of Laplacian matrix and its spectrum. Namely, it was proven that

$$Kf(G) = n \sum_{i=1}^{n-1} \frac{1}{\mu_i}.$$

The Kirchhoff index spawned a family of resistance indices, such as degree Kirchhoff index, multiplicative-degree-Kirchhoff index and additive-degree Kirchhoff index (see for example [2, 16, 17]). The degree Kirchhoff index, introduced in [4], is defined as

$$Kf^*(G) = \sum_{i < j} d_i d_j r_{ij}.$$

In analogy with the Kirchhoff index, the degree Kirchhoff index can also be represented as

$$Kf^*(G) = 2m \sum_{i=1}^{n-1} \frac{1}{\rho_i}.$$

The graph invariants $Kf(G)$ and $Kf^*(G)$ are currently much studied in the mathematical and mathematico-chemical literature; see for example [11] and the references cited therein.

In this paper, inspired by the inequality for $Kf(G)$ reported in [18], we determine an upper bound for $Kf^*(G)$ which depends on n , m and k , where k is an arbitrary real number such that $\rho_{n-1} \geq k > 0$. We also prove one general inequality that sets up a lower bound for $Kf^*(G)$ in terms of n , m and s , where s is an arbitrary real number such that $\rho_1 \geq s \geq \rho_{n-1}$. For some particular values of s we obtain some lower bounds for $Kf^*(G)$ reported in the literature.

2. PRELIMINARIES

In this section we recall some results reported in the literature on lower and upper bounds for $Kf(G)$ and $Kf^*(G)$, and an analytic inequality for real number sequences which will be used in the subsequent considerations.

The following upper bound for $Kf(G)$ that depends on parameters n , m and k , where k is an arbitrary real number with the property $\mu_{n-1} \geq k > 0$, was established in [18]

$$(2.1) \quad Kf(G) \leq \frac{(n+k)(n-1) - 2m}{k},$$

with equality if and only if $k = n$ and $G \cong K_n$, or $k = 1$ and $G \cong K_{1,n-1}$, or $k = \frac{n}{2}$ and $G \cong K_{\frac{n}{2}, \frac{n}{2}}$, or $k = n - 2$ and $G \cong K_n - e$.

In [15] (see also [1, 6, 8, 12]) the following inequality was proven

$$(2.2) \quad Kf^*(G) \geq \frac{2m(n-1)^2}{n},$$

with equality if and only if $G \cong K_n$.

In [6] the following lower bound for $Kf^*(G)$ in terms of parameters n , m , and Δ was determined

$$(2.3) \quad Kf^*(G) \geq 2m \left(\frac{\Delta}{\Delta+1} + \frac{(n-2)^2}{n-1-\frac{1}{\Delta}} \right),$$

with equality if and only if $G \cong K_n$.

In [3] it was proven that

$$(2.4) \quad Kf^*(G) \geq 2m \left(\frac{1}{P} + \frac{(n-2)^2}{n-P} \right),$$

where

$$P = 1 + \sqrt{\frac{2R_{-1}}{n(n-1)}}.$$

Equality in (2.4) is attained if and only if $G \cong K_n$.

In [10] the following upper bound for $Kf^*(G)$ was established

$$(2.5) \quad Kf^*(G) \leq 2m \frac{(n-1)(\rho_1 + \rho_{n-1}) - n}{\rho_1 \rho_{n-1}}.$$

Let $a_{k_1}, a_{k_1+1}, \dots, a_{n-k_2}$ and $b_{k_1}, b_{k_1+1}, \dots, b_{n-k_2}$ be two non-negative real number sequences of similar monotonicity, and $p_{k_1}, p_{k_1+1}, \dots, p_{n-k_2}$ be positive real number sequence, $1 \leq k_1 \leq n - k_2$, $0 \leq k_2 \leq n - 1$. Then the following inequality holds (see, for example, [13, 14])

$$(2.6) \quad \sum_{i=k_1}^{n-k_2} p_i \sum_{i=k_1}^{n-k_2} p_i a_i b_i \geq \sum_{i=k_1}^{n-k_2} p_i a_i \sum_{i=k_1}^{n-k_2} p_i b_i.$$

3. MAIN RESULTS

In the following theorem we establish an upper bound for $Kf^*(G)$ in terms of parameters n , m and k , $\rho_{n-1} \geq k > 0$. The inequality proven in this theorem is analogous to the inequality (2.1) for $Kf(G)$.

Theorem 3.1. *Let G be a simple connected graph with n , $n \geq 3$, vertices and m edges. Then, for any real k with the property $\rho_{n-1} \geq k > 0$, holds*

$$(3.1) \quad Kf^*(G) \leq m \frac{n-2+k(n-1)}{k},$$

with equality if and only if $k = \frac{n}{n-1}$ and $G \cong K_n$, or $k = 1$ and $G \cong K_{r,n-r}$, $1 \leq r \leq \lfloor \frac{n}{2} \rfloor$, or $k = \frac{3}{n-1}$ and $G \cong C_n$.

Proof. Consider the function

$$f(x) = \frac{(n-1)(x + \rho_{n-1}) - n}{x}, \quad x > 0.$$

This is an increasing function for $x > 0$. Since $0 < x = \rho_1 \leq 2$, we have $f(x) = f(\rho_1) \leq f(2)$, and according to (2.5) we get

$$(3.2) \quad Kf^*(G) \leq 2m \frac{(n-1)(2 + \rho_{n-1}) - n}{2\rho_{n-1}} = m \frac{(n-1)\rho_{n-1} + n - 2}{\rho_{n-1}}.$$

Now, consider the function

$$g(x) = \frac{n-2}{x}, \quad x > 0.$$

This is a decreasing function for $x > 0$. Since $x = \rho_{n-1} \geq k > 0$, then $g(\rho_{n-1}) \leq g(k)$, and therefore from (3.2) we get

$$Kf^*(G) \leq m \frac{n-2 + k(n-1)}{k},$$

which completes the proof. \square

Corollary 3.1. *Let G be a simple connected planar graph with n , $n \geq 3$, vertices and m edges. Then, for any real k with the property $\rho_{n-1} \geq k > 0$, holds*

$$(3.3) \quad Kf^*(G) \leq \frac{3(n-2)(n-2 + (n-1)k)}{k},$$

with equality if and only if $k = \frac{3}{2}$ and $G \cong K_3$, or $k = \frac{4}{3}$ and $G \cong K_4$.

Proof. Since for simple connected planar graphs holds $m \leq 3(n-2)$, the inequality (3.3) directly follows from (3.1). \square

In the next theorem we determine a lower bound for the degree Kirchhoff index in terms of parameters n , m and s , where s is a real number with the property $\rho_1 \geq s \geq \rho_{n-1}$.

Theorem 3.2. *Let G be a simple connected graph with n , $n \geq 3$, vertices and m edges. Then, for any real s with the property $\rho_1 \geq s \geq \rho_{n-1}$, holds*

$$(3.4) \quad Kf^*(G) \geq 2m \left(\frac{1}{s} + \frac{(n-2)^2}{n-s} \right),$$

with equality if and only if $s = \frac{n}{n-1}$ and $G \cong K_n$, or $s = 2$ and $G \cong K_{r,n-r}$, $1 \leq r \leq \lfloor \frac{n}{2} \rfloor$.

Proof. For $k_1 = 1$, $k_2 = 2$, $p_i = \rho_i$, $a_i = b_i = \frac{1}{\rho_i}$, $i = 1, 2, \dots, n-2$, the inequality (2.6) becomes

$$\sum_{i=1}^{n-2} \rho_i \sum_{i=1}^{n-2} \frac{1}{\rho_i} \geq \left(\sum_{i=1}^{n-2} 1 \right)^2,$$

i.e.

$$(n - \rho_{n-1}) \left(\sum_{i=1}^{n-1} \frac{1}{\rho_i} - \frac{1}{\rho_{n-1}} \right) \geq (n - 2)^2,$$

wherefrom follows

$$(3.5) \quad Kf^*(G) \geq 2m \left(\frac{1}{\rho_{n-1}} + \frac{(n-2)^2}{n - \rho_{n-1}} \right),$$

with equality if and only if $\rho_1 = \rho_2 = \dots = \rho_{n-2}$.

For $k_1 = 2$, $k_2 = 1$, $p_i = \rho_i$, $a_i = b_i = \frac{1}{\rho_i}$, $i = 2, 3, \dots, n-1$, the inequality (2.6) transforms into

$$\sum_{i=2}^{n-1} \rho_i \sum_{i=2}^{n-1} \frac{1}{\rho_i} \geq \left(\sum_{i=2}^{n-1} 1 \right)^2,$$

i.e.,

$$\left(\sum_{i=1}^{n-1} \rho_i - \rho_1 \right) \left(\sum_{i=1}^{n-1} \frac{1}{\rho_i} - \frac{1}{\rho_1} \right) \geq (n - 2)^2,$$

wherefrom follows

$$(3.6) \quad Kf^*(G) \geq 2m \left(\frac{1}{\rho_1} + \frac{(n-2)^2}{n - \rho_1} \right),$$

with equality if and only if $\rho_2 = \rho_3 = \dots = \rho_{n-1}$.

Consider the function

$$f(x) = \frac{1}{x} + \frac{(n-2)^2}{n-x}.$$

This is an increasing function for $x \geq \frac{n}{n-1}$, monotone decreasing for $x \leq \frac{n}{n-1}$ and has a minimum for $x = \frac{n}{n-1}$. Then for any real s , such that $\rho_1 \geq s \geq \frac{n}{n-1}$, from (3.6) follows

$$Kf^*(G) \geq 2mf(\rho_1) \geq 2mf(s) = 2m \left(\frac{1}{s} + \frac{(n-2)^2}{n-s} \right).$$

Also, for any real s with the property $\frac{n}{n-1} \geq s \geq \rho_{n-1}$, from (3.5) follows

$$Kf^*(G) \geq 2mf(\rho_{n-1}) \geq 2mf(s) = 2m \left(\frac{1}{s} + \frac{(n-2)^2}{n-s} \right).$$

Finally, from the last two inequalities we obtain (3.4). □

According to (3.5) and (3.6) we have the following corollary of Theorem 3.2.

Corollary 3.2. *Let G be a simple connected graph with n , $n \geq 3$, vertices and m edges. Then*

$$Kf^*(G) \geq 2m \max \left\{ \frac{1}{\rho_1} + \frac{(n-2)^2}{n - \rho_1}, \frac{1}{\rho_{n-1}} + \frac{(n-2)^2}{n - \rho_{n-1}} \right\},$$

with equality if and only if $G \cong K_n$, or $G \cong K_{r,n-r}$, $1 \leq r \leq \lfloor \frac{n}{2} \rfloor$.

Remark 3.1. In [3] the following inequalities were proven

$$\rho_1 \geq P \geq \frac{\Delta + 1}{\Delta} \geq \frac{n}{n-1} \geq \rho_{n-1}.$$

Now, for $s = \frac{n}{n-1}$ from (3.4) the inequality (2.2) is obtained, for $s = \frac{\Delta+1}{\Delta}$ we get (2.3), and for $s = P$ it follows (2.4).

REFERENCES

- [1] M. Bianchi, A. Cornaro, J. L. Palacios and A. Torriero, *Bounds for the Kirchhoff index via majorization techniques*, J. Math. Chem. **51** (2013), 569–587.
- [2] M. Bianchi, A. Cornaro, J. L. Palacios, J. M. Renom and A. Torriero, *Revisiting bounds for the multiplicative degree–Kirchhoff index*, MATCH Commun. Math. Comput. Chem. **75** (2016), 227–231.
- [3] S. B. Bozkurt and D. Bozkurt, *On the sum of powers of normalized Laplacian eigenvalues of graphs*, MATCH Commun. Math. Comput. Chem. **68** (2012), 917–930.
- [4] H. Chen and F. Zhang, *Resistance distance and the normalized Laplacian spectrum*, Discrete Appl. Math. **155** (2007), 654–661.
- [5] F. R. K. Chung, *Spectral Graph Theory*, AMS Press, Providence, 1997.
- [6] L. Feng, G. Yu and W. Liu, *Further results regarding the degree Kirchhoff index of graphs*, Miskolc Math. Notes **15**(1) (2014), 97–108.
- [7] I. Gutman and B. Mohar, *The quasi–Wiener and the Kirchhoff indices coincide*, Journal of Chemical Information and Modeling **36**(5) (1996), 982–985.
- [8] J. Huang and S. Li, *On the normalized Laplacian spectrum, degree–Kirchhoff index and spanning trees of graphs*, Bull. Aust. Math. Soc. **91** (2015), 353–367.
- [9] D. J. Klein and M. Randić, *Resistance distance*, J. Math. Chem. **12** (1993), 81–95.
- [10] I. Milovanović, I. Gutman and E. Milovanović, *On Kirchhoff and degree Kirchhoff indices*, Filomat **29**(8) (2015), 1869–1877.
- [11] I. Ž. Milovanović and E. I. Milovanović, *Bounds of Kirchhoff and degree Kirchhoff indices*, in: I. Gutman, B. Furtula, K. C. Das, E. Milovanović, I. Milovanović, Eds. *Bounds in Chemical Graph Theory - Mainstreams*, Mathematical Chemistry Monographs, MCM 20, University of Kragujevac, Kragujevac, (2017), pp. 93–119.
- [12] I. Ž. Milovanović, E. I. Milovanović and E. Glogić, *Lower bounds of the Kirchhoff and degree Kirchhoff indices*, Scientific Publications of the State University of Novi Pazar Series A: Applied Mathematics, Informatics and Mechanics **7**(1) (2015), 25–31.
- [13] D. S. Mitrinović and P. M. Vasić, *Analytic Inequalities*, Springer Verlag, Berlin-Heidelberg-New York, 1970.
- [14] D. S. Mitrinović and P. M. Vasić, *History, variations and generalisations of the Čebyšev inequality and the question of some priorities*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. **461–497** (1974), 1–30.
- [15] J. L. Palacios and J. M. Renom, *Broder and Karlin’s formula for hitting times and the Kirchhoff index*, Int. J. Quantum Chem. **111** (2011), 35–39.
- [16] J. L. Palacios, *A new lower bound for the multiplicative degree–Kirchhoff index*, MATCH Commun. Math. Comput. Chem. **76** (2016), 251–254.
- [17] J. L. Palacios, *Some interplay of the three Kirchhoffian indices*, MATCH Commun. Math. Comput. Chem. **75** (2016), 199–206.
- [18] S. Pirzada, H. A. Ganie and I. Gutman, *Comparison between Laplacian–energy-like invariant and Kirchhoff index*, Electron. J. Linear Algebra **31** (2016), 27–41.
- [19] M. Randić, *On characterization of molecular branching*, J. Amer. Chem. Soc. **97** (1975), 6609–6615.

- [20] H. Y. Zhu, D. J. Klein and I. Lukovits, *Extensions of the Wiener number*, Journal of Chemical Information and Modeling **36** (1996), 420–428.

FACULTY OF ELECTRONIC ENGINEERING,
UNIVERSITY OF NIŠ,
NIŠ, SERBIA
Email address: {marjan.matejic, igor, ema}@elfak.ni.ac.rs