

## SOME NEW BOUNDS ON RANDIĆ ENERGY

E. ZOGIĆ<sup>1</sup> AND B. BOROVIĆANIN<sup>2</sup>

ABSTRACT. Let  $G = (V, E)$  be a simple graph of order  $n$  with vertex set  $V = V(G) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E = E(G)$ . Let  $d_i$  be the degree of the vertex  $v_i$  in  $G$  for  $i = 1, 2, \dots, n$ . The Randić matrix  $\mathbf{R} = \mathbf{R}(G) = ||R_{ij}||_{n \times n}$  is defined by

$$R_{ij} = \begin{cases} \frac{1}{\sqrt{d_i d_j}}, & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent,} \\ 0, & \text{otherwise.} \end{cases}$$

The eigenvalues of matrix  $\mathbf{R}$ , denoted by  $\rho_1, \rho_2, \dots, \rho_n$ , are called the Randić eigenvalues of graph  $G$ . The Randić energy of graph  $G$ , denoted by  $RE$ , is defined as

$$RE = RE(G) = \sum_{i=1}^n |\rho_i|.$$

In this paper we establish some new upper and lower bounds on Randić energy.

### 1. INTRODUCTION

Let  $G = (V, E)$  be a simple graph with vertex set  $V = V(G) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E = E(G)$ . Denote by  $d_i$  the degree of vertex  $v_i \in V$ , i.e., the number of the vertices adjacent to  $v_i$ ,  $i = 1, 2, \dots, n$ . A vertex of degree zero is said to be isolated. In this paper we consider graphs without such vertices. Let  $\mathbf{A} = \mathbf{A}(G)$  be the adjacency matrix of  $G$  and denote by  $\mathbf{D} = \mathbf{D}(G)$  the diagonal matrix whose diagonal elements are vertex degrees of  $G$ .

---

*Key words and phrases.* Normalized Laplacian matrix, Randić matrix, Randić energy.  
*2010 Mathematics Subject Classification.* Primary: 05C50.  
*Received:* December 04, 2017.  
*Accepted:* December 27, 2017.

The Randić matrix of  $G$  is the  $n \times n$  matrix  $\mathbf{R} = \mathbf{R}(G) = ||R_{ij}||$ , defined by

$$R_{ij} = \begin{cases} \frac{1}{\sqrt{d_i d_j}}, & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent,} \\ 0, & \text{otherwise.} \end{cases}$$

The eigenvalues  $\rho_1, \rho_2, \dots, \rho_n$  of matrix  $\mathbf{R}$  are called the Randić eigenvalues of a graph  $G$ , and suppose they are labeled in a non-increasing order, i.e.,  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ . The Randić energy  $RE = RE(G)$  of a graph  $G$  is defined as [2, 3, 12]

$$RE = RE(G) = \sum_{i=1}^n |\rho_i|.$$

For recent results on Randić energy we refer the reader to the papers [1, 9–11, 13, 16]. Also, several lower and upper bounds on Randić energy were obtained in [2, 3, 7, 12, 14] and [15].

Randić matrix occurs in a natural way within Laplacian spectral theory and provides the non-trivial part of the so-called normalized Laplacian matrix [3]. Let  $\mathbf{D} = \mathbf{D}(G)$  be the diagonal matrix of order  $n$  whose  $i$ -th diagonal entry is  $d_i$ . Then the Laplacian matrix of  $G$  is defined as  $\mathbf{L} = \mathbf{L}(G) = \mathbf{D} - \mathbf{A}$ . If graph  $G$  has no isolated vertices, then the matrix  $\mathbf{D}^{-1/2}$  exists. For a graph without isolated vertices, the normalized Laplacian matrix can be defined [6] as

$$\tilde{\mathbf{L}} = \mathbf{D}^{-1/2} \mathbf{L} \mathbf{D}^{-1/2} = \mathbf{I} - \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2},$$

where  $\mathbf{I}$  is the unit matrix of order  $n$ .

Recall that  $\mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2} = \mathbf{R}$  for a graph without isolated vertices, implying the following relation [3] between the Randić matrix and the normalized Laplacian matrix of a graph without isolated vertices

$$(1.1) \quad \tilde{\mathbf{L}} = \mathbf{I} - \mathbf{R}.$$

The rest of the paper is organized as follows. In Section 2 we state some previously known results needed for the subsequent considerations. In Section 3 we obtain some new upper and lower bounds for the Randić energy. In addition, we determine a new lower bound on  $RE$  which is better than the lower bound obtained in [7, 15].

## 2. PRELIMINARIES

In this section we review some properties of normalized Laplacian eigenvalues, Randić eigenvalues and Randić energy. Besides, we recall some analytic inequalities for real number sequences that are of interest for the subsequent considerations.

If we denote by  $\tilde{\mu}_1 \geq \tilde{\mu}_2 \geq \dots \geq \tilde{\mu}_{n-1} \geq \tilde{\mu}_n$  the normalized Laplacian eigenvalues of a graph  $G$ , then from (1.1) it follows

$$(2.1) \quad \rho_i = 1 - \tilde{\mu}_{n-i+1}, \quad i = 1, 2, \dots, n.$$

There are numerous results for the  $\tilde{\mathbf{L}}$ -eigenvalues, see [4] for example. From [4] it follows that  $0 \leq \tilde{\mu}_i \leq 2, i = 1, 2, \dots, n$ , which implies, by (2.1),

$$(2.2) \quad -1 \leq \rho_i \leq 1, \quad i = 1, 2, \dots, n.$$

Besides, the following results for the Randić eigenvalues are known.

**Lemma 2.1.** [12, 17] *Let  $G$  be a graph on  $n$  vertices,  $n \geq 1$ , and let  $\rho_1$  be the largest eigenvalue of its Randić matrix. Then  $\rho_1 = 0$  if and only if  $G \cong \overline{K}_n$ . If  $G$  possesses at least one edge, then  $\rho_1 = 1$ .*

**Lemma 2.2.** [12] *Let  $G$  be a graph on  $n$  vertices, and let  $\mathbf{A}$  and  $\mathbf{R}$  be its adjacency and Randić matrices. If  $\mathbf{A}$  has  $n_+, n_0$ , and  $n_-$  positive, zero, and negative eigenvalues, respectively ( $n_+ + n_0 + n_- = n$ ), then  $\mathbf{R}$  has  $n_+, n_0$ , and  $n_-$  positive, zero, and negative eigenvalues, respectively.*

**Lemma 2.3.** [12] *Let  $G$  be a graph on  $n$  vertices, and let  $\mathbf{A}$  and  $\mathbf{R}$  be its adjacency and Randić matrices. If  $G$  possesses isolated vertices, then  $\det \mathbf{R} = \det \mathbf{A} = 0$ , otherwise*

$$\det \mathbf{R} = \frac{1}{d_1 d_2 \cdots d_n} \det \mathbf{A}.$$

Denote by  $\eta$  the nullity of a graph  $G$ , i.e., the multiplicity of 0 as an eigenvalue of  $\mathbf{A} = \mathbf{A}(G)$ . Lemma 2.2 implies that  $\eta$  is also the multiplicity of 0 as an eigenvalue of the matrix  $\mathbf{R}$ . In addition, as we are concerned with graphs without isolated vertices, it holds  $\eta(G) < n$ .

Denote by  $\rho_1^* \geq \rho_2^* \geq \cdots \geq \rho_n^*$  the absolute values of the Randić eigenvalues of  $G$  labeled in a non-increasing order. Then, by (2.2),  $\rho_i^* \in [0, 1]$ , for  $i = 1, 2, \dots, n$ .

In the sequel we review some known inequalities needed for subsequent considerations.

**Lemma 2.4.** [5] *Let  $a_1, a_2, \dots, a_n$  be positive real numbers such that  $0 < a_1 \leq \cdots \leq a_i \leq \cdots \leq a_k \leq \cdots \leq a_n$ . Then*

$$(2.3) \quad a_1 + a_2 + \cdots + a_n - n \sqrt[n]{a_1 a_2 \cdots a_n} \geq P(\sqrt{a_k} - \sqrt{a_i})^2,$$

where

$$P = \begin{cases} \frac{2i(n-k+1)}{n+i-k+1}, & i+k \leq n+1, \\ n-k+1, & i+k \geq n+1. \end{cases}$$

**Lemma 2.5.** [19] *Let  $a_i \in \mathbb{R}^+, i = 1, \dots, n$ . Then*

$$(2.4) \quad (n-1) \sum_{i=1}^n a_i + n \left( \prod_{i=1}^n a_i \right)^{\frac{1}{n}} \geq \left( \sum_{i=1}^n \sqrt{a_i} \right)^2 \geq \sum_{i=1}^n a_i + n(n-1) \left( \prod_{i=1}^n a_i \right)^{\frac{1}{n}}.$$

**Lemma 2.6.** [18] *Let  $a_1, \dots, a_n$  and  $b_1, \dots, b_n$  be real numbers such that  $a \leq a_i \leq A$  and  $b \leq b_i \leq B, i = 1, \dots, n$ . Then*

$$(2.5) \quad -(A - a)(B - b)n\alpha(n) \leq \sum_{i=1}^n a_i b_i - \frac{1}{n} \sum_{i=1}^n a_i \sum_{i=1}^n b_i \leq (A - a)(B - b)n\alpha(n),$$

where  $\alpha(n) = \frac{1}{4} \left(1 - \frac{(-1)^{n+1} + 1}{2n^2}\right)$ .

### 3. MAIN RESULTS

In recent papers [7, 15] the following lower bound on  $RE$  was obtained.

**Theorem 3.1.** [7, 15] *Let  $G$  be a graph on  $n$  vertices. Then*

$$(3.1) \quad RE(G) \geq 1 + (n - 1) |\det \mathbf{R}|^{\frac{1}{n-1}},$$

with equality if and only if  $G$  is a complete graph or a non-bipartite graph with three distinct Randić eigenvalues

$$\left( 1, \sqrt{\frac{2 \sum_{v_i v_j \in E(G)} \frac{1}{d_i d_j}}{n - 1}}, -\sqrt{\frac{2 \sum_{v_i v_j \in E(G)} \frac{1}{d_i d_j}}{n - 1}} \right).$$

In the sequel we obtain the lower bound on  $RE$  which improves the result (3.1).

**Theorem 3.2.** *Let  $G$  be a graph on  $n$  vertices,  $n \geq 2$ , and let  $\rho_1^* \geq \rho_2^* \geq \dots \geq \rho_n^*$  be the absolute values of its Randić eigenvalues labeled in a non-increasing order.*

(a) *If  $G$  is a non-singular graph, i.e.,  $\eta(G) = 0$ , then*

$$(3.2) \quad RE(G) \geq 1 + (n - 1) |\det \mathbf{R}|^{\frac{1}{n-1}} + (\sqrt{\rho_n^*} - \sqrt{\rho_2^*})^2.$$

(b) *If  $\eta(G) = n - j$ ,  $0 < j < n$ , then*

$$(3.3) \quad RE(G) \geq 1 + (j - 1) \left( \prod_{i=2}^{j-1} \rho_i^* \right)^{\frac{1}{j-1}} + (\sqrt{\rho_j^*} - \sqrt{\rho_2^*})^2.$$

*Proof.* (a) Since  $\eta(G) = 0$ , it follows that  $\rho_i^* \in (0, 1]$ , implying  $RE(G) = \rho_1^* + \sum_{i=2}^n \rho_i^*$ .

If we apply the inequality (2.3) to  $\sum_{i=2}^n \rho_i^*$  letting  $a_{i-1} = \rho_i^*$ ,  $i = 2, \dots, n$ , whereas  $k = n - 1$  and  $i = 1$ , we obtain that  $P = 1$ , and so

$$\begin{aligned} \sum_{i=2}^n \rho_i^* &\geq (n - 1) \sqrt[n-1]{\rho_2^* \cdots \rho_n^*} + (\sqrt{\rho_n^*} - \sqrt{\rho_2^*})^2 \\ &= (n - 1) |\det \mathbf{R}|^{\frac{1}{n-1}} + (\sqrt{\rho_n^*} - \sqrt{\rho_2^*})^2. \end{aligned}$$

By Lemma 2.1,  $\rho_1^* = 1$ , which completes the proof.

(b) We now turn to the case  $\eta(G) = n - j$ ,  $0 < j < n$ . If we apply the inequality (2.3) in the same manner as in case (a) to non-zero absolute values of Randić eigenvalues  $1 = \rho_1^* \geq \rho_2^* \geq \dots \geq \rho_j^* > 0$ , we obtain (3.3).  $\square$

*Remark 3.1.* If  $G$  is a non-singular graph, then the lower bound (3.2) is better than the lower bound (3.1) from [7, 15]. If  $\eta(G) > 0$ , then  $\det \mathbf{R} = 0$  and the lower bound (3.3) is also better than the lower bound (3.1) from [7, 15].

**Theorem 3.3.** *Let  $G$  be a non-singular graph on  $n$  vertices,  $n \geq 2$ , and let  $\rho_1^* \geq \rho_2^* \geq \dots \geq \rho_n^*$  be the absolute values of its Randić eigenvalues labeled in a non-increasing order. Then*

$$(3.4) \quad -\frac{n^2}{n-1}\alpha(n)(\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 + n|\det \mathbf{R}|^{\frac{1}{n}} \leq RE(G) \leq n^2\alpha(n)(\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 + n|\det \mathbf{R}|^{\frac{1}{n}},$$

where  $\alpha(n) = \frac{1}{4} \left(1 - \frac{(-1)^{n+1}+1}{2n^2}\right)$ .

*Proof.* In deducing the left-hand side of the inequality (3.4) we make use of the inequality (2.5) for  $A = B = \sqrt{\rho_1^*}$ ,  $a = b = \sqrt{\rho_n^*}$ ,  $a_i = b_i = \sqrt{\rho_i^*}$ ,  $i = 1, \dots, n$ . In addition, having in mind the right-hand side of the inequality (2.4), we obtain

$$\begin{aligned} RE(G) &\geq -(\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 n\alpha(n) + \frac{1}{n} \left(\sum_{i=1}^n \sqrt{\rho_i^*}\right)^2 \\ &\geq -(\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 n\alpha(n) + \frac{1}{n} \sum_{i=1}^n \rho_i^* + (n-1) \left|\prod_{i=1}^n \rho_i^*\right|^{\frac{1}{n}} \\ &= -(\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 n\alpha(n) + \frac{1}{n} RE(G) + (n-1) |\det \mathbf{R}|^{\frac{1}{n}}, \end{aligned}$$

and hence the proof follows from the last inequality.

In order to prove the right-hand side of the inequality (3.4) we will again use the inequalities (2.5) and (2.4). Letting  $A = B = \sqrt{\rho_1^*}$ ,  $a = b = \sqrt{\rho_n^*}$ ,  $a_i = b_i = \sqrt{\rho_i^*}$ ,  $i = 1, \dots, n$ , in the inequality (2.5), and then using the left-hand side of the inequality (2.4), we obtain

$$\begin{aligned} RE(G) &\leq (\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 n\alpha(n) + \frac{1}{n} \left(\sum_{i=1}^n \sqrt{\rho_i^*}\right)^2 \\ &\leq (\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 n\alpha(n) + \frac{1}{n} (n-1) \sum_{i=1}^n \rho_i^* + \left|\prod_{i=1}^n \rho_i^*\right|^{\frac{1}{n}} \\ &= (\sqrt{\rho_1^*} - \sqrt{\rho_n^*})^2 n\alpha(n) + \frac{1}{n} (n-1) RE(G) + |\det \mathbf{R}|^{\frac{1}{n}}, \end{aligned}$$

which completes the proof.  $\square$

**Acknowledgements.** The work of second author is supported by the Ministry of Education, Science and Technological Development of Republic of Serbia through the GrantNo. 174033

## REFERENCES

- [1] E. Andrade, H. Gomes and M. Robbiano, *Spectra and Randić spectra of caterpillar graphs and applications to the energy*, MATCH Commun. Math. Comput. Chem. **77** (2017), 61–75.
- [2] S. Bozkurt, A. Gungor and I. Gutman, *Randić spectral radius and Randić energy*, MATCH Commun. Math. Comput. Chem. **64** (2010), 321–334.
- [3] S. Bozkurt, A. Gungor, I. Gutman and A. Cevik, *Randić matrix and Randić energy*, MATCH Commun. Math. Comput. Chem. **64** (2010), 239–250.
- [4] F. R. K. Chung, *Spectral Graph Theory*, American Mathematical Society, Providence, 1997.
- [5] V. Cirtoaje, *The best lower bound depended on two fixed variables for Jensen's inequality with ordered variables*, J. Inequal. Appl. **2010** (2010), 1–12.
- [6] D. Cvetković, P. Rowlinson and S. K. Simić, *Introduction to the Theory of Graph Spectra*, Cambridge University Press, Cambridge, 2010.
- [7] K. C. Das and S. Sorgun, *On Randić energy of graphs*, MATCH Commun. Math. Comput. Chem. **72** (2014), 227–238.
- [8] K. C. Das, S. Sorgun and I. Gutman, *On Randić energy*, MATCH Commun. Math. Comput. Chem. **73** (2015), 81–92.
- [9] K. C. Das and S. Sun, *Extremal graphs for Randić energy*, MATCH Commun. Math. Comput. Chem. **77** (2017), 77–84.
- [10] K. C. Das, S. Sun and I. Gutman, *Normalized Laplacian eigenvalues and Randić energy of graphs*, MATCH Commun. Math. Comput. Chem **77** (2017), 45–59.
- [11] K. C. Das, S. Sorgun and K. Xu, *On Randić energy of graphs*, in: I. Gutman, X. Li (Eds.), *Graph Energies-Theory and Applications*, University of Kragujevac, Kragujevac, 2016, 111–122.
- [12] I. Gutman, B. Furtula and S. Bozkurt, *On Randić energy*, Linear Algebra Appl. **442** (2014), 50–57.
- [13] I. Gutman and X. Li, *Graph Energies-Theory and Applications*, University of Kragujevac, Kragujevac, 2016.
- [14] E. Glogić, E. Zogić and N. Glišović, *Remarks on the upper bound for the Randić energy of bipartite graphs*, Discrete Appl. Math. **221** (2017), 67–70.
- [15] J. He, Y. Liu and J. Tian, *Note on the Randić energy of graphs*, Kragujevac J. Math. **42**(2) (2018), 209–215.
- [16] X. Li and J. Wang, *Randić Energy and Randić eigenvalues*, MATCH Commun. Math. Comput. Chem. **73** (2015), 73–80.
- [17] B. Liu, Y. Huang and J. Feng, *A note on the Randić spectral radius*, MATCH Commun. Math. Comput. Chem. **68** (2012), 913–916.
- [18] D. S. Mitrinović and P. Vasić, *Analytic Inequalities*, Springer, Berlin, 1970.
- [19] B. Zhou, I. Gutman and T. Aleksić, *A note on Laplacian energy of graphs*, MATCH Commun. Math. Comput. Chem. **60** (2008), 441–446.

<sup>1</sup>DEPARTMENT OF MATHEMATICS,  
STATE UNIVERSITY OF NOVI PAZAR  
Email address: ezogic@np.ac.rs

<sup>2</sup>DEPARTMENT OF MATHEMATICS AND INFORMATICS,  
FACULTY OF SCIENCE,  
UNIVERSITY OF KRAGUJEVAC  
Email address: bojanab@kg.ac.rs