

## HERMITE-HADAMARD TYPE INEQUALITIES FOR OPERATOR GEOMETRICALLY CONVEX FUNCTIONS II

A. TAGHAVI, V. DARVISH, AND T. AZIMI ROUSHAN

ABSTRACT. In this paper, we prove some Hermite-Hadamard type inequalities for operator geometrically convex functions for non-commutative operators.

### 1. INTRODUCTION AND PRELIMINARIES

Let  $B(H)$  stand for  $C^*$ -algebra of all bounded linear operators on a complex Hilbert space  $H$  with inner product  $\langle \cdot, \cdot \rangle$ . An operator  $A \in B(H)$  is strictly positive and write  $A > 0$  if  $\langle Ax, x \rangle > 0$  for all  $x \in H$ . Let  $B(H)^{++}$  stand for all strictly positive operators on  $B(H)$ .

Let  $A$  be a self-adjoint operator in  $B(H)$ . The Gelfand map establishes a  $*$ -isometrically isomorphism  $\Phi$  between the set  $C(\text{Sp}(A))$  of all continuous functions defined on the spectrum of  $A$ , denoted  $\text{Sp}(A)$ , and the  $C^*$ -algebra  $C^*(A)$  generated by  $A$  and the identity operator  $1_H$  on  $H$  as follows.

For any  $f, g \in C(\text{Sp}(A))$  and any  $\alpha, \beta \in \mathbb{C}$  we have:

- $\Phi(\alpha f + \beta g) = \alpha\Phi(f) + \beta\Phi(g)$ ;
- $\Phi(fg) = \Phi(f)\Phi(g)$  and  $\Phi(\bar{f}) = \Phi(f)^*$ ;
- $\|\Phi(f)\| = \|f\| := \sup_{t \in \text{Sp}(A)} |f(t)|$ ;
- $\Phi(f_0) = 1_H$  and  $\Phi(f_1) = A$ , where  $f_0(t) = 1$  and  $f_1(t) = t$ , for  $t \in \text{Sp}(A)$ .

With this notation we define  $f(A) = \Phi(f)$  for all  $f \in C(\text{Sp}(A))$ , and we call it the continuous functional calculus for a self-adjoint operator  $A$ . If  $A$  is a self-adjoint operator and both  $f$  and  $g$  are real valued functions on  $\text{Sp}(A)$  then the following important property holds:  $f(t) \geq g(t)$  for any  $t \in \text{Sp}(A)$  implies that  $f(A) \geq g(A)$ ,

---

*Key words and phrases.* Operator geometrically convex function, Hermite-Hadamard inequality.  
2010 *Mathematics Subject Classification.* Primary: 47A63. Secondary: 15A60, 47B05.  
DOI 10.46793/KgJMat2101.115T

*Received:* August 19, 2018.

*Accepted:* October 09, 2018.

in the operator order of  $B(H)$ , see [12]. A real valued continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is said to be convex (concave) if

$$f(\lambda a + (1 - \lambda)b) \leq (\geq) \lambda f(a) + (1 - \lambda)f(b),$$

for  $a, b \in \mathbb{R}$  and  $\lambda \in [0, 1]$ . The following Hermite-Hadamard inequality holds for any convex function  $f$  defined on  $\mathbb{R}$

$$\begin{aligned} (b - a)f\left(\frac{a + b}{2}\right) &\leq \int_a^b f(x)dx \\ &\leq (b - a)\frac{f(a) + f(b)}{2}, \quad \text{for } a, b \in \mathbb{R}. \end{aligned}$$

The author of [8, Remark 1.9.3] gave the following refinement of Hermite-Hadamard inequalities for convex functions

$$\begin{aligned} f\left(\frac{a + b}{2}\right) &\leq \frac{1}{2} \left( f\left(\frac{3a + b}{4}\right) + f\left(\frac{a + 3b}{4}\right) \right) \\ &\leq \frac{1}{b - a} \int_a^b f(x)dx \\ &\leq \frac{1}{2} \left( f\left(\frac{a + b}{2}\right) + \frac{f(a) + f(b)}{2} \right) \\ &\leq \frac{f(a) + f(b)}{2}. \end{aligned}$$

A real valued continuous function is operator convex if

$$f(\lambda A + (1 - \lambda)B) \leq \lambda f(A) + (1 - \lambda)f(B),$$

for self-adjoint operator  $A, B \in B(H)$  and  $\lambda \in [0, 1]$ . In [2] Dragomir investigated the operator version of the Hermite-Hadamard inequality for operator convex functions. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be an operator convex function on the interval  $I$  then, for any self-adjoint operators  $A$  and  $B$  with spectra in  $I$ , the following inequalities hold

$$\begin{aligned} f\left(\frac{A + B}{2}\right) &\leq 2 \int_{\frac{1}{4}}^{\frac{3}{4}} f(tA + (1 - t)B)dt \\ &\leq \frac{1}{2} \left[ f\left(\frac{3A + B}{4}\right) + f\left(\frac{A + 3B}{4}\right) \right] \\ &\leq \int_0^1 f((1 - t)A + tB) dt \\ &\leq \frac{1}{2} \left[ f\left(\frac{A + B}{2}\right) + \frac{f(A) + f(B)}{2} \right] \\ &\leq \frac{f(A) + f(B)}{2}. \end{aligned}$$

For the first inequality in above, see [10].

A continuous function  $f : I \subseteq \mathbb{R}^+ \rightarrow \mathbb{R}^+$  ( $\mathbb{R}^+$  denoted positive real numbers) is said to be geometrically convex function (or multiplicatively convex function) if

$$f(a^\lambda b^{1-\lambda}) \leq f(a)^\lambda f(b)^{1-\lambda},$$

for  $a, b \in I$  and  $\lambda \in [0, 1]$ .

The author of [7, p. 158] showed that every polynomial  $P(x)$  with non-negative coefficients is a geometrically convex function on  $[0, \infty)$ . More generally, every real analytic function  $f(x) = \sum_{n=0}^\infty c_n x^n$  with non-negative coefficients is geometrically convex function on  $(0, R)$  where  $R$  denotes the radius of convergence. Also, see [9, 11]. In [10], the following inequalities were obtained for a geometrically convex function

$$\begin{aligned} f(\sqrt{ab}) &\leq \sqrt{\left(f(a^{\frac{3}{4}}b^{\frac{1}{4}})f(a^{\frac{1}{4}}b^{\frac{3}{4}})\right)} \\ &\leq \exp\left(\frac{1}{\log b - \log a} \int_a^b \frac{\log f(t)}{t} dt\right) \\ &\leq \sqrt{f(\sqrt{ab})} \cdot \sqrt[4]{f(a)} \cdot \sqrt[4]{f(b)} \\ &\leq \sqrt{f(a)f(b)}. \end{aligned}$$

In this paper, we prove some Hermite-Hadamard inequalities for operator geometrically convex functions. Moreover, in the final section, we present some examples and remarks.

## 2. HERMITE-HADAMARD INEQUALITIES FOR GEOMETRICALLY CONVEX FUNCTIONS

In this section, we introduce the concept of operator geometrically convex function for positive operators and prove the Hermite-Hadamard type inequalities for this function.

**Proposition 2.1.** *Let  $A, B \in B(H)^{++}$  such that  $\text{Sp}(A), \text{Sp}(B) \subseteq I$ , and  $t \in [0, 1]$ . Then  $\text{Sp}(A\sharp_t B) \subseteq I$ , where  $A\sharp_t B = A^{\frac{1}{2}}(A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^t A^{\frac{1}{2}}$  is  $t$ -geometric mean.*

*Proof.* Let  $I = [m, M]$  for some positive real numbers  $m, M$  with  $m < M$ . Since  $\text{Sp}(A), \text{Sp}(B) \subseteq I$  it is equivalent to  $m1_H \leq A \leq M1_H$  and  $m1_H \leq B \leq M1_H$ . So, by virtue of the fact that if  $a, b$  be self-adjoint operators in  $C^*$ -algebra  $\mathcal{A}$  which  $a \leq b$  and  $c \in \mathcal{A}$ , then  $c^*ac \leq c^*bc$ , and also by using the operator monotonicity property of the function  $f(x) = x^t$  on  $(0, \infty)$  for  $t \in [0, 1]$ , we get the result. □

Now, by applying Proposition 2.1, we present the following definition.

**Definition 2.1.** A continuous function  $f : I \subseteq \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is said to be operator geometrically convex if

$$f(A\sharp_t B) \leq f(A)\sharp_t f(B),$$

for  $A, B \in B(H)^{++}$  such that  $\text{Sp}(A), \text{Sp}(B) \subseteq I$  and  $t \in [0, 1]$ .

We need the following lemmas for proving our theorems.

**Lemma 2.1** ([4, 5]). *Let  $A, B \in B(H)^{++}$  and let  $t, s, u \in \mathbb{R}$ . Then*

$$(A\sharp_t B)\sharp_s(A\sharp_u B) = A\sharp_{(1-s)t+su} B.$$

**Lemma 2.2** ([4]). *Let  $A, B, C$  and  $D$  be operators in  $B(H)^{++}$  and let  $t \in \mathbb{R}$ . Then, we have*

$$A\sharp_t B \leq C\sharp_t D,$$

for  $A \leq C$  and  $B \leq D$ .

**Lemma 2.3.** *Let  $A, B \in B(H)^{++}$ . If  $f : I \subseteq \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a continuous function, then*

$$\int_0^1 f(A\sharp_t B)\sharp f(A\sharp_{1-t} B) dt \leq \left( \int_0^1 f(A\sharp_t B) dt \right)\sharp \left( \int_0^1 f(A\sharp_{1-t} B) dt \right)$$

such that  $\text{Sp}(A), \text{Sp}(B) \subseteq I$ .

*Proof.* Since the function  $t^{\frac{1}{2}}$  is operator concave, we can write

$$\begin{aligned} & \left( \left( \int_0^1 f(A\sharp_{1-u} B) du \right)^{\frac{-1}{2}} \left( \int_0^1 f(A\sharp_u B) du \right) \left( \int_0^1 f(A\sharp_{1-u} B) du \right)^{\frac{-1}{2}} \right)^{\frac{1}{2}} \\ & \quad (\text{by change of variable } v = 1 - u) \\ & = \left( \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} \left( \int_0^1 f(A\sharp_u B) du \right) \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} \right)^{\frac{1}{2}} \\ & = \left( \int_0^1 \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{1}{2}} f(A\sharp_u B) \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{1}{2}} du \right)^{\frac{1}{2}} \\ & = \left( \int_0^1 \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} (f(A\sharp_{1-u} B))^{\frac{1}{2}} \left( (f(A\sharp_{1-u} B))^{\frac{-1}{2}} f(A\sharp_u B) (f(A\sharp_{1-u} B))^{\frac{-1}{2}} \right) \right. \\ & \quad \left. \times (f(A\sharp_{1-u} B))^{\frac{1}{2}} \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} du \right)^{\frac{1}{2}} \\ & \quad (\text{by the operator Jensen inequality}) \\ & \geq \int_0^1 \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} (f(A\sharp_{1-u} B))^{\frac{1}{2}} \left( (f(A\sharp_{1-u} B))^{\frac{-1}{2}} f(A\sharp_u B) (f(A\sharp_{1-u} B))^{\frac{-1}{2}} \right)^{\frac{1}{2}} \\ & \quad \times (f(A\sharp_{1-u} B))^{\frac{1}{2}} \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} du \\ & = \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} \int_0^1 (f(A\sharp_{1-u} B))^{\frac{1}{2}} \left( (f(A\sharp_{1-u} B))^{\frac{-1}{2}} f(A\sharp_u B) (f(A\sharp_{1-u} B))^{\frac{-1}{2}} \right)^{\frac{1}{2}} \\ & \quad \times (f(A\sharp_{1-u} B))^{\frac{1}{2}} du \left( \int_0^1 f(A\sharp_v B) dv \right)^{\frac{-1}{2}} \end{aligned}$$

(by change of variable  $u = 1 - v$ )

$$= \left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{-1}{2}} \int_0^1 (f(A\sharp_{1-u}B))^{\frac{1}{2}} \left( (f(A\sharp_{1-u}B))^{\frac{-1}{2}} f(A\sharp_u B) (f(A\sharp_{1-u}B))^{\frac{-1}{2}} \right)^{\frac{1}{2}} \times (f(A\sharp_{1-u}B))^{\frac{1}{2}} du \left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{-1}{2}} .$$

So, we obtain

$$\left( \left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{-1}{2}} \left( \int_0^1 f(A\sharp_u B) du \right) \left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{-1}{2}} \right)^{\frac{1}{2}} \geq \left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{-1}{2}} \int_0^1 (f(A\sharp_{1-u}B))^{\frac{-1}{2}} \left( (f(A\sharp_{1-u}B))^{\frac{-1}{2}} f(A\sharp_u B) (f(A\sharp_{1-u}B))^{\frac{-1}{2}} \right)^{\frac{1}{2}} \times (f(A\sharp_{1-u}B))^{\frac{1}{2}} du \left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{-1}{2}} .$$

Multiplying both side of the above inequality by  $\left( \int_0^1 f(A\sharp_{1-u}B) du \right)^{\frac{1}{2}}$  we obtain

$$\left( \int_0^1 f(A\sharp_u B) du \right) \sharp \left( \int_0^1 f(A\sharp_{1-u}B) du \right) \geq \int_0^1 f(A\sharp_u B) \sharp f(A\sharp_{1-u}B) du. \quad \square$$

Before giving our theorems in this section, we mention the following remark.

*Remark 2.1.* Let  $p(x) = x^t$  and  $q(x) = x^s$  on  $[1, \infty)$ , where  $0 \leq t \leq s$ . If  $f(A) \leq f(B)$  then  $\text{Sp} \left( f(A)^{\frac{-1}{2}} (f(B)) f(A)^{\frac{-1}{2}} \right) \subseteq [1, \infty)$ . By functional calculus, we have

$$p \left( f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}} \right) \leq q \left( f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}} \right).$$

So,

$$\left( f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}} \right)^t \leq \left( f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}} \right)^s .$$

Now, we are ready to prove Hermite-Hadamard type inequality for operator geometrically convex functions.

**Theorem 2.1.** *Let  $f$  be an operator geometrically convex function. Then, we have*

$$(2.1) \quad f(A\sharp B) \leq \int_0^1 f(A\sharp_t B) dt \leq \int_0^1 f(A)\sharp_t f(B) dt.$$

Moreover, if  $f(A) \leq f(B)$ , then we have

$$(2.2) \quad \int_0^1 f(A\sharp_t B) dt \leq \int_0^1 f(A)\sharp_t f(B) dt \leq \frac{1}{2}((f(A)\sharp f(B)) + f(B)),$$

for  $A, B \in B(H)^{++}$ .

*Proof.* Let  $f$  be a geometrically convex function. Then we have

$$\begin{aligned} f(A\sharp B) &= f((A\sharp_t B)\sharp(A\sharp_{1-t} B)) \quad (\text{by Lemma 2.1}) \\ &\leq f(A\sharp_t B)\sharp f(A\sharp_{1-t} B) \quad (f \text{ is operator geometrically convex}). \end{aligned}$$

Taking integral of the both sides of the above inequalities on  $[0, 1]$ , we obtain

$$\begin{aligned} f(A\sharp B) &\leq \int_0^1 f(A\sharp_t B)\sharp f(A\sharp_{1-t} B) dt \\ &\leq \left(\int_0^1 f(A\sharp_t B) dt\right)\sharp \left(\int_0^1 f(A\sharp_{1-t} B) dt\right) \quad (\text{by Lemma 2.3}) \\ &= \int_0^1 f(A\sharp_t B) dt \\ &\leq \int_0^1 f(A)\sharp_t f(B) dt. \end{aligned}$$

For the case  $f(A) \leq f(B)$ , by applying Remark 2.1 for  $s = \frac{1}{2}$ , we have

$$\left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^t \leq \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^{\frac{1}{2}}.$$

By integrating the above inequality over  $t \in [0, \frac{1}{2}]$ , we obtain

$$\int_0^{\frac{1}{2}} \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^t dt \leq \frac{1}{2} \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^{\frac{1}{2}}.$$

Multiplying both sides of the above inequality by  $f(A)^{\frac{1}{2}}$ , we have

$$\begin{aligned} &\int_0^{\frac{1}{2}} f(A)^{\frac{1}{2}} \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^t f(A)^{\frac{1}{2}} dt \\ &\leq \frac{1}{2} \left(f(A)^{\frac{1}{2}} \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^{\frac{1}{2}} f(A)^{\frac{1}{2}}\right). \end{aligned}$$

It follows that

$$(2.3) \quad \int_0^{\frac{1}{2}} f(A)\sharp_t f(B) \leq \frac{f(A)\sharp f(B)}{2}.$$

On the other hand, by considering Remark 2.1 for  $s = 1$ , we have

$$\left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^t \leq f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}.$$

Integrating the above inequality over  $t \in [\frac{1}{2}, 1]$ , we get

$$\int_{\frac{1}{2}}^1 \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^t dt \leq \frac{1}{2} \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right).$$

By multiplying both side of the above inequality by  $f(A)^{\frac{1}{2}}$ , we have

$$\int_{\frac{1}{2}}^1 f(A)^{\frac{1}{2}} \left(f(A)^{-\frac{1}{2}}f(B)f(A)^{-\frac{1}{2}}\right)^t f(A)^{\frac{1}{2}} dt \leq \frac{f(B)}{2}.$$

It follows that

$$(2.4) \quad \int_{\frac{1}{2}}^1 f(A)\sharp_t f(B) \leq \frac{f(B)}{2}.$$

From inequalities (2.3) and (2.4) we obtain

$$\begin{aligned} \int_0^{\frac{1}{2}} f(A)\sharp_t B dt + \int_{\frac{1}{2}}^1 f(A)\sharp_t B dt &\leq \int_0^{\frac{1}{2}} f(A)\sharp_t f(B) dt + \int_{\frac{1}{2}}^1 f(A)\sharp_t f(B) dt \\ &\leq \frac{f(A)\sharp f(B)}{2} + \frac{f(B)}{2}. \end{aligned}$$

It follows that

$$\int_0^1 f(A)\sharp_t B dt \leq \int_0^1 f(A)\sharp_t f(B) dt \leq \frac{1}{2}((f(A)\sharp f(B)) + f(B)). \quad \square$$

By making use of inequalities (2.1) and (2.2), we have the following result.

**Corollary 2.1.** *Let  $f$  be an operator geometrically convex function. Then, if  $f(A) \leq f(B)$  we have*

$$f(A)\sharp B \leq \int_0^1 f(A)\sharp_t B dt \leq \frac{1}{2}((f(A)\sharp f(B)) + f(B)),$$

for  $A, B \in B(H)^{++}$ .

**Theorem 2.2.** *Let  $f$  be an operator geometrically convex function. Then, we have*

$$f(A)\sharp B \leq \int_0^1 f(A)\sharp_t B \sharp f(A)\sharp_{1-t} B dt \leq f(A)\sharp f(B),$$

for  $A, B \in B(H)^{++}$ .

*Proof.* We can write

$$\begin{aligned} f(A)\sharp B &= f((A)\sharp_t B)\sharp(A)\sharp_{1-t} B) \quad (\text{by Lemma 2.1}) \\ &\leq f(A)\sharp_t B \sharp f(A)\sharp_{1-t} B \quad (f \text{ is operator geometrically convex}) \\ &\leq (f(A)\sharp_t f(B)) \sharp (f(A)\sharp_{1-t} f(B)) \quad (\text{by Lemma 2.2}) \\ &= f(A)\sharp f(B). \end{aligned}$$

So, we obtain

$$f(A)\sharp B \leq f(A)\sharp_t B \sharp f(A)\sharp_{1-t} B \leq f(A)\sharp f(B).$$

Integrating the above inequality over  $t \in [0, 1]$  we obtain the desired result.  $\square$

We divide the interval  $[0, 1]$  to the interval  $[\nu, 1 - \nu]$  when  $\nu \in [0, \frac{1}{2})$  and to the interval  $[1 - \nu, \nu]$  when  $\nu \in (\frac{1}{2}, 1]$ . The we have the following inequalities.

**Theorem 2.3.** *Let  $A, B \in B(H)^{++}$  such that  $f(A) \leq f(B)$ . Then, we have*

(a) for  $\nu \in [0, \frac{1}{2})$

$$(2.5) \quad f(A)\sharp_\nu f(B) \leq \frac{1}{1-2\nu} \int_\nu^{1-\nu} f(A)\sharp_t f(B) dt \leq f(A)\sharp_{1-\nu} f(B);$$

(b) for  $\nu \in (\frac{1}{2}, 1]$

$$(2.6) \quad f(A)\sharp_{1-\nu}f(B) \leq \frac{1}{2\nu-1} \int_{1-\nu}^{\nu} f(A)\sharp_t f(B) dt \leq f(A)\sharp_{\nu}f(B).$$

*Proof.* Let  $\nu \in [0, \frac{1}{2})$ , then by Remark 2.1 we have

$$\begin{aligned} \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^{\nu} &\leq \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^t \\ &\leq \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^{1-\nu}, \end{aligned}$$

for  $\nu \leq t \leq 1-\nu$  and  $A, B \in B(H)^{++}$  such that  $\text{Sp}(A), \text{Sp}(B) \subseteq I$ .  
By integrating the above inequality over  $t \in [\nu, 1-\nu]$  we obtain

$$\begin{aligned} \int_{\nu}^{1-\nu} \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^{\nu} dt &\leq \int_{\nu}^{1-\nu} \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^t dt \\ &\leq \int_{\nu}^{1-\nu} \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^{1-\nu} dt. \end{aligned}$$

It follows that

$$\begin{aligned} \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^{\nu} &\leq \frac{1}{1-2\nu} \int_{\nu}^{1-\nu} \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^t dt \\ &\leq \left(f(A)^{\frac{-1}{2}} f(B) f(A)^{\frac{-1}{2}}\right)^{1-\nu}. \end{aligned}$$

Multiplying the both sides of the above inequality by  $f(A)^{\frac{1}{2}}$  gives us

$$f(A)\sharp_{\nu}f(B) \leq \frac{1}{1-2\nu} \int_{\nu}^{1-\nu} f(A)\sharp_t f(B) dt \leq f(A)\sharp_{1-\nu}f(B).$$

Also, we know that

$$\begin{aligned} \lim_{\nu \rightarrow \frac{1}{2}} f(A)\sharp_{\nu}f(B) &= \lim_{\nu \rightarrow \frac{1}{2}} \frac{1}{1-2\nu} \int_{\nu}^{1-\nu} f(A)\sharp_t f(B) dt \\ &= \lim_{\nu \rightarrow \frac{1}{2}} f(A)\sharp_{1-\nu}f(B) \\ &= f(A)\sharp f(B). \end{aligned}$$

Similarly, for  $\nu \in (\frac{1}{2}, 1]$ , by the same proof as above, we get

$$f(A)\sharp_{1-\nu}f(B) \leq \frac{1}{2\nu-1} \int_{1-\nu}^{\nu} f(A)\sharp_t f(B) dt \leq f(A)\sharp_{\nu}f(B). \quad \square$$

By definition of geometrically convex function and (2.5) we have

$$\begin{aligned} f(A\sharp_{\nu}B) &\leq \frac{1}{1-2\nu} \int_{\nu}^{1-\nu} f(A\sharp_t B) dt \\ &\leq \frac{1}{1-2\nu} \int_{\nu}^{1-\nu} f(A)\sharp_t f(B) dt \\ &\leq f(A)\sharp_{1-\nu}f(B), \end{aligned}$$



for  $\nu \in [0, \frac{1}{2})$ . We should mention here that

$$\lim_{\nu \rightarrow \frac{1}{2}} \frac{1}{1 - 2\nu} \int_{\nu}^{1-\nu} f(A\sharp_t B) dt = \lim_{\nu \rightarrow \frac{1}{2}} f(A\sharp_{\nu} B) = f(A\sharp B).$$

On the other hand, by the definition of geometrically convex function and (2.6) we have

$$\begin{aligned} f(A\sharp_{1-\nu} B) &\leq \frac{1}{2\nu - 1} \int_{1-\nu}^{\nu} f(A\sharp_t B) dt \\ &\leq \frac{1}{2\nu - 1} \int_{1-\nu}^{\nu} f(A)\sharp_t f(B) dt \\ &\leq f(A)\sharp_{\nu} f(B), \end{aligned}$$

for  $\nu \in (\frac{1}{2}, 1]$ .

### 3. EXAMPLES AND REMARKS

In this section we give some examples of the results that obtained in the previous section.

*Remark 3.1.* For positive  $A, B \in B(H)$ , Ando proved in [1] that if  $\Psi$  is a positive linear map, then we have

$$\Psi(A\sharp B) \leq \Psi(A)\sharp\Psi(B).$$

The above inequality shows that we can find some examples for Definition 2.1 when  $f$  is linear.

*Example 3.1.* It is easy to check that the function  $f(t) = t^{-1}$  is operator geometrically convex for operators in  $B(H)^{++}$ .

**Definition 3.1.** Let  $\phi$  be a map on  $C^*$ -algebra  $B(H)$ . We say that  $\phi$  is 2-positive if the  $2 \times 2$  operator matrix  $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \geq 0$ , then we have  $\begin{bmatrix} \phi(A) & \phi(B) \\ \phi(B^*) & \phi(C) \end{bmatrix} \geq 0$ .

In [6], M. Lin gave an example of a 2-positive map over contraction operators (i.e.,  $\|A\| < 1$ ). He proved that

$$(3.1) \quad \phi(t) = (1 - t)^{-1}$$

is 2-positive.

*Example 3.2.* Let  $A$  and  $B$  be two contraction operators in  $B(H)^{++}$ . Then it is easy to check  $A\sharp B$  is also a contraction and positive. Also, we know the  $2 \times 2$  operator matrix

$$\begin{bmatrix} A & A\sharp B \\ A\sharp B & B \end{bmatrix}$$

is semidefinite positive. Hence, by (3.1) we obtain

$$\begin{bmatrix} (I - A)^{-1} & (I - (A\sharp B))^{-1} \\ (I - (A\sharp B))^{-1} & (I - B)^{-1} \end{bmatrix}$$

is semidefinite positive.

On the other hand, by Ando's characterization of the geometric mean if  $X$  is a Hermitian matrix and

$$\begin{bmatrix} A & X \\ X & B \end{bmatrix} \geq 0,$$

then  $X \leq A\sharp B$ . So we conclude that  $(I - (A\sharp B))^{-1} \leq (I - A)^{-1}\sharp(I - B)^{-1}$ . Therefore, the function  $\phi(t) = (1 - t)^{-1}$  is operator geometrically convex.

Also, Lin proved that the function

$$\phi(t) = \frac{1+t}{1-t}$$

is 2-positive over contractions. By the same argument as Example 3.2 we can say the above function is operator geometrically convex too.

*Example 3.3.* In the proof of [3, Theorem 4.12], by applying Hölder-McCarthy inequality the authors showed the following inequalities

$$\begin{aligned} \langle A\sharp_{\alpha} Bx, x \rangle &= \left\langle \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right)^{\alpha} A^{\frac{1}{2}} x, A^{\frac{1}{2}} x \right\rangle \\ &\leq \left\langle \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right) A^{\frac{1}{2}} x, A^{\frac{1}{2}} x \right\rangle^{\alpha} \left\langle A^{\frac{1}{2}} x, A^{\frac{1}{2}} x \right\rangle^{1-\alpha} \\ &= \langle Ax, x \rangle^{1-\alpha} \langle Bx, x \rangle^{\alpha} \\ &= \langle Ax, x \rangle \sharp_{\alpha} \langle Bx, x \rangle, \end{aligned}$$

for  $x \in H$  and  $\alpha \in [0, 1]$ . By taking the supremum over unit vector  $x$ , we obtain that  $f(x) = \|x\|$  is geometrically convex function for usual operator norms.

By the above example and Corollary 2.1, when  $\|A\| \leq \|B\|$  we have

$$\|A\sharp B\| \leq \int_0^1 \|A\sharp_t B\| dt \leq \frac{1}{2}(\sqrt{\|A\|\|B\|} + \|B\|),$$

for  $A, B \in B(H)^{++}$ .

## REFERENCES

- [1] T. Ando, *Concavity of certain maps on positive definite matrices and applications to Hadamard products*, Linear Algebra Appl. **26** (1979), 203–241.
- [2] S. S. Dragomir, *Hermite-Hadamard's type inequalities for operator convex functions*, Appl. Math. Comput. **218** (2011), 766–772.
- [3] C. Jung, H. Lee, Y. Lim and T. Yamazaki, *Weighted geometric mean of  $n$ -operators with  $n$ -parameters*, Linear Algebra Appl. **432** (2010), 1515–1530.
- [4] F. Kubo and T. Ando, *Means of positive linear operators*, Math. Ann. **246**(3) (1980), 205–224.
- [5] J. Lawson and Y. Lim, *Weighted means and Karcher equations of positive operators*, Proc. Natl. Acad. Sci. USA **110**(39) (2013), 15626–15632.
- [6] M. Lin, *A Lewent type determinantal inequality*, Taiwanese J. Math. **17** (2013), 1303–1309.
- [7] C. P. Niculescu, *Convexity according to the geometric mean*, Math. Inequal. Appl. **3**(2) (2000), 155–167.

- [8] C. P. Niculescu and L. E. Persson, *Convex Functions and Their Applications: A Contemporary Approach*, Springer, Berlin, Heidelberg, New York, Hong Kong, London, Milan, Paris, Tokyo, 2004.
- [9] A. Taghavi, V. Darvish, H. M. Nazari and S. S. Dragomir, *Hermite-Hadamard type inequalities for operator geometrically convex functions*, Monatsh. Math. **181** (2016), 187–203.
- [10] V. Darvish, S. S. Dragomir, H. M. Nazari, and A. Taghavi, *Some inequalities associated with the Hermite-Hadamard inequalities for operator  $h$ -convex functions*, Acta Comment. Univ. Tartu. Math. **21**(2) (2017), 287-297.
- [11] A. Taghavi, T. A. Roushan and V. Darvish, *Some refinements for the arithmetic-geometric mean and Cauchy-Schwarz matrix norm interpolating inequalities*, Bulletin of the Iranian Mathematical Society **44** (2018), 927–936.
- [12] K. Zhu, *An Introduction to Operator Algebras*, CRC Press, Boca Raton, Florida, 1993.

DEPARTMENT OF MATHEMATICS,  
FACULTY OF SCIENCE,  
UNIVERSITY OF MAZANDARAN, P. O. BOX 47416-1468, BABOLSAR, IRAN  
*Email address:* taghavi@umz.ac.ir  
*Email address:* vahid.darvish@mail.com  
*Email address:* t.roushan@umz.ac.ir