

## TRIANGULAR SYSTEM OF HIGHER ORDER SINGULAR FRACTIONAL DIFFERENTIAL EQUATIONS

AMELE TAIEB<sup>1</sup> AND ZOUBIR DAHMANI<sup>2</sup>

ABSTRACT. In this paper, we introduce a high dimensional system of singular fractional differential equations. Using Schauder fixed point theorem, we prove an existence result. We also investigate the uniqueness of solution using the Banach contraction principle. Moreover, we study the Ulam-Hyers stability and the generalized-Ulam-Hyers stability of solutions. Some illustrative examples are also presented.

### 1. INTRODUCTION AND PRELIMINARIES

Recently, the fractional calculus has attracted the attention of researchers in various fields of applied sciences. For details, see [12, 16, 19, 20] and the references therein. It is important to note that some research studies deal with the existence and uniqueness of solutions for some fractional differential equations are obtained in [1, 6–9]. Other studies in [2, 3, 5, 17, 23] have been done for the singular fractional differential equations. On the other hand, the Ulam stability of fractional differential equations is quite significant in more realistic problems, numerical analysis, biology and economics. Considerable work has been done in this area, for instance, see [10, 11, 13–15, 18, 22, 24].

Let us now present some important research papers that inspired our work: We begin by [4], where C. Bai and J. Fang established the existence of solutions for the following singular fractional coupled system:

$$\begin{cases} D^\delta u(t) = f(t, v(t)), & 0 < t < 1, \\ D^\rho v(t) = g(t, u(t)), & 0 < t < 1, \end{cases}$$

---

*Key words and phrases.* Caputo derivative, fixed point, singular fractional differential equation, existence, uniqueness, Ulam-Hyers stability.

2010 *Mathematics Subject Classification.* Primary: 30C45. Secondary: 39B72, 39B82.

DOI 10.46793/KgJMat2101.081T

*Received:* March 11, 2018.

*Accepted:* October 08, 2018.

where  $0 < \delta, \rho < 1$ ,  $D^\delta, D^\rho$  are two standard Riemann-Liouville fractional derivatives,  $f, g : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$  are two given continuous functions,  $\lim_{t \rightarrow 0^+} f(t) = \infty$  and  $\lim_{t \rightarrow 0^+} g(t) = \infty$ .

In [25], A. Yang and W. Ge considered the following fractional coupled system

$$\begin{cases} D^{\alpha_1} u_1(t) + f_1(t, u_2(t), D^{\mu_1} u_2(t)) = 0, \\ \vdots \\ D^{\alpha_{n-1}} u_{n-1}(t) + f_{n-1}(t, u_n(t), D^{\mu_{n-1}} u_n(t)) = 0, \\ D^{\alpha_n} u_n(t) + f_n(t, u_1(t), D^{\mu_n} u_1(t)) = 0, \end{cases}$$

associated with the boundary conditions

$$\begin{cases} u_1(0) = u_2(0) = \dots = u_n(0) = 0, \\ u_1(1) = u_2(1) = \dots = u_n(1) = 0, \end{cases}$$

for  $1 < \alpha_j < 2$ ,  $\mu_j > 0$ ,  $\alpha_j - \mu_{j-1} > 1$ ,  $j = 1, 2, \dots, n$ ,  $\mu_0 = \mu_n$  and  $f_j : [0, 1] \times [0, \infty) \times \mathbb{R} \rightarrow [0, \infty)$  is continuous function. Some existence and multiplicity results of solutions are obtained.

In [21], A. Taïeb and Z. Dahmani established new existence and uniqueness results for the following problem:

$$\begin{cases} D^\alpha u(t) + \sum_{i=1}^m f_i(t, u(t), v(t), D^\gamma u(t), D^\rho v(t)) = 0, & t \in J, \\ D^\beta v(t) + \sum_{i=1}^m g_i(t, u(t), v(t), D^\gamma u(t), D^\rho v(t)) = 0, & t \in J, \\ u(0) = u_0^*, v(0) = v_0^*, \\ u'(0) = u''(0) = v'(0) = v''(0) = 0, \\ u'''(0) = J^r u(\tau), v'''(0) = J^\varphi v(\varsigma), \quad r > 0, \varphi > 0, \end{cases}$$

where  $\alpha, \beta \in (3, 4)$ ,  $\gamma, \rho \in (0, 3)$ ,  $\tau, \varsigma \in (0, 1)$ ,  $D^\alpha, D^\rho, D^\beta$  and  $D^\gamma$  denote the Caputo fractional derivatives and  $J^r, J^\varphi$  denote the Riemann-Liouville fractional integrals,  $J := [0, 1]$ ,  $u_0^*, v_0^* \in \mathbb{R}$ . For each  $i = 1, \dots, m$ ,  $f_i$  and  $g_i : J \times \mathbb{R}^4 \rightarrow \mathbb{R}$  are specific functions.

In this paper, we discuss the existence, uniqueness and Ulam stability of solutions for the following singular fractional coupled system:

$$(1.1) \quad \begin{cases} D^{\alpha_1} x_1(t) = f_1(t, x_1(t)), \\ D^{\alpha_2} x_2(t) = f_2(t, x_1(t), x_2(t)), \\ \vdots \\ D^{\alpha_n} x_n(t) = f_n(t, x_1(t), x_2(t), \dots, x_n(t)), \\ 0 < t \leq 1, \quad k-1 < \alpha_k < k, \quad k = 1, 2, \dots, n, \\ x_1(0) = a_0^1, \quad k = 1, \\ x_k^{(j)}(0) = a_j^k, \quad j = 0, 1, \dots, k-2, \quad k = 2, 3, \dots, n, \\ D^{\delta_{k-1}} x_k(1) = 0, \quad k-2 < \delta_{k-1} < k-1, \quad k = 2, 3, \dots, n, \end{cases}$$

where  $n \in \mathbb{N} - \{0, 1\}$ . For all  $k = 1, 2, \dots, n$ , the functions  $f_k : (0, 1] \times \mathbb{R}^k \rightarrow \mathbb{R}$  are continuous, singular at  $t = 0$ ,  $\lim_{t \rightarrow 0^+} f_k(t) = \infty$  and there exist  $\beta_k \in (0, 1)$ ,  $k = 1, 2, \dots, n$ , such that  $t^{\beta_k} f_k$ ,  $k = 1, 2, \dots, n$ , are continuous on  $[0, 1]$ .

To the best of our knowledge, there are no papers that have considered this kind of singular fractional coupled system.

We present some basic definitions and lemmas that we need to prove our main results. It can be found in [16].

**Definition 1.1.** The Riemann-Liouville fractional integral operator of order  $\alpha \geq 0$  for a continuous function  $f$  on  $[0, \infty)$  is defined as:

$$J^\alpha f(t) = \begin{cases} \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s) ds, & \alpha > 0, \\ f(t), & \alpha = 0, \end{cases}$$

where  $t \geq 0$  and  $\Gamma(\alpha) := \int_0^{+\infty} e^{-u} u^{\alpha-1} du$ .

**Definition 1.2.** The Caputo derivative of order  $\alpha$  for a function  $x : [0, +\infty) \rightarrow \mathbb{R}$ , which is at least  $k$ -times differentiable can be defined as the following:

$$D^\alpha x(t) = \frac{1}{\Gamma(k-\alpha)} \int_0^t (t-s)^{k-\alpha-1} x^{(k)}(s) ds = J^{k-\alpha} x^{(k)}(t),$$

for  $k-1 < \alpha < k$ ,  $k \in \mathbb{N} - \{0\}$ .

**Lemma 1.1.** Let  $\alpha, \beta > 0$ , and  $k-1 < \alpha < k$ ,  $k \in \mathbb{N} - \{0\}$ , and let  $j$  be a positive integer. Then

$$D^\alpha t^{\beta-1} = \frac{\Gamma(\beta)}{\Gamma(\beta-\alpha)} t^{\beta-\alpha-1}, \quad \beta > k,$$

and

$$D^\alpha t^j = 0, \quad j = 0, 1, \dots, k-1.$$

**Lemma 1.2.** Let  $q > p > 0$  and  $f \in L^1([a, b])$ . Then for all  $t \in [a, b]$ , we have

$$D^p J^q f(t) = J^{q-p} f(t), \quad t \in [a, b].$$

**Lemma 1.3.** Let  $k-1 < \alpha < k$ ,  $k \in \mathbb{N} - \{0\}$ , and let  $j$  be a positive integer. Then, the general solution of the fractional differential equation  $D^\alpha x(t) = 0$ , is given by:

$$x(t) = \sum_{j=0}^{k-1} c_j t^j, \quad (c_j)_{j=0,1,\dots,k-1} \in \mathbb{R}.$$

**Lemma 1.4.** Let  $k \in \mathbb{N} - \{0\}$ ,  $k-1 < \alpha < k$ , and let  $j$  be a positive integer. Then,

$$J^\alpha D^\alpha x(t) = x(t) + \sum_{j=0}^{k-1} c_j t^j, \quad (c_j)_{j=0,1,\dots,k-1} \in \mathbb{R}.$$

**Lemma 1.5** (Schauder fixed point theorem). Let  $(E, d)$  be a complete metric space, let  $U$  be a closed convex subset of  $E$ , and let  $T : E \rightarrow E$  be a mapping such that the set  $V := \{Tx : x \in U\}$  is relatively compact in  $E$ . Then  $T$  has at least one fixed point.

We also prove the following auxiliary result to give the integral representation of (1.1).

**Lemma 1.6.** *Assume that  $k - 1 < \alpha_k < k$ ,  $k = 1, 2, \dots, n$ ,  $n \in \mathbb{N} - \{0, 1\}$  and  $F_k \in C([0, 1], \mathbb{R})$ . Then, the following system*

$$\begin{cases} D^{\alpha_1} x_1(t) = F_1(t), \\ D^{\alpha_2} x_2(t) = F_2(t), \\ \vdots \\ D^{\alpha_n} x_n(t) = F_n(t), \end{cases}$$

associated with the conditions:

$$(1.2) \quad \begin{cases} x_1(0) = a_0^1, \\ x_k^{(j)}(0) = a_j^k, \quad k = 2, 3, \dots, n, j = 0, 1, \dots, k - 2, \\ D^{\delta_{k-1}} x_k(1) = 0, \quad k - 2 < \delta_{k-1} < k - 1, \end{cases}$$

has a unique solution  $(x_1, x_2, \dots, x_n)$ , where

$$(1.3) \quad x_k(t) = \begin{cases} \int_0^t \frac{(t-s)^{\alpha_1-1}}{\Gamma(\alpha_1)} F_1(s) ds + a_0^1, & k = 1, \\ \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} F_k(s) ds + \sum_{j=0}^{k-2} \frac{a_j^k}{j!} t^j \\ - \frac{\Gamma(k-\delta_{k-1})}{(k-1)!} t^{k-1} \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1}}{\Gamma(\alpha_k-\delta_{k-1})} F_k(s) ds, & k = 2, 3, \dots, n. \end{cases}$$

*Proof.* Using Lemma 1.4, we obtain the following integral equation:

$$(1.4) \quad x_k(t) = \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} F_k(s) ds - \sum_{j=0}^{k-1} c_j^k t^j, \quad k = 1, 2, \dots, n,$$

where

$$\begin{pmatrix} c_0^1 & 0 & \dots & \dots & \dots & 0 \\ c_0^2 & c_1^2 & 0 & \dots & \dots & 0 \\ c_0^3 & c_1^3 & c_2^3 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & 0 & 0 \\ c_0^{n-1} & c_1^{n-1} & c_2^{n-1} & \dots & c_{n-2}^{n-1} & 0 \\ c_0^n & c_1^n & c_2^n & \dots & c_{n-2}^n & c_{n-1}^n \end{pmatrix} \in M_n(\mathbb{R}).$$

Applying the conditions given in (1.2), we observe that

$$x_1(0) = -c_0^1 = a_0^1,$$

and for all  $k = 2, 3, \dots, n$ , we get

$$\begin{cases} x_k^{(j)}(0) = -j! c_j^k = a_j^k, \quad j = 0, 1, \dots, k - 2, \\ D^{\delta_{k-1}} x_k(1) = \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1}}{\Gamma(\alpha_k-\delta_{k-1})} F_k(s) ds - \frac{\Gamma(k)}{\Gamma(k-\delta_{k-1})} c_{k-1}^k = 0, \\ k - 2 < \delta_{k-1} < k - 1, \end{cases}$$

which implies that

$$(1.5) \quad c_0^1 = -a_0^1,$$

and

$$(1.6) \quad c_j^k = \begin{cases} -\frac{a_j^k}{j!}, & j = 0, 1, \dots, k-2, \\ \frac{\Gamma(k - \delta_{k-1})}{\Gamma(k)} \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1}}{\Gamma(\alpha_k - \delta_{k-1})} F_k(s) ds, & j = k-1, \end{cases}$$

where  $k = 2, 3, \dots, n$ .

Substituting (1.5) and (1.6) in (1.4), we find (1.3). The proof of Lemma 1.6 is thus achieved.  $\square$

Now, we introduce the Banach space

$$S := \{(x_1, x_2, \dots, x_n) : x_k \in C([0, 1], \mathbb{R}), k = 1, 2, \dots, n\},$$

endowed with the norm:

$$\|(x_1, x_2, \dots, x_n)\|_S = \max_{1 \leq k \leq n} \|x_k\|_\infty, \quad \|x_k\|_\infty = \max_{t \in [0, 1]} |x_k(t)|.$$

## 2. EXISTENCE AND UNIQUENESS

In this section, we try to establish sufficient conditions for the existence and uniqueness of solutions to the problem (1.1).

Define the nonlinear operator  $A : S \rightarrow S$  by

$$A(x_1, x_2, \dots, x_n)(t) := (A_1(x_1)(t), A_2(x_1, x_2)(t), \dots, A_n(x_1, x_2, \dots, x_n)(t)),$$

such that, for all  $t \in [0, 1]$ ,

$$A_k(x_1, \dots, x_k)(t) := \begin{cases} \int_0^t \frac{(t-s)^{\alpha_1-1}}{\Gamma(\alpha_1)} f_1(s) ds + a_0^1, & k = 1, \\ \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} f_k(s, \dots) ds + \sum_{j=0}^{k-2} \frac{a_j^k}{j!} t^j - \frac{\Gamma(k - \delta_{k-1})}{(k-1)!} t^{k-1} \\ \quad \times \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1}}{\Gamma(\alpha_k - \delta_{k-1})} f_k(s, \dots) ds, & k = 2, 3, \dots, n. \end{cases}$$

**Lemma 2.1.** *Let  $k-1 < \alpha_k < k$ ,  $k = 1, 2, \dots, n$ ,  $n \in \mathbb{N} - \{0, 1\}$ ,  $0 < \beta_k < 1$ ,  $T_k : (0, 1] \rightarrow \mathbb{R}$  be continuous function and  $\lim_{t \rightarrow 0^+} T_k(t) = \infty$ . Assume that  $t^{\beta_k} T_k(t)$*

is continuous on  $[0, 1]$ . Then

$$x_k(t) = \begin{cases} \int_0^t \frac{(t-s)^{\alpha_1-1}}{\Gamma(\alpha_1)} T_1(s) ds + a_0^1, & k = 1, \\ \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} T_k(s) ds + \sum_{j=0}^{k-2} \frac{a_j^k}{j!} t^j - \frac{\Gamma(k-\delta_{k-1})}{(k-1)!} t^{k-1} \\ \times \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1}}{\Gamma(\alpha_k-\delta_{k-1})} T_k(s) ds, & k = 2, 3, \dots, n, \end{cases}$$

is continuous on  $[0, 1]$ .

*Proof.* By the continuity of  $t^{\beta_k} T_k$  and

$$x_k(t) = \begin{cases} \int_0^t \frac{(t-s)^{\alpha_1-1}}{\Gamma(\alpha_1)} s^{-\beta_1} s^{\beta_1} T_1(s) ds + a_0^1, & k = 1, \\ \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} s^{-\beta_k} s^{\beta_k} T_k(s) ds + \sum_{j=0}^{k-2} \frac{a_j^k}{j!} t^j - \frac{\Gamma(k-\delta_{k-1})}{(k-1)!} t^{k-1} \\ \times \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1}}{\Gamma(\alpha_k-\delta_{k-1})} s^{-\beta_k} s^{\beta_k} T_k(s) ds, & k = 2, 3, \dots, n, \end{cases}$$

we get  $x_k(0) = a_0^k$ ,  $k = 1, 2, \dots, n$ . Then, we will divide the proof into three cases.

Case 1. For  $t_0 = 0$  and for all  $t \in (0, 1]$ , by the continuity of  $t^{\beta_k} T_k$ , there exist  $M_1, \dots, M_n > 0$ , such that for all  $t \in [0, 1]$ ,  $|t^{\beta_k} T_k(t)| \leq M_k$ . Therefore, we get

$$\begin{aligned} & |x_k(t) - x_k(0)| \\ &= \begin{cases} \left| \int_0^t \frac{(t-s)^{\alpha_1-1}}{\Gamma(\alpha_1)} s^{-\beta_1} s^{\beta_1} T_1(s) ds \right|, & k = 1, \\ \left| \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} s^{-\beta_k} s^{\beta_k} T_k(s) ds + \sum_{j=1}^{k-2} \frac{a_j^k}{j!} t^j \right. \\ \left. - \frac{\Gamma(k-\delta_{k-1})}{(k-1)!} t^{k-1} \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1}}{\Gamma(\alpha_k-\delta_{k-1})} s^{-\beta_k} s^{\beta_k} T_k(s) ds \right|, & k = 2, 3, \dots, n, \end{cases} \\ &\leq \begin{cases} \frac{M_1}{\Gamma(\alpha_1)} \int_0^t (t-s)^{\alpha_1-1} s^{-\beta_1} ds, & k = 1, \\ \frac{M_k}{\Gamma(\alpha_k)} \int_0^t (t-s)^{\alpha_k-1} s^{-\beta_k} ds + \sum_{j=1}^{k-2} \frac{|a_j^k|}{j!} t^j + \frac{\Gamma(k-\delta_{k-1}) M_k}{(k-1)! \Gamma(\alpha_k-\delta_{k-1})} t^{k-1} \\ \times \int_0^1 (1-s)^{\alpha_k-\delta_{k-1}-1} s^{-\beta_k} ds, & k = 2, 3, \dots, n. \end{cases} \end{aligned}$$

Using Beta Euler function denoted by  $B$ , we obtain

$$\begin{aligned}
 & |x_k(t) - x_k(0)| \\
 & \leq \begin{cases} \frac{M_1 t^{\alpha_1 - \beta_1}}{\Gamma(\alpha_1)} \int_0^1 (1-u)^{\alpha_1 - 1} u^{-\beta_1} du, & k = 1, \\ \frac{M_k t^{\alpha_k - \beta_k}}{\Gamma(\alpha_k)} \int_0^1 (1-u)^{\alpha_k - 1} u^{-\beta_k} du + \sum_{j=1}^{k-2} \frac{|a_j^k|}{j!} t^j \\ + \frac{\Gamma(k - \delta_{k-1}) M_k B(\alpha_k - \delta_{k-1}, 1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} t^{k-1}, & k = 2, 3, \dots, n, \end{cases} \\
 & \leq \begin{cases} \frac{M_1 B(\alpha_1, 1 - \beta_1) t^{\alpha_1 - \beta_1}}{\Gamma(\alpha_1)}, & k = 1, \\ \frac{M_k B(\alpha_k, 1 - \beta_k) t^{\alpha_k - \beta_k}}{\Gamma(\alpha_k)} + \sum_{j=1}^{k-2} \frac{|a_j^k|}{j!} t^j \\ + \frac{\Gamma(k - \delta_{k-1}) M_k B(\alpha_k - \delta_{k-1}, 1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} t^{k-1}, & k = 2, 3, \dots, n, \end{cases} \\
 & \rightarrow 0 \text{ as } t \rightarrow 0, \quad k = 1, 2, \dots, n.
 \end{aligned}$$

Case 2. For  $t_0 \in (0, 1)$  and for all  $t \in (t_0, 1]$ , we have

$$\begin{aligned}
 & |x_k(t) - x_k(t_0)| \\
 & \leq \begin{cases} \left| \int_0^t \frac{(t-s)^{\alpha_1 - 1}}{\Gamma(\alpha_1)} s^{-\beta_1} s^{\beta_1} T_1(s) ds - \int_0^{t_0} \frac{(t_0-s)^{\alpha_1 - 1}}{\Gamma(\alpha_1)} s^{-\beta_1} s^{\beta_1} T_1(s) ds \right|, & k = 1, \\ \left| \int_0^t \frac{(t-s)^{\alpha_k - 1}}{\Gamma(\alpha_k)} s^{-\beta_k} s^{\beta_k} T_k(s) ds - \int_0^{t_0} \frac{(t_0-s)^{\alpha_k - 1}}{\Gamma(\alpha_k)} s^{-\beta_k} s^{\beta_k} T_k(s) ds \right| \\ + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} (t^j - t_0^j) + \frac{\Gamma(k - \delta_{k-1})}{(k-1)!} (t^{k-1} - t_0^{k-1}) \\ \times \left| \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1}}{\Gamma(\alpha_k - \delta_{k-1})} s^{-\beta_k} s^{\beta_k} T_k(s) ds \right|, & k = 2, 3, \dots, n, \end{cases} \\
 & \leq \begin{cases} \frac{M_1}{\Gamma(\alpha_1)} \left( \int_0^t (t-s)^{\alpha_1 - 1} s^{-\beta_1} ds - \int_0^{t_0} (t_0-s)^{\alpha_1 - 1} s^{-\beta_1} ds \right), & k = 1, \\ \frac{M_k}{\Gamma(\alpha_k)} \left( \int_0^t (t-s)^{\alpha_k - 1} s^{-\beta_k} ds - \int_0^{t_0} (t_0-s)^{\alpha_k - 1} s^{-\beta_k} ds \right) \\ + \sum_{j=1}^{k-2} \frac{|a_j^k|}{j!} (t^j - t_0^j) + \frac{\Gamma(k - \delta_{k-1}) M_k}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} (t^{k-1} - t_0^{k-1}) \\ \times \int_0^1 (1-s)^{\alpha_k - \delta_{k-1} - 1} s^{-\beta_k} ds, & k = 2, 3, \dots, n. \end{cases}
 \end{aligned}$$

Therefore,

$$\begin{aligned}
& |x_k(t) - x_k(t_0)| \\
& \leq \begin{cases} \frac{M_1 (t^{\alpha_1 - \beta_1} - t_0^{\alpha_1 - \beta_1}) B(\alpha_1, 1 - \beta_1)}{\Gamma(\alpha_1)}, & k = 1, \\ \frac{M_k (t^{\alpha_k - \beta_k} - t_0^{\alpha_k - \beta_k}) B(\alpha_k, 1 - \beta_k)}{\Gamma(\alpha_k)} + \sum_{j=1}^{k-2} \frac{|a_j^k|}{j!} (t^j - t_0^j) \\ + \frac{\Gamma(k - \delta_{k-1}) M_k B(\alpha_k - \delta_{k-1}, 1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} (t^{k-1} - t_0^{k-1}), & k = 2, 3, \dots, n, \end{cases} \\
& \rightarrow 0, \text{ as } t \rightarrow t_0, \quad k = 1, 2, \dots, n.
\end{aligned}$$

Case 3. For  $t_0 \in (0, 1]$  and for all  $t \in [0, t_0]$ . Similarly, as in Case 2, it can be shown that

$$\begin{aligned}
& |x_k(t) - x_k(t_0)| \\
& \leq \begin{cases} \frac{M_1 (t_0^{\alpha_1 - \beta_1} - t^{\alpha_1 - \beta_1}) B(\alpha_1, 1 - \beta_1)}{\Gamma(\alpha_1)}, & k = 1, \\ \frac{M_k (t_0^{\alpha_k - \beta_k} - t^{\alpha_k - \beta_k}) B(\alpha_k, 1 - \beta_k)}{\Gamma(\alpha_k)} + \sum_{j=1}^{k-2} \frac{|a_j^k|}{j!} (t_0^j - t^j) \\ + \frac{\Gamma(k - \delta_{k-1}) M_k B(\alpha_k - \delta_{k-1}, 1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} (t_0^{k-1} - t^{k-1}), & k = 2, 3, \dots, n, \end{cases} \\
& \rightarrow 0, \text{ as } t \rightarrow t_0, \quad k = 1, 2, \dots, n.
\end{aligned}$$

This ends the proof.  $\square$

**Lemma 2.2.** *Let  $k - 1 < \alpha_k < k$ ,  $k = 1, 2, \dots, n$ ,  $n \in \mathbb{N} - \{0, 1\}$ ,  $0 < \beta_k < 1$ ,  $f_k : (0, 1] \times \mathbb{R}^k \rightarrow \mathbb{R}$  be continuous, and  $\lim_{t \rightarrow 0^+} f_k(t, \dots) = \infty$ . Assume that  $t^{\beta_k} f_k(t, \dots)$  is continuous on  $[0, 1] \times \mathbb{R}^k$ . Then, the operator  $A : S \rightarrow S$  is completely continuous.*

*Proof.* For all  $(x_1, \dots, x_n) \in S$ , let

$$A(x_1, x_2, \dots, x_n)(t) = (A_1(x_1), A_2(x_1, x_2), \dots, A_n(x_1, \dots, x_n))(t),$$

where

$$\begin{aligned}
& A_k(x_1, \dots, x_k)(t) \\
& := \begin{cases} \int_0^t \frac{(t-s)^{\alpha_1-1}}{\Gamma(\alpha_1)} f_1(s, x_1(s)) ds + a_0^1, & k = 1, \\ \int_0^t \frac{(t-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} f_k(s, x_1(s), \dots, x_k(s)) ds + \sum_{j=0}^{k-2} \frac{a_j^k}{j!} t^j - \frac{\Gamma(k - \delta_{k-1})}{(k-1)!} t^{k-1} \\ \times \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1}}{\Gamma(\alpha_k - \delta_{k-1})} f_k(s, x_1(s), \dots, x_k(s)) ds, & k = 2, 3, \dots, n. \end{cases}
\end{aligned}$$



By Lemma 2.1, we have  $A : S \rightarrow S$ . Let

$$(x_1^0, x_2^0, \dots, x_n^0) \in S : \|(x_1^0, x_2^0, \dots, x_n^0)\|_S = \lambda_0$$

and

$$(x_1, x_2, \dots, x_n) \in S : \|(x_1, x_2, \dots, x_n) - (x_1^0, x_2^0, \dots, x_n^0)\|_S < 1,$$

then

$$\|(x_1, x_2, \dots, x_n)\|_S < 1 + \lambda_0 = \lambda.$$

By the continuity of  $t^{\beta_k} f_k(t, x_1, \dots, x_k)$ , we know that  $t^{\beta_k} f_k(t, x_1, \dots, x_k)$  is uniformly continuous on  $[0, 1] \times [-\lambda, \lambda]^k$ .

Hence, for all  $t \in [0, 1]$  and for each  $\epsilon > 0$ , there exists  $\rho > 0$  ( $\rho < 1$ ), with

$$(2.1) \quad |t^{\beta_k} f_k(t, x_1(t), \dots, x_k(t)) - t^{\beta_k} f_k(t, x_1^0(t), \dots, x_k^0(t))| < \epsilon,$$

where  $(x_1, x_2, \dots, x_n) \in S$ , and  $\|(x_1, x_2, \dots, x_n) - (x_1^0, x_2^0, \dots, x_n^0)\|_S < \rho$ . Then

$$(2.2) \quad \begin{aligned} & \|A(x_1, x_2, \dots, x_n) - A(x_1^0, x_2^0, \dots, x_n^0)\|_S \\ &= \max_{1 \leq k \leq n} \|A_k(x_1, \dots, x_k)(t) - A_k(x_1^0, \dots, x_k^0)(t)\|_\infty. \end{aligned}$$

We have

$$\begin{aligned} & \|A_k(x_1, \dots, x_k)(t) - A_k(x_1^0, \dots, x_k^0)(t)\|_\infty \\ & \leq \begin{cases} \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_1-1} s^{-\beta_1}}{\Gamma(\alpha_1)} |s^{\beta_1} f_1(s, x_1(s)) - s^{\beta_1} f_1(s, x_1^0(s))| ds, & k=1, \\ \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} |s^{\beta_k} f_k(s, x_1(s), \dots, x_k(s)) \\ - s^{\beta_k} f_k(s, x_1^0(s), \dots, x_k^0(s))| ds + \max_{t \in [0,1]} \frac{\Gamma(k - \delta_{k-1})}{(k-1)!} t^{k-1} \\ \times \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1} s^{-\beta_k}}{\Gamma(\alpha_k - \delta_{k-1})} |s^{\beta_k} f_k(s, x_1(s), \dots, x_k(s)) \\ - s^{\beta_k} f_k(s, x_1^0(s), \dots, x_k^0(s))| ds, & k=2, 3, \dots, n. \end{cases} \end{aligned}$$

Using (2.1), we obtain

$$(2.3) \quad \begin{aligned} & \|A_k(x_1, \dots, x_k)(t) - A_k(x_1^0, \dots, x_k^0)(t)\|_\infty \\ & \leq \begin{cases} \frac{\epsilon}{\Gamma(\alpha_1)} \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_1-1} s^{-\beta_1} ds, & k=1, \\ \frac{\epsilon}{\Gamma(\alpha_k)} \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_k-1} s^{-\beta_k} ds \\ + \frac{\epsilon \Gamma(k - \delta_{k-1})}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} \max_{t \in [0,1]} t^{k-1} \int_0^1 (1-s)^{\alpha_k - \delta_{k-1} - 1} s^{-\beta_k} ds, & k=2, 3, \dots, n, \end{cases} \end{aligned}$$

$$\leq \begin{cases} \epsilon \frac{B(\alpha_1, 1 - \beta_1)}{\Gamma(\alpha_1)} \max_{t \in [0,1]} t^{\alpha_1 - \beta_1}, & k = 1, \\ \epsilon \left( \frac{B(\alpha_k, 1 - \beta_k)}{\Gamma(\alpha_k)} \max_{t \in [0,1]} t^{\alpha_k - \beta_k} + \frac{\Gamma(k - \delta_{k-1}) B(\alpha_k - \delta_{k-1}, 1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} \right), & k = 2, 3, \dots, n, \end{cases}$$

$$= \begin{cases} \epsilon \frac{\Gamma(1 - \beta_1)}{\Gamma(\alpha_1 + 1 - \beta_1)}, & k = 1, \\ \epsilon \left( \frac{\Gamma(1 - \beta_k)}{\Gamma(\alpha_k + 1 - \beta_k)} + \frac{\Gamma(k - \delta_{k-1}) \Gamma(1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1} + 1 - \beta_k)} \right), & k = 2, 3, \dots, n. \end{cases}$$

We pose:

$$(2.4) \quad \Lambda_1 := \frac{\Gamma(1 - \beta_1)}{\Gamma(\alpha_1 + 1 - \beta_1)},$$

$$\Lambda_k := \frac{\Gamma(1 - \beta_k)}{\Gamma(\alpha_k + 1 - \beta_k)} + \frac{\Gamma(k - \delta_{k-1}) \Gamma(1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1} + 1 - \beta_k)}.$$

By (2.3) and (2.4), we have

$$(2.5) \quad \|A_k(x_1, \dots, x_k)(t) - A_k(x_1^0, \dots, x_k^0)(t)\|_\infty \leq \begin{cases} \epsilon \Lambda_1, & k = 1, \\ \epsilon \Lambda_k, & k = 2, 3, \dots, n. \end{cases}$$

Thanks to (2.2) and (2.5), we get

$$\|A(x_1, x_2, \dots, x_n) - A(x_1^0, x_2^0, \dots, x_n^0)\|_S \leq \epsilon \max_{1 \leq k \leq n} \Lambda_k.$$

Therefore,

$$\|A(x_1, x_2, \dots, x_n) - A(x_1^0, x_2^0, \dots, x_n^0)\|_S \rightarrow 0,$$

as

$$\|(x_1, x_2, \dots, x_n) - (x_1^0, x_2^0, \dots, x_n^0)\|_S \rightarrow 0.$$

Hence,  $A : S \rightarrow S$  is continuous.

Let  $\theta \subset S$  be bounded. Then, there exists a positive constant  $\varsigma$  such that  $\|(x_1, x_2, \dots, x_n)\|_S \leq \varsigma$ , for all  $(x_1, x_2, \dots, x_n) \in \theta$ . Since  $t^{\beta_k} f_k(t, x_1, \dots, x_k)$ ,  $k = 1, 2, \dots, n$ , are continuous on  $[0, 1] \times [-\varsigma, \varsigma]^k$ , there exist positive constants  $L_k$ ,  $k = 1, 2, \dots, n$ , such that

$$(2.6) \quad |t^{\beta_k} f_k(t, x_1(t), \dots, x_k(t))| \leq L_k, \quad \text{for all } t \in [0, 1], \text{ for all } (x_1, x_2, \dots, x_n) \in \theta.$$

Then

$$(2.7) \quad \|A(x_1, x_2, \dots, x_n)\|_S = \max_{1 \leq k \leq n} \|A_k(x_1, \dots, x_k)\|_\infty.$$

We have

$$\|A_k(x_1, \dots, x_k)\|_\infty \leq \begin{cases} \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_1-1} s^{-\beta_1}}{\Gamma(\alpha_1)} |s^{\beta_1} f_1(s, x_1)| ds + |a_0^1|, & k=1, \\ \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} |s^{\beta_k} f_k(s, x_1, \dots, x_k)| ds \\ + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \max_{t \in [0,1]} t^j + \frac{\Gamma(k-\delta_{k-1})}{(k-1)!} \max_{t \in [0,1]} t^{k-1} \\ \times \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1} s^{-\beta_k}}{\Gamma(\alpha_k-\delta_{k-1})} |s^{\beta_k} f_k(s, x_1, \dots, x_k)| ds, \\ k=2, 3, \dots, n. \end{cases}$$

Using (2.6), we get

$$(2.8) \quad \|A_k(x_1, \dots, x_k)\|_\infty \leq \begin{cases} \frac{L_1}{\Gamma(\alpha_1)} \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_1-1} s^{-\beta_1} ds + |a_0^1|, & k=1, \\ \frac{L_k}{\Gamma(\alpha_k)} \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_k-1} s^{-\beta_k} ds + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \\ + \frac{\Gamma(k-\delta_{k-1}) L_k}{(k-1)! \Gamma(\alpha_k-\delta_{k-1})} \int_0^1 (1-s)^{\alpha_k-\delta_{k-1}-1} s^{-\beta_k} ds, & k=2, 3, \dots, n, \\ \frac{L_1 \Gamma(1-\beta_1)}{\Gamma(\alpha_1+1-\beta_1)} \max_{t \in [0,1]} t^{\alpha_1-\beta_1} + |a_0^1|, & k=1, \\ L_k \left( \frac{\Gamma(1-\beta_k)}{\Gamma(\alpha_k+1-\beta_k)} \max_{t \in [0,1]} t^{\alpha_k-\beta_k} + \frac{\Gamma(k-\delta_{k-1}) \Gamma(1-\beta_k)}{(k-1)! \Gamma(\alpha_k-\delta_{k-1}+1-\beta_k)} \right) \\ + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!}, & k=2, 3, \dots, n, \\ L_1 \Lambda_1 + |a_0^1|, & k=1, \\ L_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!}, & k=2, 3, \dots, n. \end{cases}$$

Then by (2.7) and (2.8), we get

$$\|A(x_1, x_2, \dots, x_n)\|_S \leq \max_{2 \leq k \leq n} \left\{ L_1 \Lambda_1 + |a_0^1|, L_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \right\}.$$

Thus,  $A(\theta)$  is bounded.

For all  $(x_1, x_2, \dots, x_n) \in \theta$ , and for all  $t_1, t_2 \in [0, 1]$ ,  $t_1 < t_2$ , we have:

$$\|A(x_1, x_2, \dots, x_n)(t_2) - A(x_1, x_2, \dots, x_n)(t_1)\|_S$$

$$(2.9) \quad = \max_{1 \leq k \leq n} \|A_k(x_1, \dots, x_k)(t_2) - A_k(x_1, \dots, x_k)(t_1)\|_\infty.$$

Then

$$\|A_k(x_1, \dots, x_k)(t_2) - A_k(x_1, \dots, x_k)(t_1)\|_\infty \leq \begin{cases} \max_{t \in [0,1]} \left| \int_0^{t_2} \frac{(t_2-s)^{\alpha_1-1} s^{-\beta_1}}{\Gamma(\alpha_1)} s^{\beta_1} f_1(s, x_1) ds - \int_0^{t_1} \frac{(t_1-s)^{\alpha_1-1} s^{-\beta_1}}{\Gamma(\alpha_1)} s^{\beta_1} f_1(s, x_1) ds \right|, & k=1, \\ \max_{t \in [0,1]} \left| \int_0^{t_2} \frac{(t_2-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} s^{\beta_k} f_k(s, x_1, \dots, x_k) ds - \int_0^{t_1} \frac{(t_1-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} s^{\beta_k} f_k(s, x_1, \dots, x_k) ds \right| \\ + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} (t_2^j - t_1^j) + \frac{\Gamma(k - \delta_{k-1})}{(k-1)!} (t_2^{k-1} - t_1^{k-1}) \\ \times \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1} s^{-\beta_k}}{\Gamma(\alpha_k - \delta_{k-1})} |s^{\beta_k} f_k(s, x_1, \dots, x_k)| ds, & k=2, 3, \dots, n. \end{cases}$$

Hence,

$$(2.10) \quad \|A_k(x_1, \dots, x_k)(t_2) - A_k(x_1, \dots, x_k)(t_1)\|_\infty \leq \begin{cases} \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(\alpha_1 + 1 - \beta_1)} (t_2^{\alpha_1 - \beta_1} - t_1^{\alpha_1 - \beta_1}), & k=1, \\ \frac{L_k \Gamma(1 - \beta_k) (t_2^{\alpha_k - \beta_k} - t_1^{\alpha_k - \beta_k})}{\Gamma(\alpha_k + 1 - \beta_k)} + \sum_{j=0}^{k-2} \frac{|a_j^k| (t_2^j - t_1^j)}{j!} \\ + \frac{\Gamma(k - \delta_{k-1}) L_k \Gamma(1 - \beta_k) (t_2^{k-1} - t_1^{k-1})}{(k-1)! \Gamma(\alpha_k - \delta_{k-1} + 1 - \beta_k)}, & k=2, 3, \dots, n. \end{cases}$$

Then, by (2.9) and (2.10), we obtain

$$(2.11) \quad \|A(x_1, x_2, \dots, x_n)(t_2) - A(x_1, x_2, \dots, x_n)(t_1)\|_S \leq \max \left\{ \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(\alpha_1 + 1 - \beta_1)} (t_2^{\alpha_1 - \beta_1} - t_1^{\alpha_1 - \beta_1}), \frac{L_k \Gamma(1 - \beta_k) (t_2^{\alpha_k - \beta_k} - t_1^{\alpha_k - \beta_k})}{\Gamma(\alpha_k + 1 - \beta_k)} + \sum_{j=0}^{k-2} \frac{|a_j^k| (t_2^j - t_1^j)}{j!} + \frac{\Gamma(k - \delta_{k-1}) L_k \Gamma(1 - \beta_k) (t_2^{k-1} - t_1^{k-1})}{(k-1)! \Gamma(\alpha_k - \delta_{k-1} + 1 - \beta_k)} \right\}.$$

The right-hand side of (2.11) is independent of  $(x_1, x_2, \dots, x_n)$  and tends to zero as  $t_1 \rightarrow t_2$ . Thus  $A(\theta)$  is equicontinuous. By Arzela-Ascoli theorem,  $A$  is completely continuous.  $\square$

**Theorem 2.1.** Assume that there exist nonnegative constants  $(\omega_j^k)_{j=1,\dots,k}^{k=1,\dots,n}$ , satisfying

$$(2.12) \quad t^{\beta_k} |f_k(t, x_1, \dots, x_k) - f_k(t, y_1, \dots, y_k)| \leq \sum_{j=1}^k \omega_j^k |x_j - y_j|,$$

for all  $t \in [0, 1]$  and all  $(x_1, \dots, x_k), (y_1, \dots, y_k) \in \mathbb{R}^k$ .

If

$$(2.13) \quad \Sigma := \max_{2 \leq k \leq n} \left( \omega_1^1 \Lambda_1, \sum_{j=1}^k \omega_j^k \Lambda_k \right) < 1,$$

then the system (1.1) has a unique solution on  $[0, 1]$ .

*Proof.* We will prove that  $A$  is a contractive operator on  $S$ .

Let  $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n) \in S$  and  $t \in [0, 1]$ , we have

$$(2.14) \quad \begin{aligned} & \|A(x_1, x_2, \dots, x_n) - A(y_1, y_2, \dots, y_n)\|_S \\ &= \max_{1 \leq k \leq n} \|A_k(x_1, \dots, x_k)(t) - A_k(y_1, \dots, y_k)(t)\|_\infty. \end{aligned}$$

Then

$$\begin{aligned} & \|A_k(x_1, \dots, x_k)(t) - A_k(y_1, \dots, y_k)(t)\|_\infty \\ & \leq \begin{cases} \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_1-1} s^{-\beta_1}}{\Gamma(\alpha_1)} s^{\beta_1} |f_1(s, x_1(s)) - f_1(s, y_1(s))| ds, & k=1, \\ \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} s^{\beta_k} |f_k(s, x_1(s), \dots, x_k(s)) \\ - f_k(s, y_1(s), \dots, y_k(s))| ds + \max_{t \in [0,1]} \frac{\Gamma(k - \delta_{k-1})}{(k-1)!} t^{k-1} \\ \times \int_0^1 \frac{(1-s)^{\alpha_k - \delta_{k-1} - 1} s^{-\beta_k}}{\Gamma(\alpha_k - \delta_{k-1})} s^{\beta_k} |f_k(s, x_1(s), \dots, x_k(s)) \\ - f_k(s, y_1(s), \dots, y_k(s))| ds, & k=2, 3, \dots, n. \end{cases} \end{aligned}$$

Thanks to (2.12), we can write

$$(2.15) \quad \begin{aligned} & \|A_k(x_1, \dots, x_k)(t) - A_k(y_1, \dots, y_k)(t)\|_\infty \\ & \leq \begin{cases} \frac{\omega_1^1}{\Gamma(\alpha_1)} \|x_1 - y_1\|_\infty \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_1-1} s^{-\beta_1} ds, & k=1, \\ \left( \omega_1^k \|x_1 - y_1\|_\infty + \dots + \omega_k^k \|x_k - y_k\|_\infty \right) \\ \times \left( \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} ds \right. \\ \left. + \frac{\Gamma(k - \delta_{k-1})}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} \int_0^1 (1-s)^{\alpha_k - \delta_{k-1} - 1} s^{-\beta_k} ds \right), & k=2, 3, \dots, n, \end{cases} \end{aligned}$$

$$\begin{aligned}
& \leq \begin{cases} \frac{\omega_1^1 B(\alpha_1, 1 - \beta_1) \|x_1 - y_1\|_\infty \max_{t \in [0,1]} t^{\alpha_1 - \beta_1}, & k = 1, \\ \sum_{j=1}^k \omega_j^k \max_{1 \leq k \leq n} \|x_k - y_k\|_\infty \left( \frac{B(\alpha_k, 1 - \beta_k)}{\Gamma(\alpha_k)} \max_{t \in [0,1]} t^{\alpha_k - \beta_k} \right. \\ \left. + \frac{\Gamma(k - \delta_{k-1}) B(\alpha_k - \delta_{k-1}, 1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1})} \right), & k = 2, 3, \dots, n, \end{cases} \\
& \leq \begin{cases} \frac{\omega_1^1 \Gamma(1 - \beta_1)}{\Gamma(\alpha_1 + 1 - \beta_1)} \|x_1 - y_1\|_\infty, & k = 1, \\ \sum_{j=1}^k \omega_j^k \left( \frac{\Gamma(1 - \beta_k)}{\Gamma(\alpha_k + 1 - \beta_k)} + \frac{\Gamma(k - \delta_{k-1}) \Gamma(1 - \beta_k)}{(k-1)! \Gamma(\alpha_k - \delta_{k-1} + 1 - \beta_k)} \right) \\ \times \|(x_1 - y_1, \dots, x_k - y_k)\|_S, & k = 2, 3, \dots, n. \end{cases}
\end{aligned}$$

By (2.14) and (2.15), we obtain

$$\begin{aligned}
& \|A(x_1, x_2, \dots, x_n) - A(y_1, y_2, \dots, y_n)\|_S \\
& \leq \max_{2 \leq k \leq n} \left( \omega_1^1 \Lambda_1, \sum_{j=1}^k \omega_j^k \Lambda_k \right) \|(x_1 - y_1, \dots, x_k - y_k)\|_S.
\end{aligned}$$

By (2.13), we have  $\Sigma := \max_{2 \leq k \leq n} (\omega_1^1 \Lambda_1, \sum_{j=1}^k \omega_j^k \Lambda_k) < 1$ . Hence,  $A$  is a contractive operator. Consequently, by Banach fixed point theorem,  $A$  has a fixed point which is the unique solution of system (1.1). This completes the proof.  $\square$

*Example 2.1.* Consider the following singular fractional system:

$$(2.16) \quad \begin{cases} D^{\frac{3}{4}} x_1(t) = \frac{\sin x_1(t)}{12\pi\sqrt{t}}, & 0 < t \leq 1, \\ D^{\frac{3}{2}} x_2(t) = \frac{\cos x_1(t) - \cos x_2(t)}{16\pi^3 t^{\frac{5}{7}}}, & 0 < t \leq 1, \\ D^{\frac{7}{3}} x_3(t) = \frac{(\sin x_1(t) + \sin x_2(t) + \cos x_3(t))}{24\pi t^{\frac{3}{8}}}, & 0 < t \leq 1, \\ D^{\frac{7}{2}} x_4(t) = \frac{|x_1(t) + x_2(t) + x_3(t) + x_4(t)|}{32\pi t^{\frac{1}{3}} (1 + |x_1(t) + x_2(t) + x_3(t) + x_4(t)|)}, & 0 < t \leq 1, \\ x_1(0) = 1, \\ x_2(0) = \sqrt{2}, \quad D^{\frac{1}{2}} x_2(1) = 0, \\ x_3(0) = \frac{3}{5}, \quad x_3'(0) = 2\sqrt{3}, \quad D^{\frac{4}{3}} x_3(1) = 0, \\ x_4(0) = \frac{1}{2}, \quad x_4'(0) = \sqrt{5}, \quad x_4''(0) = 1, \quad D^{\frac{5}{2}} x_4(1) = 0. \end{cases}$$

We have:

$$\begin{aligned}
n = 4, \quad \alpha_1 = \frac{3}{4}, \quad \alpha_2 = \frac{3}{2}, \quad \alpha_3 = \frac{7}{3}, \quad \alpha_4 = \frac{7}{2}, \quad \delta_1 = \frac{1}{2}, \quad \delta_2 = \frac{4}{3}, \quad \delta_3 = \frac{5}{2}, \\
a_0^1 = 1, \quad a_0^2 = \sqrt{2}, \quad a_0^3 = \frac{3}{5}, \quad a_1^3 = 2\sqrt{3}, \quad a_0^4 = \frac{1}{2}, \quad a_1^4 = \sqrt{5}, \quad a_2^4 = 1.
\end{aligned}$$

Then, for each  $t \in [0, 1]$  and  $(x_1, x_2, x_3, x_4), (y_1, y_2, y_3, y_4) \in \mathbb{R}^4$ , we have:

$$\begin{aligned} & t^{\frac{2}{3}} |f_1(t, x_1) - f_1(t, y_1)| \leq \frac{t^{\frac{1}{6}}}{12\pi} |x_1 - y_1|, \\ & t^{\frac{6}{7}} |f_2(t, x_1, x_2) - f_2(t, y_1, y_2)| \leq \frac{t^{\frac{1}{7}}}{16\pi^3} |x_1 - y_1| + \frac{t^{\frac{1}{7}}}{16\pi^3} |x_2 - y_2|, \\ & t^{\frac{7}{8}} |f_3(t, x_1, x_2, x_3) - f_3(t, y_1, y_2, y_3)| \\ & \leq \left( \frac{\sqrt{t}}{24\pi} |x_1 - y_1| + \frac{\sqrt{t}}{24\pi} |x_2 - y_2| + \frac{\sqrt{t}}{24\pi} |x_3 - y_3| \right), \\ & t^{\frac{1}{2}} |f_4(t, x_1, x_2, x_3, x_4) - f_4(t, y_1, y_2, y_3, y_4)| \\ & \leq \left( \frac{t^{\frac{1}{6}}}{32\pi} |x_1 - y_1| + \frac{t^{\frac{1}{6}}}{32\pi} |x_2 - y_2| + \frac{t^{\frac{1}{6}}}{32\pi} |x_3 - y_3| + \frac{t^{\frac{1}{6}}}{32\pi} |x_4 - y_4| \right), \end{aligned}$$

where  $\beta_1 = \frac{2}{3}, \beta_2 = \frac{6}{7}, \beta_3 = \frac{7}{8}, \beta_4 = \frac{1}{2}$ .

Moreover, we can take:

$$\begin{aligned} \omega_1^1 &= \frac{1}{12\pi}, \\ \omega_1^2 &= \omega_2^2 = \frac{1}{16\pi^3}, \quad \sum_{j=1}^2 \omega_j^2 = \frac{1}{8\pi^3}, \\ \omega_1^3 &= \omega_2^3 = \omega_3^3 = \frac{1}{24\pi}, \quad \sum_{j=1}^3 \omega_j^3 = \frac{1}{8\pi}, \\ \omega_1^4 &= \omega_2^4 = \omega_3^4 = \omega_4^4 = \frac{1}{32\pi}, \quad \sum_{j=1}^4 \omega_j^4 = \frac{1}{8\pi}. \end{aligned}$$

On the other hand, we get

$$\Lambda_1 = 2.7958, \quad \Lambda_2 = 13.4869, \quad \Lambda_3 = 9.4443, \quad \Lambda_4 = 0.5908.$$

Thus,

$$\omega_1^1 \Lambda_1 = 0.0742, \quad \sum_{j=1}^2 \omega_j^2 \Lambda_2 = 0.0544, \quad \sum_{j=1}^3 \omega_j^3 \Lambda_3 = 0.3759, \quad \sum_{j=1}^4 \omega_j^4 \Lambda_4 = 0.0235.$$

Then the singular fractional system (2.16) has a unique solution on  $[0, 1]$ .

**Theorem 2.2.** *Let  $k - 1 < \alpha_k < k$ ,  $k = 1, 2, \dots, n$ ,  $n \in \mathbb{N} - \{0, 1\}$ ,  $0 < \beta_k < 1$ . Assume that  $f_k : (0, 1] \times \mathbb{R}^k \rightarrow \mathbb{R}$  is continuous with  $\lim_{t \rightarrow 0^+} f_k(t, \dots) = \infty$  and  $t^{\beta_k} f_k(t, \dots)$  is continuous on  $[0, 1] \times \mathbb{R}^k$ . Then, the system (1.1) has at least one solution on  $[0, 1]$ .*

*Proof.* Let  $P_k = \max_{t \in [0, 1]} t^{\beta_k} |f_k(t, x_1(t), \dots, x_k(t))|$ , and define the set  $\Delta \subset S$  by

$$\Delta := \{(x_1, x_2, \dots, x_n) \in S : \|(x_1, x_2, \dots, x_n)\|_S \leq r\},$$

where

$$r = \max_{2 \leq k \leq n} \left( P_1 \Lambda_1 + |a_0^1|, P_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \right).$$

We will prove that  $A : \Delta \rightarrow \Delta$ . For  $(x_1, x_2, \dots, x_n) \in \Delta$  and  $t \in [0, 1]$ , we have

$$(2.17) \quad \|A(x_1, x_2, \dots, x_n)\|_S = \max_{1 \leq k \leq n} \|A_k(x_1, \dots, x_k)(t)\|_\infty.$$

Then

$$\begin{aligned} & \|A_k(x_1, \dots, x_k)(t)\|_\infty \\ & \leq \begin{cases} \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_1-1} s^{-\beta_1}}{\Gamma(\alpha_1)} s^{\beta_1} |f_1(s, x_1)| ds + |a_0^1|, & k=1, \\ \max_{t \in [0,1]} \int_0^t \frac{(t-s)^{\alpha_k-1} s^{-\beta_k}}{\Gamma(\alpha_k)} s^{\beta_k} |f_k(s, x_1, \dots, x_k)| ds + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \max_{t \in [0,1]} t^j \\ \quad + \frac{\Gamma(k-\delta_{k-1})}{(k-1)!} \max_{t \in [0,1]} t^{k-1} \int_0^1 \frac{(1-s)^{\alpha_k-\delta_{k-1}-1} s^{-\beta_k}}{\Gamma(\alpha_k-\delta_{k-1})} s^{\beta_k} |f_k(s, x_1, \dots, x_k)| ds, \\ & k=2, 3, \dots, n, \end{cases} \\ & \leq \begin{cases} \frac{P_1}{\Gamma(\alpha_1)} \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_1-1} s^{-\beta_1} ds + |a_0^1|, & k=1, \\ \frac{P_k}{\Gamma(\alpha_k)} \max_{t \in [0,1]} \int_0^t (t-s)^{\alpha_k-1} s^{-\beta_k} ds + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \\ \quad + \frac{\Gamma(k-\delta_{k-1}) P_k}{(k-1)! \Gamma(\alpha_k-\delta_{k-1})} \int_0^1 (1-s)^{\alpha_k-\delta_{k-1}-1} s^{-\beta_k} ds, & k=2, 3, \dots, n. \end{cases} \end{aligned}$$

Thus,

$$(2.18) \quad \begin{aligned} & \|A_k(x_1, \dots, x_k)(t)\|_\infty \\ & \leq \begin{cases} \frac{P_1 \Gamma(1-\beta_1)}{\Gamma(\alpha_1+1-\beta_1)} \max_{t \in [0,1]} t^{\alpha_1-\beta_1} + |a_0^1|, & k=1, \\ P_k \left( \frac{\Gamma(1-\beta_k)}{\Gamma(\alpha_k+1-\beta_k)} \max_{t \in [0,1]} t^{\alpha_k-\beta_k} + \frac{\Gamma(k-\delta_{k-1}) \Gamma(1-\beta_k)}{(k-1)! \Gamma(\alpha_k-\delta_{k-1}+1-\beta_k)} \right) \\ \quad + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!}, & k=2, 3, \dots, n, \end{cases} \\ & \leq \begin{cases} P_1 \Lambda_1 + |a_0^1|, & k=1, \\ P_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!}, & k=2, 3, \dots, n. \end{cases} \end{aligned}$$

Using (2.17) and (2.18), we can write

$$(2.19) \quad \|A(x_1, x_2, \dots, x_n)\|_S \leq \max_{2 \leq k \leq n} \left( P_1 \Lambda_1 + |a_0^1|, P_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!} \right).$$



Hence,  $\|A(x_1, x_2, \dots, x_n)\|_S \leq r$ . By Lemma 2.1, we have  $A(x_1, x_2, \dots, x_n)(t) \in C([0, 1])$ . Moreover, for  $(x_1, x_2, \dots, x_n) \in \Delta$ , we have  $A(x_1, x_2, \dots, x_n) \in \Delta$ . So,  $A(\Delta) \subset \Delta$ , and  $A : \Delta \rightarrow \Delta$ . Then from Lemma 2.2, we get  $A$  is completely continuous.

By Lemma 1.5, the system (1.1) has at least one solution on  $[0, 1]$ . Theorem 2.2 is thus proved.  $\square$

*Example 2.2.* Consider the following system:

$$(2.20) \left\{ \begin{array}{l} D^{\frac{1}{2}}x_1(t) = t^{-\frac{1}{3}}e^{-t} \sin x_1(t), \quad 0 < t \leq 1, \\ D^{\frac{4}{3}}x_2(t) = \frac{\cos x_1(t)}{\sqrt{t}(\pi + \sin x_2(t))}, \quad 0 < t \leq 1, \\ D^{\frac{9}{4}}x_3(t) = \frac{e^t \cos x_3}{t^{\frac{1}{5}}(4\pi + \sin(x_1 + x_2))}, \quad 0 < t \leq 1, \\ D^{\frac{7}{2}}x_4(t) = t^{-\frac{2}{9}} \frac{e^{-2t} \sin(x_1 + x_2)}{16 + \cos(x_3 + x_4)}, \quad 0 < t \leq 1, \\ D^{\frac{14}{3}}x_5(t) = \frac{\cos(u_1 + u_2 + u_3 + u_4)}{t^{\frac{1}{4}}e^t}, \quad 0 < t \leq 1, \\ x_1(0) = \sqrt{3}, \\ x_2(0) = \frac{2}{3}, \quad D^{\frac{1}{4}}x_2(1) = 0, \\ x_3(0) = -1, \quad x'_3(0) = \frac{1}{2}, \quad D^{\frac{3}{2}}x_3(1) = 0, \\ x_4(0) = \frac{\sqrt{7}}{2}, \quad x'_4(0) = \frac{1}{4}, \quad x''_4(0) = \frac{\sqrt{5}}{3}, \quad D^{\frac{11}{5}}x_4(1) = 0, \\ x_5(0) = 1, \quad x'_5(0) = \frac{4}{3}, \quad x''_5(0) = \frac{3}{7}, \quad x'''_5(0) = \frac{2\sqrt{3}}{5}, \quad D^{\frac{10}{3}}x_5(1) = 0. \end{array} \right.$$

We have:

$$\begin{aligned} n = 5, \quad \alpha_1 = \frac{1}{2}, \quad \alpha_2 = \frac{4}{3}, \quad \alpha_3 = \frac{9}{4}, \quad \alpha_4 = \frac{7}{2}, \quad \alpha_5 = \frac{14}{3}, \quad \delta_1 = \frac{1}{4}, \quad \delta_2 = \frac{3}{2}, \\ \delta_3 = \frac{11}{5}, \quad \delta_4 = \frac{10}{3}, \quad a_0^1 = \sqrt{3}, \quad a_0^2 = \frac{2}{3}, \quad a_0^3 = -1, \quad a_1^3 = \frac{1}{2}, \quad a_0^4 = \frac{\sqrt{7}}{2}, \\ a_1^4 = \frac{1}{4}, \quad a_2^4 = \frac{\sqrt{5}}{3}, \quad a_0^5 = 1, \quad a_1^5 = \frac{4}{3}, \quad a_2^5 = \frac{3}{7}, \quad a_3^5 = \frac{2\sqrt{3}}{5}. \end{aligned}$$

For  $\beta_1 = \frac{2}{3}, \beta_2 = \frac{3}{4}, \beta_3 = \frac{2}{5}, \beta_4 = \frac{4}{9}, \beta_5 = \frac{1}{2}$ , the system (2.20) has at least one solution on  $[0, 1]$ .

### 3. ULAM STABILITY

In this section, we study the Ulam-Hyers stability and the generalized Ulam-Hyers stability of solutions for system (1.1).

**Definition 3.1.** The singular fractional system (1.1) is Ulam-Hyers stable if there exists a real number  $\mu > 0$ , such that for all  $(\epsilon_1, \epsilon_2, \dots, \epsilon_n) > 0$ , and for all solution

$(x_1, x_2, \dots, x_n) \in S$  of

$$(3.1) \quad \begin{cases} |D^{\alpha_1} x_1(t) - f_1(t, x_1(t))| \leq \epsilon_1, \\ |D^{\alpha_2} x_2(t) - f_2(t, x_1(t), x_2(t))| \leq \epsilon_2, \\ \vdots \\ |D^{\alpha_n} x_n(t) - f_n(t, x_1(t), x_2(t), \dots, x_n(t))| \leq \epsilon_n, \quad 0 < t \leq 1, \end{cases}$$

there exists a solution  $(y_1, y_2, \dots, y_n) \in S$  satisfying

$$(3.2) \quad \begin{cases} D^{\alpha_1} y_1(t) = f_1(t, y_1(t)), \\ D^{\alpha_2} y_2(t) = f_2(t, y_1(t), y_2(t)), \\ \vdots \\ D^{\alpha_n} y_n(t) = f_n(t, y_1(t), y_2(t), \dots, y_n(t)), \quad 0 < t \leq 1, k = 1, \\ y_1(0) = a_0^1, \\ y_k^{(j)}(0) = a_j^k, \quad k = 2, 3, \dots, n, j = 0, 1, \dots, k-2, \\ D^{\delta_{k-1}} y_k(1) = 0, \quad k = 2, 3, \dots, n, k-2 < \delta_{k-1} < k-1, \\ k-1 < \alpha_k < k, \quad k = 1, 2, \dots, n, \end{cases}$$

with

$$\|(x_1 - y_1, \dots, x_n - y_n)\|_S \leq \mu\epsilon, \quad \epsilon > 0.$$

**Definition 3.2.** The singular fractional system (1.1) is generalized Ulam-Hyers stable if there exists  $\phi \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $\phi(0) = 0$ , such that for all  $\epsilon > 0$ , and for each solution  $(x_1, x_2, \dots, x_n) \in S$  of (3.1), there exists a solution  $(y_1, y_2, \dots, y_n) \in S$  of (3.2) with

$$\|(x_1 - y_1, \dots, x_n - y_n)\|_S \leq \phi(\epsilon), \quad \epsilon > 0.$$

**Theorem 3.1.** Let  $k-1 < \alpha_k < k$ ,  $k = 1, 2, \dots, n$ ,  $n \in \mathbb{N} - \{0, 1\}$  and  $0 < \beta_k < 1$ .

Assume that:

(H<sub>1</sub>)  $f_k : (0, 1] \times \mathbb{R}^k \rightarrow \mathbb{R}$  is continuous with  $\lim_{t \rightarrow 0^+} f_k(t, \dots) = \infty$  and  $t^{\beta_k} f_k(t, \dots)$  is continuous on  $[0, 1] \times \mathbb{R}^k$ ;

$$(H_2) \quad \|t^{\beta_k} D^{\alpha_k} x_k\|_{\infty} \geq \begin{cases} P_1 \Lambda_1 + |a_0^1|, & k = 1, \\ P_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!}, & k = 2, 3, \dots, n, \end{cases}$$

(H<sub>3</sub>) all the assumptions of Theorem 2.1 are satisfied;

$$(H_4) \quad \sum_{j=1}^k \omega_j^k < 1, \quad k = 1, 2, \dots, n.$$

Then, the singular fractional system (1.1) is generalized Ulam-Hyers stable.

*Proof.* Using (H<sub>1</sub>) we receive (2.19). Thus, for all solution  $(x_1, x_2, \dots, x_n) \in S$  of (3.1), we can write

$$(3.3) \quad \|(x_k)\|_{\infty} \leq \begin{cases} P_1 \Lambda_1 + |a_0^1|, & k = 1, \\ P_k \Lambda_k + \sum_{j=0}^{k-2} \frac{|a_j^k|}{j!}, & k = 2, 3, \dots, n. \end{cases}$$

Then, by combining  $(H_2)$  with (3.3), we get

$$(3.4) \quad \|x_k\|_\infty \leq \|t^{\beta_k} D^{\alpha_k} x_k\|_\infty.$$

On the other hand, using  $(H_3)$ , there exists a solution  $(y_1, y_2, \dots, y_n) \in S$  satisfying (3.2). Therefore, by (3.4) we can write:

$$\begin{aligned} \|x_k - y_k\|_\infty &\leq \|t^{\beta_k} D^{\alpha_k} (x_k - y_k)\|_\infty \\ &\leq \|t^{\beta_k} (D^{\alpha_k} x_k - f_k(t, x_1, \dots, x_k)) - t^{\beta_k} (D^{\alpha_k} y_k - f_k(t, y_1, \dots, y_k)) \\ &\quad + t^{\beta_k} (f_k(t, x_1, \dots, x_k) - f_k(t, y_1, \dots, y_k))\|_\infty \\ &\leq \|t^{\beta_k} (D^{\alpha_k} x_k - f_k(t, x_1, \dots, x_k))\|_\infty + \|t^{\beta_k} (D^{\alpha_k} y_k - f_k(t, y_1, \dots, y_k))\|_\infty \\ &\quad + \|t^{\beta_k} (f_k(t, x_1, \dots, x_k) - f_k(t, y_1, \dots, y_k))\|_\infty \\ &\leq \|t^{\beta_k}\|_\infty \| (D^{\alpha_k} x_k - f_k(t, x_1, \dots, x_k)) \|_\infty \\ &\quad + \|t^{\beta_k}\|_\infty \| (D^{\alpha_k} y_k - f_k(t, y_1, \dots, y_k)) \|_\infty \\ &\quad + \|t^{\beta_k} (f_k(t, x_1, \dots, x_k) - f_k(t, y_1, \dots, y_k))\|_\infty \end{aligned}$$

From (2.12), (3.1) and (3.2), we obtain

$$\|(x_k - y_k)\|_\infty \leq \epsilon_k + \sum_{j=1}^k \omega_j^k \max_{1 \leq k \leq n} \|(x_k - y_k)\|_\infty.$$

Then

$$\max_{1 \leq k \leq n} \|(x_k - y_k)\|_\infty \leq \frac{\epsilon}{1 - \sum_{j=1}^k \omega_j^k} := \mu\epsilon, \quad \epsilon = \max_{1 \leq k \leq n} \epsilon_k, \quad \mu = \frac{1}{1 - \sum_{j=1}^k \omega_j^k}.$$

Hence,

$$\|(x_1 - y_1, \dots, x_n - y_n)\|_S \leq \mu\epsilon.$$

Using  $(H_4)$ , we get  $\mu > 0$ . Thus, system (1.1) is Ulam-Hyers stable. Taking  $\phi(\epsilon) = \mu\epsilon$ , we get system (1.1) is generalized Ulam-Hyers stable. This ends the proof.  $\square$

### REFERENCES

- [1] M. A. Abdellaoui, Z. Dahmani and N. BedjaouiI, *New existence results for a coupled system on nonlinear differential equations of arbitrary order*, IJNAA **6**(2) (2015), 65–75.
- [2] R. P. Agarwal and D. O'Regan, *Existence theory for singular initial and boundary value problems: a fixed point approach*, Appl. Anal. **81** (2002), 391–434.
- [3] R. P. Agarwal, D. O'Regan and S. Stanek, *Positive solutions for Dirichlet problems of singular nonlinear fractional differential equations*, J. Math. Anal. Appl. **371** (2010), 57–68.
- [4] C. Bai and J. Fang, *The existence of a positive solution for a singular coupled system of nonlinear fractional differential equations*, Appl. Math. Comput. **150**(3) (2004), 611–621.
- [5] D. Baleanu, S. Z. Nazemi and S. Rezapour, *The existence of positive solutions for a new coupled system of multiterm singular fractional integrodifferential boundary value problems*, Abstr. Appl. Anal. (2013), Article ID 368659, 15 pages.

- [6] Z. Dahmani and A. Taïeb, *Solvability of a coupled system of fractional differential equations with periodic and antiperiodic boundary conditions*, Pure and Applied Mathematics Letters **1** (2015), 29–36.
- [7] Z. Dahmani and A. Taïeb, *New existence and uniqueness results for high dimensional fractional differential systems*, Facta Univ. Ser. Math. Inform. **30**(3) (2015), 281–293.
- [8] Z. Dahmani and A. Taïeb, *Solvability for high dimensional fractional differential systems with high arbitrary orders*, Journal of Advanced Scientific Research in Dynamical and Control Systems **7**(4) (2015), 51–64.
- [9] Z. Dahmani and A. Taïeb, *A coupled system of fractional differential equations involving two fractional orders*, ROMAI J. **11**(2) (2015), 141–177.
- [10] Z. Dahmani, A. Taïeb and N. Bedjaoui, *Solvability and stability for nonlinear fractional integro-differential systems of high fractional orders*, Facta Univ. Ser. Math. Inform. **31**(3) (2016), 629–644.
- [11] M. Fečkan, J. Wang and Y. Zhou, *Ulam’s type stability of impulsive ordinary differential equations*, J. Math. Anal. Appl. **395** (2012), 258–264.
- [12] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, River Edge, New Jersey, 2000.
- [13] R. W. Ibrahim, *Stability of a fractional differential equation*, International Journal of Mathematical, Computational, Physical and Quantum Engineering **7**(3) (2013), 300–305.
- [14] R. W. Ibrahim, *Ulam stability of boundary value problem*, Kragujevac J. Math. **37**(2) (2013), 287–297.
- [15] S. M. Jung and T. M. Rassias, *Generalized Hyers-Ulam stability of Riccati differential equation*, Math. Inequal. Appl. **11** (2008), 777–782.
- [16] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North Holland Mathematics Studies **204**, Elsevier, Amsterdam, 2006.
- [17] R. Li, *Existence of solutions for nonlinear singular fractional differential equations with fractional derivative condition*, Adv. Difference Equ. **2014**(292) (2014), 12 pages.
- [18] Z. Lin, W. Wei and J. R. Wang, *Existence and stability results for impulsive integro-differential equations*, Ser. Math. Inform. **29**(2) (2014), 119–130.
- [19] F. Mainardi, *Fractional calculus: some basic problems in continuum and statistical mechanics*, in: A. Carpinteri and F. Mainardi (Eds.), *Fractals and Fractional Calculus in Continuum Mechanics*, Springer, New York, 1997, 291–348.
- [20] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [21] A. Taïeb and Z. Dahmani, *A coupled system of nonlinear differential equations involving  $m$  nonlinear terms*, Georgian Math. J. **23**(3) (2016), 447–458.
- [22] A. Taïeb and Z. Dahmani, *The high order Lane-Emden fractional differential system: existence, uniqueness and Ulam stabilities*, Kragujevac J. Math. **40**(2) (2016), 238–259.
- [23] A. Taïeb and Z. Dahmani, *A new problem of singular fractional differential equations*, J. Dyn. Syst. Geom. Theor. **14**(2) (2016), 161–183.
- [24] A. Taïeb and Z. Dahmani, *On singular fractional differential systems and Ulam-Hyers stabilities*, International Journal of Modern Mathematical Sciences **14**(3) (2016), 262–282.
- [25] A. Yang and W. Ge, *Positive solutions for boundary value problems of  $n$ -dimension nonlinear fractional differential system*, Bound. Value Probl. (2008), Article ID 437453, 15 pages.

<sup>1</sup>LPAM, FACULTY ST,  
UMAB MOSTAGANEM,  
ALGERIA  
*Email address:* taieb5555@yahoo.com

<sup>2</sup>LPAM, FACULTY SEI,  
UMAB MOSTAGANEM,  
ALGERIA  
*Email address:* zzdahmani@yahoo.fr