

SPECTRA OF THE LOWER TRIANGULAR MATRIX
 $\mathbb{B}(r_1, \dots, r_l; s_1, \dots, s_{l'})$ OVER c_0

SANJAY KUMAR MAHTO^{1,2}, ARNAB PATRA¹, AND P. D. SRIVASTAVA³

ABSTRACT. The infinite lower triangular matrix $\mathbb{B}(r_1, \dots, r_l; s_1, \dots, s_{l'})$ is considered over the sequence space c_0 , where l and l' are positive integers. The diagonal and sub-diagonal entries of the matrix consist of the oscillatory sequences $r = (r_{k(\bmod l)+1})$ and $s = (s_{k(\bmod l')+1})$, respectively. The rest of the entries of the matrix are zero. It is shown that the matrix represents a bounded linear operator. Then the spectrum of the matrix is evaluated and partitioned into its fine structures: point spectrum, continuous spectrum, residual spectrum, etc. In particular, the spectra of the matrix $\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6)$ are determined. Finally, an example is taken in support of the results.

1. INTRODUCTION

The study of the spectrum of a bounded linear operator has received much attention in recent years due to its wide application in functional analysis, classical quantum mechanics, etc. Let A be an infinite matrix that is bounded and linear in a Banach space U . Then many dynamical systems can be reformulated as the system of linear equations $Ax = \lambda x$, where λ is a complex number and x is a nonzero vector in U . The stability of this system can be explained by the spectrum of A . In this course, spectrum localization of an infinite matrix over a sequence space is viewed as an important problem by many authors [10, 14–16, 23, 26]. An extensive study of most of the research done in this direction can be found in the review articles [25] and [17].

Key words and phrases. Fine spectra, sequence space, lower triangular infinite matrix, point spectrum, continuous spectrum, residual spectrum.

2010 *Mathematics Subject Classification.* Primary: 47A10. Secondary: 47B37.

DOI 10.46793/KgJMat2203.369M

Received: September 05, 2019.

Accepted: December 27, 2019.

For a sequence $x = (x_k)$, the backward difference operator Δ is defined by $\Delta x = x_k - x_{k-1}$, where $x_{-1} = 0$. The matrix representation of this operator is as follows:

$$\Delta = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdots \\ -1 & 1 & 0 & 0 & \cdots \\ 0 & -1 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

In short, Δ is an infinite matrix whose diagonal entries and subdiagonal entries are the constant sequences $(1, 1, \dots)$ and $(-1, -1, \dots)$, respectively. Akhmedov and Başar [1] determined the spectral decompositions of this operator over bv_p ($1 \leq p < \infty$), whereas Altay and Başar [3] evaluated the spectra of the same operator over the spaces c and c_0 . Altay and Başar [4] then considered the difference operator $B(r, s)$ over c_0 and c , which is a generalization of the operator Δ . The diagonal and subdiagonal entries of $B(r, s)$ contain the sequences (r, r, \dots) and (s, s, \dots) , where r and $s \neq 0$ are real numbers. Furkan and Bilgiç studied $B(r, s)$ in the same direction over ℓ_p and bv_p in [6]. For more study, we refer [2, 7, 8, 12, 13, 18, 19, 22, 24] etc. Now if one considers the more generalized difference matrix whose diagonal and subdiagonal entries are the oscillatory sequences $(r_1, r_2, \dots, r_l, r_1, \dots)$ and $(s_1, s_2, \dots, s_{l'}, s_1, \dots)$, where l and l' are some positive integers, then the number of limit points of both the sequences will be different and it will be interesting to study the spectral property of the matrix.

In this paper, we have determined the spectra and fine spectra of the generalized difference matrix $\mathbb{B}(r_1, \dots, r_l; s_1, \dots, s_{l'})$ in which the diagonal entries consist of a sequence whose terms are oscillating between the points r_1, r_2, \dots, r_l and the subdiagonal entries consist of an oscillatory sequence whose terms are oscillating between the points $s_1, s_2, \dots, s_{l'}$. Furthermore, the spectra and fine spectra of the matrix $\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6)$ are also discussed.

2. PRELIMINARIES

Let U and V be Banach spaces. Then the space of all bounded linear operators from U into V is denoted by $B(U, V)$. If $U = V$, then the space is denoted by $B(U)$. Let $L \in B(U)$ and U^* be dual of U . Then the adjoint $L^* \in B(U^*)$ of L is defined by $(L^*f)(t) = f(Lt)$ for all $f \in U^*$. Let $J : D(J) \rightarrow U$ be a linear operator defined over a subset $D(J)$ of U . Then the operator $(J - \lambda I)^{-1}$ is called the resolvent operator of J , where λ is a complex number and I is the identity operator.

A complex number λ is said to be a *regular value* [11] of a linear operator $J : D(J) \rightarrow U$ if and only if the operator $(J - \lambda I)^{-1}$ exists, bounded and is defined on a set which is dense in U . The set of all regular values of the linear operator J is called *resolvent set* and is denoted by $\rho(J)$. The complement $\sigma(J) = \mathbb{C} - \rho(J)$ is called the *spectrum* of J . The spectrum $\sigma(J)$ is further partitioned into the following three disjoint sets.

- (a) $\sigma_p(J) = \{\lambda \in \mathbb{C} : (J - \lambda I)^{-1} \text{ does not exist}\}$. This set is called the *point spectrum (discrete spectrum)* of the operator J . The members of this set are called *eigenvalues* of J .
- (b) $\sigma_c(T)$, which is defined as the set of all complex numbers λ for which $(J - \lambda I)^{-1}$ exists and defined on a set which is dense in U , but it is not a bounded operator in U . This set is called *continuous spectrum* of J .
- (c) $\sigma_r(T)$, which contains all those complex numbers for which $(J - \lambda I)^{-1}$ exists, defined on a set which is not dense in U . This set is called the *residual spectrum* of J .

Let $R(J - \lambda I)$ denotes the range of the operator $J - \lambda I$. Goldberg [9] has classified the spectrum using the following six properties of $R(J - \lambda I)$ and $(J - \lambda I)^{-1}$:

- (I) $R(J - \lambda I) = U$;
- (II) $R(J - \lambda I) \neq U$ but $\overline{R(J - \lambda I)} = U$;
- (III) $\overline{R(J - \lambda I)} \neq U$

and

- (1) $(J - \lambda I)^{-1}$ exists and it is bounded;
- (2) $(J - \lambda I)^{-1}$ exists but it is not bounded;
- (3) $(J - \lambda I)^{-1}$ does not exist.

Based on the above six properties, the Goldberg’s classification of the spectrum can be given as shown in the Table 1.

TABLE 1. Subdivisions of spectrum of a bounded linear operator

| | (I) | (II) | (III) |
|---|------------------|------------------|------------------|
| 1 | $\rho(J, U)$ | — | $\sigma_r(J, U)$ |
| 2 | $\sigma_c(J, U)$ | $\sigma_c(J, U)$ | $\sigma_r(J, U)$ |
| 3 | $\sigma_p(J, U)$ | $\sigma_p(J, U)$ | $\sigma_p(J, U)$ |

Theorem 2.1 ([21]). *Let L be a bounded linear operator on a normed linear space U . Then L has a bounded inverse if and only if L^* is onto.*

Lemma 2.1 ([20]). *An infinite matrix $A = (a_{nk}) \in B(c_0)$ if and only if*

- (a) $(a_{nk})_k \in \ell_1$ for all n and $\sup_n \sum_k |a_{nk}| < \infty$;
- (b) $(a_{nk})_n \in c_0$ for all k .

Moreover, the norm $\|A\| = \sup_n \sum_k |a_{nk}|$.

Throughout the paper, we denote the set of natural numbers by \mathbb{N} , the set of complex numbers by \mathbb{C} and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. We assume that $x_{-n} = 0$ for all $n \in \mathbb{N}$.

3. MAIN RESULTS

Let l and l' be two natural numbers. Suppose that H is the least common multiple of l and l' . Let $r_i, i = 1, \dots, l$, and $s_i \neq 0, i = 1, \dots, l'$, be complex numbers. Then

$$(3.4) \quad \begin{cases} n - k = mH + \zeta, \\ \zeta = m'l + \zeta', \\ \zeta = m''l' + \zeta'', \end{cases}$$

for some non-negative integers m, m' and m'' .

Lemma 3.1. *If $(|\lambda - r_1| \cdots |\lambda - r_l|)^{1/l} > (|s_1| \cdots |s_{l'}|)^{1/l'}$, then $(\mathbb{B} - \lambda I)^{-1} \in B(c_0)$.*

Proof. Since $(|\lambda - r_1| \cdots |\lambda - r_l|)^{1/l} > (|s_1| \cdots |s_{l'}|)^{1/l'}$ and $s_1, s_2, \dots, s_{l'}$ are non-zero, therefore $\lambda \neq r_i$ for all $i = 1, 2, \dots, l$. Then the matrix $\mathbb{B} - \lambda I$ is a triangle and hence $(\mathbb{B} - \lambda I)^{-1} = (z_{nk})$ exists, which is given by (3.3). We first consider a row of $(\mathbb{B} - \lambda I)^{-1}$ which is a multiple of H , that is $n = \tilde{m}H$ for some $\tilde{m} \in \mathbb{N}_0$. Now, let $k = \hat{m}H$ for $\hat{m} = 0, 1, \dots, \tilde{m}$. Then (3.4) implies that $n - k = (\tilde{m} - \hat{m})H$ and $m' = m'' = \zeta = \zeta' = \zeta'' = 0$. Thus, from (3.3), we have

$$z_{nk} = \frac{(-1)^{n-k}}{r_{\alpha(k)} - \lambda} \left\{ \frac{(s_1 \cdots s_{l'})^{\frac{H}{l'}}}{\{(r_1 - \lambda) \cdots (r_l - \lambda)\}^{\frac{H}{l}}} \right\}^{\tilde{m} - \hat{m}},$$

for all $\hat{m} = 0, 1, \dots, \tilde{m}$. Therefore,

$$\sum_{k \in [0_H]} |z_{nk}| = \frac{1}{|r_{\alpha(k)} - \lambda|} \sum_{j=0}^{\tilde{m}} \left\{ \frac{(|s_1| \cdots |s_{l'}|)^{\frac{H}{l'}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{\frac{H}{l}}} \right\}^j,$$

where $[0_H]$ denotes the set of all non-negative integers which are multiple of H . For the same row, if we consider $k = \hat{m}H + 1$ for $\hat{m} = 0, 1, \dots, \tilde{m} - 1$, then $n - k = (\tilde{m} - \hat{m} - 1)H + H - 1$. Let m'_1 and m''_1 be quotients and ζ'_1 and ζ''_1 be remainders when $H - 1$ is divided by l and l' respectively, that is

$$\begin{aligned} H - 1 &= m'_1 l + \zeta'_1, \\ H - 1 &= m''_1 l' + \zeta''_1. \end{aligned}$$

Then, from (3.3), we obtain that

$$\begin{aligned} z_{nk} &= \frac{(-1)^{n-k} s_{\beta(k)} \cdots s_{\beta(k+\zeta''_1-1)}}{(r_{\alpha(k)} - \lambda) \cdots (r_{\alpha(k+\zeta'_1)} - \lambda)} \cdot \frac{(s_1 \cdots s_{l'})^{m''_1}}{\{(r_1 - \lambda) \cdots (r_l - \lambda)\}^{m'_1}} \\ &\quad \times \left\{ \frac{(s_1 \cdots s_{l'})^{\frac{H}{l'}}}{\{(r_1 - \lambda) \cdots (r_l - \lambda)\}^{\frac{H}{l}}} \right\}^{\tilde{m} - \hat{m} - 1}, \end{aligned}$$

for all $\hat{m} = 0, 1, \dots, \tilde{m} - 1$. Hence,

$$\begin{aligned} \sum_{k \in [1_H]} |z_{nk}| &= \frac{|s_{\beta(k)}| \cdots |s_{\beta(k+\zeta''_1-1)}|}{|r_{\alpha(k)} - \lambda| \cdots |r_{\alpha(k+\zeta'_1)} - \lambda|} \cdot \frac{(|s_1| \cdots |s_{l'}|)^{m''_1}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{m'_1}} \\ &\quad \times \sum_{j=0}^{\tilde{m}-1} \left\{ \frac{(|s_1| \cdots |s_{l'}|)^{\frac{H}{l'}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{\frac{H}{l}}} \right\}^j, \end{aligned}$$

where $[1_H]$ denotes the set of all nonnegative integers x such that H divides $x - 1$. Similarly, for $k = \hat{m}H + 2, \dots, \hat{m}H + H - 1$, we have

$$\begin{aligned} \sum_{k \in [2_L]} |z_{nk}| &= \frac{|s_{\beta(k)}| \cdots |s_{\beta(k+\zeta''_2-1)}|}{|r_{\alpha(k)} - \lambda| \cdots |r_{\alpha(k+\zeta'_2)} - \lambda|} \cdot \frac{(|s_1| \cdots |s_{l'}|)^{m'_2}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{m'_2}} \\ &\quad \times \sum_{j=0}^{\tilde{m}-1} \left\{ \frac{(|s_1| \cdots |s_{l'}|)^{\frac{H}{l'}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{\frac{H}{l'}}} \right\}^j, \\ &\quad \vdots \\ \sum_{k \in [(H-1)_L]} |z_{nk}| &= \frac{|s_{\beta(k)}| \cdots |s_{\beta(k+\zeta''_{H-1}-1)}|}{|r_{\alpha(k)} - \lambda| \cdots |r_{\alpha(k+\zeta'_{H-1})} - \lambda|} \cdot \frac{(|s_1| \cdots |s_{l'}|)^{m''_{H-1}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{m'_{H-1}}} \\ &\quad \times \sum_{j=0}^{\tilde{m}-1} \left\{ \frac{(|s_1| \cdots |s_{l'}|)^{\frac{H}{l'}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{\frac{H}{l'}}} \right\}^j, \end{aligned}$$

for some integers $\zeta'_i, \zeta''_i, m'_i$ and m''_i for all $i \in \{2, 3, \dots, H - 1\}$. Thus,

$$\begin{aligned} \sum_{k=0}^{\infty} |z_{nk}| &= \frac{1}{|r_{\alpha(k)} - \lambda|} \sum_{j=0}^{\tilde{m}} \left\{ \frac{(|s_1| \cdots |s_{l'}|)^{\frac{H}{l'}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{\frac{H}{l'}}} \right\}^j \\ (3.5) \quad &\quad + M \sum_{j=0}^{\tilde{m}-1} \left\{ \frac{(|s_1| \cdots |s_{l'}|)^{\frac{H}{l'}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{\frac{H}{l'}}} \right\}^j, \end{aligned}$$

where

$$\begin{aligned} M &= \frac{|s_{\beta(k)}| \cdots |s_{\beta(k+\zeta''_1-1)}|}{|r_{\alpha(k)} - \lambda| \cdots |r_{\alpha(k+\zeta'_1)} - \lambda|} \cdot \frac{(|s_1| \cdots |s_{l'}|)^{m'_1}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{m'_1}} \\ &\quad + \cdots + \frac{|s_{\beta(k)}| \cdots |s_{\beta(k+\zeta''_{L-1}-1)}|}{|r_{\alpha(k)} - \lambda| \cdots |r_{\alpha(k+\zeta'_{L-1})} - \lambda|} \cdot \frac{(|s_1| \cdots |s_{l'}|)^{m''_{H-1}}}{\{|r_1 - \lambda| \cdots |r_l - \lambda|\}^{m'_{H-1}}}. \end{aligned}$$

Let $M_0 = \max \left\{ \frac{1}{|r_{\alpha(k)} - \lambda|}, M \right\}$. Then

$$\sum_{k=0}^{\infty} |z_{nk}| \leq \frac{2M_0(|r_1 - \lambda||r_2 - \lambda| \cdots |r_l - \lambda|)^{\frac{H}{l'}}}{(|r_1 - \lambda||r_2 - \lambda| \cdots |r_l - \lambda|)^{\frac{H}{l'}} - (|s_1||s_2| \cdots |s_{l'}|)^{\frac{H}{l'}}}.$$

Therefore, $\sup_{n \in [0_H]} \sum_{k=0}^{\infty} |z_{nk}| < \infty$. Similarly, we prove that

$$\sup_{n \in [1_H]} \sum_{k=0}^{\infty} |z_{nk}| < \infty,$$

$$\sup_{n \in [2_H]} \sum_{k=0}^{\infty} |z_{nk}| < \infty,$$

⋮

$$\sup_{n \in [(H-1)_H]} \sum_{k=0}^{\infty} |z_{nk}| < \infty.$$

Thus,

$$\sup_n \sum_{k=0}^{\infty} |z_{nk}| = \max \left\{ \sup_{n \in [0_H]} \sum_{k=0}^{\infty} |z_{nk}|, \sup_{n \in [1_H]} \sum_{k=0}^{\infty} |z_{nk}|, \dots, \sup_{n \in [(H-1)_H]} \sum_{k=0}^{\infty} |z_{nk}| \right\}.$$

This implies that $\sup_n \sum_{k=0}^{\infty} |z_{nk}| < \infty$. Likewise, for an arbitrary column of $(\mathbb{B} - \lambda I)^{-1}$, adding the entries separately whose rows n belong to $[0_H], [1_H], \dots, [(H-1)_H]$ respectively, we get $\sum_{n=0}^{\infty} |z_{nk}| < \infty$. Therefore, $\lim_{n \rightarrow \infty} |z_{nk}| = 0$ for all $k \in \mathbb{N}_0$. Hence, by Lemma 2.1, the matrix $(\mathbb{B} - \lambda I)^{-1} \in B(c_0)$. \square

Consider the set $S = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} \leq (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}} \right\}$. Then we have the following theorem.

Theorem 3.2. $\sigma(\mathbb{B}, c_0) = S$.

Proof. First, we prove that $\sigma(\mathbb{B}, c_0) \subseteq S$. Let λ be a complex number that does not belong to S . Then $(|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} > (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}}$. In that case, from Lemma 3.1, it follows that $(\mathbb{B} - \lambda I)^{-1} \notin B(c_0)$. That is, $\lambda \notin \sigma(\mathbb{B}, c_0)$. Hence, $\sigma(\mathbb{B}, c_0) \subseteq S$.

Next, we show that $S \subseteq \sigma(\mathbb{B}, c_0)$. Let $\lambda \in S$. Then, $(|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} \leq (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}}$. If λ equals any of the r_i for all $i \in \{1, 2, \dots, l\}$, then the range of the operator $\mathbb{B} - \lambda I$ is not dense in c_0 , and hence $\lambda \in \sigma(\mathbb{B}, c_0)$. Therefore, we assume that $\lambda \neq r_i$ for all $i \in \{1, 2, \dots, l\}$. In that case, $\mathbb{B} - \lambda I$ is a triangle and $(\mathbb{B} - \lambda I)^{-1} = (z_{nk})$ exists, which is given by (3.3). Let $y = (1, 0, 0, \dots) \in c_0$ and let $x = (x_k)$ be the sequence such that $(\mathbb{B} - \lambda I)^{-1}y = x$. It follows, from (3.3), that

$$(3.6) \quad x_{nH} = z_{nH,0} = \frac{(-1)^{nH}}{r_1 - \lambda} \left\{ \frac{(s_1 s_2 \cdots s_{l'})^{\frac{H}{l'}}}{\{(r_1 - \lambda) \cdots (r_l - \lambda)\}^{\frac{H}{l}}} \right\}^n,$$

for all $n \in \mathbb{N}_0$. Since $\{(r_1 - \lambda) \cdots (r_l - \lambda)\}^{\frac{1}{l}} \leq (s_1 \cdots s_{l'})^{\frac{1}{l'}}$, the subsequence (x_{nH}) of x does not converge to 0. Consequently, the sequence $x = (x_k) \notin c_0$. Therefore, $(\mathbb{B} - \lambda I)^{-1} \notin B(c_0)$. Thus, $\lambda \in \sigma(\mathbb{B}, c_0)$ and hence $S \subseteq \sigma(\mathbb{B}, c_0)$. This proves the theorem. \square

Theorem 3.3. $\sigma_p(\mathbb{B}, c_0) = \emptyset$.

Proof. Let $\lambda \in \sigma_p(\mathbb{B}, c_0)$. Then there exists a nonzero sequence $x = (x_k)$ such that $\mathbb{B}x = \lambda x$. This implies that

$$(3.7) \quad s_{(k-1)(\text{mod } l')+1} x_{k-1} + r_{k(\text{mod } l)+1} x_k = \lambda x_k.$$

Let x_{k_0} be the first non-zero term of the sequence $x = (x_k)$. Then from the relation (3.7), we find that $\lambda = r_{k_0(\text{mod } l)+1}$. Next, for $k = k_0 + l$, (3.7) becomes

$$s_{(k_0+l-1)(\text{mod } l')+1} x_{k_0+l-1} + r_{(k_0+l)(\text{mod } l)+1} x_{k_0+l} = \lambda x_{k_0+l}.$$

$$\times \left\{ \frac{((\lambda - r_1) \cdots (\lambda - r_l))^{\frac{H}{l}}}{(s_1 \cdots s_{l'})^{\frac{H}{l'}}} \right\}^k x_0.$$

Thus,

$$\begin{aligned} \sum_{k=0}^{\infty} |x_k| &= \sum_{k=0}^{\infty} |x_{kH}| + \sum_{k=0}^{\infty} |x_{kH+1}| + \cdots + \sum_{n=0}^{\infty} |x_{kH+H-1}| \\ &= \left(1 + \left| \frac{\lambda - r_1}{s_1} \right| + \cdots + \left| \frac{(\lambda - r_1)^{\frac{H}{l}} \cdots (\lambda - r_{l-1})^{\frac{H}{l}} (\lambda - r_l)^{\frac{H}{l}-1}}{s_1^{\frac{H}{l'}} \cdots s_{l'-1}^{\frac{H}{l'}} s_{l'}^{\frac{H}{l}-1}} \right| \right) \\ &\quad \times \sum_{k=0}^{\infty} \left| \frac{((\lambda - r_1) \cdots (\lambda - r_l))^{\frac{H}{l}}}{(s_1 \cdots s_{l'})^{\frac{H}{l'}}} \right|^k |x_0|. \end{aligned}$$

Clearly, the sequence $x = (x_k) \in \ell_1$ if and only if $(|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} < (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}}$. This proves the theorem. \square

Theorem 3.5. $\sigma_r(\mathbb{B}, c_0) = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} < (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}} \right\}$.

Proof. The residual spectrum of a bounded linear operator L on a Banach space U is given by the relation $\sigma_r(L, U) = \sigma_p(L^*, U^*) \setminus \sigma_p(L, U)$. Therefore, $\sigma_r(\mathbb{B}, c_0) = \sigma_p(\mathbb{B}^*, c_0^*) \setminus \sigma_p(\mathbb{B}, c_0)$. Then the proof of this theorem is an easy consequence of the Theorems 3.3 and 3.4. \square

Theorem 3.6. $\sigma_c(\mathbb{B}, c_0) = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} = (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}} \right\}$.

Proof. Since spectrum of an operator on a Banach space is disjoint union of point, residual and continuous spectrum, therefore from Theorems 3.2, 3.3 and 3.5, we deduce that

$$\sigma_c(\mathbb{B}, c_0) = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| \cdots |\lambda - r_l|)^{\frac{1}{l}} = (|s_1| \cdots |s_{l'}|)^{\frac{1}{l'}} \right\}. \quad \square$$

Theorem 3.7. $\{r_1, r_2, \dots, r_l\} \subseteq III_1(\mathbb{B}, c_0)$.

Proof. Theorem 3.5 shows that $r_1 \in \sigma_r(\mathbb{B}, c_0)$. However, $\sigma_r(\mathbb{B}, c_0) = III_1(\mathbb{B}, c_0) \cup III_2(\mathbb{B}, c_0)$. Therefore, to prove $r_1 \in III_1\sigma(\mathbb{B}, c_0)$, we shall show that the matrix $\mathbb{B} - r_1 I$ has bounded inverse and from Theorem 2.1, it will be sufficient to show that $(\mathbb{B} - r_1 I)^*$ is onto. For this, let $y = (y_k) \in \ell_1$. Then $(\mathbb{B} - r_1 I)^* x = y$ implies that

$$(3.10) \quad (r_{i(\text{mod } l)+1} - r_1)x_i + s_{i(\text{mod } l')+1}x_{i+1} = y_i,$$

for all $i \in \mathbb{N}_0$. Solving (3.10) for $x = (x_i)$, we obtain that

$$(3.11) \quad x_{mH+k} = \sum_{j=0}^{k-2} \frac{1}{s_{j(\text{mod } l')+1}} \prod_{i=j+1}^{k-1} \frac{r_1 - r_{i(\text{mod } l)+1}}{s_{i(\text{mod } l')+1}} y_{mH+j} + \frac{y_{mH+k-1}}{s_{(k-1)(\text{mod } l')+1}},$$

for $k = 1, \dots, H$, and $m = 0, \dots, \infty$. Let

$$C_j = \frac{1}{s_{j(\text{mod } l')+1}} \prod_{i=j+1}^{k-1} \frac{r_1 - r_{i(\text{mod } l)+1}}{s_{i(\text{mod } l')+1}},$$

for $j = 0, \dots, k-2$, and

$$C_{k-1} = \frac{1}{s^{(k-1)(\text{mod } l)+1}}.$$

Then (3.11) can be written as

$$(3.12) \quad x_{mH+k} = C_0 y_{mH} + C_1 y_{mH+1} + \dots + C_{k-1} y_{mH+k-1}.$$

Taking summation from $m = 0$ to ∞ of the absolute values of x_{mH+k} , we obtain

$$(3.13) \quad \sum_{m=0}^{\infty} |x_{mH+k}| \leq |C_0| \sum_{m=0}^{\infty} |y_{mH}| + |C_1| \sum_{m=0}^{\infty} |y_{mH+1}| + \dots + |C_{k-1}| \sum_{m=0}^{\infty} |y_{mH+k-1}|.$$

Since $y = (y_k) \in \ell_1$, therefore the right hand side of the inequality (3.13) is a sum of k finite terms. Thus, $\sum_{m=0}^{\infty} |x_{mH+k}| < \infty$ for $k \in \{1, 2, \dots, H\}$. This implies that the series

$$(3.14) \quad \sum_i |x_i| = |x_0| + \sum_{m=0}^{\infty} |x_{mH+1}| + \sum_{m=0}^{\infty} |x_{mH+2}| + \dots + \sum_{m=0}^{\infty} |x_{mH+H}|$$

is a sum of $H+1$ finite terms. Hence, $x = (x_i) \in \ell_1$. We have shown that for every $y = (y_i) \in \ell_1$ there exists a sequence $x = (x_i) \in \ell_1$ such that $(\mathbb{B} - r_1 I)^* x = y$. That is, $(\mathbb{B} - r_1 I)^*$ is onto. Similarly, we can show that $r_i \in III_1(\mathbb{B}, c_0)$ for $i = 2, \dots, l$. This proves the theorem. \square

Theorem 3.8. $\sigma_r(\mathbb{B}, c_0) \setminus \{r_1, r_2, \dots, r_l\} \subseteq III_2(\mathbb{B}, c_0)$.

Proof. Let λ belongs to the set $\sigma_r(\mathbb{B}, c_0) \setminus \{r_1, r_2, \dots, r_l\}$. Then $(|\lambda - r_1| \dots |\lambda - r_l|)^{\frac{1}{l}} < (|s_1| \dots |s_l|)^{\frac{1}{l}}$ and $\lambda \notin r_i$ for all $i \in \{1, 2, \dots, l\}$. This inequality shows that the series $\sum_{k=0}^{\infty} |z_{nk}|$ in (3.5) is not convergent when n goes to infinity. In that case, $\mathbb{B} - \lambda I$ does not have bounded inverse. Then from Table 1, we find that $\lambda \in III_2(\mathbb{B}, c_0)$. Hence $\sigma_r(\mathbb{B}, c_0) \setminus \{r_1, r_2, \dots, r_l\} \subseteq III_2(\mathbb{B}, c_0)$. \square

Theorem 3.9. $III_1(\mathbb{B}, c_0) = \{r_1, r_2, \dots, r_l\}$.

Proof. From Table 1, we have $\sigma_r(\mathbb{B}, c_0) = III_1(\mathbb{B}, c_0) \cup III_2(\mathbb{B}, c_0)$ and the union is disjoint. Then taking complement of the inclusion of Theorem 3.8 in $\sigma_r(\mathbb{B}, c_0)$, we obtain that $\sigma_r(\mathbb{B}, c_0) \setminus III_2(\mathbb{B}, c_0) \subseteq \{r_1, r_2, \dots, r_l\}$. That is, $III_1(\mathbb{B}, c_0) \subseteq \{r_1, r_2, \dots, r_l\}$. This inclusion together with Theorem 3.7 implies that $III_1(\mathbb{B}, c_0) = \{r_1, r_2, \dots, r_l\}$. \square

Theorem 3.10. $III_2(\mathbb{B}, c_0) = \sigma_r(\mathbb{B}, c_0) \setminus \{r_1, r_2, \dots, r_l\}$.

Proof. Taking complement of the result of Theorem 3.9 in $\sigma_r(\mathbb{B}, c_0)$, we obtain that $III_2(\mathbb{B}, c_0) = \sigma_r(\mathbb{B}, c_0) \setminus \{r_1, r_2, \dots, r_l\}$. \square

4. FINE SPECTRA OF THE MATRIX $\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6)$

We consider the matrix

$$\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6) = \begin{bmatrix} r_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ s_1 & r_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & s_2 & r_3 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & s_3 & r_4 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & s_4 & r_1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & s_5 & r_2 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & s_6 & r_3 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & s_1 & r_4 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_2 & r_1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots \end{bmatrix}.$$

Now, consider the following sets:

$$D = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| |\lambda - r_2| |\lambda - r_3| |\lambda - r_4|)^{\frac{1}{4}} \leq (|s_1| |s_2| |s_3| |s_4| |s_5| |s_6|)^{\frac{1}{6}} \right\},$$

$$D_1 = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| |\lambda - r_2| |\lambda - r_3| |\lambda - r_4|)^{\frac{1}{4}} < (|s_1| |s_2| |s_3| |s_4| |s_5| |s_6|)^{\frac{1}{6}} \right\},$$

$$D_2 = \left\{ \lambda \in \mathbb{C} : (|\lambda - r_1| |\lambda - r_2| |\lambda - r_3| |\lambda - r_4|)^{\frac{1}{4}} = (|s_1| |s_2| |s_3| |s_4| |s_5| |s_6|)^{\frac{1}{6}} \right\}.$$

From the discussion of the previous section, we deduce the following results:

- (a) $\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6) \in B(c_0)$;
- (b) $\|\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6)\|_{c_0} \leq \max_{i,j} \{|r_i| + |s_j|\} : 1 \leq i \leq 4, 1 \leq j \leq 6$;
- (c) $\sigma(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = D$;
- (d) $\sigma_p(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = \emptyset$;
- (e) $\sigma_p(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6)^*, c_0^* \cong \ell_1) = D_1$;
- (f) $\sigma_r(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = D_1$;
- (g) $\sigma_c(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = D_2$;
- (h) $III_1\sigma(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = \{r_1, r_2, r_3, r_4\}$;
- (i) $III_2\sigma(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = D_1 \setminus \{r_1, r_2, r_3, r_4\}$.

In particular, if we take $r_1 = 1 - i$, $r_2 = -i$, $r_3 = -1.5$, $r_4 = -i$ and $s_1 = i$, $s_2 = 1 + i$, $s_3 = -2$, $s_4 = -1.5$, $s_5 = 1 - i$, $s_6 = -1$, then the spectrum is given by

$$\sigma(\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6), c_0) = \left\{ \lambda \in \mathbb{C} : (|\lambda - 1 + i| |\lambda + i|^2 |\lambda + 1.5|)^{\frac{1}{4}} \leq 6^{\frac{1}{6}} \right\},$$

which is shown by the shaded region in Figure 1.

5. CONCLUSIONS

We have studied the spectral decomposition of the matrix $\mathbb{B}(r_1, \dots, r_l; s_1, \dots, s_{l'})$, which generalizes the following matrices.

- The backward difference operator Δ [3] for $l = 1$, $l' = 1$, $r_1 = 1$ and $s_1 = -1$.
- The Right shift operator for $l = 1$, $l' = 1$, $r_1 = 0$ and $s_1 = 1$.




FIGURE 1. Spectrum of $\mathbb{B}(r_1, \dots, r_4; s_1, \dots, s_6)$.

- The Zweier matrix [5] for $l = 1$, $l' = 1$, $r_1 = s$ and $s_1 = 1 - s$ for some complex numbers $s \neq 0, 1$.
- The generalized difference operator $B(r, s)$ [4] for $l = 1$, $l' = 1$, $r_1 = r$ and $s_1 = s$ for some complex numbers r and $s \neq 0$.

Acknowledgements. The authors gratefully acknowledge the reviewers for their helpful comments.

REFERENCES

- [1] A. M. Akhmedov and F. Başar, *The fine spectra of the difference operator Δ over the sequence space bv_p ($1 \leq p < \infty$)*, Acta Math. Sin. (Engl. Ser.) **23** (2007), 1757–1768.
- [2] A. M. Akhmedov and S. R. El-Shabrawy, *Spectra and fine spectra of lower triangular double-band matrices as operators on ℓ_p ($1 \leq p < \infty$)*, Math. Slovaca **65** (2015), 1137–1152.
- [3] B. Altay and F. Başar, *On the fine spectrum of the difference operator Δ on c_0 and c* , Inform. Sci. **168** (2004), 217–224.
- [4] B. Altay and F. Başar, *On the fine spectrum of the generalized difference operator $B(r, s)$ over the sequence spaces c_0 and c* , Int. J. Math. Math. Sci. **2005** (2005), 3005–3013.
- [5] B. Altay and M. Karakuş, *On the spectrum and the fine spectrum of the zweier matrix as an operator on some sequence spaces*, Thai J. Math. **3** (2005), 153–162.
- [6] H. Bilgiç and H. Furkan, *On the fine spectrum of the generalized difference operator $B(r, s)$ over the sequence spaces ℓ_p and bv_p ($1 < p < \infty$)*, Nonlinear Anal. **68** (2008), 499–506.
- [7] R. Birbonshi and P. D. Srivastava, *On some study of the fine spectra of n -th band triangular matrices*, Complex Anal. Oper. Theory **11** (2017), 739–753.
- [8] E. Dündar and F. Başar, *On the fine spectrum of the upper triangle double band matrix Δ^+ on the sequence space c_0* , Math. Commun. **18** (2013), 337–348.

- [9] S. Goldberg, *Unbounded linear operators: Theory and applications*, Courier Corporation, Mineola, New York, 2006.
- [10] M. Gonzalez, *The fine spectrum of the Cesàro operator in ℓ_p ($1 < p < \infty$)*, Arch. Math. (Basel) **44** (1985), 355–358.
- [11] E. Kreyszig, *Introductory functional analysis with applications*, Wiley, New York, 1989.
- [12] H. S. Özarslan and E. Yavuz, *A new note on absolute matrix summability*, J. Inequal. Appl. **2013** (2013), Paper ID 474, 7 pages.
- [13] A. Patra, R. Birbonshi and P. D. Srivastava, *On some study of the fine spectra of triangular band matrices*, Complex Anal. Oper. Theory **13** (2019), 615–635.
- [14] J. B. Reade, *On the spectrum of the Cesàro operator*, Bull. Lond. Math. Soc. **17** (1985), 263–267.
- [15] B. E. Rhoades, *The fine spectra for weighted mean operators*, Pacific J. Math. **104** (1983), 219–230.
- [16] B. E. Rhoades, *The fine spectra for weighted mean operators in $B(\ell^p)$* , Integral Equations Operator Theory **12** (1989), 82–98.
- [17] P. D. Srivastava, *Spectrum and fine spectrum of the generalized difference operators acting on some Banach sequence spaces - A review*, Ganita **67** (2017), 57–68.
- [18] P. D. Srivastava and S. Kumar, *Fine spectrum of the generalized difference operator Δ_v on sequence space ℓ_1* , Thai J. Math. **8** (2010), 221–233.
- [19] P. D. Srivastava and S. Kumar, *Fine spectrum of the generalized difference operator Δ_{uv} on sequence space ℓ_1* , Appl. Math. Comput. **218** (2012), 6407–6414.
- [20] M. Stieglitz and H. Tietz, *Matrixtransformationen von folgenräumen eine ergebnisübersicht*, Math. Z. **154** (1977), 1–16.
- [21] A. E. Taylor and C. J. A. Halberg Jr, *General theorems about a bounded linear operator and its conjugate*, J. Reine Angew. Math. **198** (1957), 93–111.
- [22] B. C. Tripathy and P. Saikia, *On the spectrum of the Cesàro operator C_1 on $\overline{bv}_0 \cap \ell_\infty$* , Math. Slovaca **63** (2013), 563–572.
- [23] R. B. Wenger, *The fine spectra of the Hölder summability operators*, Indian J. Pure Appl. Math. **6** (1975), 695–712.
- [24] M. Yeşilkayagil and F. Başar, *On the fine spectrum of the operator defined by the lambda matrix over the spaces of null and convergent sequences*, Abstr. Appl. Anal. **2013** (2013), Article ID 687393, 13 pages.
- [25] M. Yeşilkayagil and F. Başar, *A survey for the spectrum of triangles over sequence spaces*, Numer. Funct. Anal. Optim. (2019) (accepted).
- [26] M. Yildirim, *On the spectrum and fine spectrum of the compact Rhalay operators*, Indian J. Pure Appl. Math. **27** (1996), 779–784.

¹DEPARTMENT OF MATHEMATICS,
 INDIAN INSTITUTE OF TECHNOLOGY KHARAGPUR,
 WEST BENGAL 721302 INDIA
 Email address: skmahto0777@gmail.com
 Email address: arnptr1991@gmail.com

²DEPARTMENT OF MATHEMATICS,
 R. N. A. R. COLLEGE SAMASTIPUR,
 BIHAR 848101 INDIA

³DEPARTMENT OF MATHEMATICS,
 INDIAN INSTITUTE OF TECHNOLOGY BHILAI,
 CHATTISGARH 492015 INDIA
 Email address: pds@maths.iitkgp.ernet.in