

## EXISTENCE AND STABILITY OF NONLOCAL INITIAL VALUE PROBLEMS INVOLVING GENERALIZED KATUGAMPOLA DERIVATIVE

ARIF S. BAGWAN<sup>1</sup> AND DEEPAK B. PACHPATTE<sup>2</sup>

**ABSTRACT.** In this paper, the existence results for the solutions to nonlocal initial value problems involving generalized Katugampola derivative are established. Some fixed point theorem techniques are used to derive the existence results. In the sequel, we investigate the generalized Ulam-Hyers-Rassias stability corresponding to our problem. Some examples are given to illustrate our main results.

### 1. INTRODUCTION

In recent decades, the theory of continuous fractional calculus and their applications have remains a centre of attraction in many mathematical research. Indeed, fractional differential equations (FDEs) have grabbed desired attention by many authors. One can see [1–5, 7–13, 20, 21, 23, 26, 27, 33, 34] and references therein. Several definitions of fractional derivatives and integrals have been introduced during the theoretical development of fractional calculus. See [1, 2, 5, 7, 8, 16, 20–22, 25, 27] and references therein.

Initially, Hilfer et al. [16, 17] have proposed linear differential equations involving new fractional operator. They applied operational method to solve such FDEs. Further, Furati et al. [14, 15] investigated non-linear problems and discussed existence and non-existence results for FDEs with Hilfer derivative operator. Benchohra et al. [6, 7], U. N. Katugampola [20, 21], D. B. Dhaigude et al. [8, 9], Kou et al. [23], J. Wang et al. [32, 33] and many more authors, see [1, 2, 5, 19, 29, 31] and references therein, have established the existence results for FDEs with several fractional derivative operators.

---

*Key words and phrases.* Generalized Katugampola derivative, nonlocal initial value problem, Existence, Ulam-Hyers-Rassias stability.

2010 *Mathematics Subject Classification.* Primary: 26A33, 34A08. Secondary: 34A12.

DOI 10.46793/KgJMat2203.443B

*Received:* November 28, 2018.

*Accepted:* January 08, 2020.

Recently, D. S. Oliveira et al. [27] in their article proposed a new fractional differential operator: Hilfer-Katugampola fractional derivative (also known as generalized Katugampola derivative). Further, they established the existence and uniqueness results for the FDEs with generalized Katugampola derivative.

The theory of Ulam stability is also evolved as one of the most interesting field of research. Initially, Ulam [30] established the results on the stability of functional equations. Thereafter, remarkable interest have been shown by authors towards the study of Ulam-Hyers stability and Ulam-Hyers-Rassias stability for various FDEs, see [1, 6, 7, 18, 24, 31, 33] and references therein.

In this paper, we studied the existence and stability of nonlocal initial value problem (IVP) involving generalized Katugampola derivative of the form:

$$(1.1) \quad {}^\rho D_{a^+}^{\mu, \nu} u(t) = f(t, u(t)), \quad \mu \in (0, 1), \nu \in [0, 1], t \in (a, b],$$

$$(1.2) \quad {}^\rho I_{a^+}^{1-\beta} u(a) = \sum_{i=1}^m \lambda_i u(\kappa_i), \quad \mu \leq \beta = \mu + \nu(1 - \mu) < 1, \kappa_i \in (a, b],$$

where  $f$  is a given function such that  $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $0 < \rho$ . The operator  ${}^\rho D_{a^+}^{\mu, \nu}$  is the generalized Katugampola fractional derivative of order  $\mu$  and type  $\nu$  and the operator  ${}^\rho I_{a^+}^{1-\beta} u(a)$  is the Katugampola fractional integral of order  $1 - \beta$ , with  $a > 0$ ,  $\kappa_i, i = 1, 2, \dots, m$ , are prefixed points satisfying  $a < \kappa_1 \leq \kappa_2 \leq \dots \leq \kappa_m < b$ .

Furthermore, the paper is arranged as follows. In Section 2, we recall some basic definitions, important results and preliminary facts. We establish the equivalent mixed type Volterra integral equation for the IVP (1.1)–(1.2). In Section 3, we present existence of solution using the Krasnoselskii fixed point theorem. Further, we present the generalized Ulam-Hyers-Rassias stability to our problem. An illustrative example is given at the end of the main results.

## 2. PRELIMINARY RESULTS

In this section, we provide some basic definitions of generalized fractional integrals and derivatives, some important results and preliminary facts that are very useful to us in the sequel.

Let  $0 < a < b < \infty$  be a finite interval on  $\mathbb{R}^+$  and  $C[a, b]$  be the Banach space of all continuous functions  $h : [a, b] \rightarrow \mathbb{R}$  with the norm

$$\|h\|_C = \max \{|h(t)| : t \in [a, b]\}.$$

For  $0 \leq \beta < 1$  and the parameter  $\rho > 0$  we define the weighted space of continuous functions  $h$  on  $(a, b]$  by

$$C_{\beta, \rho}[a, b] = \left\{ h : (a, b] \rightarrow \mathbb{R} : \left( \frac{t^\rho - a^\rho}{\rho} \right)^\beta h(t) \in C[a, b] \right\},$$

with the norm

$$\|h\|_{C_{\beta,\rho}} = \left\| \left( \frac{t^\rho - a^\rho}{\rho} \right)^\beta h(t) \right\|_C = \max_{t \in [a,b]} \left| \left( \frac{t^\rho - a^\rho}{\rho} \right)^\beta h(t) \right|.$$

It is obvious that  $C_{0,\rho}[a, b] = C[a, b]$ .

Let  $\delta_\rho = \left( t^{\rho-1} \frac{d}{dt} \right)$ . We define the Banach space of continuously differentiable functions  $h$  on  $[a, b]$  by

$$C_{\delta_\rho,\beta}^1[a, b] = \{h : [a, b] \rightarrow \mathbb{R} : \delta_\rho h \in C_{\beta,\rho}[a, b]\},$$

with the norms

$$\|h\|_{C_{\delta_\rho,\beta}^1} = \|h\|_C + \|\delta_\rho h\|_{C_{\beta,\rho}}$$

and

$$\|h\|_{C_{\delta_\rho,\beta}^0} = \max \{|\delta_\rho h(t)| : t \in [a, b]\}.$$

Note that  $C_{\delta_\rho,\beta}^0[a, b] = C_{\beta,\rho}[a, b]$ .

**Definition 2.1** (Katugampola fractional integral [20, 27]). Let  $\mu, c \in \mathbb{R}$ , with  $\mu > 0$ ,  $u \in Z_c^\mu(a, b)$ , where  $Z_c^\mu(a, b)$  is the space of Lebesgue measurable functions with complex values. The left-sided Katugampola fractional integral of order  $\mu$  is defined by

$$(2.1) \quad ({}^\rho I_{a^+}^\mu u)(t) = \frac{\rho^{1-\mu}}{\Gamma(\mu)} \int_a^t \frac{x^{\rho-1} u(x)}{(t^\rho - x^\rho)^{1-\mu}} dx, \quad t > a.$$

**Definition 2.2** (Katugampola fractional derivative [21, 27]). Let  $\mu, \rho \in \mathbb{R}$  be such that  $\mu \notin \mathbb{N}$ ,  $0 < \mu, \rho$ . The left-sided Katugampola fractional derivative of order  $\mu$  is defined by

$$(2.2) \quad ({}^\rho D_{a^+}^\mu u)(t) = \delta_\rho^n ({}^\rho I_{a^+}^{n-\mu} u)(t) = \frac{\rho^{1-n+\mu}}{\Gamma(n-\mu)} \left( t^{1-\rho} \frac{d}{dt} \right)^n \int_a^t \frac{x^{\rho-1} u(x)}{(t^\rho - x^\rho)^{1-n+\mu}} dx,$$

where  $n = [\mu] + 1$  is such that  $[\mu]$  is the integer part of  $\mu$ .

**Definition 2.3** (Generalized Katugampola fractional derivative [27]). Let  $0 < \mu \leq 1$  and  $0 \leq \nu \leq 1$ . The generalized Katugampola fractional derivative (of order  $\mu$  and type  $\nu$ ) with respect to  $t$  is defined by

$$(2.3) \quad \begin{aligned} ({}^\rho D_{a^\pm}^{\mu,\nu} u)(t) &= \left\{ \pm {}^\rho I_{a^\pm}^{\nu(1-\mu)} \left( t^{\rho-1} \frac{d}{dt} \right)^1 {}^\rho I_{a^\pm}^{(1-\nu)(1-\mu)} u \right\} (t) \\ &= \left\{ \pm {}^\rho I_{a^\pm}^{\nu(1-\mu)} \delta_\rho {}^\rho I_{a^\pm}^{(1-\nu)(1-\mu)} u \right\} (t), \end{aligned}$$

where  $\rho > 0$ ,  $u \in C_{1-\beta,\rho}[0, 1]$  and  $I$  is Katugampola fractional integral defined in (2.1).

*Remark 2.1.* ([27]). For  $\beta = \mu + \nu(1 - \mu)$ , the generalized Katugampola fractional derivative operator  ${}^\rho D_{a^+}^{\mu, \nu}$  can be expressed as

$$(2.4) \quad {}^\rho D_{a^+}^{\mu, \nu} = {}^\rho I_{a^+}^{\nu(1-\mu)} \delta_\rho {}^\rho I_{a^+}^{1-\beta} = {}^\rho I_{a^+}^{\nu(1-\mu)} {}^\rho D_{a^+}^\beta.$$

**Lemma 2.1** ([27]). *Let  $\mu > 0$ ,  $0 \leq \beta < 1$  and  $u \in C_{\beta, \rho}[a, b]$ . Then*

$$({}^\rho D_{a^+}^\mu {}^\rho I_{a^+}^\mu u)(t) = u(t), \quad \text{for all } t \in (a, b].$$

**Lemma 2.2** (Semigroup property [27]). *Let  $\mu > 0$ ,  $\nu > 0$ ,  $1 \leq q \leq \infty$ ,  $a, b \in (0, \infty)$  such that  $a < b$  and  $\rho, s \in \mathbb{R}$ ,  $s \leq \rho$ . Then the following property holds true*

$$({}^\rho I_{a^+}^\mu {}^\rho I_{a^+}^\nu u)(t) = ({}^\rho I_{a^+}^{\mu+\nu} u)(t),$$

for all  $u \in Z_s^q(a, b)$ .

**Lemma 2.3** ([27]). *Let  $t > a$  and for  $\mu \geq 0$  and  $\nu > 0$ , we have*

$$\begin{aligned} \left[ {}^\rho D_{a^+}^\mu \left( \frac{x^\rho - a^\rho}{\rho} \right)^{\mu-1} \right] (t) &= 0, \quad 0 < \mu < 1, \\ \left[ {}^\rho I_{a^+}^\mu \left( \frac{x^\rho - a^\rho}{\rho} \right)^{\nu-1} \right] (t) &= \frac{\Gamma(\nu)}{\Gamma(\mu + \nu)} \left( \frac{x^\rho - a^\rho}{\rho} \right)^{\mu+\nu-1}. \end{aligned}$$

**Lemma 2.4** ([27]). *Let  $\mu > 0$ ,  $0 \leq \beta < 1$  and  $a, b \in (0, \infty)$  such that  $a < b$  and  $u \in C_{\beta, \rho}[a, b]$ . Then*

$$({}^\rho I_{a^+}^\mu u)(a) = \lim_{t \rightarrow a^+} ({}^\rho I_{a^+}^\mu u)(t) = 0,$$

and  ${}^\rho I_{a^+}^\mu u$  is continuous on  $[a, b]$  if  $\beta < \mu$ .

**Lemma 2.5** ([27]). *Let  $\mu \in (0, 1)$ ,  $\nu \in [0, 1]$  and  $\beta = \mu + \nu - \mu\nu$ . If  $u \in C_{1-\beta}^\beta[a, b]$  then*

$${}^\rho I_{a^+}^\beta {}^\rho D_{a^+}^\beta u = {}^\rho I_{a^+}^\mu {}^\rho D_{a^+}^{\mu, \nu} u$$

and

$${}^\rho D_{a^+}^\beta {}^\rho I_{a^+}^\mu u = {}^\rho D_{a^+}^{\nu(1-\mu)} u.$$

**Lemma 2.6** ([27]). *Let  $\mu \in (0, 1)$ ,  $0 \leq \beta < 1$ . If  $u \in C_\beta[a, b]$  and  ${}^\rho I_{a^+}^{1-\mu} u \in C_\beta^1[a, b]$ , then for all  $t \in (a, b]$*

$$({}^\rho I_{a^+}^\mu {}^\rho D_{a^+}^\mu u)(t) = - \left( \frac{t^\rho - a^\rho}{\rho} \right)^{\mu-1} \frac{({}^\rho I_{a^+}^{1-\beta} u)(a)}{\Gamma(\mu)} + u(t).$$

**Lemma 2.7** ([27]). *Let  $u \in L^1(a, b)$ . If  ${}^\rho D_{a^+}^{\nu(1-\mu)} u$  exists on  $L^1(a, b)$ , then*

$${}^\rho D_{a^+}^{\mu, \nu} {}^\rho I_{a^+}^\mu u = {}^\rho I_{a^+}^{\nu(1-\mu)} {}^\rho D_{a^+}^{\nu(1-\mu)} u.$$

**Lemma 2.8** ([27]). *Let  $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  be a function where  $f(\cdot, u(\cdot)) \in C_{1-\beta}[a, b]$ . A function  $u \in C_{1-\beta}^\beta[a, b]$  is a solution of fractional IVP*

$$D_{a^+}^{\mu, \nu} u(t) = f(t, u(t)), \quad \mu \in (0, 1), \nu \in [0, 1],$$

$$I_{a^+}^{1-\beta} u(a^+) = u_0, \quad \beta = \mu + \nu - \mu\nu,$$

if and only if  $u$  satisfies the integral equation of Volterra type:

$$u(t) = \frac{u_0(t-a)^{\beta-1}}{\Gamma(\beta)} + \frac{1}{\Gamma(\mu)} \int_a^t (t-x)^{\mu-1} f(x, u(x)) dx.$$

**Definition 2.4** (Volterra integral equation). A linear Volterra integral equation of the second kind has the form of

$$u(t) = u_0(t) + \int_a^t K(t, x)u(x) dx,$$

where  $K$  is a kernel.

**Theorem 2.1** (Krasnoselskii fixed point theorem [28]). *Let  $E$  be a nonempty closed, bounded and convex subset of a Banach space  $(\mathcal{B}, \|\cdot\|)$ . Further, assume that  $F$  and  $G$  are two operators defined on  $E$  which map  $E$  into  $\mathcal{B}$  such that*

- (a)  $F(x) + G(y) \in E$  for all  $x, y \in E$ ;
- (b)  $F$  is a contraction;
- (c)  $G$  is continuous and compact.

Then  $F + G$  has a fixed point in  $E$ .

Using the above fundamental results, the following theorem yields the equivalence between the IVP (1.1)–(1.2) and an improved mixed type Volterra integral equation.

**Theorem 2.2.** *Let  $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  be a function such that for any  $u \in C_{1-\beta}[a, b]$   $f(\cdot, u(\cdot)) \in C_{1-\beta}[a, b]$ , where  $\beta = \mu + \nu - \mu\nu$ , with  $0 < \mu \leq 1$ ,  $0 \leq \nu \leq 1$ . Function  $u \in C_{1-\beta}^\beta[a, b]$  is a solution of IVP (1.1)–(1.2) if and only if it satisfies the following mixed type Volterra integral equation*

$$(2.5) \quad u(t) = \frac{K}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx$$

$$+ \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx,$$

where  $K = \left\{ \Gamma(\beta) - \sum_{i=1}^m \lambda_i \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} \right\}^{-1}$ .

*Proof.* Let  $u \in C_{1-\beta}^\beta[a, b]$  be a solution of IVP (1.1)–(1.2). Then by the Lemma 2.8 the solution of IVP (1.1)–(1.2) can be written as

$$(2.6) \quad u(t) = \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \frac{({}^\rho I_{a^+}^{1-\beta} u)(a)}{\Gamma(\beta)} + \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx.$$

Now, substitute  $t = \kappa_i$  in the above equation

$$u(\kappa_i) = \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} \frac{({}^\rho I_{a^+}^{1-\beta} u)(a)}{\Gamma(\beta)} + \frac{1}{\Gamma(\mu)} \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx.$$

Multiplying by  $\lambda_i$  the both hand sides, we get

$$\lambda_i u(\kappa_i) = \lambda_i \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} \frac{({}^\rho I_{a^+}^{1-\beta} u)(a)}{\Gamma(\beta)} + \frac{\lambda_i}{\Gamma(\mu)} \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx.$$

Thus, we have

$$\begin{aligned} {}^\rho I_{a^+}^{1-\beta} u(a) &= \sum_{i=1}^m \lambda_i u(\kappa_i), \\ &= \frac{({}^\rho I_{a^+}^{1-\beta} u)(a)}{\Gamma(\beta)} \sum_{i=1}^m \lambda_i \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} \\ &\quad + \frac{1}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx, \end{aligned}$$

which implies

$$(2.7) \quad ({}^\rho I_{a^+}^{1-\beta} u)(a) = \frac{\Gamma(\beta)}{\Gamma(\mu)} K \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx.$$

Substituting (2.7) in (2.6) we get (2.5), which proved that  $u$  also satisfies integral equation (2.5) when it satisfies IVP (1.1)–(1.2). This proved the necessity. Now, we prove the sufficiency by applying  ${}^\rho I_{a^+}^{1-\beta}$  to both hand sides of the integral equation (2.5), we have

$$\begin{aligned} {}^\rho I_{a^+}^{1-\beta} u(t) &= {}^\rho I_{a^+}^{1-\beta} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \frac{K}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \\ &\quad + {}^\rho I_{a^+}^{1-\beta} {}^\rho I_{a^+}^\mu f(x, u(x)). \end{aligned}$$

By using Lemma 2.2, Lemma 2.1 and Lemma 2.3, we have

$${}^\rho I_{a^+}^{1-\beta} u(t) = \frac{\Gamma(\beta)}{\Gamma(\mu)} K \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx + {}^\rho I_{a^+}^{1-\nu(1-\mu)} f(t, u(t)).$$

Since  $1 - \nu(1 - \mu) > 1 - \beta$ , by taking the limit as  $t \rightarrow a$  and using Lemma 2.4, we have

$$(2.8) \quad ({}^\rho I_{a^+}^{1-\beta} u)(a) = \frac{\Gamma(\beta)}{\Gamma(\mu)} K \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx.$$

Now, substituting  $t = \kappa_i$  in (2.5), we have

$$\begin{aligned} u(\kappa_i) &= \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} \frac{K}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \\ &\quad + \frac{1}{\Gamma(\mu)} \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx. \end{aligned}$$

Then we have

$$\begin{aligned} \sum_{i=1}^m \lambda_i u(\kappa_i) &= \frac{K}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \\ &\quad + \frac{1}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \\ &= \frac{1}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \\ &\quad \times \left\{ K \sum_{i=1}^m \lambda_i \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\beta-1} + 1 \right\} \\ (2.9) \quad &= \frac{\Gamma(\beta)}{\Gamma(\mu)} K \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx. \end{aligned}$$

It follows from (2.8) and (2.9), that

$${}^\rho I_{a^+}^{1-\beta} u(a) = \sum_{i=1}^m \lambda_i u(\kappa_i).$$

It follows from Lemma 2.3 and Lemma 2.5 and by applying  ${}^\rho D_{a^+}^\beta$  to both hand sides of (2.5) that

$$(2.10) \quad {}^\rho D_{a^+}^\beta u(t) = {}^\rho D_{a^+}^{\nu(1-\mu)} f(t, u(t)).$$

Since  $u \in C_{1-\beta}^\beta[a, b]$  and by the definition of  $C_{1-\beta}^\beta[a, b]$ , we have  ${}^\rho D_{a^+}^\beta u \in C_{1-\beta}^\beta[a, b]$ . Then  ${}^\rho D_{a^+}^{\nu(1-\mu)} f = {}^\rho D^\rho I_{a^+}^{1-\nu(1-\mu)} f \in C_{1-\beta}[a, b]$ . It is obvious that for any  $f \in C_{1-\beta}[a, b]$ ,  ${}^\rho I_{a^+}^{1-\nu(1-\mu)} f \in C_{1-\beta}[a, b]$ , then  ${}^\rho I_{a^+}^{1-\nu(1-\mu)} f \in C_{1-\beta}^1[a, b]$ . Thus,  $f$  and  ${}^\rho I_{a^+}^{1-\nu(1-\mu)} f$  satisfy both the conditions of Lemma 2.6.

Now, it follows from Lemma 2.6, by applying  ${}^\rho I_{a^+}^{\nu(1-\mu)}$  on both sides of (2.10), that

$$(2.11) \quad ({}^\rho D_{a^+}^{\mu,\nu} u)(t) = - \left( \frac{t^\rho - a^\rho}{\rho} \right)^{\nu(1-\mu)-1} \frac{{}^\rho I_{a^+}^{1-\nu(1-\mu)} f(a)}{\Gamma(\nu(1-\mu))} + f(t, u(t)).$$

By Lemma 2.4, it implies that  ${}^\rho I_{a^+}^{1-\nu(1-\mu)} f(a) = 0$ . Hence, (2.11) reduces to

$$({}^\rho D_{a^+}^{\mu,\nu} u)(t) = f(t, u(t)).$$

This completes the proof. □

### 3. MAIN RESULT

In the sequel, let us introduce the following hypothesis.

[Q1] Let  $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function such that for any  $u \in C_{1-\beta}[a, b]$   $f(\cdot, u(\cdot)) \in C_{1-\beta}^{\nu(1-\mu)}[a, b]$ . For all  $u, v \in \mathbb{R}$  there exists a positive constant  $J > 0$  such that

$$|f(t, u) - f(t, v)| \leq J|u - v|.$$

[Q2] The constant

$$(3.1) \quad \sigma := \frac{JB(\mu, \beta)}{\Gamma(\mu)} \left\{ |K| \sum_{i=1}^m \lambda_i \left( \frac{\kappa_i^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} + \left( \frac{b^\rho - a^\rho}{\rho} \right)^\mu \right\} < 1,$$

where  $K$  is defined in the Theorem 2.2.

Now, we will establish our main existence result for IVP (1.1)–(1.2) using Krasnoselskii fixed point theorem.

**Theorem 3.1.** *Assume that the hypothesis [Q1] and [Q2] are satisfied. Then IVP (1.1)–(1.2) has at least one solution in  $C_{1-\beta}^\beta[a, b]$ .*

*Proof.* According to Theorem 2.2, it is sufficient to prove the existence result for the mixed type integral equation (2.5).

Now, define the operator  $\Delta$  by

$$(3.2) \quad \begin{aligned} (\Delta u)(t) &= \frac{K}{\Gamma(\mu)} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left( \frac{\kappa_i^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \\ &+ \frac{1}{\Gamma(\mu)} \int_a^t \left( \frac{t^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx. \end{aligned}$$

It is obvious that the operator  $\Delta$  is well defined and maps  $C_{1-\beta}[a, b]$  into  $C_{1-\beta}[a, b]$ . Let  $\hat{f}(x) = f(x, 0)$  and

$$(3.3) \quad \eta := \frac{B(\mu, \beta)}{\Gamma(\mu)} \left\{ |K| \sum_{i=1}^m \lambda_i \left( \frac{\kappa_i^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} + \left( \frac{b^\rho - a^\rho}{\rho} \right)^\mu \right\} \|\hat{f}\|_{C_{1-\beta}}.$$

Consider a ball  $B_s := \{u \in C_{1-\beta}[a, b] : \|u\|_{C_{1-\beta}} \leq s\}$ , with  $\frac{\eta}{1-\sigma} \leq s, \sigma < 1$ .



Now, let us subdivide the operator  $\Delta$  into two operators  $F$  and  $G$  on  $B_s$  as follows:

$$(Fu)(t) = \frac{K}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx$$

and

$$(Gu)(t) = \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx.$$

The proof is divided into following steps.

Step I. For every  $u, v \in B_s$ ,  $Fu + Gv \in B_s$ . For the operator  $F$

$$(Fu)(t) \left(\frac{t^\rho - a^\rho}{\rho}\right)^{1-\beta} = \frac{K}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx, \quad t \in (a, b],$$

we have

$$\begin{aligned} \left| (Fu)(t) \left(\frac{t^\rho - a^\rho}{\rho}\right)^{1-\beta} \right| &\leq \frac{|K|}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} |f(x, u(x))| dx \\ &\leq \frac{|K|}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} (|f(x, u(x)) - f(x, 0)| \\ &\quad + |f(x, 0)|) dx \\ &\leq \frac{|K|}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} (J|u(x)| + |\hat{f}(x)|) dx. \end{aligned}$$

Here we use the fact that

$$\begin{aligned} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} |u(x)| dx &\leq \left\{ \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} \left(\frac{x^\rho - a^\rho}{\rho}\right)^{\beta-1} x^{\rho-1} dx \right\} \\ &\quad \times \|u(x)\|_{C_{1-\beta}} \\ (3.4) \qquad \qquad \qquad &= \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\mu+\beta-1} B(\mu, \beta) \|u(x)\|_{C_{1-\beta}}. \end{aligned}$$

Thus, we have

$$\begin{aligned} \left| (Fu)(t) \left(\frac{t^\rho - a^\rho}{\rho}\right)^{1-\beta} \right| &\leq \frac{|K|}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \left\{ \left(\frac{\kappa_i^\rho - a^\rho}{\rho}\right)^{\mu+\beta-1} B(\mu, \beta) \right. \\ &\quad \left. \times (J\|u(x)\|_{C_{1-\beta}} + \|\hat{f}(x)\|_{C_{1-\beta}}) \right\}, \end{aligned}$$

which gives

$$(3.5) \quad \|Fu\|_{C_{1-\beta}} \leq \frac{|K| B(\mu, \beta)}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \left\{ \left( \frac{\kappa_i^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} \left( J\|u(x)\|_{C_{1-\beta}} + \|\hat{f}(x)\|_{C_{1-\beta}} \right) \right\}.$$

For  $t \in (a, b]$  and the operator  $G$

$$(Gu)(t) \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} = \frac{1}{\Gamma(\mu)} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \int_a^t \left( \frac{t^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx,$$

we have

$$\begin{aligned} \left| (Gu)(t) \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \right| &\leq \frac{1}{\Gamma(\mu)} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \int_a^t \left( \frac{t^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} |f(x, u(x))| dx \\ &\leq \frac{1}{\Gamma(\mu)} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \\ &\quad \times \int_a^t \left( \frac{t^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} (J|u(x)| + |\hat{f}(x)|) dx. \end{aligned}$$

Again, by using (3.4), we have

$$\begin{aligned} \left| (Gu)(t) \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \right| &\leq \frac{1}{\Gamma(\mu)} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \left\{ \left( \frac{t^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} \right. \\ &\quad \left. \times B(\mu, \beta) \left( J\|u(x)\|_{C_{1-\beta}} + \|\hat{f}(x)\|_{C_{1-\beta}} \right) \right\} \\ &\leq \frac{B(\mu, \beta)}{\Gamma(\mu)} \left( \frac{b^\rho - a^\rho}{\rho} \right)^\mu \left( J\|u(x)\|_{C_{1-\beta}} + \|\hat{f}(x)\|_{C_{1-\beta}} \right), \end{aligned}$$

which gives

$$(3.6) \quad \|(Gu)\|_{C_{1-\beta}} \leq \frac{B(\mu, \beta)}{\Gamma(\mu)} \left( \frac{b^\rho - a^\rho}{\rho} \right)^\mu \left( J\|u(x)\|_{C_{1-\beta}} + \|\hat{f}(x)\|_{C_{1-\beta}} \right).$$

Combining (3.5) and (3.6) for every  $u, v \in B_s$  we have

$$\|Fu + Gv\|_{C_{1-\beta}} \leq \|Fu\|_{C_{1-\beta}} + \|(Gv)\|_{C_{1-\beta}} \leq \sigma s + \eta \leq s,$$

which implies that  $Fu + Gv \in B_s$ .

Step II. The operator  $F$  is contraction mapping.

For any  $u, v \in B_s$  and the operator  $F$

$$\{(Fu)(t) - (Fv)(t)\} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta}$$

$$= \frac{K}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left( \frac{\kappa_i^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} [f(x, u(x)) - f(x, v(x))] dx$$

we have

$$\begin{aligned} \left| \{(Fu)(t) - (Fv)(t)\} \left( \frac{t^\rho - a^\rho}{\rho} \right)^{1-\beta} \right| &\leq \frac{|K|}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left( \frac{\kappa_i^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} \\ &\quad \times |f(x, u(x)) - f(x, v(x))| dx \\ &\leq \frac{|K|}{\Gamma(\mu)} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left( \frac{\kappa_i^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} \\ &\quad \times J |u(x) - v(x)| dx \\ &\leq \frac{J|K|}{\Gamma(\mu)} B(\mu, \beta) \sum_{i=1}^m \lambda_i \left( \frac{\kappa_i^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} \\ &\quad \times \|u - v\|_{C_{1-\beta}}, \end{aligned}$$

which gives

$$\|Fu - Fv\|_{C_{1-\beta}} \leq \frac{J|K|}{\Gamma(\mu)} B(\mu, \beta) \sum_{i=1}^m \lambda_i \left( \frac{\kappa_i^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} \|u - v\|_{C_{1-\beta}} \leq \sigma \|u - v\|_{C_{1-\beta}}.$$

Hence, by the hypothesis [Q2] the operator  $F$  is a contraction mapping.

Step III. The operator  $G$  is compact and continuous.

Since the function  $f \in C_{1-\beta}[a, b]$ , it is obvious from the definition of  $C_{1-\beta}[a, b]$  that the operator  $G$  is continuous.

From the equation (3.6) of Step I clearly,  $G$  is uniformly bounded on  $B_s$ . Next we prove the compactness.

For any  $a < t_1 < t_2 \leq b$  we have

$$\begin{aligned} |(Gu)(t_1) - (Gu)(t_2)| &= \left| \frac{1}{\Gamma(\mu)} \int_a^{t_1} \left( \frac{t_1^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \right. \\ &\quad \left. - \frac{1}{\Gamma(\mu)} \int_a^{t_2} \left( \frac{t_2^\rho - x^\rho}{\rho} \right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \right| \\ &\leq \frac{\|f\|_{C_{1-\beta}}}{\Gamma(\mu)} \left| \int_a^{t_1} \left( \frac{t_1^\rho - x^\rho}{\rho} \right)^{\mu-1} \left( \frac{x^\rho - a^\rho}{\rho} \right)^{\beta-1} x^{\rho-1} dx \right. \\ &\quad \left. - \int_a^{t_2} \left( \frac{t_2^\rho - x^\rho}{\rho} \right)^{\mu-1} \left( \frac{x^\rho - a^\rho}{\rho} \right)^{\beta-1} x^{\rho-1} dx \right| \\ &\leq \frac{\|f\|_{C_{1-\beta}} B(\mu, \beta)}{\Gamma(\mu)} \left| \left( \frac{t_1^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} - \left( \frac{t_2^\rho - a^\rho}{\rho} \right)^{\mu+\beta-1} \right| \end{aligned}$$

tending to zero as  $t_2 \rightarrow t_1$ , whether  $\mu + \beta - 1 \geq 0$  or  $\mu + \beta - 1 < 0$ . Thus,  $G$  is equicontinuous. Hence, by Arzel-Ascoli Theorem, the operator  $G$  is compact on  $B_s$ .

It follows from Krasnoselskii fixed point theorem that the IVP (1.1)–(1.2) has at least one solution  $u \in C_{1-\beta}[a, b]$ . Using the Lemma 2.7 and repeating the process of proof in Theorem 2.2, one can show that this solution is actually in  $C_{1-\beta}^\beta[a, b]$ . This completes the proof.  $\square$

**3.1. Ulam-Hyers-Rassias stability.** In this section, we discuss the Ulam stability results for the solution of IVP (1.1)–(1.2).

**Definition 3.1** ([1]). The solution of IVP (1.1)–(1.2) is said to be Ulam-Hyers stable if there exists a real number  $\psi > 0$  such that for every  $\varepsilon > 0$  and for each solution  $u \in C_{\beta,\rho}$  of the inequality

$$(3.7) \quad |({}^\rho D_{a^+}^{\mu,\nu} u)(t) - f(t, u(t))| \leq \varepsilon, \quad t \in (a, b],$$

there exists  $v \in C_{\beta,\rho}$ , a solution of IVP (1.1)–(1.2) satisfying

$$|u(t) - v(t)| \leq \varepsilon\psi, \quad t \in (a, b].$$

**Definition 3.2** ([1]). The solution of IVP (1.1)–(1.2) is said to be generalized Ulam-Hyers stable if there exists a continuous function  $\psi_f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ , with  $\psi_f(0) = 0$  such that for every solution  $u \in C_{\beta,\rho}$  of the inequality (3.7) there exists  $v \in C_{\beta,\rho}$ , a solution of IVP (1.1)–(1.2) satisfying

$$|u(t) - v(t)| \leq \psi_f(\varepsilon), \quad t \in (a, b].$$

**Definition 3.3** ([1]). The solution of IVP (1.1)–(1.2) is said to be Ulam-Hyers-Rassias stable with respect to  $\Psi \in C_{\beta,\rho}((a, b], \mathbb{R}_+)$  if there exists a real number  $0 < \psi_\theta$  such that for every  $0 < \varepsilon$  and for every solution  $u \in C_{\beta,\rho}$  of the inequality

$$(3.8) \quad |({}^\rho D_{a^+}^{\mu,\nu} u)(t) - f(t, u(t))| \leq \varepsilon\Psi(t), \quad t \in (a, b],$$

there exists  $v \in C_{\beta,\rho}$  a solution of IVP (1.1)–(1.2) satisfying

$$|u(t) - v(t)| \leq \varepsilon\psi_\theta\Psi(t), \quad t \in (a, b].$$

**Definition 3.4** ([1]). The solution of IVP (1.1)–(1.2) is said to be generalized Ulam-Hyers-Rassias stable with respect to  $\Psi \in C_{\beta,\rho}((a, b], \mathbb{R}_+)$  if there exists a real number  $0 < \psi_\theta$  such that for every solution  $u \in C_{\beta,\rho}$  of the inequality

$$(3.9) \quad |({}^\rho D_{a^+}^{\mu,\nu} u)(t) - f(t, u(t))| \leq \Psi(t), \quad t \in (a, b],$$

there exists  $v \in C_{\beta,\rho}$  a solution of IVP (1.1)–(1.2) satisfying

$$|u(t) - v(t)| \leq \psi_\theta\Psi(t), \quad t \in (a, b].$$

*Remark 3.1* ([1]). Clearly

- (a) from Definition 3.1 follows Definition 3.2;
- (b) from Definition 3.3 follows Definition 3.4;
- (c) from Definition 3.3 for  $\Psi(\cdot) = 1$  follows Definition 3.2.

Now, we establish the results on generalized Ulam-Hyers-Rassias stability of the IVP (1.1)–(1.2).

**Theorem 3.2.** *Assume that [Q1] and following hypothesis hold.*

[Q3] *There exists  $\omega_\theta > 0$  such that for each  $t \in (a, b]$  we have*

$${}^\rho I_{a^+}^\mu \Psi(t) \leq \omega_\theta \Psi(t).$$

[Q4] *There exists a function  $p \in C[(a, b], [0, \infty)]$  such that for each  $t \in (a, b]$*

$$|f(t, u(t))| \leq \frac{p(t) \Psi(t)}{1 + |u|} |u|.$$

*Then the solution of IVP (1.1)–(1.2) satisfies the generalized Ulam-Hyers-Rassias stability with respect to  $\Psi$ .*

*Proof.* Let  $u$  be a solution of the inequality (3.9) and let  $v$  be a solution of IVP (1.1)–(1.2). Then we have

$$\begin{aligned} v(t) &= \frac{K}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, v(x)) dx \\ &\quad + \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, v(x)) dx \\ &= \Phi_v + \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, v(x)) dx, \end{aligned}$$

where

$$\Phi_v = \frac{K}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, v(x)) dx.$$

On the other hand, if  $\sum_{i=1}^m \lambda_i u(\kappa_i) = \sum_{i=1}^m \lambda_i v(\kappa_i)$  and  ${}^\rho I_{a^+}^{1-\beta} u(a) = {}^\rho I_{a^+}^{1-\beta} v(a)$ , then  $\Phi_u = \Phi_v$ . Indeed,

$$\begin{aligned} |\Phi_u - \Phi_v| &\leq \frac{|K|}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} \\ &\quad \times |f(x, u(x)) - f(x, v(x))| dx \\ &\leq \frac{|K|}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i \int_a^{\kappa_i} \left(\frac{\kappa_i^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} J |u - v| dx \\ &\leq \frac{J|K|}{\Gamma(\mu)} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta-1} \sum_{i=1}^m \lambda_i {}^\rho I_{a^+}^{1-\beta} |u(\kappa_i) - v(\kappa_i)| \\ &= 0. \end{aligned}$$

Hence,  $\Phi_u = \Phi_v$ . Then we have

$$v(t) = \Phi_u + \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, v(x)) dx.$$

From inequality (3.9) and [Q3] for each  $t \in (a, b]$  we have

$$\left| u(t) - \Phi_u - \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \right| \leq {}^\rho I_{a^+}^\mu \Psi(t) \leq \omega_\theta \Psi(t).$$

Set  $\tilde{p} = \sup_{t \in (a, b]} p(t)$ . From the hypothesis [Q3] and [Q4] for each  $t \in (a, b]$  we have

$$\begin{aligned} |u(t) - v(t)| &\leq \left| u(t) - \Phi_u - \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} f(x, u(x)) dx \right| \\ &\quad + \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} |f(x, u(x)) - f(x, v(x))| dx \\ &\leq \omega_\theta \Psi(t) + \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} 2\tilde{p}\Psi(x) dx \\ &\leq \omega_\theta \Psi(t) + 2\tilde{p} ({}^\rho I_{a^+}^\mu \Psi)(t) \\ &\leq (1 + 2\tilde{p}) \omega_\theta \Psi(t) \\ &:= \psi_\theta \Psi(t). \end{aligned}$$

Thus, the IVP (1.1)–(1.2) is generalized Ulam-Hyers-Rassias stable with respect to  $\Psi$ . This completes the proof. □

Following theorem will be useful in the progress of our next result.

**Theorem 3.3** ([1]). *Let  $(\Omega, d)$  be a generalized complete metric space and a strictly contractive operator  $\Phi : \Omega \rightarrow \Omega$ , with a Lipschitz constant  $E < 1$ . If there exists a non negative integer  $j$  such that  $d(\Phi^{j+1}u, \Phi^{j+1}u) < \infty$  for some  $u \in \Omega$ , then the following propositions hold true:*

- A:**  $\{\Phi^j u\}_{n \in \mathbb{N}}$  converges to a fixed point  $u^*$  of  $\Phi$ ;
- B:**  $u^*$  is a unique fixed point of  $\Phi$  in  $\Omega^* = \{v \in \Omega : d(\Phi^*u, v) < \infty\}$ ;
- C:** if  $v \in \Omega^*$ , then  $d(v, u^*) \leq \frac{1}{1-E} d(v, \Phi u)$ .

Let  $Z = Z(I, \mathbb{R})$  be the metric space with the metric

$$d(u, v) = \sup_{t \in (a, b]} \frac{\left(\frac{t^\rho - a^\rho}{\rho}\right)^{1-\beta} |u(t) - v(t)|}{\Psi(t)}.$$

**Theorem 3.4.** *Assume that [Q3] and the following assumption hold.*

[Q5] *There exists  $\phi \in C((a, b], [0, \infty))$  such that for every  $u, v \in \mathbb{R}$  and for each  $t \in (a, b]$ , we have*

$$|f(t, u) - f(t, v)| \leq \left(\frac{t^\rho - a^\rho}{\rho}\right)^{1-\beta} \phi(t) \Psi(t) |u - v|.$$

If

$$E := \left(\frac{G^\rho - a^\rho}{\rho}\right)^{1-\beta} \phi^* \omega_\theta < 1,$$

where  $\phi^* = \sup_{t \in (a, b]} \phi(t)$ , then there exists a unique solution  $u_0$  of the IVP (1.1)–(1.2) and IVP (1.1)–(1.2) is generalized Ulam-Hyers-Rassias stable. Moreover,

$$|u(t) - u_0(t)| \leq \frac{\Psi(t)}{1 - E}.$$

*Proof.* Let the operator  $\Delta : C_{\beta, \rho} \rightarrow C_{\beta, \rho}$  be defined in (3.2). By using Theorem 3.3, we have

$$\begin{aligned} |(\Delta u)(t) - (\Delta v)(t)| &\leq \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} |f(x, u(x)) - f(x, v(x))| dx \\ &\leq \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} \phi(x) \Psi(x) \\ &\quad \times \left| \left(\frac{x^\rho - a^\rho}{\rho}\right)^{1-\beta} u(x) - \left(\frac{x^\rho - a^\rho}{\rho}\right)^{1-\beta} v(x) \right| dx \\ &\leq \frac{1}{\Gamma(\mu)} \int_a^t \left(\frac{t^\rho - x^\rho}{\rho}\right)^{\mu-1} x^{\rho-1} \phi^*(x) \Psi(x) \|u - v\|_{C_{1-\beta}} dx \\ &\leq \phi^* ({}^\rho I_{a+}^\mu) \Psi(t) \|u - v\|_C \\ &\leq \phi^* \omega_\theta \Psi(t) \|u - v\|_C. \end{aligned}$$

Hence,

$$\begin{aligned} \left| \left(\frac{t^\rho - x^\rho}{\rho}\right)^{1-\beta} (\Delta u)(t) - \left(\frac{t^\rho - x^\rho}{\rho}\right)^{1-\beta} (\Delta v)(t) \right| &\leq \left(\frac{G^\rho - a^\rho}{\rho}\right)^{1-\beta} \phi^* \omega_\theta \\ &\quad \times \Psi(t) \|u - v\|_C. \end{aligned}$$

Thus, we have

$$d(\Delta u, \Delta v) = \sup_{t \in (a, b]} \frac{\|(\Delta u)(t) - (\Delta v)(t)\|_C}{\Psi(t)} \leq E \|u - v\|_C.$$

This completes the theorem. □

3.2. **Examples.**

*Example 3.1.* Consider the following IVP:

$$(3.10) \quad {}^\rho D_{0^+}^{\mu, \nu} u(t) = \frac{|u(t)|}{50e^{t+5}(1+|u(t)|)}, \quad t \in (0, 1],$$

$$(3.11) \quad {}^\rho I_{0^+}^{1-\beta} u(0) = 5u\left(\frac{1}{2}\right) + 3u\left(\frac{3}{4}\right), \quad \beta = \mu + \nu(1 - \mu),$$

where  $\mu = \frac{1}{2}$ ,  $\nu = \frac{2}{3}$  and  $\beta = \frac{5}{6}$ . Set  $f(t, u) = \frac{|u|}{50e^{t+5}(1+|u|)}$ ,  $t \in (0, 1]$ .

It is obvious that the function  $f$  is continuous. For any  $u, v \in \mathbb{R}$  and  $t \in (0, 1]$

$$|f(t, u) - f(t, v)| \leq \frac{1}{50e^5} |u - v|.$$

Thus, the condition [Q1] of Theorem 3.1 is satisfied, with  $J = \frac{1}{50e^5}$ . Moreover, with some elementary computation for  $\rho > 0$  we have

$$|K| = \left| \left\{ \Gamma\left(\frac{5}{6}\right) - \left[ 5 \left( \frac{(1/2)^\rho - 0^\rho}{\rho} \right)^{-1/6} + 3 \left( \frac{(3/4)^\rho - 0^\rho}{\rho} \right)^{-1/6} \right] \right\}^{-1} \right| < 1$$

and

$$\begin{aligned} \sigma &= \frac{1}{50e^5} \cdot \frac{B(1/2, 5/6)}{\Gamma(1/2)} \left\{ |K| \left[ 5 \left( \frac{(1/2)^\rho - 0^\rho}{\rho} \right)^{1/3} + 3 \left( \frac{(3/4)^\rho - 0^\rho}{\rho} \right)^{1/3} \right] \right. \\ &\quad \left. + \left( \frac{1^\rho - 0^\rho}{\rho} \right)^{1/2} \right\} < 1. \end{aligned}$$

Hence, the condition [Q2] of Theorem 3.1 is satisfied.

It follows, from Theorem 3.1, that the IVP (3.10)–(3.11) has at least one solution in  $C_{1/6}[0, 1]$ .

Now, let  $\Psi(t) = \frac{1}{t^{2\rho-4}}$  and  $p(t) = \frac{1}{50e^{t+5}}$ , then

$$|f(t, u(t))| \leq \frac{1}{50e^{t+5}} \cdot \frac{1}{t^{2\rho-4}} \cdot \frac{|u(t)|}{(1+|u(t)|)}.$$

Thus, the condition [Q4] of Theorem 3.2 is satisfied and with the obvious elementary computation, we have

$${}^\rho I_{0^+}^\mu \Psi(t) = \frac{\rho^{1-\mu}}{\Gamma(\mu)} \int_a^t \frac{x^{\rho-1} \Psi(x)}{(t^\rho - x^\rho)^{1-\mu}} dx \leq \frac{1}{\rho^\mu \Gamma(\mu)} B\left(\mu, \frac{4}{\rho} - 1\right) \Psi(t) \leq \omega_\theta \Psi(t).$$

Hence, the condition [Q4] of Theorem 3.2 is satisfied with  $\omega_\theta = \frac{1}{\rho^\mu \Gamma(\mu)} B\left(\mu, \frac{4}{\rho} - 1\right)$ . It follows from the Theorem 3.2 that the IVP (3.10)–(3.11) is generalized Ulam-Hyers-Rassias stable.



## 4. CONCLUSION

We have investigated the sufficient conditions for the existence of solutions to the nonlocal initial value problems involving generalized Katugampola derivative. We have used Krasnoselskii fixed point theorem to develop the existence results. Further, we established some conditions for the generalized Ulam-Hyers-Rassias stability corresponding to the considered problem. Finally, as an application, a suitable example is given to demonstrate our main results.

## REFERENCES

- [1] S. Abbas, M. Benchohra, J. Lagreg, A. Alsaedi and Y. Zhou, *Existence and Ulam stability for fractional differential equations of Hilfer-Hadamard type*, Adv. Difference Equ. **180** (2017), DOI 10.1186/s13662-017-1231-1.
- [2] B. Ahmad and S. K. Ntouyas, *Initial value problem of fractional order Hadamard-type functional differential equations*, Electron. J. Differential Equations **2015** (2015), 1–9.
- [3] A. O. Akdemir, A. Ekinici and E. Set, *Conformable fractional integrals and related new integral inequalities*, J. Nonlinear Convex Anal. **18** (2017), 661–674.
- [4] A. O. Akdemir, E. Set and A. Ekinici, *On new conformable fractional integral inequalities for product of different kinds of convexity*, TWMS Journal of Applied and Engineering Mathematics **9** (2019), 142–150.
- [5] R. Almeida, A. B. Malinowska and T. Odziejewicz, *Fractional differential equations with dependence on the Caputo-Katugampola derivative*, Journal of Computational and Nonlinear Dynamics **11** (2016), 11 pages.
- [6] M. Benchohra and J. E. Lazreg, *On stability for nonlinear implicit fractional differential equations*, Matematiche (Catania) **70** (2014), 49–61.
- [7] M. Benchohra and J. E. Lazreg, *Existence and Ulam stability for nonlinear implicit fractional differential equations with Hadamard derivative*, Stud. Univ. Babeş-Bolyai Math. **62** (2017), 27–38.
- [8] D. B. Dhaigude and S. P. Bhairat, *Existence and uniqueness of solution of Cauchy type problem for Hilfer fractional differential equations*, Communications in Applied Analysis **22** (2018), 121–134.
- [9] D. B. Dhaigude and S. P. Bhairat, *Existence and continuation of solutions of Hilfer fractional differential equations*, J. Math. Model. **7** (2019), 1–20.
- [10] M. A. Dokuyucu, D. Baleanu and E. Celik, *Analysis of the fractional Keller-Segel model*, Filomat **32**(16) (2018), 5633–5643.
- [11] M. A. Dokuyucu, E. Celik, H. Bulut and H. M. Baskonus, *Cancer treatment model with the Caputo-Fabrizio fractional derivative*, Eur. Phys. J. Plus **133** (2018), Paper ID 92, DOI 10.1140/epjp/i2018-11950-y.
- [12] A. Ekinici and N. Eroğlu, *New generalizations for convex functions via conformable fractional integrals*, Filomat **33**(14) (2019), 4525–4534.
- [13] A. Ekinici and M. E. Özdemir, *Some new integral inequalities via Riemann-Liouville integral operators*, Appl. Comput. Math. **3** (2019), 288–295.
- [14] K. M. Furati, M. D. Kassim and N. e. Tatar, *Existence and uniqueness for a problem involving Hilfer fractional derivative*, Comput. Math. Appl. **64** (2012), 1616–1626.
- [15] K. M. Furati, M. D. Kassim and N. e. Tatar, *Non-existence of global solutions for a differential equations involving Hilfer fractional derivative*, Electron. J. Differential Equations **2013** (2013), 1–10.
- [16] R. Hilfer, *Fractional Time Evolution: Applications in Fractional Calculus in Physics*, World Scientific, London, 2000.

- [17] R. Hilfer, Y. Luckho and Z. Tomovski, *Operational method for the solution of fractional differential equations with generalized Riemann-Liouville fractional derivatives*, *Fract. Calc. Appl. Anal.* **12** (2009), 289–318.
- [18] S. M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, 2001.
- [19] M. D. Kassim, K. M. Furati and N. -E. Tatar, *On a differential equation involving Hilfer-Hadamard fractional derivative*, *Abstr. Appl. Anal.* **2012** (2012), Article ID 391062, 17 pages.
- [20] U. N. Katugampola, *New approach to a generalized fractional integral*, *Appl. Math. Comput.* **218** (2011), 860–865.
- [21] U. N. Katugampola, *A new approach to generalized fractional derivatives*, *Bull. Math. Anal. Appl.* **6** (2014), 1–15.
- [22] A. A. Kilbas, *Hadamard-type fractional calculus*, *J. Korean Math. Soc.* **38** (2001), 1191–1204.
- [23] C. Kou, H. Zhou and M. Medved, *Existence and continuation theorems of Riemann-Liouville type fractional differential equations*, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **22** (2011), Article ID 1250077, 12 pages.
- [24] K. D. Kucche and S. T. Sutar, *Stability via successive approximation for nonlinear implicit fractional differential equations*, *Moroccan Journal of Pure and Applied Analysis* **3** (2017), 36–55.
- [25] K. S. Miller and B. Ross, *A Introduction to the Fractional Calculus and Differential Equations*, John Wiley, New York, 1993.
- [26] I. Mumcu, E. Set and A. O. Akdemir, *Hermite-Hadamard type inequalities for harmonically convex functions via Katugampola fractional integrals*, *Miskolc Math. Notes* **20** (2019), 409–424.
- [27] D. S. Oliveira and E. C. de Oliveira, *Hilfer-Katugampola fractional derivative*, *Comput. Appl. Math.* **37** (2018), 3672–3690.
- [28] D. R. Smart, *Fixed Point Theorems*, Cambridge University Press, Cambridge, 1973.
- [29] R. Subashini, K. Jothimani, S. Saranya and C. Ravichandran, *On the results of Hilfer fractional derivatives with nonlocal conditions*, *International Journal of Pure and Applied Mathematics* **118** (2018), 277–289.
- [30] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience, New York, 1968.
- [31] D. Vivek, K. Kanagrajan and E. M. Elsayed, *Nonlocal initial value problems for implicit differential equations with Hilfer-Hadamard fractional derivative*, *Nonlinear Analysis: Modelling and Control* **23** (2018), 341–360.
- [32] J. Wang and Y. Zhang, *Nonlocal initial value problems for differential equations with Hilfer fractional derivative*, *Appl. Math. Comput.* **266** (2015), 850–859.
- [33] J. Wang, Y. Zhou and M. Medved, *Existence and stability of fractional differential equations with Hadamard derivative*, *Topol. Methods Nonlinear Anal.* **41** (2013), 113–133.
- [34] H. Yaldiz and A. O. Akdemir, *Katugampola fractional integrals within the class of convex functions*, *Turkish Journal of Science* **3** (2018), 40–50.

<sup>1</sup>DEPARTMENT OF FIRST YEAR ENGINEERING,  
PIMPRI CHINCHWAD COLLEGE OF ENGINEERING,  
NIGDI, PUNE-411044, INDIA  
*Email address:* arif.bagwan@gmail.com

<sup>2</sup>DEPARTMENT OF MATHEMATICS,  
DR. BABASAHEB AMBEDKAR MARATHWADA UNIVERSITY,  
AURANGABAD-431004, INDIA  
*Email address:* pachpatte@gmail.com