

QUOTIENT HOOPS INDUCED BY QUASI-VALUATION MAPS

R. A. BORZOOEI¹, G. R. REZAEI², M. AALY KOLOGANI³, AND Y. B. JUN⁴

ABSTRACT. In this paper, our aim was making a metric space on hoop algebras, because of that, we introduced the notion of valuation maps from F -quasi-valuation map based on hoops and related properties of them are investigated. By using these notions, we introduced a quasi-metric space. The continuity of operations of a hoop is studied with topology induced by a quasi-valuation. Also, we studied hoop homomorphism and investigated that under which condition this homomorphism is an F -quasi-valuation map. Moreover, we wanted to find a congruence relation on hoops in a new way and study about the quotient structure that is made by it. So, we defined a congruence relation by F -quasi-valuation map and proved that the quotient is a hoop.

1. INTRODUCTION

Non-classical logic has become a considerable formal tool for computer science and artificial intelligence to deal with fuzzy information and uncertainty information. Many-valued logic, a great extension and development of classical logic, has always been a crucial direction in non-classical logic. In order to research the many-valued logical system whose propositional value is given in a lattice, Bosbach in [14, 15], proposed the concept of hoops, and discussed their some properties. Hoops are naturally ordered commutative residuated integral monoids. In the last years, hoops theory and related structures was enriched with deep structure theorems [1, 3–10, 12, 13, 16–18, 22, 24, 27]. Many of these results have a strong impact with fuzzy logic. Particularly, from the structure theorem of finite basic hoops one obtains an elegant short proof of the completeness theorem for propositional basic logic, introduced

Key words and phrases. Hoop, quasi-valuation map, S_{\odot} -quasi-valuation map, S_{\rightarrow} -quasi-valuation map, F -quasi-valuation map, (pseudo) metric space.

2010 *Mathematics Subject Classification.* Primary: 06B30. Secondary: 03G25, 06D20.

DOI 10.46793/KgJMat2205.743B

Received: July 27, 2019.

Accepted: March 17, 2020.

by Hájek in [21]. The algebraic structures corresponding to Hájek's propositional (fuzzy) basic logic, BL-algebras, are particular cases of hoops. The main example of BL-algebras in interval $[0, 1]$ endowed with the structure induced by a t-norm. MV-algebras, product algebras and Gödel algebras are the most known classes of BL-algebras. Recent investigations are concerned with non-commutative generalizations for these structures. During these years, many researchers study on hoops in different way, and got some results on hoops [11, 20, 23, 26]. Algebra and topology, the two fundamental domains of mathematics, play complementary roles. Topology studies continuity and convergence and provides a general framework to study the concept of a limit. Algebra studies all kinds of operations and provides a basis for algorithms and calculations. Many of the most important objects of mathematics represent a blend of algebraic and of topological structures. Topological function spaces and linear topological spaces in general, topological groups and topological fields and topological lattices are objects of this kind. Very often an algebraic structure and a topology come naturally together. The rules that describe the relationship between a topology and algebraic operation are almost always transparent and natural the operation has to be continuous, jointly continuous, jointly or separately. In the 20th century many topologists and algebraists have contributed to topological algebra. Song, Roh and Jun, in [25] introduced the notion of quasi-valuation maps based on a subalgebra and an ideal in BCK/BCI-algebras, and then they investigated several properties. They provided relations between a quasi-valuation map based on a subalgebra and a quasi-valuation map based on an ideal. In a BCI-algebra, they gave a condition for a quasi-valuation map based on an ideal to be a quasi-valuation map based on a subalgebra, and found conditions for a real-valued function on a BCK/BCI-algebra to be a quasi-valuation map based on an ideal. Using the notion of a quasi-valuation map based on an ideal, they constructed (pseudo) metric spaces, and shew that the binary operation \star in BCK-algebras is uniformly continuous. In [2], Aaly and Rezaei, introduced the notion of quasi-valuation maps such as $(S_{\circ}, S_{\rightarrow})$ S -quasi-valuation maps and F -quasi-valuation map based on subalgebras and filters and related properties of them are investigated. Also, they studied the relation between them and proved that every F -quasi-valuation map is an S -quasi-valuation map. Finally, by using the notion F -quasi-valuation map, they introduced a metric space and proved that if λ is an F -quasi-valuation map of hoop H then all operation of H are continuous.

In this paper, our aim was making a metric space on hoop algebras, because of that, we introduced the notion of valuation maps from F -quasi-valuation map based on hoops and related properties of them are investigated. By using these notions, we introduced a quasi-metric space. The continuity of operations of a hoop is studied with topology induced by a quasi-valuation. Also, we studied hoop homomorphism and investigated that under which condition these homomorphism is an F -quasi-valuation map. Moreover, we wanted to find a congruence relation on hoops in a new way and

study about the quotient structure that is made by it. So, we defined a congruence relation by F -quasi-valuation map and proved that the quotient is a hoop.

2. PRELIMINARIES

By a *hoop* we mean an algebra $(H, \odot, \rightarrow, 1)$ in which $(H, \odot, 1)$ is a commutative monoid and for all $x, y, z \in H$ the following assertions are valid:

- (H1) $x \rightarrow x = 1$;
- (H2) $x \odot (x \rightarrow y) = y \odot (y \rightarrow x)$;
- (H3) $x \rightarrow (y \rightarrow z) = (x \odot y) \rightarrow z$.

We define a relation " \leq " on a hoop H by

$$(\forall x, y \in H)(x \leq y \Leftrightarrow x \rightarrow y = 1).$$

It is easy to see that (H, \leq) is a poset. A hoop H is *bounded* if there is an element $0 \in H$ such that, for all $x \in H$, $0 \leq x$. Let $x^0 = 1$ and $x^n = x^{n-1} \odot x$, for any $n \in \mathbb{N}$. If H is a bounded hoop, then we define a negation " $'$ " on H such that, for all $x \in H$, $x' = x \rightarrow 0$. A nonempty subset S of H is called a *subhoop* of H if it satisfies:

$$(\forall x, y \in S)(x \odot y \in S, x \rightarrow y \in S).$$

Note that every subhoop contains the element 1.

Proposition 2.1 ([19]). *Let $(H, \odot, \rightarrow, 1)$ be a hoop. For any $x, y, z \in H$, the following conditions hold:*

- (a1) (H, \leq) is a meet-semilattice with $x \wedge y = x \odot (x \rightarrow y)$;
- (a2) $x \odot y \leq z$ if and only if $x \leq y \rightarrow z$;
- (a3) $x \odot y \leq x, y$ and $x^n \leq x$ for any $n \in \mathbb{N}$;
- (a4) $x \leq y \rightarrow x$;
- (a5) $1 \rightarrow x = x$ and $x \rightarrow 1 = 1$;
- (a6) $x \odot (x \rightarrow y) \leq y$ and $x \odot y \leq x \wedge y \leq x \rightarrow y$;
- (a7) $x \rightarrow y \leq (y \rightarrow z) \rightarrow (x \rightarrow z)$;
- (a8) $x \leq y$ implies $x \odot z \leq y \odot z$, $z \rightarrow x \leq z \rightarrow y$ and $y \rightarrow z \leq x \rightarrow z$;
- (a9) $x \rightarrow (y \rightarrow z) = (x \odot y) \rightarrow z = y \rightarrow (x \rightarrow z)$.

A nonempty subset F of a hoop H is called a *filter* of H (see [19]) if the following assertions are valid:

$$(2.1) \quad (\forall x, y \in H)(x, y \in F \Rightarrow x \odot y \in F),$$

$$(2.2) \quad (\forall x, y \in H)(x \in F, x \leq y \Rightarrow y \in F).$$

Note that the conditions (2.1) and (2.2) mean that F is closed under the operation \odot and F is upward closed, respectively.

Note that a subset F of a hoop H is a filter of H if and only if the following assertions are valid (see [19]):

$$1 \in F, \\ (\forall x, y \in H)(x \rightarrow y \in F, x \in F \Rightarrow y \in F).$$

Definition 2.1 ([2]). A real valued function λ of H is called

- an S_{\odot} -quasi-valuation map of H if

$$(\forall x, y \in H)(\lambda(x \odot y) \geq \lambda(x) + \lambda(y));$$

- an S_{\rightarrow} -quasi-valuation map of H if

$$(\forall x, y \in H)(\lambda(x \rightarrow y) \geq \lambda(x) + \lambda(y));$$

- an S -quasi-valuation map of H if it is an S_{\odot} -quasi-valuation map and an S_{\rightarrow} -quasi-valuation map of H .

Definition 2.2 ([2]). A real valued function λ of H is called an F -quasi-valuation map of H if

$$\lambda(1) = 0, \\ (\forall x, y \in H)(\lambda(y) \geq \lambda(x) + \lambda(x \rightarrow y)).$$

Proposition 2.2 ([2]). Let λ be an F -quasi-valuation map on H . Then the following statements hold:

- (i) λ is an S -quasi-valuation map on H ;
- (ii) λ is an order preserving map;
- (iii) for any $x \in H$, $\lambda(x) \leq 0$.

Theorem 2.1 ([2]). If an F -quasi-valuation map λ of H satisfies the following condition

$$(\forall x \in H)(\lambda(x) = 0 \Rightarrow x = 1),$$

then (H, d_{λ}) is a metric space.

Note. In what follows, let H denote a hoop unless otherwise specified.

3. QUASI-VALUATION MAPS ON HOOPS

In this section, we introduce the notion of valuation maps from F -quasi-valuation map based on hoops and related properties of them are investigated. By using these notions, we introduce a quasi-metric space. The continuity of operations of a hoop was studied with topology induced by a quasi-valuation.

If a F -quasi-valuation map λ of H satisfies:

$$(\forall x \in H)(x \neq 1 \Rightarrow \lambda(x) \neq 0),$$

then we say that λ is an F -valuation map of H .

TABLE 1. Cayley table for the binary operation “ \odot ”

\odot	0	a	b	1
0	0	0	0	0
a	0	0	a	a
b	0	a	b	b
1	0	a	b	1

TABLE 2. Cayley table for the binary operation “ \rightarrow ”

\rightarrow	0	a	b	1
0	1	1	1	1
a	a	1	1	1
b	0	a	1	1
1	0	a	b	1

Example 3.1. Let $H = \{0, a, b, 1\}$ be a set with Cayley tables (Table 1 and 2). Then $(H, \odot, \rightarrow, 1)$ is a bounded hoop. Define a map λ on H as follows:

$$\lambda : H \rightarrow \mathbb{R}, \quad x \mapsto \begin{cases} -30, & \text{if } x = 0, \\ -25, & \text{if } x = a, \\ -20, & \text{if } x = b, \\ 0, & \text{if } x = 1. \end{cases}$$

It is routine to verify that λ is an F -valuation map of H .

For any non-empty subset F of H and a negative real number k , define a real valued function λ_F on H as follows:

$$(3.1) \quad \lambda_F : H \rightarrow \mathbb{R}, \quad x \mapsto \begin{cases} 0, & \text{if } x \in F, \\ k, & \text{otherwise.} \end{cases}$$

Lemma 3.1 ([2]). *If F is a filter of H , then the function λ_F in (3.1) is an F -quasi-valuation map of H and $F_{\lambda_F} = F$.*

Theorem 3.1. *The function λ_F in (3.1) is an F -valuation map of H if and only if F is the trivial filter of H , that is $F = \{1\}$.*

Proof. Straightforward. □

In the following, we introduce quasi-metric space by using the notion of valuation maps from F -quasi-valuation map based on hoops. The continuity of operations of a hoop will study with topology induced by a quasi-valuation.

Definition 3.1. A function $d : H \times H \rightarrow \mathbb{R}$ is called a *quasi-metric* on H if it satisfies:

$$\begin{aligned} (\forall x, y \in H) (d(x, y) \leq 0, d(x, x) = 0), \\ (\forall x, y \in H) (d(x, y) = d(y, x)), \\ (\forall x, y, z \in H) (d(x, z) \geq d(x, y) + d(y, z)). \end{aligned}$$

We say that the pair (H, d) is a *quasi-metric space*.

Theorem 3.2. *If λ is an F -quasi-valuation map of H , then (H, d_λ) is a quasi-metric space which is called the quasi-metric space induced by λ , where*

$$d_\lambda : H \times H \rightarrow \mathbb{R}, \quad (x, y) \mapsto \lambda(x \rightarrow y) + \lambda(y \rightarrow x).$$

Proof. Since λ is an F -quasi-valuation map of H , by Proposition 2.2, λ is order preserving and for any $x \in H$, $\lambda(x) \leq 0$. Thus, according to definition of d_λ , it is clear that, for any $x, y \in H$, $d_\lambda(x, y) \leq 0$. Let $x \in H$. Then $d_\lambda(x, x) = \lambda(x \rightarrow x) = \lambda(1) = 0$. Also, for any $x, y \in H$,

$$d_\lambda(x, y) = \lambda(x \rightarrow y) + \lambda(y \rightarrow x) = \lambda(y \rightarrow x) + \lambda(x \rightarrow y) = d_\lambda(y, x).$$

Moreover, by Proposition 2.1 (a7), for any $x, y, z \in H$, $(x \rightarrow y) \odot (y \rightarrow z) \leq x \rightarrow z$. Since λ is an F -quasi-valuation map of H , by Proposition 2.2, λ is order preserving and S -quasi-valuation map of H . Then

$$\lambda(x \rightarrow y) + \lambda(y \rightarrow z) \leq \lambda((x \rightarrow y) \odot (y \rightarrow z)) \leq \lambda(x \rightarrow z).$$

By the similar way, $\lambda(z \rightarrow y) + \lambda(y \rightarrow x) \leq \lambda((z \rightarrow y) \odot (y \rightarrow x)) \leq \lambda(z \rightarrow x)$. Hence,

$$\begin{aligned} d_\lambda(x, y) + d_\lambda(y, z) &= \lambda(x \rightarrow y) + \lambda(y \rightarrow x) + \lambda(y \rightarrow z) + \lambda(z \rightarrow y) \\ &\leq \lambda(x \rightarrow z) + \lambda(z \rightarrow x) = d_\lambda(x, z). \end{aligned}$$

Therefore, (H, d_λ) is a quasi-metric space which is called the quasi-metric space induced by λ . \square

Proposition 3.1. *Every quasi-metric space (H, d_λ) induced by an F -quasi-valuation map λ of H satisfies:*

$$\begin{aligned} d_\lambda(x, y) &\leq \min\{d_\lambda(x \rightarrow a, y \rightarrow a), d_\lambda(a \rightarrow x, a \rightarrow y)\}, \\ d_\lambda(x \rightarrow y, a \rightarrow b) &\geq d_\lambda(x \rightarrow y, a \rightarrow y) + d_\lambda(a \rightarrow y, a \rightarrow b), \\ d_\lambda(x \odot y, a \odot b) &\geq d_\lambda(x \odot y, a \odot y) + d_\lambda(a \odot y, a \odot b), \end{aligned}$$

for all $a, b, x, y \in H$.

Proof. Let (H, d_λ) be a quasi-metric space. By Proposition 2.1 (a7) for any $x, y, z \in H$, $x \rightarrow y \leq (y \rightarrow a) \rightarrow (x \rightarrow a)$ and $y \rightarrow x \leq (x \rightarrow a) \rightarrow (y \rightarrow a)$. Since λ is an F -quasi-valuation map of H , by Proposition 2.2, λ is order preserving and S -quasi-valuation map of H . Then

$$\begin{aligned} d_\lambda(x, y) &\leq \lambda(x \rightarrow y) + \lambda(y \rightarrow x) \leq \lambda((y \rightarrow a) \rightarrow (x \rightarrow a)) + \lambda((x \rightarrow a) \rightarrow (y \rightarrow a)) \\ &= d_\lambda(x \rightarrow a, y \rightarrow a). \end{aligned}$$

By the similar way, $d_\lambda(x, y) \leq d_\lambda(a \rightarrow x, a \rightarrow y)$. Hence,

$$d_\lambda(x, y) \leq \min\{d_\lambda(x \rightarrow a, y \rightarrow a), d_\lambda(a \rightarrow x, a \rightarrow y)\}.$$

Now, let $x, y, a \in H$. Then by Proposition 2.1 (a7), we have

$$((x \rightarrow y) \rightarrow (y \rightarrow a)) \odot ((y \rightarrow a) \rightarrow (a \rightarrow b)) \leq (x \rightarrow y) \rightarrow (a \rightarrow b).$$

By the similar way,

$$((a \rightarrow b) \rightarrow (a \rightarrow y)) \odot ((a \rightarrow y) \rightarrow (x \rightarrow y)) \leq (a \rightarrow b) \rightarrow (x \rightarrow y).$$

Since λ is an F -quasi-valuation map of H , by Proposition 2.2, λ is order preserving and S -quasi-valuation map of H . Then it is clear that

$$d_\lambda(x \rightarrow y, a \rightarrow y) + d_\lambda(a \rightarrow y, a \rightarrow b) \leq d_\lambda(x \rightarrow y, a \rightarrow b).$$

Also, since $y \odot a \leq y \odot a$, by Proposition 2.1 (a2) and (a8), we have $y \leq a \rightarrow (y \odot a)$, and so $x \rightarrow y \leq (x \odot a) \rightarrow (y \odot a)$. Then, by Proposition 2.2, λ is order preserving, thus, $\lambda(x \rightarrow y) \leq \lambda((x \odot a) \rightarrow (y \odot a))$. By the similar way, $\lambda(y \rightarrow x) \leq \lambda((y \odot a) \rightarrow (x \odot a))$. Hence,

$$\begin{aligned} d_\lambda(x, y) &= \lambda(x \rightarrow y) + \lambda(y \rightarrow x) \leq \lambda((x \odot a) \rightarrow (y \odot a)) + \lambda((y \odot a) \rightarrow (x \odot a)) \\ &= d_\lambda(x \odot a, y \odot a). \end{aligned}$$

Then, for any $x, y, a, b \in H$, since (H, d_λ) is a quasi-metric space, we have,

$$d_\lambda(x \odot y, a \odot y) + d_\lambda(a \odot y, a \odot b) \leq d_\lambda(x \odot y, a \odot b). \quad \square$$

Theorem 3.3. *If λ is an F -valuation map of H , then the quasi-metric space induced by λ satisfies the following assertion,*

$$(3.2) \quad (\forall x, y \in H) (d_\lambda(x, y) = 0 \Rightarrow x = y).$$

Proof. Let λ be an F -valuation map of H . Then λ is an F -quasi-valuation map of H . Thus, by Theorem 3.2, $d_\lambda(x, y)$ is quasi-metric. Now, for any $x, y \in H$, if $d_\lambda(x, y) = 0$, then $\lambda(x \rightarrow y) + \lambda(y \rightarrow x) = 0$. Since λ is an F -quasi-valuation map of H , by Proposition 2.2, for any $x \in H$, $\lambda(x) \leq 0$. So, by routine calculations, it is clear that $\lambda(x \rightarrow y) = \lambda(y \rightarrow x) = 0$, and so by Theorem 2.1, $x \rightarrow y = 1$ and $y \rightarrow x = 1$. Therefore, $x = y$. □

We consider conditions for an F -quasi-valuation map to be an F -valuation map.

Theorem 3.4. *If the quasi-metric space (H, d_λ) induced by an F -quasi-valuation map λ of H satisfies the condition (3.2), then λ is an F -valuation map of H .*

Proof. Let λ be an F -quasi-valuation map of H and there exists $1 \neq x \in H$ such that $\lambda(x) = 0$. Since λ is an F -quasi-valuation map of H that satisfying the condition (3.2), we have

$$d_\lambda(1, x) = \lambda(1 \rightarrow x) + \lambda(x \rightarrow 1) = \lambda(x) + \lambda(1) = 0.$$

Then $d_\lambda(1, x) = 0$. Since (H, d_λ) is a quasi metric, we have $x = 1$, which is a contradiction. Hence, for any $1 \neq x \in H$, $\lambda(x) \neq 0$. Therefore, λ is an F -valuation map of H . \square

Note. If (H, d_λ) is a quasi-metric space, then for any $x \in H$ and $\varepsilon < 0$ the set $B_\varepsilon(x) = \{y \in H \mid d_\lambda(x, y) > \varepsilon\}$ is called a *ball of radius $|\varepsilon|$ with center at x* . The set $U \subseteq H$ is open in (H, d_λ) if, for any $x \in U$, there is an $\varepsilon < 0$ such that $B_\varepsilon(x) \subseteq U$. The topology \mathcal{T}_{d_λ} induced by d_λ is the collection of all open sets in (H, d_λ) .

Theorem 3.5. *If \mathcal{T}_λ is an induced topology by d_λ , then $(H, \odot, \rightarrow, \mathcal{T}_\lambda)$ is a topological hoop.*

Proof. By Theorem 3.3, (H, d_λ) is a quasi-metric space. Let $x, y \in H$ such that $x \rightarrow y \in B_\varepsilon(x \rightarrow y)$ for any $\varepsilon < 0$. We claim that $B_\varepsilon(x) \rightarrow B_\varepsilon(y) \subseteq B_\varepsilon(x \rightarrow y)$. For this, suppose $z \in B_\varepsilon(x) \rightarrow B_\varepsilon(y)$. Then there exist $p \in B_\varepsilon(x)$ and $q \in B_\varepsilon(y)$ such that $z = p \rightarrow q$. Thus, $d_\lambda(x, p) \geq \frac{\varepsilon}{2}$ and $d_\lambda(y, q) \geq \frac{\varepsilon}{2}$. By Proposition 3.1, it is clear that $d_\lambda(x \rightarrow y, p \rightarrow q) \geq d_\lambda(x, p)$ and $d_\lambda(p \rightarrow y, p \rightarrow q) \geq d_\lambda(y, q)$. Thus,

$$d_\lambda(x \rightarrow y, p \rightarrow q) \geq d_\lambda(x \rightarrow y, p \rightarrow y) + d_\lambda(p \rightarrow y, p \rightarrow q) \geq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

So, $d_\lambda(x \rightarrow y, p \rightarrow q) \geq \varepsilon$. Hence, $z \in B_\varepsilon(x \rightarrow y)$ and so, $(H, \rightarrow, \mathcal{T}_\lambda)$ is a topological hoop. By the similar way, we can prove that $(H, \odot, \mathcal{T}_\lambda)$ is a topological hoop. Therefore, $(H, \odot, \rightarrow, \mathcal{T}_\lambda)$ is a topological hoop. \square

Theorem 3.6. *For any F -quasi-valuation map λ of H , if we define a relation R_λ on H as follows:*

$$(\forall x, y) ((x, y) \in R_\lambda \Leftrightarrow d_\lambda(x, y) = 0),$$

then R_λ is a congruence relation on H .

We say that R_λ is a congruence relation on H induced by λ .

Proof. Let $x, y, z \in H$. For proving that R_λ is a congruence relation on H , first of all we have to prove that R_λ is an equivalence relation on H . It is clear that R_λ is reflexive and symmetric relation on H . Suppose $(x, y) \in R_\lambda$ and $(y, z) \in R_\lambda$. Then $d_\lambda(x, y) = 0$ and $d_\lambda(y, z) = 0$. By Proposition 2.1 (a7), for any $x, y, z \in H$, we have $(x \rightarrow y) \odot (y \rightarrow z) \leq x \rightarrow z$. Since λ is an F -quasi-valuation map of H , by Proposition 2.2, λ is order preserving and S -quasi-valuation map of H , then $\lambda(x \rightarrow y) + \lambda(y \rightarrow z) \leq \lambda(x \rightarrow z)$. By the similar way, $\lambda(z \rightarrow y) + \lambda(y \rightarrow x) \leq \lambda(z \rightarrow x)$. Hence,

$$0 = \lambda(x \rightarrow y) + \lambda(y \rightarrow x) + \lambda(y \rightarrow z) + \lambda(z \rightarrow y) \leq \lambda(x \rightarrow z) + \lambda(z \rightarrow x) = d_\lambda(x, z).$$

Since λ is an F -quasi-valuation map of H , by Proposition 2.2, for any $x \in H$, $\lambda(x) \leq 0$. Then $d_\lambda(x, z) = 0$ and so $(x, z) \in R_\lambda$. Therefore, R_λ is a transitive relation on H . Now, we prove that R_λ is a congruence relation on H . For any $x, y, z \in H$ such that $(x, y) \in R_\lambda$. Since $y \leq z \rightarrow (y \odot z)$, by Proposition 2.1 (a8),

$$x \rightarrow y \leq x \rightarrow (z \rightarrow (y \odot z)) = (x \odot z) \rightarrow (y \odot z).$$

Since λ is an F -quasi-valuation map of H , by Proposition 2.2, λ is order preserving, then $\lambda(x \rightarrow y) \leq \lambda((x \odot z) \rightarrow (y \odot z))$. By the similar way, it is clear that

$$\lambda(y \rightarrow x) \leq \lambda((y \odot z) \rightarrow (x \odot z)).$$

Hence,

$$0 = d_\lambda(x, y) \leq \lambda((x \odot z) \rightarrow (y \odot z)) + \lambda((y \odot z) \rightarrow (x \odot z)) = d_\lambda(x \odot z, y \odot z).$$

Since λ is an F -quasi-valuation map of H , by Proposition 2.2, for any $x \in H$, $\lambda(x) \leq 0$. Then $d_\lambda(x \odot z, y \odot z) = 0$ and so $(x \odot z, y \odot z) \in R_\lambda$. Moreover, if $(x, y) \in R_\lambda$, then by Proposition 3.1, it is clear that $(x \rightarrow z, y \rightarrow z) \in R_\lambda$ and $(z \rightarrow x, z \rightarrow y) \in R_\lambda$. Therefore, R_λ is a congruence relation on H induced by λ . \square

For any congruence relation R_λ induced by F -quasi-valuation map λ of H , let H_λ denote the set of all equivalence classes, that is,

$$H_\lambda := \{x_\lambda \mid x \in H\},$$

where $x_\lambda := \{y \in H \mid (x, y) \in R_\lambda\}$.

Theorem 3.7. *If λ is an F -quasi-valuation map of H , then $(H_\lambda, \square, \rightarrow, 1_\lambda)$ is a hoop, where*

$$(\forall x_\lambda, y_\lambda \in H_\lambda) (x_\lambda \square y_\lambda = (x \odot y)_\lambda, x_\lambda \rightarrow y_\lambda = (x \rightarrow y)_\lambda).$$

Proof. Let $x \in H$. Then it is clear that $x_\lambda \rightarrow x_\lambda = (x \rightarrow x)_\lambda = 1_\lambda$ and $(H_\lambda, \square, 1_\lambda)$ is a commutative monoid. Suppose $x_\lambda, y_\lambda, z_\lambda \in H_\lambda$. Then

$$\begin{aligned} (x_\lambda \square y_\lambda) \rightarrow z_\lambda &= (x \odot y)_\lambda \rightarrow z_\lambda = ((x \odot y) \rightarrow z)_\lambda = (x \rightarrow (y \rightarrow z))_\lambda \\ &= x_\lambda \rightarrow (y_\lambda \rightarrow z_\lambda). \end{aligned}$$

Moreover, by routine calculations, we have

$$x_\lambda \square (x_\lambda \rightarrow y_\lambda) = (x \odot (x \rightarrow y))_\lambda = (y \odot (y \rightarrow x))_\lambda = y_\lambda \square (y_\lambda \rightarrow x_\lambda).$$

Therefore, $(H_\lambda, \square, \rightarrow, 1_\lambda)$ is a hoop. \square

Theorem 3.7 is illustrated by the following example.

Example 3.2. According to Example 3.1, $H_\lambda = \{1_\lambda, x_\lambda, y_\lambda, 0_\lambda\}$.

Lemma 3.2 ([2]). *If $\lambda : H \rightarrow \mathbb{R}$ is an F -quasi-valuation map of H , then the set*

$$F_\lambda := \{x \in H \mid \lambda(x) = 0\}$$

is a filter of H .

Proposition 3.2. *If λ is an F -quasi-valuation map of H , then $F_\lambda = 1_\lambda$.*

Proof. Let λ be an F -quasi-valuation map of H . Then, by Lemma 3.2, we have

$$\begin{aligned} F_\lambda &= \{x \in H \mid \lambda(x) = 0\} = \{x \in H \mid \lambda(1 \rightarrow x) + \lambda(x \rightarrow 1) = 0\} \\ &= \{x \in H \mid d_\lambda(1, x) = 0\} \\ &= \{x \in H \mid (x, 1) \in R_\lambda\} \\ &= 1_\lambda. \end{aligned} \quad \square$$

For any filter F of H , let η_F be a relation on H defined by

$$(\forall x, y \in H) ((x, y) \in \eta_F \Leftrightarrow x \rightarrow y \in F, y \rightarrow x \in F).$$

Then η_F is a congruence relation on H (induced by F). Denote by H/F the set of all equivalence classes, that is,

$$H/F := \{[x] \mid x \in H\},$$

where $[x] = \{y \in H \mid (x, y) \in \eta_F\}$.

Theorem 3.8. *If λ is an F -quasi-valuation map of H , then $\eta_{F_\lambda} = R_\lambda$.*

Proof. Let $x, y \in H$. Then

$$\begin{aligned} (x, y) \in \eta_{F_\lambda} &\Leftrightarrow x \rightarrow y \in F_\lambda \text{ and } y \rightarrow x \in F_\lambda \\ &\Leftrightarrow \lambda(x \rightarrow y) = \lambda(y \rightarrow x) = 0 \\ &\Leftrightarrow \lambda(x \rightarrow y) + \lambda(y \rightarrow x) = 0 \\ &\Leftrightarrow d_\lambda(x, y) = 0 \\ &\Leftrightarrow (x, y) \in R_\lambda. \end{aligned} \quad \square$$

Theorem 3.9. *Let λ and g be F -quasi-valuation maps of H with $\lambda \neq g$. If $1_\lambda = 1_g$, then R_λ and R_g coincide and so $H_\lambda = H_g$.*

Proof. By routine calculations, we can see that $1_\lambda = \{x \in H \mid \lambda(x) = 0\}$. Suppose $x, y \in H$ such that $(x, y) \in R_\lambda$. Then $d_\lambda(x, y) = 0$ and so $\lambda(x \rightarrow y) + \lambda(y \rightarrow x) = 0$. Thus, $\lambda(x \rightarrow y) \geq -\lambda(y \rightarrow x)$. Since λ is an F -quasi-valuation map of H , by Proposition 2.2, we get that $\lambda(x \rightarrow y) = \lambda(y \rightarrow x) = 0$. Thus, $x \rightarrow y, y \rightarrow x \in 1_\lambda$. By assumption, $1_\lambda = 1_g$ we get that $x \rightarrow y, y \rightarrow x \in 1_g$, and so $g(x \rightarrow y) = g(y \rightarrow x) = 0$. Hence, $g(x \rightarrow y) + g(y \rightarrow x) = 0$, and so $d_g(x, y) = 0$. So $(x, y) \in R_g$. The proof of converse is similar. Therefore, R_λ and R_g coincide and so $H_\lambda = H_g$. \square

Theorem 3.10. *For any filter F and any F -quasi-valuation map λ of H such that $1_\lambda \subseteq F$ consider the set*

$$\overline{F}_\lambda := \{x_\lambda \mid x \in F\}.$$

Then the following assertions are valid:

- (1) $(\forall x \in H) (x \in F \Leftrightarrow x_\lambda \in \overline{F}_\lambda)$;
- (2) F_λ is a filter of H_λ .

Proof. (1) It is clear that if $x \in F$, then $x_\lambda \in \overline{F_\lambda}$. Suppose $x_\lambda \in \overline{F_\lambda}$. Then there exists $y \in F$ such that $x_\lambda = y_\lambda$. Thus, $(x, y) \in R_\lambda$. Since R_λ is a congruence relation on H , we have $(y \rightarrow x, 1) = (y \rightarrow x, y \rightarrow y) \in R_\lambda$. Hence, $y \rightarrow x \in 1_\lambda$. Since $1_\lambda \subseteq F$, we have $y \rightarrow x \in F$. Moreover, $y \in F$ and F is a filter of H , then $x \in F$.

(2) Since F is a filter of H , $1 \in F$, and so $1_\lambda \in \overline{F_\lambda}$. Suppose $x_\lambda, x_\lambda \rightarrow y_\lambda \in \overline{F_\lambda}$. Then by (1), $x \in F$ and $x \rightarrow y \in F$. Since F is a filter of H , $y \in F$. Thus, by (1), $y_\lambda \in \overline{F_\lambda}$. Therefore, $\overline{F_\lambda}$ is a filter of H_λ . \square

Proposition 3.3. *For any F -quasi-valuation map λ of H , let F^* be a filter of F_λ . Then the set*

$$F := \{x \in H \mid x_\lambda \in F^*\}$$

is a filter of H and $1_\lambda \subseteq F$.

Proof. Since F^* is a filter of $\overline{F_\lambda}$, $1_\lambda \in F^*$ and so $1 \in F$. Now, suppose $x, x \rightarrow y \in F$. Then $x_\lambda, (x \rightarrow y)_\lambda \in F^*$. Since F^* is a filter of $\overline{F_\lambda}$, we have $y_\lambda \in F^*$ and so $y \in F$. Hence, F is a filter of H . \square

Let $\mathcal{F}(H_\lambda)$ denote the set of all filters of F_λ and let $\mathcal{F}(H, \lambda)$ denote the set of all filters of H containing 1_λ . Then there exists a bijection between $\mathcal{F}(H_\lambda)$ and $\mathcal{F}(H, \lambda)$, that is,

$$f : \mathcal{F}(H_\lambda) \rightarrow \mathcal{F}(H, \lambda), \quad F \mapsto F_\lambda$$

is a bijection.

Theorem 3.11. *Let $g : H \rightarrow G$ be a homomorphism of hoops. Then the following hold.*

- (1) *If λ is an F -quasi-valuation map of G , then the composition $\lambda \circ g$ of λ and g is an F -quasi-valuation map of H .*
- (2) *If g is an isomorphism and if λ is an F -quasi-valuation map of G , then $H_{\lambda \circ g}$ and G_λ are isomorphic.*

Proof. (1) Since g is a homomorphism of hoops, we have $(\lambda \circ g)(1) = \lambda(g(1)) = \lambda(1)$. Since λ is an F -quasi-valuation map of H , we have $\lambda(1) = 0$ and so $(\lambda \circ g)(1) = 0$. Now, suppose $x, y \in H$. Since λ is an F -quasi-valuation map of H , we have

$$\begin{aligned} (\lambda \circ g)(x \rightarrow y) + (\lambda \circ g)(x) &= \lambda(g(x \rightarrow y)) + \lambda(g(x)) \\ &= \lambda(g(x) \rightarrow g(y)) + \lambda(g(x)) \\ &\leq \lambda(g(y)) = (\lambda \circ g)(y). \end{aligned}$$

(2) Let define the map $\phi : H_{\lambda \circ g} \rightarrow G_\lambda$ such that, for any $x_{\lambda \circ g} \in H_{\lambda \circ g}$, $\phi(x_{\lambda \circ g}) = (g(x))_\lambda$. Now, we prove that ϕ is an isomorphism. For this, let $x_{\lambda \circ g}, y_{\lambda \circ g} \in H_{\lambda \circ g}$. Then

$$\begin{aligned} \phi(x_{\lambda \circ g} \boxtimes y_{\lambda \circ g}) &= \phi((x \odot y)_{\lambda \circ g}) \\ &= (g(x \odot y))_\lambda = (g(x) \odot g(y))_\lambda \\ &= (g(x))_\lambda \boxtimes (g(y))_\lambda = \phi(x_{\lambda \circ g}) \boxtimes \phi(y_{\lambda \circ g}) \end{aligned}$$

and

$$\begin{aligned}\phi(x_{\lambda \circ g} \twoheadrightarrow y_{\lambda \circ g}) &= \phi((x \rightarrow y)_{\lambda \circ g}) \\ &= (g(x \rightarrow y))_{\lambda} = (g(x) \rightarrow g(y))_{\lambda} \\ &= (g(x))_{\lambda} \twoheadrightarrow (g(y))_{\lambda} = \phi(x_{\lambda \circ g}) \twoheadrightarrow \phi(y_{\lambda \circ g}),\end{aligned}$$

Hence, ϕ is a homomorphism of hoop.

Let $x, y \in H$ such that $\phi(x) = \phi(y)$. Then $\lambda \circ g(x) = \lambda \circ g(y)$. Thus, $(g(x))_{\lambda} = (g(y))_{\lambda}$, and so $(g(x), g(y)) \in R_{\lambda}$. Hence, $d_{\lambda}(g(x), g(y)) = 0$. Since d_{λ} is a quasi-metric and g is an isomorphism, we have $g(x) = g(y)$ and so $x = y$. Hence, ϕ is a one to one homomorphism.

Let $x_{\lambda} \in G_{\lambda}$. Since g is onto, there exists $y \in G$, such that $(g(y))_{\lambda} = x_{\lambda}$. Then $(\lambda \circ g)(y) = x_{\lambda}$, thus, $\phi(y) = x_{\lambda}$. Hence, ϕ is an isomorphism and so $H_{\lambda \circ g}$ and G_{λ} are isomorphic. \square

Theorem 3.12. *For any F -quasi-valuation map λ of H , we have the following assertions.*

- (1) *The map $\pi : H \rightarrow H_{\lambda}$, $x \mapsto x_{\lambda}$, is an onto homomorphism.*
- (2) *For any F -quasi-valuation map ϕ^* of H_{λ} , there exists an F -quasi-valuation map ϕ of H such that $\phi = \phi^* \circ \pi$.*
- (3) *The map $\lambda^* : H_{\lambda} \rightarrow \mathbb{R}$, $x_{\lambda} \mapsto \lambda(x)$, is an F -quasi-valuation map of H_{λ} .*

Proof. (1) By definition of H_{λ} , the proof is clear.

(2) Let define $\phi = \phi^*(x_{\lambda})$. We show that ϕ is an F -quasi-valuation map of H . For this, since $1 \in H$, we have $\phi(1) = \phi^*(1_{\lambda})$. Moreover, ϕ^* is an F -quasi-valuation map of H , $\phi(1) = 0$. Suppose $x, y \in H$ such that

$$\phi(x) + \phi(x \rightarrow y) = \phi^*(x_{\lambda}) + \phi^*(x_{\lambda} \twoheadrightarrow y_{\lambda}) \leq \phi^*(y_{\lambda}).$$

Since ϕ^* is an F -quasi-valuation map of H , we have $\phi(x) + \phi(x \rightarrow y) \leq \phi(y)$.

(3) Let $x_{\lambda} \in H_{\lambda}$. Since λ is an F -quasi-valuation map of H , we have

$$\lambda^*(x_{\lambda}) + \lambda^*(x_{\lambda} \twoheadrightarrow y_{\lambda}) = \lambda(x) + \lambda(x \rightarrow y) \leq \lambda(y) = \lambda^*(y_{\lambda}). \quad \square$$

Proposition 3.4. *Let H and G be two hoops and $\lambda : H \rightarrow \mathbb{R}$ and $\gamma : G \rightarrow \mathbb{R}$ be quasi-valuations. If $f : H \rightarrow G$ is a homomorphism, then the following statements are equivalent:*

- (i) *f is a quasi-valuation preserving;*
- (ii) *f is an isometry.*

Proof. (i) \Rightarrow (ii) Let f be a quasi-valuation preserving. Then, for any $x \in H$, define $\gamma(f(x)) = \lambda(x)$. For any $x, y \in H$, we have

$$\begin{aligned} d_\gamma(f(x), f(y)) &= \gamma(f(x) \rightarrow f(y)) + \gamma(f(y) \rightarrow f(x)) \\ &= \gamma(f(x \rightarrow y)) + \gamma(f(y \rightarrow x)) \\ &= \gamma \circ f(x \rightarrow y) + \gamma \circ f(y \rightarrow x) \\ &= \lambda(x \rightarrow y) + \lambda(y \rightarrow x) \\ &= d_\lambda(x, y). \end{aligned}$$

Hence, f is an isometry.

(ii) \Rightarrow (i) Let f be an isometry. Then, for any $x \in H$,

$$\begin{aligned} \lambda(x) = d_\lambda(x, 1) &= d_\gamma(f(x), f(1)) = \gamma(f(x) \rightarrow f(1)) + \gamma(f(1) \rightarrow f(x)) = \gamma(f(x)) \\ &= \gamma \circ f(x). \end{aligned}$$

Hence, f is a quasi-valuation preserving. □

Proposition 3.5. *Let $f : H \rightarrow G$ be a hoop isomorphism. If λ is a quasi-valuation on H , then $\gamma : G \rightarrow \mathbb{R}$ that, for any $y \in G$, is defined by $\gamma(y) = \lambda \circ f^{-1}(y)$ is a quasi-valuation. Moreover, if λ is an F -quasi-valuation on H , then γ is an F -quasi-valuation on G .*

Proof. Let $y_1, y_2 \in G$. Since f is an isomorphism, there exist $x_1, x_2 \in H$ such that $f(x_1) = y_1$ and $f(x_2) = y_2$. Then

$$\begin{aligned} \gamma(y_1 \rightarrow y_2) &= \lambda \circ f^{-1}(y_1 \rightarrow y_2) = \lambda(f^{-1}(y_1 \rightarrow y_2)) \\ &= \lambda(f^{-1}(y_1) \rightarrow f^{-1}(y_2)) = \lambda(x_1 \rightarrow x_2) \\ &\geq \lambda(x_1) + \lambda(x_2) = \lambda(f^{-1}(y_1)) + \lambda(f^{-1}(y_2)) \\ &= \gamma(y_1) + \gamma(y_2). \end{aligned}$$

By the similar way, we can prove that $\gamma(y_1 \odot y_2) \geq \gamma(y_1) + \gamma(y_2)$. Hence, γ is a quasi-valuation.

Since f is a hoop isomorphism, it is clear that $f(1_H) = 1_G$. Since λ is an F -quasi-valuation on H , we have $\gamma(1_G) = \lambda \circ f^{-1}(1_G) = \lambda(f^{-1}(1_G)) = \lambda(1_H) = 0$, and so $\gamma(1_G) = 0$. Let $x, y \in H$ and λ be an F -quasi-valuation on H . Since f is an isomorphism, there exist $a, b \in H$ such that $f(a) = x$ and $f(b) = y$. Then

$$\begin{aligned} \gamma(y) &= \lambda \circ f^{-1}(y) = \lambda(f^{-1}(y)) = \lambda(b) \\ &\geq \lambda(a \rightarrow b) + \lambda(a) = \lambda(f^{-1}(x)) + \lambda(f^{-1}(x) \rightarrow f^{-1}(y)) \\ &= \lambda(f^{-1}(x)) + \lambda(f^{-1}(x \rightarrow y)) = \lambda \circ f^{-1}(x) + \lambda \circ f^{-1}(x \rightarrow y) \\ &= \gamma(x) + \gamma(x \rightarrow y). \end{aligned} \quad \square$$

4. CONCLUSIONS AND FUTURE WORKS

In this paper, our aim was making a metric space on hoop algebras, because of that we introduced the notion of valuation maps from F -quasi-valuation map based on hoops and related properties of them are investigated. By using these notions, we introduce a quasi-metric space. The continuity of operations of a hoop was studied with topology induced by a quasi-valuation. Also, we study hoop homomorphism and investigate that under which condition these homomorphism is an F -quasi-valuation map. Moreover, we wanted to find a congruence relation on hoops in a new way and study about the quotient structure that is made by it. Because of that, we define a congruence relation by F -quasi-valuation map and prove that the quotient is a hoop. In our future work, we want to study about the product of finite number of this quasi-metric space and investigate that the quotient space of hoop has a quasi-metric or not. Finally we study the completion of this pace.

REFERENCES

- [1] M. Aaly Kologani and R. A. Borzooei, *On ideal theory of hoops*, Math. Bohem. (2019), DOI 10.21136/MB.2019.0140-17.
- [2] M. Aaly Kologani, G. R. Rezaei, R. A. Borzooei and Y. B. Jun, *Hoops with quasi-valuation maps*, J. Algebr. Syst. (to appear).
- [3] M. Aaly Kologani, M. Mohseni Takallo, and H. S. Kim, *Fuzzy filters of hoop based on fuzzy points*, Mathematics **7**(5) (2019), DOI 10.3390/math7050430.
- [4] M. Al Tahan, S. Hoskova-Mayerova and B. Davvaz, *An overview of topological hypergroupoids*, Journal of Intelligent and Fuzzy Systems **34**(3) (2018), 1907–1916.
- [5] S. Z. Alavi, R. A. Borzooei and M. Aaly Kologani, *Filter theory of pseudo hoop-algebras*, Italian Journal of Pure and Applied Mathematics **37** (2017), 619–632.
- [6] M. J. Atteya, C. Haetinger and D. I. Rasen, *(σ, τ) -derivation of semiprime rings*, Kragujevac J. Math. **4**(2) (2019), 239–246.
- [7] W. J. Blok and I. M. A. Ferreirim, *On the structure of hoops*, Algebra Universalis **43** (2000), 233–257.
- [8] W. J. Blok and I. M. A. Ferreirim, *Hoops and their implicational reducts*, Logic in Computer Science **28** (1993), 219–230.
- [9] H. Bordbar, M. M. Zahedi, and Y. B. Jun, *Ideals of is-algebras based on N -structures*, Kragujevac J. Math. **42**(4) (2018), 631–641.
- [10] R. A. Borzooei and M. Aaly Kologani, *Results on hoops*, Journal of Algebraic Hyperstructures and Logical Algebras **1**(1) (2019), 61–77.
- [11] R. A. Borzooei and M. Aaly Kologani, *Filter theory of hoop-algebras*, Journal of Advanced Research in Pure Mathematics **6**(4) (2014), 72–86.
- [12] R. A. Borzooei and M. Aaly Kologani, *Local and perfect semihoops*, Journal of Intelligent and Fuzzy Systems **29**(1) (2015), 223–234.
- [13] R. A. Borzooei, H. R. Varasteh and K. Borna, *Fundamental hoop-algebras*, Ratio Mathematica **29** (2015), 25–40.
- [14] B. Boshach, *Komplementäre halbgruppen axiomatik und arithmetik*, Fund. Math. **64**(3) (1969), 257–287.
- [15] B. Boshach, *Komplementäre halbgruppen kongruenzen und quotienten*, Fund. Math. **69**(1) (1970), 1–14.

- [16] A. Broumand Saeid, C. Flaut, S. Hoskova-Mayerovaa, M. Afshar and M. Kuchaki Rafsanjani, *Some connections between BCK-algebras and n -ary block codes*, Soft Computing **22** (2018), 41–46.
- [17] I. Cristea and S. Hoskova, *Fuzzy topological hypergroupoids*, Iran. J. Fuzzy Syst. **6**(4) (2009), 13–21.
- [18] S. Ebrahimi Atani and M. Sedghi Shanbeh Bazari, *Decomposable filters of lattices*, Kragujevac J. Math. **43**(1) (2019), 59–73.
- [19] G. Georgescu, L. Leustean, and V. Preoteasa, *Pseudo-hoops*, J. Mult.-Valued Logic Soft Comput. **11**(1–2) (2005), 153–184.
- [20] S. Ghorbani, *Localization of hoop-algebras*, Journal Advanced Research in Pure Mathematics **5**(3) (2013), 1–13.
- [21] P. Hájek, *Metamathematics of Fuzzy Logic*, Springer-Verlag, New York, 1998.
- [22] S. Hoskova, *Topological hypergroupoids*, Comput. Appl. Math. **64**(9) (2012), 2845–2849.
- [23] C. F. Luo, X. L. Xin and P. He, *n -Fold (positive) implicative filters of hoops*, Italian Journal of Pure and Applied Mathematics **38** (2017), 631–642.
- [24] E. Mohammadzadeh and R. A. Borzooei, *Results on (engel, solvable, nilpotent) fuzzy subpolygroups*, Kragujevac J. Math. **45**(5) (2021), 667–684.
- [25] S. Z. Song, E. H. Roh and Y. B. Jun, *Quasi-valuation maps on BCK/BCI-algebras*, Kyungpook Math. J. **55** (2015), 859–870.
- [26] R. Tayebi Khorami and A. Borumand Saeid, *Some unitary operators on hoop-algebras*, Fuzzy Information and Engineering **9**(2) (2017), 205–223.
- [27] M. A. Zander, *Decomposability of the finitely generated free hoop residuation algebra*, Studia Logica **88**(2) (2008), 233–246.

¹DEPARTMENT OF MATHEMATICS,
SHAHID BEHESHTI UNIVERSITY,
TEHRAN, IRAN
Email address: borzooei@sbu.ac.ir

²DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF SISTAN AND BALUCHESTAN,
ZAHEDAN, IRAN
Email address: grezaei@math.usb.ac.ir

³HATEF HIGHER EDUCATION INSTITUTE,
ZAHEDAN, IRAN
Email address: mona4011@gmail.com

⁴DEPARTMENT OF MATHEMATICS EDUCATION,
GYEONGSANG NATIONAL UNIVERSITY,
JINJU 52828, KOREA
Email address: skywine@gmail.com