

NEW MIXED RECURRENCE RELATIONS OF TWO-VARIABLE ORTHOGONAL POLYNOMIALS VIA DIFFERENTIAL OPERATORS

MOSAED M. MAKKY¹ AND MOHAMMAD SHADAB²

ABSTRACT. In this paper, we derive new recurrence relations for two-variable orthogonal polynomials for example Jacobi polynomial, Bateman's polynomial and Legendre polynomial via two different differential operators $\Xi = \left(\frac{\partial}{\partial z} + \sqrt{w}\frac{\partial}{\partial w}\right)$ and $\Delta = \left(\frac{1}{w}\frac{\partial}{\partial z} + \frac{1}{z}\frac{\partial}{\partial w}\right)$. We also derive some special cases of our main results.

1. INTRODUCTION AND PRELIMINARIES

In recent decades, the study of the multi-variable orthogonal polynomials has been substantially developed by many authors [3, 5, 15]. The properties of the multi-variable orthogonal polynomials have been analyzed by different approaches. The analytical properties of two-variable orthogonal polynomials like generating functions, recurrence relations, partial differential equations, and orthogonality have remained the main attraction of the topic due to its wide range of applications in different research areas [1, 4, 7, 10, 16].

Some new classes of two-variables analogues of the Jacobi polynomials have been introduced from Jacobi weights by Koornwinder [9]. These all classes are introduced by means of two different partial differential operators D_1 and D_2 , where D_1 has order two, and D_2 may have any arbitrary order. Koornwinder constructed bases of orthogonal polynomials in two-variables by using a tool given by Agahanov [2].

In 2017, M. Marriaga et al. [11] derived some new recurrence relations involving two-variable orthogonal polynomials in a different way. In 2019, G. V. Milovanović et

Key words and phrases. Jacobi polynomials, Legendre polynomials, Bateman's polynomials, differential operators.

2020 *Mathematics Subject Classification.* Primary: 33C45, 33C47. Secondary: 11B37.

DOI 10.46793/KgJMat2403.383M

Received: September 21, 2020.

Accepted: April 28, 2021.

al. [12] presented the study of various recurrence relations, generating functions and series expansion formulas for two families of orthogonal polynomials in two-variables. Motivated by these two studies, we present here some recurrence relations of two-variables orthogonal polynomials via differential operators.

The generalized hypergeometric function [14, p. 42–43] can be defined as

$${}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} z \right] = \sum_{n=0}^{\infty} \frac{(\alpha_1)_n \cdots (\alpha_p)_n}{(\beta_1)_n \cdots (\beta_q)_n} \cdot \frac{z^n}{n!},$$

with certain convergence conditions given in [14, p. 43].

The Pochhammer symbol $(\lambda)_\nu$ ($\lambda, \nu \in \mathbb{C}$) [13, p. 22, (1)], is defined by

$$(\lambda)_\nu := \frac{\Gamma(\lambda + \nu)}{\Gamma(\lambda)} = \begin{cases} 1, & \nu = 0, \lambda \in \mathbb{C} \setminus \{0\}, \\ \lambda(\lambda + 1) \cdots (\lambda + \nu - 1), & \nu = n \in \mathbb{N}, \lambda \in \mathbb{C}, \end{cases}$$

being understood *conventionally* that $(0)_0 = 1$ and assumed *tacitly* that the Γ quotient exists.

The classical Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ of degree n , $n = 0, 1, 2, \dots$, [13, p. 254 (1)] is defined as

$$P_n^{(\alpha, \beta)}(x) = \frac{(1 + \alpha)_n}{n!} {}_2F_1 \left(-n, 1 + \alpha + \beta + n; 1 + \alpha; \frac{1 - x}{2} \right), \\ \operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, x \in (-1, 1).$$

The generating function of the Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ of degree n [13, p. 270, (2)] is defined by

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x) t^n = F_4 \left(1 + \beta, 1 + \alpha; 1 + \alpha, 1 + \beta; \frac{1}{2}t(x - 1), \frac{1}{2}t(x + 1) \right),$$

where

$$F_4 \left(1 + \beta, 1 + \alpha; 1 + \alpha, 1 + \beta; \frac{1}{2}t(x - 1), \frac{1}{2}t(x + 1) \right),$$

is an Appell polynomial [14, p. 53, (7)].

An elementary generating function of the Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ [13, p. 271, (6)] can be presented in the form

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x) t^n = \rho^{-1} \left(\frac{2}{1 + t + \rho} \right)^\beta \left(\frac{2}{1 - t + \rho} \right)^\alpha$$

or

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x) t^n = 2^{\alpha + \beta} \rho^{-1} (1 + t + \rho)^{-\beta} (1 - t + \rho)^{-\alpha},$$

where $\rho = (1 - 2xt + t^2)^{\frac{1}{2}}$ and on setting $\alpha = \beta = 0$, the Jacobi polynomial reduce to the Legendre polynomial.

Recently, R. Khan et al. [8] introduced generalization of two-variable Jacobi polynomial

$$(1.1) \quad P_n^{(\alpha, \beta)}(x, y) = \sum_{k=0}^n \frac{(1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n - k)!(1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{x - \sqrt{y}}{2} \right)^k, \\ n = 0, 1, 2, \dots, \operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, x, y \in (-1, 1),$$

which can be presented in the alternate form

$$P_n^{(\alpha, \beta)}(x, y) = \sum_{n, k=0}^{\infty} \frac{(1 + \alpha)_n (1 + \beta)_n}{k!(n - k)!(1 + \alpha)_k (1 + \beta)_{n-k}} \left(\frac{x - \sqrt{y}}{2} \right)^k \left(\frac{x + \sqrt{y}}{2} \right)^{n-k}$$

and

$$P_n^{(\alpha, \beta)}(x, y) = \frac{(1 + \alpha)_n}{n!} \left(\frac{x + \sqrt{y}}{2} \right)^n {}_2F_1 \left(-n, -\beta - n; 1 + \alpha; \frac{x - \sqrt{y}}{x + \sqrt{y}} \right)$$

or

$$P_n^{(\alpha, \beta)}(x, y) = \frac{(1 + \alpha)_n}{n!} {}_2F_1 \left(-n, 1 + \alpha + \beta + n; 1 + \alpha; \frac{\sqrt{y} - x}{2} \right).$$

The generating functions of generalized Jacobi polynomial of two-variables $P_n^{(\alpha, \beta)}(x, y)$ [8] can be presented as follows

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x, y) t^n = \mu^{-1} \left(\frac{2}{1 + \sqrt{y}t + \mu} \right)^{\beta} \left(\frac{2}{1 - \sqrt{y}t + \mu} \right)^{\alpha}$$

or

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x, y) t^n = 2^{\alpha+\beta} \mu^{-1} (1 + \sqrt{y}t + \mu)^{-\beta} (1 - \sqrt{y}t + \mu)^{-\alpha},$$

where $\mu = (1 - 2xt + y t^2)^{\frac{1}{2}}$.

In another way, the generating function of generalized Jacobi polynomials of two variables $P_n^{(\alpha, \beta)}(x, y)$ [8] can be presented as follows

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x, y) t^n = F_4 \left(1 + \beta, 1 + \alpha; 1 + \alpha, 1 + \beta; \frac{1}{2}t(x - \sqrt{y}), \frac{1}{2}t(x + \sqrt{y}) \right),$$

which can be written in the form

$$\sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x, y) t^n = \sum_{n, k=0}^{\infty} \frac{(1 + \alpha)_{n+k} (1 + \beta)_{n+k} \frac{1}{2} (x - \sqrt{y})^k \frac{1}{2} (x + \sqrt{y})^n t^n}{k!n! (1 + \alpha)_k (1 + \beta)_n}.$$

Bateman's generating function for $P_n^{(\alpha, \beta)}(x, y)$ [8] can be presented as follows

$$B_n^{(\alpha, \beta)}(x, y; t) = \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2} (x - \sqrt{y})^n t^n}{n! (1 + \alpha)_n} \right] \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2} (x + \sqrt{y})^n t^n}{n! (1 + \beta)_n} \right], \\ \operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |x| < 1, |y| < 1,$$

where

$$B_n^{(\alpha, \beta)}(x, y; t) = \sum_{n=0}^{\infty} \frac{P_n^{(\alpha, \beta)}(x, y) t^n}{(1 + \alpha)_n (1 + \beta)_n}.$$

The generalized Jacobi polynomial of two-variables $P_n^{(\alpha, \beta)}(x, y)$ reduces to the Legendre polynomial of two variables $P_n(x, y)$ for $\alpha = \beta = 0$ in (1.1)

$$P_n(x, y) = \sum_{k=0}^n \frac{(n+k)!}{(k!)^2 (n-k)!} \left(\frac{x - \sqrt{y}}{2} \right)^k,$$

and its generating function can be given by

$$\sum_{n=0}^{\infty} P_n(x, y) t^n = (1 - 2xt + yt^2)^{-\frac{1}{2}}.$$

Also, Khan and Abukhammash [6] defined the Legendre Polynomials of two-variables $P_n(x, y)$ as

$$P_n(x, y) = \sum_{k=0}^{[n/2]} \frac{(-y)^k \left(\frac{1}{2}\right)_{n-k} (2x)^{n-2k}}{k!(n-k)!}$$

and the generating function for $P_n(x, y)$ is given by

$$\sum_{k=0}^n P_n(x, y) t^n = (1 - 2xt + y t^2)^{\frac{1}{2}}.$$

2. RECURRENCE RELATIONS FOR JACOBI POLYNOMIALS

In this section, we will study the action of the following differential operator

$$(2.1) \quad \Xi = \left(\frac{\partial}{\partial z} + \sqrt{w} \frac{\partial}{\partial w} \right)$$

on complex bivariate Jacobi polynomial $P_n^{(\alpha, \beta)}(z, w)$ (2.2) to obtain the desired results.

Now, we present complex bivariate Jacobi polynomial by replacing $x, y \in \mathbb{R}$ by $z, w \in \mathbb{C}$ such that

$$(2.2) \quad P_n^{(\alpha, \beta)}(z, w) = \sum_{k=0}^n \frac{(1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n-k)! (1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{z - \sqrt{w}}{2} \right)^k,$$

$$\operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |z| < 1, |w| < 1.$$

Following conjugate relations will be use frequently in the paper.

$$\begin{aligned} (1 + \alpha + \beta)_{n+k+1} &= (1 + \alpha + \beta) (2 + \alpha + \beta) [1 + (1 + \alpha) + (1 + \beta)]_{(n-1)+k}, \\ (1 + \alpha)_{k+1} &= (1 + \alpha) (1 + (1 + \alpha))_k, \\ (1 + \alpha)_n &= (1 + \alpha) (1 + (1 + \alpha))_{n-1}, \\ (1 + \alpha + \beta)_{n+1} &= (1 + \alpha + \beta)_n (1 + \alpha + \beta + n). \end{aligned}$$

Theorem 2.1. *Following recurrence relation for the Jacobi Polynomial $P_n^{(\alpha,\beta)}(z, w)$, it holds true*

$$\begin{aligned} & \frac{\partial}{\partial z} P_n^{(\alpha,\beta)}(z, w) + \sqrt{w} \frac{\partial}{\partial w} P_n^{(\alpha,\beta)}(z, w) \\ & - \frac{(1 + \alpha + \beta + n)}{4} \left(\frac{z - \sqrt{w}}{2} \right) P_{n-1}^{(1+\alpha),(1+\beta)}(z, w) = 0, \\ & \text{Re}(\alpha) > -1, \text{Re}(\beta) > -1, |z| < 1, |w| < 1. \end{aligned}$$

Proof. On applying the operator (2.1) in (2.2), we get

$$\begin{aligned} & \left(\frac{\partial}{\partial z} + \sqrt{w} \frac{\partial}{\partial w} \right) P_n^{(\alpha,\beta)}(z, w) \\ & = \left(\frac{\partial}{\partial z} + \sqrt{w} \frac{\partial}{\partial w} \right) \sum_{k=0}^n \frac{(1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n - k)! (1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{z - \sqrt{w}}{2} \right)^k \\ & = \sum_{k=0}^n \frac{k (1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n - k)! (1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{z - \sqrt{w}}{2} \right)^{k-1} \left(\frac{1}{2} - \frac{1}{4} \right) \\ & = \frac{1}{4} \sum_{k=0}^n \frac{k (1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n - k)! (1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{z - \sqrt{w}}{2} \right)^{k-1}. \end{aligned}$$

Now, on replacing $k \rightarrow k + 1$ and simplifications, we get

$$\begin{aligned} & \frac{1}{4} \sum_{k=0}^n \frac{(1 + \alpha)_n (1 + \alpha + \beta)_{n+k+1}}{k! [n - (k + 1)]! (1 + \alpha)_{k+1} (1 + \alpha + \beta)_n} \left(\frac{z - \sqrt{w}}{2} \right)^k \\ & = \frac{1}{4} \sum_{k=0}^n \frac{(1 + \alpha) (1 + (1 + \alpha))_{n-1} (1 + \alpha + \beta) (2 + \alpha + \beta)}{k! ((n - 1) - k)! (1 + \alpha) (1 + (1 + \alpha))_k} \\ & \quad \times \frac{(1 + (1 + \alpha) + (1 + \beta))_{(n-1)+k} (1 + \alpha + \beta + n)}{(1 + (1 + \alpha) + (1 + \beta))_{(n-1)} (1 + \alpha + \beta) (2 + \alpha + \beta)} \left(\frac{z - \sqrt{w}}{2} \right)^k \\ & = \frac{1 + (\alpha + \beta + n)}{4} \left(\frac{z - \sqrt{w}}{2} \right) \\ & \quad \times \sum_{k=0}^n \frac{(1 + (1 + \alpha))_{n-1} [1 + (1 + \alpha) + (1 + \beta)]_{(n-1)+k}}{k! ((n - 1) - k)! (1 + (1 + \alpha))_k [1 + (1 + \alpha) + (1 + \beta)]_{n-1}} \left(\frac{z - \sqrt{w}}{2} \right)^{k-1} \\ & = \frac{(1 + \alpha + \beta + n)}{4} \left(\frac{z - \sqrt{w}}{2} \right) P_{n-1}^{(1+\alpha),(1+\beta)}(z, w). \end{aligned}$$

Therefore, we get the desired result. □

Corollary 2.1. *Following recurrence relation for the Jacobi Polynomial $P_n^{(\alpha,\beta)}(z, 1)$, it holds true*

$$\frac{\partial}{\partial z} P_n^{(\alpha,\beta)}(z, 1) - \frac{(1 + \alpha + \beta + n)}{2} \left(\frac{z - 1}{2} \right) P_{n-1}^{(1+\alpha),(1+\beta)}(z, 1) = 0,$$

$$\operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |z| < 1.$$

Proof. First, put $w = 1$ in (2.2) we consider the Jacobi polynomials

$$P_n^{(\alpha,\beta)}(z, 1) = \sum_{k=0}^n \frac{(1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n - k)!(1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{z - 1}{2}\right)^k.$$

Taking differential operator $\Xi_z = \left(\frac{\partial}{\partial z}\right)$ then following the same process used in the above theorem leads to the desired result. \square

Corollary 2.2. *Following recurrence relation for the Jacobi Polynomial $P_n^{(\alpha,\beta)}(1, w)$, it holds true*

$$\left(\sqrt{w} \frac{\partial}{\partial w}\right) P_n^{(\alpha,\beta)}(1, w) + \frac{(1 + \alpha + \beta + n)}{4} \left(\frac{1 - \sqrt{w}}{2}\right) P_{n-1}^{(1+\alpha),(1+\beta)}(1, w) = 0,$$

$$\operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |w| < 1.$$

Proof. By putting $z = 1$ in (2.2) we get

$$P_n^{(\alpha,\beta)}(1, w) = \sum_{k=0}^n \frac{(1 + \alpha)_n (1 + \alpha + \beta)_{n+k}}{k!(n - k)!(1 + \alpha)_k (1 + \alpha + \beta)_n} \left(\frac{1 - \sqrt{w}}{2}\right)^k.$$

Taking differential operator $\Xi_w = \left(\sqrt{w} \frac{\partial}{\partial w}\right)$ then following the same process used in the above theorem leads to the desired result. \square

3. RECURRENCE RELATIONS FOR BATEMAN’S GENERATING FUNCTION

Now, we present complex bivariate Bateman’s generating function by replacing $x, y \in \mathbb{R}$ by $z, w \in \mathbb{C}$ such that

$$(3.1) \quad B_n^{(\alpha,\beta)}(z, w; t) = \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z - \sqrt{w})^n t^n}{n!(1 + \alpha)_n} \right] \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z + \sqrt{w})^n t^n}{n!(1 + \beta)_n} \right]$$

and

$$B_n^{(\alpha,\beta)}(z, w; t) = \sum_{n=0}^{\infty} \frac{P_n^{(\alpha,\beta)}(z, w) t^n}{(1 + \alpha)_n (1 + \beta)_n}, \quad \operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |z| < 1, |w| < 1.$$

We can also write the conjugate relationships for the purpose to use these relations in this section

$$(1 + \alpha)_{n+1} = (1 + \alpha) (1 + (1 + \alpha))_n,$$

$$(1 + \beta)_{n+1} = (1 + \beta) (1 + (1 + \beta))_n.$$

Theorem 3.1. *Following recurrence relation for the Bateman’s generating function $B_n^{(\alpha,\beta)}(z, w; t)$, it holds true*

$$\frac{\partial}{\partial z} B_n^{(\alpha,\beta)}(z, w; t) + \sqrt{w} \frac{\partial}{\partial w} B_n^{(\alpha,\beta)}(z, w; t) - \frac{t}{2(1 + \alpha)} B_n^{[(1+\alpha),\beta]}(z, w; t)$$

$$- \frac{3t}{2(1 + \beta)} B_n^{[\alpha,(1+\beta)]}(z, w; t) = 0, \quad \operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |z| < 1, |w| < 1.$$

Proof. Using the differential operator Ξ for the Bateman's generating function $B_n^{(\alpha,\beta)}(z, w; t)$, we see that

$$\begin{aligned} & \Xi B_n^{(\alpha,\beta)}(z, w; t) \\ &= \left(\frac{\partial}{\partial z} + \sqrt{w} \frac{\partial}{\partial w} \right) \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z - \sqrt{w})^n t^n}{n! (1 + \alpha)_n} \right] \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z + \sqrt{w})^n t^n}{n! (1 + \beta)_n} \right] \\ &= \frac{t}{2(1 + \alpha)} \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z - \sqrt{w})^n t^n}{n! [1 + (1 + \alpha)]_n} \right] \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z + \sqrt{w})^n t^n}{n! (1 + \beta)_n} \right] \\ &\quad + \frac{3t}{2(1 + \beta)} \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z - \sqrt{w})^n t^n}{n! (1 + \alpha)_n} \right] \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(z + \sqrt{w})^n t^n}{n! [1 + (1 + \beta)]_n} \right] \\ &= \frac{t}{2(1 + \alpha)} B_n^{[(1+\alpha),\beta]}(z, w; t) - \frac{3t}{2(1 + \beta)} B_n^{[\alpha,(1+\beta)]}(z, w; t). \end{aligned}$$

Therefore, we get the desired result. □

Corollary 3.1. *Following recurrence relation for the Bateman's generating function $B_n^{(\alpha,\beta)}(z, 1)$, it holds true*

$$\begin{aligned} & \frac{\partial}{\partial z} B_n^{(\alpha,\beta)}(z, 1; t) - \frac{t}{(1 + \alpha)} B_n^{[(1+\alpha),\beta]}(z, 1; t) - \frac{t}{(1 + \beta)} B_n^{[\alpha,(1+\beta)]}(z, 1; t) = 0, \\ & \text{Re}(\alpha) > -1, \text{Re}(\beta) > -1, |z| < 1. \end{aligned}$$

Proof. First, substitute $w=1$ in the Bateman's generating function (3.1), we have

$$B_n^{(\alpha,\beta)}(z, 1; t) = \sum_{n=0}^{\infty} \frac{\frac{1}{2}(z - 1)^n t^n}{n! (1 + \alpha)_n} \sum_{n=0}^{\infty} \frac{\frac{1}{2}(z + 1)^n t^n}{n! (1 + \beta)_n}.$$

Taking differential operator $\Xi_z = \left(\frac{\partial}{\partial z} \right)$ then following the same process used in the above theorem leads to the desired result. □

Corollary 3.2. *Following recurrence relation for the Bateman's generating function $B_n^{(\alpha,\beta)}(1, w)$, it holds true*

$$\begin{aligned} & \sqrt{w} \frac{\partial}{\partial w} B_n^{(\alpha,\beta)}(1, w; t) + \frac{t}{2(1 + \alpha)} B_n^{[(1+\alpha),\beta]}(1, w; t) - \frac{t}{2(1 + \beta)} B_n^{[\alpha,(1+\beta)]}(1, w; t) = 0, \\ & \text{Re}(\alpha) > -1, \text{Re}(\beta) > -1, |w| < 1. \end{aligned}$$

Proof. Put $z = 1$ in the Bateman's generating function (3.1), we get

$$B_n^{(\alpha,\beta)}(1, w; t) = \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(1 - \sqrt{w})^n t^n}{n! (1 + \alpha)_n} \right] \left[\sum_{n=0}^{\infty} \frac{\frac{1}{2}(1 + \sqrt{w})^n t^n}{n! (1 + \beta)_n} \right].$$

Taking differential operator $\Xi_w = \left(\sqrt{w} \frac{\partial}{\partial w} \right)$ then following the same process used in the above theorem leads to the desired result. □

4. RECURRENCE RELATIONS FOR LEGENDRE POLYNOMIALS

In this sections, we will study the action of the following differential operator

$$\Delta = \left(\frac{1}{w} \frac{\partial}{\partial z} + \frac{1}{z} \frac{\partial}{\partial w} \right),$$

on complex bivariate Legendre polynomial $P_n(z, w)$ (2.2) to obtain the desired results.

Now, we present complex bivariate Legendre polynomial by replacing $x, y \in \mathbb{R}$ by $z, w \in \mathbb{C}$ such that

$$(4.1) \quad P_n(z, w) = \sum_{k=0}^{[n/2]} \frac{(-w)^k \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k}}{k!(n-k)!},$$

where $\text{Re}(\alpha) > -1, \text{Re}(\beta) > -1, |z| < 1, |w| < 1$.

Theorem 4.1. *Following recurrence relation for the Legendre polynomials $P_n(z, w)$, it holds true*

$$\frac{1}{w} \frac{\partial}{\partial z} P_n(z, w) + \left(\frac{1}{z} \frac{\partial}{\partial w} \right) P_n(z, w) - \left(\frac{n}{zw} \right) P_n(z, w) + \left(\frac{1}{2z^2} \right) P_{n-1}(z, w) = 0,$$

$\text{Re}(\alpha) > -1, \text{Re}(\beta) > -1, |z| < 1, |w| < 1$.

Proof. For Legendre polynomials (4.1) of two variables $P_n(z, w)$ we see that

$$\Delta P_n(z, w) = \left(\frac{1}{w} \frac{\partial}{\partial z} + \frac{1}{z} \frac{\partial}{\partial w} \right) \sum_{k=0}^{[n/2]} \frac{(-w)^k \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k}}{k!(n-k)!}.$$

After applying the differential operator, we get

$$\begin{aligned} & -2 \sum_{k=0}^{[n/2]} \frac{(-w)^{k-1} \left(\frac{1}{2}\right)_{n-k} (n-2k) (2z)^{n-2k-1}}{k!(n-k)!} - 2 \sum_{k=0}^{[n/2]} \frac{k (-w)^{k-1} \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k-1}}{k!(n-k)!} \\ &= -2 \sum_{k=0}^{[n/2]} \frac{(n-k) (-w)^{k-1} \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k-1}}{k!(n-k)!}. \\ &= -2n \sum_{k=0}^{[n/2]} \frac{(-w)^{k-1} \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k-1}}{k!(n-k)!} - 2 \sum_{k=0}^{[n/2]} \frac{k (-w)^{k-1} \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k-1}}{k!(n-k)!} \\ &= \left(\frac{n}{zw} \right) \sum_{k=0}^{[n/2]} \frac{(-w)^k \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k}}{k!(n-k)!} \\ & \quad - 2 \left(\frac{1}{2z} \right)^2 \sum_{k=0}^{[n/2]} \frac{(-w)^k \left(\frac{1}{2}\right)_{(n-1)-k} (2z)^{[(n-1)-2(k-1)]}}{k! [(n-1)-k]!}. \\ &= \left(\frac{n}{zw} \right) P_n(z, w) - \left(\frac{1}{2z^2} \right) P_{n-1}(z, w). \end{aligned}$$

Now, on some simplification, we get our desired result. □

Corollary 4.1. *Following recurrence relation for the Legendre polynomials $P_n(z, 1)$, it holds true*

$$\frac{1}{w} \frac{\partial}{\partial z} P_n(z, 1) - \left(\frac{n}{z}\right) P_n(z, 1) + \left(\frac{1}{2z^2}\right) P_{n-1}(z, 1) = 0,$$

$$\operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |z| < 1.$$

Proof. First, substitute $w = 1$ in equation (4.1) we get

$$P_n(z, 1) = \sum_{k=0}^{[n/2]} \frac{(-1)^k \left(\frac{1}{2}\right)_{n-k} (2z)^{n-2k}}{k!(n-k)!}.$$

Taking differential operator $\Delta_z = \left(\frac{1}{w} \frac{\partial}{\partial z}\right)$ then following the same process used in the above theorem leads to the desired result. \square

Corollary 4.2. *Following recurrence relation for the Legendre polynomials $P_n(1, w)$, it holds true*

$$\frac{1}{z} \frac{\partial}{\partial w} P_n(1, w) - \left(\frac{n}{w}\right) P_n(1, w) + \left(\frac{1}{2}\right) P_{n-1}(1, w) = 0,$$

$$\operatorname{Re}(\alpha) > -1, \operatorname{Re}(\beta) > -1, |w| < 1.$$

Proof. Put $z = 1$ in equation (4.1) we get

$$P_n(1, w) = \sum_{k=0}^{[n/2]} \frac{(-w)^k \left(\frac{1}{2}\right)_{n-k} (2)^{n-2k}}{k!(n-k)!}.$$

Taking differential operator $\Delta_w = \left(\frac{1}{z} \frac{\partial}{\partial w}\right)$ then following the same process used in the above theorem leads to the desired result. \square

Acknowledgements. The authors are highly thankful to the editor and anonymous referees for their efforts which could shape the manuscript in the present form.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Fathematical Functions*, National Bureau of Standards, New York, 1970.
- [2] S. A. Agahanov, *A method of constructing orthogonal polynomials of two variables for a certain class of weight functions* (Russian), Vestnik Leningrad University **20** (1965), 5–10.
- [3] G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999. <https://doi.org/10.1017/CB09781107325937>.
- [4] E. H. Doha, W. M. Abd-elhameed and H. M. Ahmed, *The coefficients of differentiated expansions of double and triple Jacobi polynomials*, Bull. Iran. Math. Soc. **38**(3) (2012), 739–765.
- [5] C. F. Dunkl and Y. Xu, *Orthogonal Polynomials of Several Variables*, Cambridge University Press, Cambridge, 2001. <https://doi.org/10.1017/CB09781107786134>
- [6] M. A. Khan and G. S. Abukhamash, *On a new class of polynomial set suggested by Legendre polynomials*, Acta Ciencia Indica **4**(857) (2003), XXIXM.

- [7] M. A. Khan, A. H. Khan and S. M. Abbas, *A note on pseudo two variables Jacobi polynomials*, Ain Shams Engineering Journal **3** (2013), 127–131. <https://doi.org/10.1016/j.asej.2012.06.003>
- [8] R. Khan, N. Kumar and R. Qamar, *Two variables generalization of Jacobi polynomials*, Global Journal of Pure and Applied Mathematics **13**(5) (2017), 1387–1399.
- [9] T. H. Koornwinder, *Two-variable analogues of the classical orthogonal polynomials*, in: R. A. Askey (Ed.), *Theory and Application of Special Functions*, Proceedings of an Advanced Seminar Sponsored by the Mathematics Research Center, The University of Wisconsin-Madison, Academic Press, New York, 1975, 435–495. <https://doi.org/10.1016/B978-0-12-064850-4.50015-X>
- [10] F. Marcellan, S. Jabee and M. Shadab, *Analytic properties of Touchard based hybrid polynomials via operational techniques*, Bull. Malaysian Math. Sci. Soc. **44** (2020), 223–242. <https://doi.org/10.1007/s40840-020-00945-4>
- [11] M. Marriaga, T. E. Perez and M. A. Pinar, *Three term relations for a class of bivariate orthogonal polynomials*, Mediterr. J. Math. **14**(54) (2017), 25 pages. <https://doi.org/10.1007/s00009-017-0859-0>
- [12] G. V. Milovanović, G. Ozturk and R. Aktas, *Properties of some of two-variable orthogonal polynomials*, Bull. Malaysian Math. Sci. Soc. **43** (2019), 1403–1431. <https://doi.org/10.1007/s40840-019-00750-8>
- [13] E. D. Rainville, *Special Functions*, The Macmillan Co. Inc., New York, 1960, reprinted by Chelsea Publ. Co. Bronx, New York, 1971.
- [14] H. M. Srivastava and H. L. Manocha, *A Treatise on Generating functions*, Halsted Press (Ellis Horwood Ltd., Chichester, U.K.), John Wiley and Sons, New York, Chichester, Brisbane, Toronto, 1984. <https://doi.org/10.1137/1028045>.
- [15] G. Szego, *Orthogonal Polynomials*, 4th Edn., Colloquium Publications **23**, Amer. Math. Soc., Providence, 1978.
- [16] M. I. Qureshi, M. Shadab and M. S. Baboo, *Evaluation of some novel integrals involving Legendre function of second kind using hypergeometric approach*, Palestine J. Math. **6**(1) (2017), 68–75.

¹DEPARTEMENT OF MATHEMATICS,
FACULTY OF SCIENCE, SOUTH VALLEY UNIVERSITY,
(QENA-EGYPT)
Email address: mosaed_makky11@yahoo.com, mosaed_makky@sci.svu.edu.eg

²DEPARTMENT OF NATURAL AND APPLIED SCIENCES,
SCHOOL OF SCIENCE AND TECHNOLOGY, GLOCAL UNIVERSITY,
SAHARANPUR 247121, INDIA
Email address: shadabmohd786@gmail.com