

LOGARITHMICALLY COMPLETE MONOTONICITY OF RECIPROCAL ARCTAN FUNCTION

VLADIMIR JOVANOVIĆ¹ AND MILANKA TREML¹

ABSTRACT. We prove the conjecture stated in F. Qi and R. Agarwal, *On complete monotonicity for several classes of functions related to ratios of gamma functions*, J. Inequal. Appl. (2019), that the function $1/\arctan$ is logarithmically completely monotonic on $(0, \infty)$, but not a Stieltjes transform.

1. INTRODUCTION

By a *completely monotonic function* (shortly CM) we mean here an infinitely differentiable function $f : (0, \infty) \rightarrow \mathbb{R}$, such that

$$(-1)^n f^{(n)} \geq 0, \quad n = 0, 1, 2, \dots$$

If f' is completely monotonic and $f \geq 0$, then we call f a *Bernstein function*. Here we are mostly interested in *logarithmically completely monotonic functions*, that is, infinitely differentiable functions $f : (0, \infty) \rightarrow (0, \infty)$ with the property

$$(-1)^n (\log f)^{(n)} \geq 0, \quad n = 1, 2, 3, \dots$$

A basic fact concerning CM - functions is the Bernstein theorem: a function f is CM if and only if there exists a non-decreasing function α on $(0, \infty)$ satisfying

$$f(x) = \int_0^\infty e^{-xt} d\alpha(t),$$

for all $x > 0$ (see [9, p. 161]). In some occasions it has been proven a stronger property which leads to complete monotonicity of a function f , namely that there exist $a \geq 0$

Key words and phrases. Complete monotonicity, Stieltjes transform.

2020 *Mathematics Subject Classification.* Primary: 26A48. Secondary: 30E20.

DOI 10.46793/KgJMat2501.105J

Received: November 29, 2021.

Accepted: December 27, 2021.

and a non-negative Borel measure μ on $[0, \infty)$ for which the equality

$$f(x) = a + \int_0^\infty \frac{d\mu(t)}{x+t}$$

holds for $x > 0$, where the measure μ fulfills the condition

$$\int_0^\infty \frac{d\mu(t)}{1+t} < \infty.$$

Such functions are called *Stieltjes transforms*. We recall that all Stieltjes transforms are logarithmically completely monotonic (see [2] and further generalizations [3]), and the latter are CM (see [5], but also [7] and [8]).

In [6] the authors set the conjecture that the function $f(x) = \frac{1}{\arctan x}$ is logarithmically completely monotonic on $(0, \infty)$, but not a Stieltjes transform. The aim of this paper is to justify these assertions. We will do it in the next section.

2. FORMULATIONS AND PROOFS

Theorem 2.1. *The function $f(x) = \frac{1}{\arctan x}$ is logarithmically completely monotonic on $(0, \infty)$.*

The idea of the proof of Theorem 2.1 is based on the Remark 1 in [1], where the authors suggest employing the residue theorem in an attempt to obtain integral representations of functions under consideration.

Proof. It suffices to prove that

$$g(x) = -(\log f(x))' = \frac{1}{(x^2 + 1) \arctan x}$$

is CM on $(0, \infty)$. In what follows we always assume that \log denotes the principle value of logarithm, i.e., $\log z = \ln |z| + i \arg z$, with $\arg z \in (-\pi, \pi]$.

Let us consider the integral $\int_{\Gamma_{R,r}} G(z) dz$, over the "keyhole" contour $\Gamma_{R,r}$ given in Figure 1, where

$$G(z) = \frac{z+1}{z(z-z_0)\log z}$$


and $z_0 = \frac{i-x}{i+x}$ for $x > 0$.

We assume $R > 1$ and $r < 1$. Note that $|z_0| = 1$ and that 1, z_0 are the only singularities of G lying inside $\Gamma_{R,r}$. From the residue theorem, we have

$$\int_{\Gamma_{R,r}} G(z) dz = 2\pi i (\text{Res}(G(z); z_0) + \text{Res}(G(z); 1)).$$

Since z_0 is a first-order pole, it follows

$$\text{Res}(G(z); z_0) = \frac{1+z_0}{z_0 \log z_0} = \frac{1 + \frac{i-x}{i+x}}{\frac{i-x}{i+x} \log \frac{i-x}{i+x}} = \frac{2i}{(i-x)2i \arctan x} = -\frac{(i+x)}{(x^2+1) \arctan x},$$


 FIGURE 1. Keyhole contour $\Gamma_{R,r}$

where we used the fact that $\arctan x = \frac{1}{2i} \log \frac{1+ix}{1-ix}$, for $x > 0$. Similarly,

$$\operatorname{Res}(G(z); 1) = \lim_{z \rightarrow 1} (z-1) \frac{1+z}{z(z-z_0) \log z} = \frac{2}{1-z_0} = \frac{2}{1 - \frac{i-x}{i+x}} = \frac{i+x}{x},$$

whence,

$$(2.1) \quad g(x) = \frac{1}{x} - \frac{1}{2\pi i(x+i)} \int_{\Gamma_{R,r}} G(z) dz.$$

Now, it remains to calculate the integral $\int_{\Gamma_{R,r}} G(z) dz$. In order to accomplish it, we start from the relation

$$(2.2) \quad \int_{\Gamma_{R,r}} G(z) dz = \int_{\Gamma_R} G(z) dz + \int_{\Gamma_r} G(z) dz + \int_{\Gamma_{R,r}^+} G(z) dz + \int_{\Gamma_{R,r}^-} G(z) dz.$$

The first two integrals vanish as $R \rightarrow \infty$ and $r \rightarrow 0+$. It follows from the estimates

$$\left| \int_{\Gamma_R} G(z) dz \right| \leq 2R\pi \max_{|z|=R} \frac{|z+1|}{|z| |\log z| |z-z_0|} \leq 2\pi \frac{R+1}{(\ln R - 2\pi)(R-1)}$$

and

$$\left| \int_{\Gamma_r} G(z) dz \right| \leq 2r\pi \max_{|z|=r} \frac{|z+1|}{|z| |\log z| |z-z_0|} \leq 2\pi \frac{1+r}{(-\ln r - 2\pi)(1-r)}.$$

We also have for $t < 0$

$$\lim_{\substack{z \rightarrow t \\ \Im z > 0}} G(z) = \frac{t+1}{t(\ln(-t) + \pi i)(t-z_0)} = G^+(t)$$

and

$$\lim_{\substack{z \rightarrow t \\ \Im z < 0}} G(z) = \frac{t+1}{t(\ln(-t) - \pi i)(t - z_0)} = G^-(t).$$

Consequently,

$$(2.3) \quad \int_{\Gamma_{R,r}^+} G(z) dz + \int_{\Gamma_{R,r}^-} G(z) dz = \int_{-R}^{-r} [G^+(t) - G^-(t)] dt.$$

Let us denote $I = \lim_{\substack{R \rightarrow \infty \\ r \rightarrow 0^+}} \int_{\Gamma_{R,r}} G(z) dz$. From (2.2) and (2.3) we obtain

$$\begin{aligned} I &= \int_{-\infty}^0 [G^+(t) - G^-(t)] dt \\ &= \int_{-\infty}^0 \frac{2\pi i(t+1) dt}{t(\log^2(-t) + \pi^2)(t - z_0)} \\ &= 2\pi i \int_0^{\infty} \frac{(1-t) dt}{t(\log^2 t + \pi^2)(t + z_0)}. \end{aligned}$$

Using $z_0 = \frac{i-x}{i+x}$, we have

$$\begin{aligned} I &= \int_0^{\infty} \frac{2\pi i(1-t) dt}{t(\log^2 t + \pi^2)(t + \frac{i-x}{i+x})} \\ &= \int_0^{\infty} \frac{2\pi i(i+x)(1-t) dt}{t(\log^2 t + \pi^2)(x(t-1) + i(t+1))} \\ &= -2\pi i(i+x) \int_0^{\infty} \frac{((1-t)^2 x + i(1-t^2)) dt}{t(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)}. \end{aligned}$$

Note that (2.1) implies

$$(2.4) \quad g(x) = \frac{1}{x} - \frac{1}{2\pi i(x+i)} I$$

and since $\frac{1}{2\pi i(x+i)} I$ is real, we conclude that

$$\int_0^{\infty} \frac{(1-t^2) dt}{t(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)} = 0.$$

Therefore, from (2.4), it follows

$$(2.5) \quad g(x) = \frac{1}{x} + \int_0^{\infty} \frac{(1-t)^2 x dt}{t(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)}.$$

Employing

$$\frac{1}{x} = \int_0^{\infty} \frac{dt}{xt(\log^2 t + \pi^2)},$$

we get

$$g(x) = \int_0^{\infty} \frac{(2(1-t)^2 x + (1+t)^2) dt}{xt(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)}.$$

The substitution $t \mapsto \frac{1}{t}$ implies

$$\int_0^1 \frac{(2(1-t)^2x + (1+t)^2) dt}{xt(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)} = \int_1^\infty \frac{(2(1-t)^2x + (1+t)^2) dt}{xt(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)}.$$

Hence,

$$(2.6) \quad g(x) = 2 \int_0^1 \frac{(2(1-t)^2x + (1+t)^2) dt}{xt(x^2(1-t)^2 + (1+t)^2)(\log^2 t + \pi^2)}.$$

For $a, b, x > 0$ it is

$$\frac{2a^2x^2 + b^2}{x(a^2x^2 + b^2)} = \frac{1}{x} + \frac{1}{2} \left(\frac{1}{x + \frac{bi}{a}} + \frac{1}{x - \frac{bi}{a}} \right)$$

and using

$$\frac{1}{x} = \int_0^\infty e^{-xs} ds, \quad \frac{1}{x + \frac{bi}{a}} = \int_0^\infty e^{-xs} e^{-\frac{bi}{a}s} ds, \quad \frac{1}{x - \frac{bi}{a}} = \int_0^\infty e^{-xs} e^{\frac{bi}{a}s} ds,$$

one obtains

$$\frac{2a^2x^2 + b^2}{x(a^2x^2 + b^2)} = \int_0^\infty e^{-xs} \left(1 + \cos \frac{bs}{a} \right) ds.$$

Setting $a = 1 - t$ and $b = 1 + t$ yields

$$\frac{2(1-t)^2x + (1+t)^2}{x(x^2(1-t)^2 + (1+t)^2)} = \int_0^\infty e^{-xs} \left(1 + \cos \frac{1+t}{1-t}s \right) ds.$$

From (2.6), we have

$$g(x) = 2 \int_0^1 \left(\int_0^\infty \frac{e^{-xs} (1 + \cos \frac{1+t}{1-t}s) ds}{t(\ln^2 t + \pi^2)} \right) dt,$$

and, finally, after interchanging integration order, we obtain

$$(2.7) \quad g(x) = \int_0^\infty \left(\int_0^1 \frac{2(1 + \cos \frac{1+t}{1-t}s) dt}{t(\ln^2 t + \pi^2)} \right) e^{-xs} ds.$$

Now, it is evident that (2.7) implies complete monotonicity of g . \square

Theorem 2.2. *The function $f(x) = \frac{1}{\arctan x}$ is not a Stieltjes transform on $(0, \infty)$.*

For the proof of this theorem, we use the following result on Stieltjes transforms from [4].

Proposition 2.1. *If $f \neq 0$ is a Stieltjes transform, then $\frac{1}{f}$ is a Bernstein function.*

Proof of Theorem 2.2. The function $h(x) = \frac{1}{f(x)} = \arctan x$ is not a Bernstein function, since

$$h^{(3)}(x) = -2 \frac{3x^2 - 1}{(1 + x^2)^3}$$

changes its sign on $(0, \infty)$. Therefore, according to Proposition 2.1, f is not a Stieltjes transform. \square

REFERENCES

- [1] H. Alzer and C. Berg, *Some classes of completely monotonic functions*, Annales Academiae Scientiarum Fennicae. Mathematica **27** (2002), 445–460.
- [2] C. Berg, *Integral representation of some functions related to the gamma function*, Mediterr. J. Math. **1** (2004), 433–439. <https://doi.org/10.1007/s00009-004-0022-6>
- [3] C. Berg, S. Koumandos and H. L. Pedersen, *Nielsen's beta function and some infinitely divisible distributions*, Math. Nachr. **294** (2021), 426–429. <https://doi.org/10.1002/mana.201900217>
- [4] C. Berg and H. L. Pedersen, *A completely monotone function related to the gamma function*, J. Comput. Appl. Math. **133** (2001), 219–230. [https://doi.org/10.1016/S0377-0427\(00\)00644-0](https://doi.org/10.1016/S0377-0427(00)00644-0)
- [5] R. Horn, *On infinitely divisible matrices, kernels and functions*, Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete **8** (1967), 219–230. <https://doi.org/10.1007/BF00531524>
- [6] F. Qi and R. Agarwal, *On complete monotonicity for several classes of functions related to ratios of gamma functions*, J. Inequal. Appl. (2019), 1–42. <https://doi.org/10.1186/s13660-019-1976-z>
- [7] F. Qi and C.-P. Chen, *A complete monotonicity property of the gamma function*, J. Math. Anal. Appl. **296** (2004), 603–607. <https://doi.org/10.1016/j.jmaa.2004.04.026>
- [8] R. L. Schilling, R. Song and Z. Vondraček, *Bernstein Functions*, De Gruyter, 2012. <https://doi.org/10.1515/9783110269338>
- [9] D. V. Widder, *The Laplace Transform*, Princeton University Press, 1946.

¹FACULTY OF SCIENCES AND MATHEMATICS,
UNIVERSITY OF BANJA LUKA,
MLADENA STOJANOVIĆA 2
BANJA LUKA
REPUBLIC OF SRPSKA
BOSNIA AND HERZEGOVINA
Email address: vladimir.jovanovic@pmf.unibl.org
ORCID iD: <https://orcid.org/0009-0006-7839-8575>
Email address: milanka.treml@pmf.unibl.org