

WEAVING CONTINUOUS CONTROLLED K - g -FUSION FRAMES IN HILBERT SPACES

PRASENJIT GHOSH¹ AND TAPAS K. SAMANTA²

ABSTRACT. We introduce the notion of weaving continuous controlled K - g -fusion frame in Hilbert space. Some characterizations of weaving continuous controlled K - g -fusion frame have been presented. We extend some of the recent results of woven K - g -fusion frame and controlled K - g -fusion frame to woven continuous controlled K - g -fusion frame. Finally, a perturbation result of woven continuous controlled K - g -fusion frame has been studied.

1. INTRODUCTION AND PRELIMINARIES

Duffin and Schaeffer [13] introduced frame for Hilbert space to study some fundamental problems in non-harmonic Fourier series. Later on, after some decades, frame theory was popularized by Daubechies et al. [11]. At present, frame theory has been widely used in signal and image processing, filter bank theory, coding and communications, system modeling and so on.

Let H be a separable Hilbert space associated with the inner product $\langle \cdot, \cdot \rangle$. Frame for Hilbert space was defined as a sequence of basis-like elements in Hilbert space. A sequence $\{f_i\}_{i=1}^{+\infty} \subset H$ is called a frame for H , if there exist positive constants $0 < A \leq B < +\infty$ such that

$$A\|f\|^2 \leq \sum_{i=1}^{+\infty} |\langle f, f_i \rangle|^2 \leq B\|f\|^2, \quad \text{for all } f \in H.$$

The constants A and B are called lower and upper bounds, respectively.

Key words and phrases. Frame, g -fusion frame, continuous g -fusion frame, controlled frame, woven frame.

2020 *Mathematics Subject Classification.* Primary: 42C15. Secondary: 94A12, 46C07.
<https://doi.org/10.46793/KgJMat2601.115G>

Received: June 13, 2022.

Accepted: June 19, 2023.

Throughout this paper, H is considered to be a separable Hilbert space with associated inner product $\langle \cdot, \cdot \rangle$ and \mathbb{H} is the collection of all closed subspaces of H . (X, μ) denotes abstract measure space with positive measure μ . I_H is the identity operator on H . $\mathcal{B}(H_1, H_2)$ is a collection of all bounded linear operators from H_1 to H_2 . In particular, $\mathcal{B}(H)$ denotes the space of all bounded linear operators on H . For $S \in \mathcal{B}(H)$, we denote $\mathcal{N}(S)$ and $\mathcal{R}(S)$ for null space and range of S , respectively. Also, $P_M \in \mathcal{B}(H)$ is the orthonormal projection of H onto a closed subspace $M \subset H$. The set $\mathcal{S}(H)$ of all self-adjoint operators on H is a partially ordered set with respect to the partial order \leq which is defined as for $R, S \in \mathcal{S}(H)$

$$R \leq S \Leftrightarrow \langle Rf, f \rangle \leq \langle Sf, f \rangle, \quad \text{for all } f \in H.$$

$\mathcal{GB}(H)$ denotes the set of all bounded linear operators which have bounded inverse. If $S, R \in \mathcal{GB}(H)$, then R^*, R^{-1} and SR also belongs to $\mathcal{GB}(H)$. An operator $U \in \mathcal{B}(H)$ is called positive if $\langle Uf, f \rangle \geq 0$ for all $f \in H$. In notation, we can write $U \geq 0$. If $V \in \mathcal{B}(H)$ is positive then there exists a unique positive U such that $V^2 = U$. This will be denoted by $V = U^{1/2}$. Moreover, if an operator V commutes with U then V commutes with every operator in the C^* -algebra generated by U and I , specially V commutes with $U^{1/2}$. $\mathcal{GB}^+(H)$ is the set of all positive operators in $\mathcal{GB}(H)$ and T, U are invertible operators in $\mathcal{GB}(H)$. For each $m > 1$, we define $[m] = \{1, 2, \dots, m\}$.

We present some theorems in operator theory which will be needed throughout this paper.

Theorem 1.1 (Douglas' factorization theorem [12]). *Let $S, V \in \mathcal{B}(H)$. Then the following conditions are equivalent.*

- (i) $\mathcal{R}(S) \subseteq \mathcal{R}(V)$.
- (ii) $SS^* \leq \lambda^2 VV^*$ for some $\lambda > 0$.
- (iii) $S = VW$ for some bounded linear operator W on H .

Theorem 1.2 ([15]). *Let $M \subset H$ be a closed subspace and $T \in \mathcal{B}(H)$. Then $P_M T^* = P_M T^* P_{\overline{TM}}$. If T is an unitary operator (i.e., $T^*T = I_H$), then $P_{\overline{TM}} T = T P_M$.*

Theorem 1.3 ([8]). *Let H_1, H_2 be two Hilbert spaces and $U : H_1 \rightarrow H_2$ be a bounded linear operator with closed range \mathcal{R}_U . Then, there exists a bounded linear operator $U^\dagger : H_2 \rightarrow H_1$ such that $UU^\dagger x = x$ for all $x \in \mathcal{R}_U$.*

1.1. K - g -fusion frame. Construction of K - g -fusion frames and their dual were presented by Sadri and Rahimi [1] to generalize the theory of K -frame [16], fusion frame [9], and g -frame [35].

Definition 1.1 ([1]). Let $\{W_j\}_{j \in J}$ be a collection of closed subspaces of H and $\{v_j\}_{j \in J}$ be a collection of positive weights, $\{H_j\}_{j \in J}$ be a sequence of Hilbert spaces. Suppose $\Lambda_j \in \mathcal{B}(H, H_j)$ for each $j \in J$ and $K \in \mathcal{B}(H)$. Then $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$ is called a K - g -fusion frame for H respect to $\{H_j\}_{j \in J}$ if there exist constants $0 < A \leq B < +\infty$

such that

$$A \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B \|f\|^2,$$

for all $f \in H$. The constants A and B are called the lower and upper bounds of K - g -fusion frame, respectively. If $K = I_H$ then the family is called g -fusion frame and it has been widely studied in [18–20, 31].

Define the space

$$\ell^2(\{H_j\}_{j \in J}) = \left\{ \{f_j\}_{j \in J} : f_j \in H_j, \sum_{j \in J} \|f_j\|^2 < +\infty \right\},$$

with inner product given by

$$\langle \{f_j\}_{j \in J}, \{g_j\}_{j \in J} \rangle = \sum_{j \in J} \langle f_j, g_j \rangle_{H_j}.$$

Clearly, $\ell^2(\{H_j\}_{j \in J})$ is a Hilbert space with the pointwise operations [1].

1.2. Controlled K - g -fusion frame. Controlled frame is one of the newest generalization of frame. P. Balaz et al. [6] introduced controlled frame to improve the numerical efficiency of interactive algorithms for inverting the frame operator. In recent times, several generalizations of controlled frame namely, controlled K -frame [26], controlled g -frame [27], controlled fusion frame [23], controlled g -fusion frame [34], controlled K - g -fusion frame [28] etc. have been appeared.

Definition 1.2 ([28]). Let $K \in \mathcal{B}(H)$ and $\{W_j\}_{j \in J}$ be a collection of closed subspaces of H and $\{v_j\}_{j \in J}$ be a collection of positive weights. Let $\{H_j\}_{j \in J}$ be a sequence of Hilbert spaces, $T, U \in \mathcal{GB}(H)$ and $\Lambda_j \in \mathcal{B}(H, H_j)$ for each $j \in J$. Then the family $\Lambda_{TU} = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$ is a (T, U) -controlled K - g -fusion frame for H if there exist constants $0 < A \leq B < +\infty$ such that

$$(1.1) \quad A \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \langle \Lambda_j P_{W_j} U f, \Lambda_j P_{W_j} T f \rangle \leq B \|f\|^2,$$

for all $f \in H$. If Λ_{TU} satisfies only the right inequality of (1.1) it is called a (T, U) -controlled g -fusion Bessel sequence in H .

Let Λ_{TU} be a (T, U) -controlled g -fusion Bessel sequence in H with a bound B . The synthesis operator $T_C : \mathcal{K}_{\Lambda_j} \rightarrow H$ is defined as

$$T_C \left(\left\{ v_j \left(T^* P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} U \right)^{1/2} f \right\}_{j \in J} \right) = \sum_{j \in J} v_j^2 T^* P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} U f,$$

for all $f \in H$ and the analysis operator $T_C^* : H \rightarrow \mathcal{K}_{\Lambda_j}$ is given by

$$T_C^* f = \left\{ v_j \left(T^* P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} U \right)^{1/2} f \right\}_{j \in J}, \quad \text{for all } f \in H,$$

where

$$\mathcal{K}_{\Lambda_j} = \left\{ \left\{ v_j \left(T^* P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} U \right)^{1/2} f \right\}_{j \in J} : f \in H \right\} \subset \ell^2 \left(\{H_j\}_{j \in J} \right).$$

The frame operator $S_C : H \rightarrow H$ is defined as follows:

$$S_C f = T_C T_C^* f = \sum_{j \in J} v_j^2 T^* P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} U f,$$

for all $f \in H$ and it is easy to verify that

$$\langle S_C f, f \rangle = \sum_{j \in J} v_j^2 \langle \Lambda_j P_{W_j} U f, \Lambda_j P_{W_j} T f \rangle,$$

for all $f \in H$. Furthermore, if Λ_{TU} is a (T, U) -controlled K - g -fusion frame with bounds A and B , then $AKK^* \leq S_C \leq BI_H$.

1.3. Continuous controlled g -fusion frame. In recent times, controlled frames and their generalizations are also studied in continuous case by many researchers. P. Ghosh and T. K. Samanta studied continuous version of controlled g -fusion frame in [21].

Definition 1.3 ([21]). Let $F : X \rightarrow \mathbb{H}$ be a mapping, $v : X \rightarrow \mathbb{R}^+$ be a measurable function and $\{K_x\}_{x \in X}$ be a collection of Hilbert spaces. For each $x \in X$, suppose that $\Lambda_x \in \mathcal{B}(F(x), K_x)$ and $T, U \in \mathcal{GB}^+(H)$. Then $\Lambda_{TU} = \{(F(x), \Lambda_x, v(x))\}_{x \in X}$ is called a continuous (T, U) -controlled generalized fusion frame or continuous (T, U) -controlled g -fusion frame for H with respect to (X, μ) and v , if

(i) for each $f \in H$, the mapping $x \mapsto P_{F(x)}(f)$ is measurable (i.e., is weakly measurable);

(ii) there exist constants $0 < A \leq B < +\infty$ such that

$$(1.2) \quad A \|f\|^2 \leq \int_X v^2(x) \langle \Lambda_x P_{F(x)} U f, \Lambda_x P_{F(x)} T f \rangle d\mu_x \leq B \|f\|^2,$$

for all $f \in H$, where $P_{F(x)}$ is the orthogonal projection of H onto the subspace $F(x)$. The constants A, B are called the frame bounds. If only the right inequality of (1.2) holds then Λ_{TU} is called a continuous (T, U) -controlled g -fusion Bessel family for H .

Let Λ_{TU} be a continuous (T, U) -controlled g -fusion Bessel family for H . Then the operator $S_C : H \rightarrow H$ defined by

$$\langle S_C f, g \rangle = \int_X v^2(x) \langle T^* P_{F(x)} \Lambda_x^* \Lambda_x P_{F(x)} U f, g \rangle d\mu_x,$$

for all $f, g \in H$, is called the frame operator. If Λ_{TU} is a continuous (T, U) -controlled g -fusion frame for H , then from (1.2), we get

$$A \langle f, f \rangle \leq \langle S_C f, f \rangle \leq B \langle f, f \rangle, \quad \text{for all } f \in H.$$

The bounded linear operator $T_C : L^2(X, K) \rightarrow H$ defined by

$$\langle T_C \Phi, g \rangle = \int_X v^2(x) \langle T^* P_{F(x)} \Lambda_x^* \Lambda_x P_{F(x)} U f, g \rangle d\mu_x,$$

where for all $f \in H$, $\Phi = \left\{ v(x) \left(T^* P_{F(x)} \Lambda_x^* \Lambda_x P_{F(x)} U \right)^{1/2} f \right\}_{x \in X}$ and $g \in H$, is called synthesis operator and its adjoint operator is called analysis operator.

1.4. Weaving frame. Woven frame is a new notion in frame theory which has been introduced by Bemrose et al. [7]. Two frames $\{f_i\}_{i \in I}$ and $\{g_i\}_{i \in I}$ for H are called woven if there exist constants $0 < A \leq B < +\infty$ such that for any subset $\sigma \subset I$ the family $\{f_i\}_{i \in \sigma} \cup \{g_i\}_{i \in \sigma^c}$ is a frame for H . This frame has been generalized for the discrete as well as the continuous case such as woven fusion frame [17], woven g -frame [24], woven g -fusion frame [25], woven K - g -fusion frame [32], continuous weaving frame [36], continuous weaving fusion frame [33], continuous weaving g -frames [3], weaving continuous K - g -frames [5], controlled weaving frames [29], continuous controlled K - g -frames [30] etc.

In this paper, woven continuous controlled K - g -fusion frame in Hilbert spaces is presented and some of their properties are going to be established. We discuss sufficient conditions for weaving continuous controlled K - g -fusion frame. Construction of woven continuous controlled K - g -fusion frame by bounded linear operator is given. At the end, we discuss a perturbation result of woven continuous controlled K - g -fusion frame.

2. WEAVING CONTINUOUS CONTROLLED K - g -FUSION FRAME

In this section, we first give the continuous version of controlled K - g -fusion frame for H and then present weaving continuous controlled K - g -fusion frame for H .

Definition 2.1. Let $K \in \mathcal{B}(H)$ and $F : X \rightarrow \mathbb{H}$ be a mapping, $v : X \rightarrow \mathbb{R}^+$ be a measurable function and $\{K_x\}_{x \in X}$ be a collection of Hilbert spaces. For each $x \in X$, suppose that $\Lambda(x) \in \mathcal{B}(F(x), K_x)$ and $T, U \in \mathcal{GB}^+(H)$. Then $\Lambda_{TU} = \{(F(x), \Lambda(x), v(x))\}_{x \in X}$ is called a continuous (T, U) -controlled K - g -fusion frame for H with respect to (X, μ) and v , if

(i) for each $f \in H$, the mapping $x \mapsto P_{F(x)}(f)$ is measurable (i.e., is weakly measurable);

(ii) there exist constants $0 < A \leq B < +\infty$ such that

$$(2.1) \quad A \|K^* f\|^2 \leq \int_X v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \leq B \|f\|^2,$$

for all $f \in H$, where $P_{F(x)}$ is the orthogonal projection of H onto the subspace $F(x)$. The constants A, B are called the frame bounds.

Now, we consider the following cases.

(i) If only the right inequality of (2.1) holds, then Λ_{TU} is called a continuous (T, U) -controlled K - g -fusion Bessel family for H .

- (ii) If $U = I_H$, then Λ_{TU} is called a continuous (T, I_H) -controlled K - g -fusion frame for H .
- (iii) If $T = U = I_H$, then Λ_{TU} is called a continuous K - g -fusion frame for H (for more details, refer to [4]).
- (iv) If $K = I_H$, then Λ_{TU} is called a continuous (T, U) -controlled g -fusion frame for H .

Remark 2.1. If the measure space $X = \mathbb{N}$ and μ is the counting measure then a continuous (T, U) -controlled K - g -fusion frame will be the discrete (T, U) -controlled K - g -fusion frame.

2.0.1. *Example.* Let $H = \mathbb{R}^3$ and $\{e_1, e_2, e_3\}$ be an standard orthonormal basis for H . Consider

$$\mathcal{B} = \{x \in \mathbb{R}^3 : \|x\| \leq 1\}.$$

Then it is a measure space equipped with the Lebesgue measure μ . Let us now consider that $\{B_1, B_2, B_3\}$ is a partition of \mathcal{B} where $\mu(B_1) \geq \mu(B_2) \geq \mu(B_3) > 1$. Let $\mathbb{H} = \{W_1, W_2, W_3\}$, where $W_1 = \overline{\text{Span}}\{e_1, e_2\}$, $W_2 = \overline{\text{Span}}\{e_2, e_3\}$ and $W_3 = \overline{\text{Span}}\{e_1, e_3\}$. Define $F : \mathcal{B} \rightarrow \mathbb{H}$ by

$$F(x) = \begin{cases} W_1, & \text{if } x \in B_1, \\ W_2, & \text{if } x \in B_2, \\ W_3, & \text{if } x \in B_3, \end{cases}$$

and $v : \mathcal{B} \rightarrow [0, +\infty)$ by

$$v(x) = \begin{cases} 1, & \text{if } x \in B_1, \\ 2, & \text{if } x \in B_2, \\ -1, & \text{if } x \in B_3. \end{cases}$$

It is easy to verify that F and v are measurable functions. For each $x \in \mathcal{B}$, define the operators

$$\Lambda(x)(f) = \frac{1}{\sqrt{\mu(B_k)}} \langle f, e_k \rangle e_k,$$

$f \in H$, where k is such that $x \in \mathcal{B}_k$ and $K : H \rightarrow H$ by

$$Ke_1 = e_1, \quad Ke_2 = e_2, \quad Ke_3 = 0.$$

It is easy to verify that $K^*e_1 = e_1$, $K^*e_2 = e_2$, $K^*e_3 = 0$. Now, for any $f \in H$, we have

$$\|K^*f\|^2 = \left\| \sum_{i=1}^3 \langle f, e_i \rangle K^*e_i \right\|^2 = |\langle f, e_1 \rangle|^2 + |\langle f, e_2 \rangle|^2 \leq \|f\|^2.$$

Let $T(f_1, f_2, f_3) = (5f_1, 4f_2, 5f_3)$ and $U(f_1, f_2, f_3) = \left(\frac{f_1}{6}, \frac{f_2}{3}, \frac{f_3}{6}\right)$ be two operators on H . Then it is easy to verify that $T, U \in \mathcal{GB}^+(H)$ and $TU = UT$. Now, for any

$f = (f_1, f_2, f_3) \in H$, we have

$$\begin{aligned} & \int_{\mathbb{B}} v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ &= \sum_{i=1}^3 \int_{\mathbb{B}_i} v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ &= \frac{5}{6}f_1^2 + \frac{16}{3}f_2^2 + \frac{5}{6}f_3^2. \end{aligned}$$

This implies that

$$\frac{5}{6} \|K^*f\|^2 \leq \int_{\mathbb{B}} v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \leq \frac{16}{3} \|f\|^2.$$

Thus, Λ_{TU} be a continuous (T, U) -controlled K - g -fusion frame for \mathbb{R}^3 .

Now, we present woven continuous controlled K - g -fusion frame for H .

Definition 2.2. A family of continuous (T, U) -controlled K - g -fusion frames given by $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in X}$ for H is said to be woven continuous (T, U) -controlled K - g -fusion frame if there exist universal positive constants $0 < A \leq B < +\infty$ such that for each partition $\{\sigma_i\}_{i \in [m]}$ of X , the family $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ is a continuous (T, U) -controlled K - g -fusion frame for H with bounds A and B .

Each family $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ is called a weaving continuous (T, U) -controlled K - g -fusion frame. For abbreviation, we use W. C. C. K. G. F. F. instead of the statement of woven continuous (T, U) -controlled K - g -fusion frame.

In the following proposition, we will see that every woven continuous controlled K - g -fusion frame has a universal upper bound.

Proposition 2.1. Suppose for each $i \in [m]$, $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{x \in X}$ be a continuous (T, U) -controlled K - g -fusion Bessel family for H with bound B_i . Then for any partition $\{\sigma_i\}_{i \in [m]}$ of X , the family $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ is a continuous (T, U) -controlled K - g -fusion Bessel family for H .

Proof. Let $\{\sigma_i\}_{i \in [m]}$ be an arbitrary partition of X . For each $f \in H$, we have

$$\begin{aligned} & \sum_{i \in [m]} \int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}Uf, \Lambda_i(x)P_{F_i(x)}Tf \rangle d\mu_x \\ & \leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}Uf, \Lambda_i(x)P_{F_i(x)}Tf \rangle d\mu_x \\ & \leq \left(\sum_{i \in [m]} B_i \right) \|f\|^2. \end{aligned}$$

This completes the proof. □

Next, we give a characterization of W. C. C. K. G. F. F. for H in terms of an operator.

Theorem 2.1. *Let the families given by $\Lambda = \{(F(x), \Lambda(x), v(x))\}_{x \in X}$ and $\Gamma = \{(G(x), \Lambda(x), v(x))\}_{x \in X}$ be continuous (T, U) -controlled K - g -fusion frames for H . The following statements are equivalent.*

- (i) Λ and Γ are W. C. C. K. G. F. F. for H .
- (ii) For each partition σ of X , there exist $\alpha > 0$ and a bounded linear operator $\Theta_\sigma : L_\sigma^2(X, K) \rightarrow H$ defined by

$$\begin{aligned} \langle \Theta_\sigma \Phi, g \rangle &= \int_\sigma v^2(x) \langle T^* P_{F(x)} \Lambda(x)^* \Lambda(x) P_{F(x)} U f, g \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle T^* P_{G(x)} \Gamma(x)^* \Gamma(x) P_{G(x)} U f, g \rangle d\mu_x, \end{aligned}$$

$g \in H$ such that $\alpha K K^* \leq \Theta_\sigma \Theta_\sigma^*$, where

$$L_\sigma^2(X, K) = \left\{ \Phi = \phi \cup \psi : \int_X \|\Phi\|^2 d\mu < +\infty \right\},$$

where for all $f \in H$,

$$\phi = \left\{ v(x) \left(T^* P_{F(x)} \Lambda(x)^* \Lambda(x) P_{F(x)} U \right)^{1/2} f \right\}_{x \in \sigma}$$

and

$$\psi = \left\{ v(x) \left(T^* P_{G(x)} \Gamma(x)^* \Gamma(x) P_{G(x)} U \right)^{1/2} f \right\}_{x \in \sigma^c}.$$

Proof. (i) \Rightarrow (ii) Suppose that A and B are the universal lower and upper bounds for Λ and Γ . Take $\Theta_\sigma = T_C^\sigma$, for every partition σ of X , where T_C^σ is the synthesis operator of

$$\{(F(x), \Lambda(x), v(x))\}_{x \in \sigma} \cup \{(G(x), \Lambda(x), v(x))\}_{x \in \sigma^c}.$$

Thus, for each $\Phi \in L_\sigma^2(X, K)$, we have

$$\begin{aligned} \langle \Theta_\sigma \Phi, g \rangle &= \langle T_C^\sigma \Phi, g \rangle \\ &= \int_\sigma v^2(x) \langle T^* P_{F(x)} \Lambda(x)^* \Lambda(x) P_{F(x)} U f, g \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle T^* P_{G(x)} \Gamma(x)^* \Gamma(x) P_{G(x)} U f, g \rangle d\mu_x, \quad g \in H. \end{aligned}$$

Since Λ and Γ are woven, for each $f \in H$, we have

$$A \|K^* f\|^2 \leq \|(T_C^\sigma)^* f\|^2 = \|\Theta_\sigma^* f\|^2.$$

Thus, $\alpha K K^* \leq \Theta_\sigma \Theta_\sigma^*$, $\alpha = A$.

(ii) \Rightarrow (i) Let σ be a partition of X and $f \in H$. Now it is easy to verify that

$$\Theta_\sigma^* f = \left\{ v(x) \left(T^* P_{F(x)} \Lambda(x)^* \Lambda(x) P_{F(x)} U \right)^{1/2} f \right\}_{x \in \sigma}$$

$$\cup \left\{ v(x) \left(T^* P_{G(x)} \Gamma(x)^* \Gamma(x) P_{G(x)} U \right)^{1/2} f \right\}_{x \in \sigma^c}.$$

Thus, for each $f \in H$, we have

$$\begin{aligned} \alpha \|K^* f\|^2 &\leq \|\Theta_\sigma^* f\|^2 = \int_\sigma v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x. \end{aligned}$$

Hence, Λ and Γ are W. C. C. K. G. F. F. for H . This completes the proof. \square

In the following theorem, we will construct W. C. C. K. G. F. F. for H by using a bounded linear operator.

Theorem 2.2. *Let $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ be a W. C. C. K. G. F. F. for H with universal bounds A and B . If $V \in \mathcal{B}(H)$ is invertible such that V^* commutes with T, U and V commutes with K , then $\{(VF_i(x), \Lambda_i(x) P_{F_i(x)} V^*, v_i(x))\}_{i \in [m], x \in \sigma_i}$ is a W. C. C. K. G. F. F. for H .*

Proof. Since $P_{F_i(x)} V^* = P_{F_i(x)} V^* P_{V F_i(x)}$ for all $x \in \sigma_i$ and $i \in [m]$, the mapping $x \mapsto P_{V F_i(x)}$ is weakly measurable. For each $f \in H$, we have

$$\begin{aligned} &\sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} V^* P_{V F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} V^* P_{V F_i(x)} T f \rangle d\mu_x \\ &= \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} V^* U f, \Lambda_i(x) P_{F_i(x)} V^* T f \rangle d\mu_x \\ &= \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U V^* f, \Lambda_i(x) P_{F_i(x)} T V^* f \rangle d\mu_x \\ &\leq B \|V^* f\|^2 \leq B \|V\|^2 \|f\|^2. \end{aligned}$$

On the other hand, for each $f \in H$, we have

$$\begin{aligned} &\sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} V^* P_{V F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} V^* P_{V F_i(x)} T f \rangle d\mu_x \\ &\geq A \|K^* V^* f\|^2 = A \|V^* K^* f\|^2 \geq A \|V^{-1}\|^{-2} \|K^* f\|^2. \end{aligned}$$

This completes the proof. \square

Corollary 2.1. *Let $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ be a W. C. C. K. G. F. F. for H with universal bounds A and B . If $V \in \mathcal{B}(H)$ is invertible such that V^* commutes with T, U and V commutes with K , then $\{(VF_i(x), \Lambda_i(x) P_{F_i(x)} V^*, v_i(x))\}_{i \in [m], x \in \sigma_i}$ is a W. C. C. $V K V^*$. G. F. F. for H .*

Proof. According to the proof of Theorem 2.2, universal upper bounds is $B\|V\|^2$. On the other hand, for each $f \in H$, we have

$$\begin{aligned} & \frac{A}{\|V\|^2} \|(VKV^*)^* f\|^2 = \frac{A}{\|V\|^2} \|VK^*V^*f\|^2 \leq A \|K^*V^*f\|^2 \\ & \leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}UV^*f, \Lambda_i(x)P_{F_i(x)}TV^*f \rangle d\mu_x \\ & = \sum_{i \in [m]} \int_X v_i^2(x) \langle \Gamma_i(x)P_{V F_i(x)}Uf, \Gamma_i(x)P_{V F_i(x)}Tf \rangle d\mu_x, \end{aligned}$$

where $\Lambda_i(x)P_{F_i(x)}V^* = \Gamma_i(x)$. This completes the proof. \square

Theorem 2.3. *Let $V \in \mathcal{B}(H)$ be invertible operator such that V^* , $(V^{-1})^*$ commutes with T and U . Suppose $\{(VF_i(x), \Lambda_i(x)P_{F_i(x)}V^*, v_i(x))\}_{i \in [m], x \in \sigma_i}$ is a W. C. C. K. G. F. F. for H with universal bounds A and B . Then $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ be a W. C. C. $V^{-1}KV$. G. F. F. for H .*

Proof. Now, for each $f \in H$, using Theorem 1.2, and taking $\Lambda_i(x)P_{F_i(x)}V^* = \Gamma_i(x)$, we have

$$\begin{aligned} & \frac{A}{\|V\|^2} \|(V^{-1}KV)^* f\|^2 = \frac{A}{\|V\|^2} \|V^*K^*(V^{-1})^*f\|^2 \\ & \leq A \|K^*(V^{-1})^*f\|^2 \\ & \leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Gamma_i(x)P_{V F_i(x)}U(V^{-1})^*f, \Gamma_i(x)P_{V F_i(x)}T(V^{-1})^*f \rangle d\mu_x \\ & \leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Gamma_i(x)U(V^{-1})^*f, \Gamma_i(x)T(V^{-1})^*f \rangle d\mu_x \\ & = \sum_{i \in [m]} \int_X v_i^2(x) \langle \Gamma_i(x)(V^{-1})^*Uf, \Gamma_i(x)(V^{-1})^*Tf \rangle d\mu_x \\ & = \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}Uf, \Lambda_i(x)P_{F_i(x)}Tf \rangle d\mu_x. \end{aligned}$$

On the other hand, for each $f \in H$, it is easy to verify that

$$\sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}Uf, \Lambda_i(x)P_{F_i(x)}Tf \rangle d\mu_x \leq B \|V^{-1}\|^2 \|f\|^2.$$

This completes the proof. \square

Next, we will see that the intersection of components of a W. C. C. K. G. F. F. with a closed subspace is a W. C. C. K. G. F. F. for the smaller space.

Theorem 2.4. *Let $\{F(x), \Lambda(x), v(x)\}_{x \in X}$ and $\{G(x), \Gamma(x), w(x)\}_{x \in X}$ be W. C. C. K. G. F. F. for H and W be a closed subspace of H . Then the families given by*

$\{F(x) \cap W, \Lambda(x), v(x)\}_{x \in X}$ and $\{G(x) \cap W, \Gamma(x), w(x)\}_{x \in X}$ are W. C. C. K. G. F. F. for W .

Proof. The operators $P_{F(x) \cap W} = P_{F(x)}(P_W)$ and $P_{G(x) \cap W} = P_{G(x)}(P_W)$ are orthogonal projections of H onto $F(x) \cap W$ and $G(x) \cap W$, respectively. Let σ be a measurable subset of X . Then for each $f \in W$, we have

$$\begin{aligned} & \int_{\sigma} v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ & + \int_{\sigma^c} w^2(x) \langle \Gamma(x)P_{G(x)}Uf, \Gamma(x)P_{G(x)}Tf \rangle d\mu_x \\ & = \int_{\sigma} v^2(x) \langle \Lambda(x)P_{F(x)}P_WUf, \Lambda(x)P_{F(x)}P_WTf \rangle d\mu_x \\ & + \int_{\sigma^c} w^2(x) \langle \Gamma(x)P_{G(x)}P_WUf, \Gamma(x)P_{G(x)}P_WTf \rangle d\mu_x \\ & = \int_{\sigma} v^2(x) \langle \Lambda(x)P_{F(x) \cap W}Uf, \Lambda(x)P_{F(x) \cap W}Tf \rangle d\mu_x \\ & + \int_{\sigma^c} w^2(x) \langle \Gamma(x)P_{G(x) \cap W}Uf, \Gamma(x)P_{G(x) \cap W}Tf \rangle d\mu_x. \end{aligned}$$

This completes the proof. \square

The following theorem states the equivalence between W. C. C. K. G. F. F. and a bounded linear operator.

Theorem 2.5. *Let $V \in \mathcal{B}(H)$ be an invertible operator such that V^* commutes with T, U . Suppose K be a bounded linear operator on H which have closed range. Let $\Lambda_{TU} = \{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ be a W. C. C. K. G. F. F. for H with universal bounds A and B . Then the family given by*

$$\Delta_{TU} = \left\{ (VF_i(x), \Lambda_i(x)P_{F_i(x)}V^*, v_i(x)) \right\}_{i \in [m], x \in \sigma_i}$$

*is a W. C. C. K. G. F. F. for H if and only if there exists a $\delta > 0$ such that for each $f \in H$, we have $\|V^*f\| \geq \delta \|K^*f\|$.*

Proof. Suppose that Δ_{TU} is a W. C. C. K. G. F. F. for H with bounds C and D . Then for each $f \in H$, using the Theorem 1.2, and taking $\Lambda_i(x)P_{F_i(x)}V^* = \Gamma_i(x)$, we have

$$\begin{aligned} C \|K^*f\|^2 & \leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Gamma_i(x)P_{V_{F_i(x)}}Uf, \Gamma_i(x)P_{V_{F_i(x)}}Tf \rangle d\mu_x \\ & = \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}V^*Uf, \Lambda_i(x)P_{F_i(x)}V^*Tf \rangle d\mu_x \\ & = \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x)P_{F_i(x)}UV^*f, \Lambda_i(x)P_{F_i(x)}TV^*f \rangle d\mu_x \end{aligned}$$

$$\leq B \|V^* f\|^2.$$

Thus,

$$\|V^* f\| \geq \sqrt{C/B} \|K^* f\|, \quad \text{for all } f \in H.$$

Conversely, suppose $\|V^* f\| \geq \delta \|K^* f\|$ for all $f \in H$. Since K have a closed range, by Theorem 1.3, for all $f \in H$, we get

$$\|V^* f\| = \|(K^\dagger)^* K^* V^* f\| \leq \|K^\dagger\| \|K^* V^* f\|.$$

Now, for $f \in H$, we have

$$\begin{aligned} & \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} V^* P_{V_{F_i(x)}} U f, \Lambda_i(x) P_{F_i(x)} V^* P_{V_{F_i(x)}} T f \rangle d\mu_x \\ &= \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U V^* f, \Lambda_i(x) P_{F_i(x)} T V^* f \rangle d\mu_x \\ &\geq A \|K^* V^* f\|^2 \geq A \|K^\dagger\|^{-2} \|V^* f\|^2 \geq A \delta^2 \|K^\dagger\|^{-2} \|K^* f\|^2. \end{aligned}$$

This completes the proof. \square

The next theorem shows that it is enough to check continuous weaving controlled K - g -fusion woven on smaller measurable space than the original.

Theorem 2.6. *Suppose for each $i \in [m]$, $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{x \in X}$ be a continuous (T, U) -controlled K - g -fusion frame for H with universal bounds A_i and B_i . If there exists a measurable subset $Y \subset X$ such that the family of continuous (T, U) -controlled K - g -fusion frame $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in Y}$ is a W. C. C. K. G. F. F. for H with universal frame bounds A and B . Then the family given by $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in X}$ is a W. C. C. K. G. F. F. for H with universal frame bounds A and $\sum_{i \in [m]} B_i$.*

Proof. Let $\{\sigma_i\}_{i \in [m]}$ be an arbitrary partition of X . For each $f \in H$, we define $\varphi : X \rightarrow \mathbb{C}$ by

$$\varphi(x) = \sum_{i \in [m]} \chi_{\sigma_i}(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle.$$

Then φ is measurable. Now, for each $f \in H$, we have

$$\begin{aligned} & \sum_{i \in [m]} \int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \\ &\leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \\ &\leq \left(\sum_{i \in [m]} B_i \right) \|f\|^2. \end{aligned}$$

It is easy to verify that $\{\sigma_i \cap Y\}_{i \in [m]}$ is a partitions of Y . Thus, the family given by $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i \cap Y}$ is a continuous (T, U) -controlled K - g -fusion frame for H with lowest frame bound A . Therefore,

$$\begin{aligned} & \sum_{i \in [m]} \int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \\ & \geq \sum_{i \in [m]} \int_{\sigma_i \cap Y} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \\ & \geq A \|K^* f\|^2. \end{aligned}$$

This completes the proof. \square

In the following theorem, we show that it is possible to remove vectors from continuous controlled K - g -fusion frames and still be left with woven frames.

Theorem 2.7. *Let $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i}$ be a W. C. C. K. G. F. F. for H with universal bounds A and B . If there exists $0 < D < A$ and a measurable subset $Y \subset X$ and $n \in [m]$ such that for $f \in H$*

$$\sum_{i \in [m] \setminus \{n\}} \int_{X \setminus Y} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \leq D \|K^* f\|^2,$$

then the family $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in Y}$ is a W. C. C. K. G. F. F. for H with frame bounds $A - D$ and B .

Proof. Suppose that $\{\sigma_i\}_{i \in [m]}$ and $\{\gamma_i\}_{i \in [m]}$ are partitions of Y and $X \setminus Y$, respectively. For a given $f \in H$, we define $\varphi : Y \rightarrow \mathbb{C}$ by

$$\varphi(x) = \sum_{i \in [m]} \chi_{\sigma_i}(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle,$$

and $\phi : X \rightarrow \mathbb{C}$ by

$$\phi(x) = \sum_{i \in [m]} \chi_{\sigma_i \cup \gamma_i}(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle.$$

Since $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{i \in [m], x \in \sigma_i \cup \gamma_i}$ is a continuous (T, U) -controlled K - g -fusion frame for H and $\varphi = \phi|_Y$, φ and ϕ are measurable. So, for each $f \in H$, we have

$$\begin{aligned} & \sum_{i \in [m]} \int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \\ & \leq \sum_{i \in [m]} \int_{\sigma_i \cup \gamma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \leq B \|f\|^2. \end{aligned}$$

Now, we assume that $\{\xi_i\}_{i \in [m]}$ such that $\xi_n = \theta$. Then $\{\xi_i \cup \sigma_i\}_{i \in [m]}$ is a partition of X and so for any $f \in H$, we have

$$\sum_{i \in [m]} \int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x$$

$$\begin{aligned}
&= \sum_{i \in [m] \setminus \{n\}} \left[\int_{\xi_i \cup \sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \right. \\
&\quad - \int_{\xi_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \\
&\quad \left. + \int_{\sigma_n} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \right] \\
&\geq \sum_{i \in [m] \setminus \{n\}} \left[\int_{\xi_i \cup \sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \right. \\
&\quad - \int_{X \setminus Y} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \\
&\quad \left. + \int_{\sigma_n} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \right] \\
&= \sum_{i \in [m] \setminus \{n\}} \int_{\xi_i \cup \sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \\
&\quad - \sum_{i \in [m] \setminus \{n\}} \int_{X \setminus Y} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} U f, \Lambda_i(x) P_{F_i(x)} T f \rangle d\mu_x \\
&\geq (A - D) \|K^* f\|^2.
\end{aligned}$$

This completes the proof. \square

Proposition 2.2. *Let $K \in \mathcal{B}(H)$ be a closed range operator, $V \in \mathcal{B}(H)$ be a unitary operator and $\{(F(x), \Lambda(x), v(x))\}_{x \in X}$ be a continuous (T, U) -controlled K -g-fusion frame for H with bounds A, B . If $\|I_H - V\|^2 \|K^\dagger\|^2 \leq A/B$ and V commutes with T, U , then*

$$\Lambda = \{(F(x), \Lambda(x), v(x))\}_{x \in X}, \quad \Lambda' = \{(V^{-1}F(x), \Lambda(x)V, v(x))\}_{x \in X}$$

are W. C. C. K. G. F. F. for \mathcal{R}_K .

Proof. Let σ be a partition of X . Since $K \in \mathcal{B}(H)$ has a closed range, for $f \in \mathcal{R}_K$, we have $\|f\|^2 \leq \|K^\dagger\|^2 \|K^* f\|^2$. Now, for each $f \in \mathcal{R}_K$, we have

$$\begin{aligned}
&\int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\
&\quad + \int_{\sigma^c} v^2(x) \langle \Lambda(x) V P_{V^{-1}F(x)} U f, \Lambda(x) V P_{V^{-1}F(x)} T f \rangle d\mu_x \\
&= \int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x
\end{aligned}$$

$$\begin{aligned}
& + \int_{\sigma^c} v^2(x) \langle \Lambda(x)P_{F(x)}UVf, \Lambda(x)P_{F(x)}TVf \rangle d\mu_x \\
& \geq \int_X v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\
& \quad - \int_{\sigma^c} v^2(x) \langle \Lambda(x)P_{F(x)}U(I_H - V)f, \Lambda(x)P_{F(x)}T(I_H - V)f \rangle d\mu_x \\
& \geq A\|K^*f\|^2 - B\|I_H - V\|^2\|f\|^2 \\
& \geq A\|K^*f\|^2 - B\|I_H - V\|^2\|K^\dagger\|^2\|K^*f\|^2 \\
& = \left(A - B\|I_H - V\|^2\|K^\dagger\|^2 \right) \|K^*f\|^2.
\end{aligned}$$

Hence, the families Λ and Λ' are W. C. C. K. G. F. F. for \mathcal{R}_K . \square

Next, we will see that under some sufficient conditions sum of two continuous (T, U) -controlled K - g -fusion frames is woven with itself.

Theorem 2.8. *Let $K \in \mathcal{B}(H)$ be an invertible operator, the families given by $\Lambda = \{(F(x), \Lambda(x), v(x))\}_{x \in X}$ and $\Gamma = \{(G(x), \Lambda(x), v(x))\}_{x \in X}$ be continuous (T, U) -controlled K - g -fusion frames for H with bounds A, B and C, D , respectively. Suppose for each $x \in X$*

- (i) $F(x) \subset G(x)^\perp$;
- (ii) $\Lambda(x)P_{F(x)}\mathcal{R}(U) \perp \Lambda(x)P_{G(x)}\mathcal{R}(T)$;
- (iii) $\Lambda(x)P_{F(x)}\mathcal{R}(T) \perp \Lambda(x)P_{G(x)}\mathcal{R}(U)$.

If for any partition σ of X , $(T_\Gamma^\sigma)^*$ is bounded below then

$$\Delta = \{(F(x) + G(x), \Lambda(x), v(x))\}_{x \in X},$$

and Λ are W. C. C. K. G. F. F. for H .

Proof. Since for each $x \in X$, $F(x) \subset G(x)^\perp$, we have $P_{F(x)+G(x)} = P_{F(x)} + P_{G(x)}$. Now, for each $x \in X$, using the given conditions (ii) and (iii), we have

$$\begin{aligned}
& \int_X v^2(x) \langle \Lambda(x)P_{F(x)+G(x)}Uf, \Lambda(x)P_{F(x)+G(x)}Tf \rangle d\mu_x \\
& = \int_X v^2(x) \langle \Lambda(x) (P_{F(x)} + P_{G(x)})Uf, \Lambda(x) (P_{F(x)} + P_{G(x)})Tf \rangle d\mu_x \\
& = \int_X v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\
(2.2) \quad & + \int_X v^2(x) \langle \Lambda(x)P_{G(x)}Uf, \Lambda(x)P_{G(x)}Tf \rangle d\mu_x \\
& \leq (B + D)\|f\|^2.
\end{aligned}$$

On the other hand, from (2.2), we get

$$\int_X v^2(x) \langle \Lambda(x)P_{F(x)+G(x)}Uf, \Lambda(x)P_{F(x)+G(x)}Tf \rangle d\mu_x \geq (A + C) \|K^*f\|^2,$$

for all $f \in H$. Thus, Δ is a continuous (T, U) -controlled K - g -fusion frame for H with bounds $(A + C)$ and $(B + D)$.

Furthermore, since K is a invertible operator and for any partition σ of X , $(T_\Gamma^\sigma)^*$ is bounded below, for each $f \in H$, there exists $M > 0$ such that

$$\|(T_\Gamma^\sigma)^* f\|^2 \geq M^2 \|f\|^2 \geq \frac{M^2}{\|K\|^2} \|K^*f\|^2.$$

Now, for each $f \in H$, we have

$$\begin{aligned} & \int_\sigma v^2(x) \langle \Lambda(x)P_{F(x)+G(x)}Uf, \Lambda(x)P_{F(x)+G(x)}Tf \rangle d\mu_x \\ & + \int_{\sigma^c} v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ & = \int_X v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ & \quad - \int_\sigma v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ & \quad + \int_\sigma v^2(x) \langle \Lambda(x) (P_{F(x)} + P_{G(x)}) Uf, \Lambda(x) (P_{F(x)} + P_{G(x)}) Tf \rangle d\mu_x \\ & = \int_X v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ & \quad + \int_\sigma v^2(x) \langle \Lambda(x)P_{G(x)}Uf, \Lambda(x)P_{G(x)}Tf \rangle d\mu_x \\ & \geq A \|K^*f\|^2 + \|(T_\Gamma^\sigma)^* f\|^2 \geq \left(A + \frac{M^2}{\|K\|^2} \right) \|K^*f\|^2. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \int_\sigma v^2(x) \langle \Lambda(x)P_{F(x)+G(x)}Uf, \Lambda(x)P_{F(x)+G(x)}Tf \rangle d\mu_x \\ & \quad + \int_{\sigma^c} v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \\ & \leq \int_X v^2(x) \langle \Lambda(x)P_{F(x)+G(x)}Uf, \Lambda(x)P_{F(x)+G(x)}Tf \rangle d\mu_x \\ & \quad + \int_X v^2(x) \langle \Lambda(x)P_{F(x)}Uf, \Lambda(x)P_{F(x)}Tf \rangle d\mu_x \end{aligned}$$

$$\leq (2B + D)\|f\|^2.$$

Thus, Δ and Λ are W. C. C. K. G. F. F. for H . Similarly, it can be shown that Δ and Γ are W. C. C. K. G. F. F. for H . This completes the proof. \square

In the following theorem, we present a sufficient condition for weaving continuous controlled K - g -fusion frame in terms of positive operators associated with given continuous controlled K - g -fusion frame.

Theorem 2.9. *Let the families given by $\Lambda = \{(F(x), \Lambda(x), v(x))\}_{x \in X}$ and $\Gamma = \{(G(x), \Lambda(x), v(x))\}_{x \in X}$ be continuous (T, U) -controlled K - g -fusion frames for H . Suppose for each $x \in X$, the operator $U_x : H \rightarrow H$ defined by*

$$\langle U_x(f), g \rangle = \int_X v^2(x) \langle T^* \Delta(x) U f, g \rangle d\mu_x,$$

$f, g \in H$, where $\Delta(x) = P_{G(x)} \Gamma^*(x) \Gamma(x) P_{G(x)} - P_{F(x)} \Lambda^*(x) \Lambda(x) P_{F(x)}$, is a positive operator. Then Λ and Γ are W. C. C. K. G. F. F. for H .

Proof. Let A, B and C, D be frame bounds of Λ and Γ , respectively. Take σ be any partition of X . Then for each $f \in H$, we have

$$\begin{aligned} A \|K^* f\|^2 &\leq \int_X v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &= \int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle T^* P_{F(x)} \Lambda(x)^* \Lambda(x) P_{F(x)} U f, f \rangle d\mu_x \\ &= \int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\quad - \int_{\sigma^c} v^2(x) \langle T^* \Delta(x) U f, f \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle T^* P_{G(x)} \Gamma(x)^* \Gamma(x) P_{G(x)} U f, f \rangle d\mu_x \\ &\leq \int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x \\ &\leq (B + D) \|f\|^2. \end{aligned}$$

Thus, Λ and Γ are W. C. C. K. G. F. F. for H with universal bounds A and $B + D$. \square

Theorem 2.10. *Suppose for each $i \in [m]$, $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{x \in X}$ be a continuous (T, U) -controlled K - g -fusion frame for H with bounds A_i and B_i . Suppose Y be*

measurable subset X and there exists $N > 0$ such that for all $i, k \in [m]$ with $i \neq k$

$$0 \leq \int_Y \langle \Gamma_{i,k} Uf, \Gamma_{i,k} Tf \rangle d\mu_x \leq N \min\{\Theta, \Omega\}, \quad f \in H,$$

where

$$\begin{aligned} \Gamma_{i,k} &= v_i^2(x) \Lambda_i(x) P_{F_i(x)} - v_k^2(x) \Lambda_k(x) P_{F_k(x)}, \\ \Theta &= \int_Y v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x, \\ \Omega &= \int_Y v_k^2(x) \langle \Lambda_i(x) P_{F_k(x)} Uf, \Lambda_k(x) P_{F_k(x)} Tf \rangle d\mu_x. \end{aligned}$$

Then the family $\{(F_i(x), \Lambda_i(x), v_i(x))\}_{x \in X, i \in [m]}$ is *W. C. C. K. G. F. F.* for H with universal bounds $\frac{A}{(m-1)(N+1)+1}$ and B , where $A = \sum_{i \in [m]} A_i$ and $B = \sum_{i \in [m]} B_i$.

Proof. Let $\{\sigma_i\}_{i \in [m]}$ be a partition of X . Then for $f \in H$, we have

$$\begin{aligned} \sum_{i \in [m]} A_i \|K^* f\|^2 &\leq \sum_{i \in [m]} \int_X v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \\ &= \sum_{i \in [m]} \sum_{k \in [m], k \neq i} \int_{\sigma_k} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \\ &\leq \sum_{i \in [m]} \left[\int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \right. \\ &\quad + \sum_{k \in [m], k \neq i} \int_{\sigma_k} \langle \Gamma_{i,k} Uf, \Gamma_{i,k} Tf \rangle d\mu_x \\ &\quad \left. + \sum_{k \in [m], k \neq i} \int_{\sigma_k} v_k^2(x) \langle \Lambda_k(x) P_{F_k(x)} Uf, \Lambda_k(x) P_{F_k(x)} Tf \rangle d\mu_x \right], \\ \Gamma_{i,k} &= v_i^2(x) \Lambda_i(x) P_{F_i(x)} - v_k^2(x) \Lambda_k(x) P_{F_k(x)} \\ &\leq \sum_{i \in [m]} \left[\int_{\sigma_i} v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \right. \\ &\quad \left. + \sum_{k \in [m], k \neq i} (N+1) \int_{\sigma_k} v_k^2(x) \langle \Lambda_k(x) P_{F_k(x)} Uf, \Lambda_k(x) P_{F_k(x)} Tf \rangle d\mu_x \right], \\ &= D \sum_{i \in [m], \sigma_i} \int v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x, \end{aligned}$$

where $D = \{(m-1)(N+1)+1\}$. Thus, for each $f \in H$, we have

$$\begin{aligned} \frac{A}{(m-1)(N+1)+1} \|K^* f\|^2 &\leq \sum_{i \in [m], \sigma_i} \int v_i^2(x) \langle \Lambda_i(x) P_{F_i(x)} Uf, \Lambda_i(x) P_{F_i(x)} Tf \rangle d\mu_x \\ &\leq B \|f\|^2. \end{aligned}$$

This completes the proof. \square

3. PERTURBATION OF WOVEN CONTINUOUS CONTROLLED g -FUSION FRAME

In frame theory, one of the most important problem is the stability of frame under some perturbation. P. Casazza and Chirstensen [10] have been generalized the Paley-Wiener perturbation theorem to perturbation of frame in Hilbert space. P. Ghosh and T. K. Samanta have studied perturbation of dual g -fusion frame and continuous controlled g -fusion frame in [18, 21]. In this section, we will see that under some small perturbations, continuous controlled K - g -fusion frames constitute woven continuous controlled K - g -fusion frame.

Theorem 3.1. *Let the families given by $\Lambda = \{(F(x), \Lambda(x), v(x))\}_{x \in X}$ and $\Gamma = \{(G(x), \Gamma(x), v(x))\}_{x \in X}$ be continuous (T, U) -controlled K - g -fusion frames for H with bounds A, B and C, D , respectively. Suppose that there exist non-negative constants λ_1, λ_2 and μ with $0 < \lambda_1 < 1$, $\mu < (1 - \lambda_1)A - \lambda_2B$ such that for each $f \in H$, we have*

$$\begin{aligned} 0 &\leq \int_X v^2(x) \langle T^* \Delta(x) U f, f \rangle d\mu_x \\ &\leq \lambda_1 \int_X v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\quad + \lambda_2 \int_X v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x + \mu \|K^* f\|^2, \end{aligned}$$

where $\Delta(x) = (P_{F(x)} \Lambda(x)^* \Lambda(x) P_{F(x)} - P_{G(x)} \Gamma(x)^* \Gamma(x) P_{G(x)})$. Then, Λ and Γ are W. C. C. K. G. F. F. for H .

Proof. Let σ be a partition of X . Now, for each $f \in H$, we have

$$\begin{aligned} &\int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x + \int_{\sigma^c} v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x \\ &\geq \int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x - \int_{\sigma^c} v^2(x) \langle T^* \Delta(x) U f, f \rangle d\mu_x \\ &\quad + \int_{\sigma^c} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\geq \int_X v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x - \int_X v^2(x) \langle T^* \Delta(x) U f, f \rangle d\mu_x \\ &\geq (1 - \lambda_1) \int_X v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ &\quad - \lambda_2 \int_X v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x - \mu \|K^* f\|^2 \end{aligned}$$

$$\geq ((1 - \lambda_1) A - \lambda_2 B - \mu) \|K^* f\|^2.$$

On the other hand,

$$\begin{aligned} & \int_{\sigma} v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ & + \int_{\sigma^c} v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x \\ & \leq \int_X v^2(x) \langle \Lambda(x) P_{F(x)} U f, \Lambda(x) P_{F(x)} T f \rangle d\mu_x \\ & \quad + \int_X v^2(x) \langle \Gamma(x) P_{G(x)} U f, \Gamma(x) P_{G(x)} T f \rangle d\mu_x \\ & \leq (B + D) \|f\|^2. \end{aligned}$$

This completes the proof. \square

Acknowledgement. The authors would like to thank the editor and the referees for their helpful suggestions and comments to improve this paper.

REFERENCES

- [1] R. Ahmadi, G. Rahimlou, V. Sadri and R. Z. Farfar, *Constructions of K - g fusion frames and their duals in Hilbert spaces*, Bull. Transilv. Univ. Brasov Ser. III. Math. Comput. Sci. **13**(62) (2020), 17–32. <https://doi.org/10.31926/but.mif.2020.12.61.1.2>
- [2] S. T. Ali, J. P. Antonie and J. P. Gazeau, *Continuous frames in Hilbert spaces*, Annals of Physics **222** (1993), 1–37. <https://doi.org/10.1006/aphy.1993.1016>
- [3] E. Alizadeh and V. Sadri, *On continuous weaving G -frames in Hilbert spaces*, Wavelets and Linear Algebra **7**(1) (2020), 23–36. <https://doi.org/10.22072/wala.2020.114423.1248>
- [4] E. Alizadeh, A. Rahimi, E. Osgooei and M. Rahman, *Continuous K - G -fusion frames in Hilbert spaces*, TWMS J. Pure Appl. Math. **11**(1) (2021), 44–55.
- [5] E. Alizadeh and V. Sadri, *Construction of weaving continuous g -frames for operators in Hilbert spaces*, Probl. Anal. Issues Anal. **10**(2) (2021), 3–17. <https://doi.org/10.15393/j3.art.2021.9310>
- [6] P. Balazs, J. P. Antonie and A. Grybos, *Weighted and controlled frames: Mutual relationship and first numerical properties*, Int. J. Wavelets Multiresolut. Inf. Process. **14**(1) (2010), 109–132. <https://doi.org/10.1142/S0219691310003377>
- [7] T. Bemrose, P. G. Casazza, K. Grochenic, M. C. Lammers and R. G. Lynch, *Weaving frames*, Operators and Matrices **10**(4) (2016), 1093–1116. <https://doi.org/10.7153/oam-10-61>
- [8] O. Christensen, *An Introduction to Frames and Riesz Bases*, Birkhauser, 2008.
- [9] P. Casazza and G. Kutyniok, *Frames of subspaces*, Contemp. Math. **345** (2004), 87–114. <https://doi.org/10.1090/conm/345/06242>
- [10] P. Casazza and O. Christensen, *Perturbation of operators and applications to frame theory*, J. Fourier Anal. Appl. **3** (1997), 543–557. <https://doi.org/10.1007/BF02648883>
- [11] I. Daubechies, A. Grossmann and Y. Mayer, *Painless nonorthogonal expansions*, J. Math. Phys. **27**(5) (1986), 1271–1283. <https://doi.org/10.1063/1.527388>
- [12] R. G. Douglas, *On majorization, factorization, and range inclusion of operators on Hilbert space*. Proc. Amer. Math. Soc. **17** (1966), 413–415. <https://doi.org/10.1080/03081087.2017.1402859>

- [13] R. J. Duffin and A. C. Schaeffer, *A class of nonharmonic Fourier series*, Trans. Amer. Math. Soc. **72** (1952), 341–366.
- [14] M. H. Faroughi, A. Rahimi and R. Ahmadi, *GC-fusion frames*, Methods Funct. Anal. Topology **16**(2) (2010), 112–119.
- [15] P. Gavruta, *On the duality of fusion frames*, J. Math. Anal. Appl. **333** (2007), 871–879. <https://doi.org/10.1016/j.jmaa.2006.11.052>
- [16] L. Gavruta, *Frames for operator*, Appl. Comput. Harmon. Anal. **32**(1) (2012), 139–144. <https://doi.org/10.1016/j.acha.2011.07.006>
- [17] S. Garg, K. L. Vashisht and G. Verma, *On weaving fusion frames for Hilbert spaces*, International Conference on Sampling Theory and Applications (SampTA) (2017), 381–385. <https://doi.org/10.1109/SAMPSTA.2017.8024363>
- [18] P. Ghosh and T. K. Samanta, *Stability of dual g -fusion frame in Hilbert spaces*, Methods Funct. Anal. Topology **26**(3) (2020), 227–240.
- [19] P. Ghosh and T. K. Samanta, *Generalized atomic subspaces for operators in Hilbert spaces*, Math. Bohem. **147**(2) (2022), 325–345. <https://doi.org/10.21136/MB.2021.0130-20>
- [20] P. Ghosh and T. K. Samanta, *Generalized fusion frame in tensor product of Hilbert spaces*, J. Indian Math. Soc. **89** (1–2) (2022), 58–71. <https://doi.org/10.18311/jims/2022/29307>
- [21] P. Ghosh and T. K. Samanta, *Continuous controlled generalized fusion frames in Hilbert spaces*, J. Indian Math. Soc. (to appear).
- [22] G. Kaiser, *A Friendly Guide to Wavelets*, Birkhauser, 1994.
- [23] A. Khosravi and K. Musazadeh, *Controlled fusion frames*, Methods Funct. Anal. Topology **18**(3) (2012), 256–265.
- [24] D. Li, J. Leng and T. Huang, *On weaving g -frames for Hilbert spaces*, Complex Anal. Oper. Theory **14**(33) (2020). <https://doi.org/10.1007/s11785-020-00991-7>
- [25] M. Mohammadrezaee, M. Rashidi-Kouchi, A. Nazari and A. Oloomi, *Woven g -fusion frames in Hilbert spaces*, Sahand Communications in Mathematical Analysis **18**(3) (2021), 133–151. <https://doi.org/10.22130/scma.2021.137940.870>
- [26] M. Nouri, A. Rahimi and Sh. Najafizadeh, *Controlled K -frames in Hilbert spaces*, Int. J. Anal. Appl. **4**(2) (2015), 39–50.
- [27] A. Rahimi and A. Fereydooni, *Controlled g -frames and their g -multipliers in Hilbert spaces*, An. Stiint. Univ. “Ovidius” Constanta Ser. Mat. **21**(2) (2013), 223–236. <https://doi.org/10.2478/auom-2013-0035>
- [28] G. Rahimlou, V. Sadri and R. Ahmadi, *Construction of controlled K - g -fusion frame in Hilbert spaces*, UPB Scientific Bulletin, Series A **82**(1) (2020).
- [29] R. Rezapour, A. Rahimi, E. Osgooei and H. Dehghan, *Controlled weaving frames in Hilbert spaces*, Infinite Dimensional Analysis Quantum Probability and Related Topics **22**(1) (2019), Paper ID 1950003. <https://doi.org/10.1142/S0219025719500036>
- [30] R. Rezapour, A. Rahimi, E. Osgooei and H. Dehghan, *Continuous controlled K - g -frames in Hilbert spaces*, Indian J. Pure Appl. Math. **50** (2019), 863–875. <https://doi.org/10.1007/s13226-019-0359-y>
- [31] V. Sadri, Gh. Rahimlou, R. Ahmadi and R. Zarghami Farfar, *Generalized fusion frames in Hilbert spaces*, Infinite Dimensional Analysis Quantum Probability and Related Topics **23**(2) (2020), Paper ID 2050015. <https://doi.org/10.1142/S0219025720500150>
- [32] V. Sadri, G. Rahimlou and R. Ahmadi, *K - g -fusion woven in Hilbert spaces*, TWMS J. Pure Appl. Math. **11**(3) (2021), 947–958.
- [33] V. Sadri, R. Ahmadi and G. Rahimlou, *On continuous weaving fusion frames in Hilbert spaces*, Int. J. Wavelets Multiresolut. Inf. Process. **18**(5) (2020), Paper ID 2050035, 17 pages. <https://doi.org/10.1142/S0219691320500356>
- [34] H. Shakoory, R. Ahmadi, N. Behzadi and S. Nami, *(C, C') -Controlled g -fusion frames*, Iran. J. Math. Sci. **18**(1) (2023), 179–191. <https://doi.org/10.52547/ijmsi.18.1.179>

- [35] W. Sun, *G-frames and G-Riesz bases*, J. Math. Anal. **322**(1) (2006), 437–452. <https://doi.org/10.1016/j.jmaa.2005.09.039>
- [36] L. K. Vashisht and Deepshikha, *On continuous weaving frames*, Adv. Pure Appl. Math. **8**(1) (2017), 15–31. <https://doi.org/10.1515/apam-2015-0077>

¹DEPARTMENT OF PURE MATHEMATICS,
UNIVERSITY OF CALCUTTA,
35, BALLYGUNGE CIRCULAR ROAD, KOLKATA, 700019, WEST BENGAL, INDIA
Email address: prasenjitpuremath@gmail.com

²DEPARTMENT OF MATHEMATICS,
ULUBERIA COLLEGE,
ULUBERIA, HOWRAH, 711315, WEST BENGAL, INDIA
Email address: mumpu_tapas5@yahoo.co.in