

ON HADAMARD-CAPUTO IMPLICIT FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS WITH BOUNDARY FRACTIONAL CONDITIONS

AHMED A. HAMOUD^{1*}, CHAHRA KECHAR², ABDELOUAHEB ARDJOUNI²,
HOMAN EMADIFAR^{3,4,5}, ATUL KUMAR⁶, AND LAITH ABUALIGAH^{7,8,9,10,11,12}

ABSTRACT. The purpose of this paper is to investigate the existence and uniqueness of solutions for nonlinear fractional implicit integro-differential equations of Hadamard-Caputo type with fractional boundary conditions. The reasoning is inspired by diverse classical fixed point theory, such as the Schauder and Banach fixed point theorems. The theoretical findings are illustrated through an example.

1. INTRODUCTION

In mathematical analysis, fractional calculus (FC) is a subject that studies different approaches of defining non-integer order derivatives (i.e., fractional differential calculus (FDC)) and integrals (i.e., fractional integral calculus (FIC)). Fractional calculus is widely and efficiently used to describe many phenomena arising in physics, engineering, bioengineering and biomedical sciences, finance, viscoelasticity, control theory, stochastic processes and economy. Recently, fractional differential equations (FDEs) have attracted many authors (see for example [1–3, 8, 13] and references therein).

By flipping the differential and integral sections of the Hadamard derivative, a novel method known as the Hadamard-Caputo derivative is created. The primary distinction between the Hadamard fractional derivative and the Hadamard-Caputo fractional derivative, notwithstanding the various demands placed on the function itself, is that the Hadamard-Caputo derivative of a constant is zero [24]. The most

Key words and phrases. Fixed point theorems, existence, uniqueness, Hadamard-Caputo fractional derivative, integro-differential equation.

2020 *Mathematics Subject Classification.* Primary: 34A09. Secondary: 26A33, 34B10.
<https://doi.org/10.46793/KgJMat2603.491H>

Received: August 29, 2023.

Accepted: February 10, 2024.

significant benefit of the Hadamard-Caputo derivative is that it gave rise to a new concept that can be used to establish the integer order beginning conditions for fractional.

For more details and properties of Hadamard-Caputo derivative and Hadamard fractional derivatives, integrals see [10, 11, 26].

The implicit fractional differential equations (IFDEs) are a very important class of fractional differential equations. This type of equation is derived from the implicit ordinary differential equation (IODE) of the following form

$$H(\varrho, G(\varrho), G'(\varrho), \dots, G^{(n-1)}(\varrho)) = 0,$$

with different kind of initial or boundary conditions, for more details see [6, 14, 15, 23].

Benchohra et al. [7, 9] and Nieto et al. [28, 30] have initiated the study of implicit fractional differential equations (IFDEs) of the form

$$D^\alpha G(\varrho) = H(\varrho, G(\varrho), D^\alpha G(\varrho)),$$

with different kind of initial or boundary conditions. This sort of equation is crucial in many different fields of science and engineering [34].

In [33], Vivek et al. showed that a class of boundary value systems for nonlinear IFDEs with complex order have a solution and are stable

$$\begin{aligned} {}^c D^\theta G(\varrho) &= H(\varrho, G(\varrho), {}^c D^\theta G(\varrho)), \quad \theta = m + i\alpha, \quad \varrho \in Y := [0, \chi], \\ aG(0) + bG(\chi) &= c, \end{aligned}$$

where $\theta \in \mathbb{C}$, ${}^c D^\theta$ is the Caputo fractional derivative. Suppose $m \in (0, 1]$, $\alpha \in \mathbb{R}_+$, $0 < \alpha < 1$, $H : Y \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function. Further $a, b, c \in \mathbb{R}$ with $a + b \neq 0$. The results are based upon the Schaefer's fixed point theorem and Banach contraction principle.

In [27] Karthikeyan and Arul studied the uniqueness of integral BVP for IFDEs involving Hadamard-Caputo fractional derivative

$$\begin{aligned} {}^{CH} D^\theta \zeta(\varrho) &= H(\varrho, \zeta(\varrho), {}^{CH} D^\theta \zeta(\varrho)), \quad \varrho \in \xi := [n, \chi], \\ \zeta(n) &= 0, \quad \zeta(\chi) = \Omega \int_n^\sigma \zeta(\omega) d\omega, \quad n < \sigma < \chi, \end{aligned}$$

where $\Omega \in \mathbb{R}$, $n < \sigma < \chi$, $1 < \theta \leq 2$, ${}^{CH} D^\theta$ is the Hadamard-Caputo fractional derivative, and $H : \xi \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function.

In [12] N. Dardar established the uniqueness of solutions for the system:

$$\begin{aligned} {}^C_H D^r v(\varrho) &= W(\varrho, v(\varrho), {}^C_H D^r v(\varrho)), \\ v(1) &= 0, \quad \alpha {}_H I^q v(\eta) + \beta {}^C_H D^\gamma v(\Psi) = \lambda, \end{aligned}$$

by using different fixed point theorem.

Recently, many authors focus on the development of techniques for discussing the solutions of FIDEs.

Balachandran and Trujillo [5], investigated the existence of a unique solution for FIDEs with boundary value conditions.

$${}^C\mathcal{D}^\alpha f(v) = W(v, f(v)) + \int_{v_0}^v K(v, \varrho, f(\varrho)) d\varrho, \quad 0 < \alpha \leq 1.$$

In [32] the authors investigated the uniqueness of solution for iterative integro-differential system:

$$\begin{aligned} \mathcal{D}^\alpha w(v) &= Q(v) + \int_0^\varrho H(v, s) w(\lambda w(s)) ds, \\ w(0) &= w_0. \end{aligned}$$

In [17] A. A. Hamoud established the uniqueness and stability for fractional nonlinear Fredholm–Volterra system:

$$\begin{aligned} {}^C\mathcal{D}^\alpha w(\varrho) &= H(\varrho) + \int_0^\varrho \vartheta(\varrho, s) w(w(s)) ds + \int_0^\Psi \theta(\varrho, s) w(w(s)) ds, \\ aw(0) + bw(\Psi) &= c, \quad a, b, c, \in \mathbb{R}. \end{aligned}$$

For some other results on FIDE, see [4, 5, 16, 18–22, 25, 29].

Motivated by the above papers and the reference [12], we study the theoretical analysis of solutions for a class of system for nonlinear implicit FIDEs of Hadamard-Caputo type with fractional boundary conditions

$$(1.1) \quad \begin{aligned} {}^C_H D^r v(\varrho) &= f\left(\varrho, v(\varrho), {}^C_H D^r v(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right), \\ v(1) &= 0, \quad \alpha {}_H I^q v(\eta) + \beta {}^C_H D^\gamma v(\Psi) = \lambda, \end{aligned}$$

where ${}_H I^q$ is the standard Hadamard fractional integral, ${}^C_H D^r$ is the Hadamard-Caputo fractional derivative, $f : \xi \times \mathbb{R}^3 \rightarrow \mathbb{R}$, $K : \xi \times \xi \times \mathbb{R} \rightarrow \mathbb{R}$ are given functions, $\eta \in \xi =: (1, \Psi)$, $\Psi > 1$ and α, λ, β are real constants.

2. PRELIMINARIES AND BACKGROUND MATERIALS

Let us introduce some necessary notations and definitions which will be utilised throughout the entire process [28, 29, 31, 33–35].

We represent by the symbol $C(\xi, \mathbb{R})$ the space of all continuous functions $v : \xi \rightarrow \mathbb{R}$ in Banach space with the supremum norm

$$\|v\|_\infty = \sup \{|v(\varrho)| : \varrho \in \xi\}.$$

Let now $[b, c]$, $-\infty < b < c < +\infty$, is finite interval and we suppose $AC([b, c], \mathbb{R})$ is the space of functions $\psi : [b, c] \rightarrow \mathbb{R}$ that are absolutely continuous.

Assume $\delta = \varrho \frac{d}{dt} d\varrho$ is the Hadamard derivative, $\delta^n = \delta(\delta^{n-1})$, we consider the set of functions:

$$AC_\delta^n([b, c], \mathbb{R}) = \left\{ \psi : [b, c] \rightarrow \mathbb{R} : \delta^{n-1} \psi(\varrho) \in AC([b, c], \mathbb{R}) \right\}.$$

Definition 2.1 ([28]). The Hadamard fractional integral of order $\alpha > 0$ for a continuous function $\psi : [1, +\infty) \rightarrow \mathbb{R}$ is given by

$${}_H I_1^\alpha \psi(\varrho) = \frac{1}{\Gamma(\alpha)} \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\alpha-1} \psi(s) \frac{ds}{s},$$

where $\log(\cdot) = \log_e(\cdot)$ and $\Gamma(\cdot)$ is Gamma function.

Definition 2.2 ([31]). For a function $\psi \in AC_\delta^n([b, c], \mathbb{R})$, the Hadamard-Caputo fractional derivative of order α is given by

$${}^C D_1^\alpha \psi(\varrho) = \frac{1}{\Gamma(n-\alpha)} \left(t \frac{d}{dt}\right)^n \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{n-\alpha-1} \psi(s) \frac{ds}{s}, \quad n-1 < \alpha < n,$$

where $\delta^n = \left(\varrho \frac{d}{d\varrho}\right)^n$, $n = [\alpha] + 1$, and $[\alpha]$ denotes the integer part of α .

Lemma 2.1 ([24]). Let $\psi \in AC_\delta^n[b, c]$ or $\psi \in C_\delta^n[b, c]$ and $\alpha \in \mathbb{C}$. Then,

$${}_H I_b^\alpha \left({}^C D_b^\alpha \psi\right)(\varrho) = \psi(\varrho) - \sum_{k=0}^{n-1} \frac{\delta^{(k)} \psi(b)}{k!} \left(\log \frac{\varrho}{b}\right)^k.$$

Proposition 2.1 ([24]). Let $\alpha > 0$, $\beta > 0$, $n = [\alpha] + 1$, and $b > 0$. Then,

$$\begin{aligned} \left({}_H I_{b^+}^\alpha \left(\log \frac{v}{b}\right)^{\beta-1}\right)(v) &= \frac{\Gamma(\beta)}{\Gamma(\beta+\alpha)} \left(\log \frac{v}{b}\right)^{\beta+\alpha-1}, \\ \left({}^C D_{b^+}^\alpha \left(\log \frac{v}{b}\right)^{\beta-1}\right)(v) &= \frac{\Gamma(\beta)}{\Gamma(\beta-\alpha)} \left(\log \frac{v}{b}\right)^{\beta-\alpha-1}, \quad \alpha < \beta. \end{aligned}$$

Theorem 2.1. [24] Let $v(\varrho) \in AC_\delta^n[b, c]$, $0 < b < c < +\infty$ and $\alpha \geq 0$, $\beta \geq 0$. Then,

$$\begin{aligned} {}^C D_b^\alpha \left(I^\beta v\right)(\varrho) &= \left(I^{\beta-\alpha} v\right)(\varrho), \\ {}^C D^\alpha \left({}^C D^\beta\right)(\varrho) &= {}^C D^{\alpha+\beta}(\varrho). \end{aligned}$$

Theorem 2.2 (Schauder’s fixed point [35]). Suppose that E be a Banach space, and P be a nonempty, convex and closed subset of E . Assume that $\mathcal{A} : P \rightarrow P$ be a continuous mapping and $\mathcal{A}(P)$ is a relatively compact subset of E . Then \mathcal{A} admits at least one fixed point in P .

3. MAIN RESULTS

Definition 3.1. A function $v \in AC_\delta^2(\xi, \mathbb{R})$ is said to be a solution of the system (1.1) if v satisfies the equation ${}^C D_H^\kappa v(\varrho) = f\left(\varrho, v(\varrho), {}^C D_H^\kappa v(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right)$, and satisfies the conditions $v(1) = 0$, $\alpha {}_H I^q v(\eta) + \beta {}^C D_H^\gamma v(\Psi) = \lambda$.

In what follows, we present the following lemma to show the existence of solutions of the system (1.1).

Lemma 3.1 ([12]). *Suppose that $h : [0, +\infty) \rightarrow \mathbb{R}$ is a continuous. A function v is a solution of the following system*

$$v(\varrho) = \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} h(s) \frac{ds}{s} + \frac{\log \varrho}{\Lambda} \left[\lambda - \frac{\alpha}{\Gamma(\kappa+q)} \int_1^\eta \left(\log \frac{\eta}{s}\right)^{\kappa+q-1} h(s) \frac{ds}{s} - \frac{\beta}{\Gamma(\kappa-\gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s}\right)^{\kappa-\gamma-1} h(s) \frac{ds}{s} \right],$$

where

$$\Lambda = \frac{\alpha (\log \eta)^{q+1}}{\Gamma(q+2)} + \frac{\beta (\log \Psi)^{1-\gamma}}{\Gamma(2-\gamma)},$$

is equivalent to v is a solution of the following problem

$${}^C_H D^\kappa v(\varrho) = h(\varrho) \\ v(1) = 0, \quad \alpha {}^C_H I^q v(\eta) + \beta {}^C_H D^\gamma v(\Psi) = \lambda, \quad q, \gamma \in [0, 1].$$

Now, we prove the existence of a solution of the system (1.1).

Our hypotheses are as follows.

(H1) $f : \xi \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is continuous.

(H2) There exist three constants $L_1 > 0$, $0 < L_2 < 1$ and $L_3 > 0$ as follows

$$|f(\varrho, \varsigma, \varphi, w) - f(\varrho, \bar{\varsigma}, \bar{\varphi}, \bar{w})| \leq L_1 |\varsigma - \bar{\varsigma}| + L_2 |\varphi - \bar{\varphi}| + L_3 |w - \bar{w}|,$$

for each $\varsigma, \varphi, w, \bar{\varsigma}, \bar{\varphi}$ and $\bar{w} \in \mathbb{R}$ for a.e. $\varrho \in \xi$.

(H3) There exists a function $k(\varrho, s) \in C[0, 1]$, as follows:

$$|K(\varrho, s, v(s)) - K(\varrho, s, y(s))| \leq k(\varrho, s) |v(s) - y(s)|.$$

Also, we denote

$$\sigma_k = \sup_{\varrho \in \xi} \int_1^\varrho |K(\varrho, s, 0)| ds, \\ \sigma_k^* = \sup_{\varrho \in \xi} \sigma_k(\varrho), \\ \beta_k = \sup_{\varrho \in \xi} \int_1^\varrho |k(\varrho, s)| ds.$$

Theorem 3.1. *Let the assumptions (H1)-(H3) be true. If*

$$(3.1) \quad \rho := \frac{L_1 + \beta_k L_3}{1 - L_2} \left[\frac{(\log \Psi)^\kappa}{\Gamma(\kappa+1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa+q+1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa-\gamma+1)} \right] < 1,$$

then the system (1.1) has a unique solution $v \in AC_\delta^2(\xi, \mathbb{R})$ on ξ .

Proof. Let $F : C(\xi, \mathbb{R}) \rightarrow C(\xi, \mathbb{R})$ be defined as

$$Fv(\varrho) = \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} \sigma_v(s) \frac{ds}{s}$$

$$(3.2) \quad + \frac{\log \varrho}{\Lambda} \left[\lambda - \frac{\alpha}{\Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s} \right)^{\kappa+q-1} \sigma_v(s) \frac{ds}{s} - \frac{\beta}{\Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s} \right)^{\kappa-\gamma-1} \sigma_v(s) \frac{ds}{s} \right],$$

where

$$\sigma_v(s) = f \left(s, v(s), D^\kappa v(s), \int_1^s K(s, \tau, v(\tau)) d\tau \right).$$

Clearly, the fixed points of F are solutions of the system (1.1).

Let $v, y \in AC_\delta^2(\xi, \mathbb{R})$. Then for each $\varrho \in \xi$ we obtain

$$(3.3) \quad \begin{aligned} |(Fv)(\varrho) - (Fy)(\varrho)| &= \left| \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{t}{s} \right)^{\kappa-1} \sigma_v(s) \frac{ds}{s} + \frac{\log \varrho}{\Lambda} \left[\lambda - \frac{\alpha}{\Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s} \right)^{\kappa+q-1} \sigma_v(s) \frac{ds}{s} - \frac{\beta}{\Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s} \right)^{\kappa-\gamma-1} \sigma_v(s) \frac{ds}{s} \right] - \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s} \right)^{\kappa-1} \sigma_y(s) \frac{ds}{s} - \frac{\log \varrho}{\Lambda} \left[\lambda - \frac{\alpha}{\Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s} \right)^{\kappa+q-1} \sigma_y(s) \frac{ds}{s} - \frac{\beta}{\Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s} \right)^{\kappa-\gamma-1} \sigma_y(s) \frac{ds}{s} \right] \right| \\ &\leq \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s} \right)^{\kappa-1} |\sigma_v(s) - \sigma_y(s)| \frac{ds}{s} + \frac{|\alpha| \log \varrho}{|\Lambda| \Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s} \right)^{\kappa+q-1} |\sigma_v(s) - \sigma_y(s)| \frac{ds}{s} + \frac{|\beta| \log \varrho}{|\Lambda| \Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s} \right)^{\kappa-\gamma-1} |\sigma_v(s) - \sigma_y(s)| \frac{ds}{s}, \end{aligned}$$

with

$$\sigma_v(\varrho) = f \left(\varrho, v(\varrho), \sigma_v(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds \right)$$

and

$$\sigma_y(\varrho) = f \left(\varrho, y(\varrho), \sigma_y(\varrho), \int_1^\varrho K(\varrho, s, y(s)) ds \right).$$

By using (H2), we find

$$\begin{aligned} \left| \int_1^\varrho K(\varrho, s, v(s)) ds - \int_1^\varrho K(\varrho, s, y(s)) ds \right| &\leq \int_1^\varrho k(\varrho, s) |v(\varrho) - y(\varrho)| ds \\ &\leq \beta_k \|v - y\| \end{aligned}$$

and

$$\begin{aligned} |\sigma_v(\varrho) - \sigma_y(\varrho)| &= \left\| f\left(\varrho, v(\varrho), \sigma_v(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right) \right. \\ &\quad \left. - f\left(\varrho, y(\varrho), \sigma_y(\varrho), \int_1^\varrho K(\varrho, s, y(s)) ds\right) \right\| \\ &\leq L_1 |v(\varrho) - y(\varrho)| + L_2 |\sigma_v(\varrho) - \sigma_y(\varrho)| + L_3 \beta_k \|v - y\|. \end{aligned}$$

Thus,

$$(3.4) \quad |\sigma_v(\varrho) - \sigma_y(\varrho)| \leq \frac{L_1 + \beta_k L_3}{1 - L_2} \|v - y\|.$$

Replacing (3.4) in (3.3), we obtain

$$\begin{aligned} & |(Fv)(\varrho) - (Fy)(\varrho)| \\ & \leq \frac{1}{\Gamma(\kappa)} \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} |v(s) - y(s)| \frac{ds}{s} \\ & \quad + \frac{|\alpha| \log \varrho}{|\Lambda| \Gamma(\kappa + q)} \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \int_1^\eta \left(\log \frac{\eta}{s}\right)^{\kappa+q-1} |v(s) - y(s)| \frac{ds}{s} \\ & \quad + \frac{|\beta| \log \varrho}{|\Lambda| \Gamma(\kappa - \gamma)} \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \int_1^\Psi \left(\log \frac{\Psi}{s}\right)^{\kappa-\gamma-1} |v(s) - y(s)| \frac{ds}{s} \\ & \leq \left[\frac{1}{\Gamma(\kappa)} \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} \frac{ds}{s} \right. \\ & \quad + \frac{|\alpha| \log \varrho}{|\Lambda| \Gamma(\kappa + q)} \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \int_1^\eta \left(\log \frac{\eta}{s}\right)^{\kappa+q-1} \frac{ds}{s} \\ & \quad \left. + \frac{|\beta| \log \varrho}{|\Lambda| \Gamma(\kappa - \gamma)} \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \int_1^\Psi \left(\log \frac{\Psi}{s}\right)^{\kappa-\gamma-1} \frac{ds}{s} \right] |v(s) - y(s)| \\ & \leq \left(\frac{L_1 + \beta_k L_3}{1 - L_2}\right) \\ & \quad \times \left[\frac{(\log \Psi)^\kappa}{\Gamma(\kappa + 1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa + q + 1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa - \gamma + 1)} \right] |v(s) - y(s)|. \end{aligned}$$

Hence,

$$\|(Fv)(\varrho) - (Fy)(\varrho)\|_\infty \leq \rho \|v - y\|_\infty,$$

for $v, y \in AC_\delta^2(\xi, \mathbb{R})$, where

$$\rho := \frac{L_1 + \beta_k L_3}{1 - L_2} \left[\frac{(\log \Psi)^\kappa}{\Gamma(\kappa + 1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa + q + 1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa - \gamma + 1)} \right].$$

Consequently by (3.1), F is a contraction. As a consequence of Banach's contraction principle, we conclude that F admits a unique fixed point which is the solution of the system (1.1). □

Next, we study the second result, by using the fixed point theorem of Schauder.

(H4) There exist $p, \varphi, w, z \in C(\xi, \mathbb{R}_+)$, with $z^* = \sup_{\varrho \in \xi} z(\varrho) < 1$, $\omega^* = \sup_{\varrho \in \xi} \omega(\varrho) < 1$, $\varphi^* = \sup_{\varrho \in \xi} \varphi(\varrho) < 1$ and $p^* = \sup_{\varrho \in \xi} p(\varrho) < 1$, such that

$$f(\varrho, \varsigma, \varphi, w) \leq p(\varrho) + \varphi(\varrho) |\varsigma| + \omega(\varrho) |\varphi| + z(\varrho) |w|,$$

for any $\varsigma, \varphi, w \in \mathbb{R}$ for a.e. $\varrho \in \xi$.

Theorem 3.2. *Assume that (H1), (H3) and (H4) are true. Moreover if*

$$(3.5) \quad \omega^* + M(v^* + \beta_k z^*) < 1,$$

with

$$M := \frac{(\log \Psi)^\kappa}{\Gamma(\alpha + 1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa + q + 1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa - \gamma + 1)},$$

then the system (1.1) admits at least one solution.

Proof. Let

$$R \geq \frac{M(p^* + \sigma_k^* z^*) + \frac{|\lambda|(1-\omega^*) \log \Psi}{|\Lambda|}}{1 - (\omega^* + M(\varphi^* + \beta_k z^*))},$$

and consider $\Delta_R = \{v \in C(\xi, \mathbb{R}) : \|v\|_\infty \leq R\}$. It is clear that the subset Δ_R is closed, convex and bounded. We will use Schauder's fixed point theorem to demonstrate that F defined by (3.2) admits a fixed point.

This could be proved through three steps.

Step 1: F is a continuous mapping.

Let $\{v_n\}$ be a sequence as follows $v_n \rightarrow v$ in $AC_\delta^2(\xi, \mathbb{R})$. Then for any $\varrho \in \xi$

$$\begin{aligned} & |(Fv_n)(\varrho) - (Fv)(\varrho)| \\ & \leq \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} |\psi_n(s) - \psi(s)| \frac{ds}{s} \\ & \quad + \frac{|\alpha| \log \varrho}{|\Lambda| \Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s}\right)^{\kappa+q-1} |\psi_n(s) - \psi(s)| \frac{ds}{s} \\ & \quad + \frac{|\beta| \log \varrho}{|\Lambda| \Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s}\right)^{\kappa-\gamma-1} |\psi_n(s) - \psi(s)| \frac{ds}{s} \\ & \leq \left[\frac{(\log \Psi)^\kappa}{\Gamma(\alpha + 1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa + q + 1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa - \gamma + 1)} \right] |\psi_n(s) - \psi(s)|, \end{aligned}$$

where $\psi, \psi_n \in C(\xi, \mathbb{R})$ are

$$\begin{aligned} \psi(\varrho) &= f\left(\varrho, v(\varrho), \psi(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right), \\ \psi_n(\varrho) &= f\left(\varrho, v(\varrho), \psi_n(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right). \end{aligned}$$

Since ψ is a continuous functions (i.e., f is continuous), then from the Lebesgue theorem of dominated convergence, we get

$$\|F(v_n)(\varrho) - (Fv)(\varrho)\|_\infty \rightarrow 0, \quad \text{as } n \rightarrow +\infty$$

Hence, $F(v_n)(\varrho) \rightarrow (Fv)(\varrho)$ as $n \rightarrow +\infty$ which implies that F is continuous.

Step 2: $F(\Delta_R) \subset \Delta_R$.

Let $v \in \Delta_R$. We demonstrate $F(v) \in \Delta_R$, for all $\varrho \in \xi$, we find

$$\begin{aligned} |Fv(\varrho)| &\leq \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} |\psi(s)| \frac{ds}{s} \\ &\quad + \frac{|\alpha| \log \varrho}{|\Lambda| \Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s}\right)^{\kappa+q-1} |\psi(s)| \frac{ds}{s} \\ (3.6) \quad &\quad + \frac{|\beta| \log \varrho}{|\Lambda| \Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s}\right)^{\kappa-\gamma-1} |\psi(s)| \frac{ds}{s} + \frac{|\lambda| \log \varrho}{|\Lambda|}, \end{aligned}$$

where $\psi \in C(\xi, \mathbb{R})$ is

$$\psi(\varrho) = f\left(\varrho, v(\varrho), \psi(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right).$$

From (H3), we get

$$\begin{aligned} &\left| \int_1^\varrho K(\varrho, s, v(s)) ds - \int_1^\varrho K(\varrho, s, 0) ds + \int_1^\varrho K(\varrho, s, 0) ds \right| \\ &\leq \int_1^\varrho |k(\varrho, s)| |v(\varrho)| ds + \int_1^\varrho |K(\varrho, s, 0)| ds \\ &\leq \beta_k \|v\|_\infty + \sigma_k(\varrho). \end{aligned}$$

From (H4), for $\varrho \in \xi$ we get

$$\begin{aligned} |\psi(\varrho)| &= \left| f\left(\varrho, v(\varrho), \psi(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds\right) \right| \\ &\leq p(\varrho) + \varphi(\varrho) |v(\varrho)| + \omega(\varrho) |\psi(\varrho)| + \beta_k z(\varrho) \|v\|_\infty + \sigma_k(\varrho) z(\varrho) \\ &\leq p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) \|v\|_\infty + \omega^* |\psi(\varrho)|. \end{aligned}$$

Hence,

$$(3.7) \quad |\psi(\varrho)| \leq \frac{p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R}{1 - \omega^*}.$$

In the inequality (3.6), we obtain by substituting (3.7)

$$\begin{aligned} |Fv(\varrho)| &\leq \frac{1}{\Gamma(\kappa)} \int_1^\varrho \left(\log \frac{\varrho}{s}\right)^{\kappa-1} |\psi(s)| \frac{ds}{s} + \frac{|\alpha| \log \varrho}{|\Lambda| \Gamma(\kappa + q)} \int_1^\eta \left(\log \frac{\eta}{s}\right)^{\kappa+q-1} |\psi(s)| \frac{ds}{s} \\ &\quad + \frac{|\beta| \log \varrho}{|\Lambda| \Gamma(\kappa - \gamma)} \int_1^\Psi \left(\log \frac{\Psi}{s}\right)^{\kappa-\gamma-1} |\psi(s)| \frac{ds}{s} + \frac{|\lambda| \log \varrho}{|\Lambda|} \\ &\leq \left(\frac{p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R}{1 - \omega^*} \right) \end{aligned}$$

$$\begin{aligned}
& \times \left[\frac{(\log \Psi)^\kappa}{\Gamma(\kappa+1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa+q+1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa-\gamma+1)} \right] + \frac{|\lambda| \log \Psi}{|\Lambda|} \\
& = \frac{M(p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R)}{1 - \omega^*} + \frac{|\lambda| \log \Psi}{|\Lambda|} \\
& \leq R.
\end{aligned}$$

Step 3: We demonstrate that the expression $F(\Delta_R)$ is equicontinuous.

It is clear from step 2 that $F(\Delta_R) \subset \Delta_R$ is bounded. For the $F(\Delta_R)$ equicontinuity.

Let $\mu_1, \mu_2 \in (1, \Psi]$, $\mu_1 < \mu_2$ and let $v \in \Delta_R$. Then

$$\begin{aligned}
& |(Fv)(\mu_2) - (Fv)(\mu_1)| \\
& = \left| \frac{1}{\Gamma(\kappa)} \int_1^{\mu_1} \left(\log \frac{\mu_2}{s} \right)^{\kappa-1} \psi(s) \frac{ds}{s} - \frac{1}{\Gamma(\kappa)} \int_1^{\mu_2} \left(\log \frac{\mu_1}{s} \right)^{\kappa-1} \psi(s) \frac{ds}{s} \right| \\
& = \left| \frac{1}{\Gamma(\kappa)} \int_1^{\mu_1} \left[\left(\log \frac{\mu_2}{s} \right)^{\kappa-1} - \left(\log \frac{\mu_1}{s} \right)^{\kappa-1} \right] \psi(s) \frac{ds}{s} \right. \\
& \quad \left. + \frac{1}{\Gamma(\kappa)} \int_{\mu_1}^{\mu_2} \left(\log \frac{\mu_2}{s} \right)^{\kappa-1} \psi(s) \frac{ds}{s} \right| \\
& \leq \frac{|\psi(s)|}{\Gamma(\kappa)} \left| \int_1^{\mu_1} \left[\left(\log \frac{\mu_2}{s} \right)^{\kappa-1} - \left(\log \frac{\mu_1}{s} \right)^{\kappa-1} \right] \frac{ds}{s} \right| \\
& \quad + \frac{|\psi(s)|}{\Gamma(\kappa)} \left| \int_{\mu_1}^{\mu_2} \left(\log \frac{\mu_2}{s} \right)^{\kappa-1} \frac{ds}{s} \right| \\
& \leq \frac{p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R}{(1 - \omega^*) \Gamma(\kappa)} \left| \int_1^{\mu_1} \left[\left(\log \frac{\mu_2}{s} \right)^{\kappa-1} - \left(\log \frac{\mu_1}{s} \right)^{\kappa-1} \right] \frac{ds}{s} \right| \\
& \quad + \frac{p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R}{(1 - \omega^*) \Gamma(\kappa)} \left| \int_{\mu_1}^{\mu_2} \left(\log \frac{\mu_2}{s} \right)^{\kappa-1} \frac{ds}{s} \right| \\
& \leq \frac{p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R}{(1 - \omega^*) \Gamma(\kappa+1)} \left[\left| (\log \mu_1)^\kappa + \left(\log \frac{\mu_2}{\mu_1} \right)^\kappa - (\log \mu_2)^\kappa \right| + \left| \left(\log \frac{\mu_2}{\mu_1} \right)^\kappa \right| \right] \\
& \leq \frac{p^* + \sigma_k^* z^* + (\varphi^* + \beta_k z^*) R}{(1 - \omega^*) \Gamma(\kappa+1)} |(\log \mu_1)^\kappa - (\log \mu_2)^\kappa| \\
& \rightarrow 0, \quad \text{as } \mu_1 \rightarrow \mu_2.
\end{aligned}$$

The Arzela-Ascoli theorem shows that F is relatively compact in both scenarios, and Schauder's fixed point theorem states that F has a fixed point. Then, F is a solution of the system (1.1). \square

4. APPLICATIONS

Example 4.1. Assume that the nonlinear system is,

$${}^C_H D^{\frac{3}{2}} v(\varrho) = f \left(\varrho, v(\varrho), {}^C_H D^{\frac{7}{5}} v(\varrho), \int_1^\varrho K(\varrho, s, v(s)) ds \right), \quad \varrho \in [1, e],$$

$$v(1) = 0, \quad \alpha {}_H I^{\frac{2}{3}} v(\eta) + \beta {}_H^C D^{\frac{3}{7}} v(\Psi) = \lambda.$$

We see that $\kappa = \frac{3}{2}, q = \frac{2}{3}, \gamma = \frac{3}{7}, \eta = 2, \alpha = \frac{1}{3}, \beta = 3, \lambda = \frac{4}{5}, \Psi = e$, and

$$f(\varrho, \varsigma, \varphi, w) = \frac{\varrho^4}{11 + \varrho^2} + \frac{|\varsigma|}{7\varrho^4(|\varsigma| + 3)} + \frac{3}{7\varrho^2} \cos \varphi + \frac{1}{\varrho^2} w, \quad \varrho \in [1, e], \varsigma, \varphi, w \in \mathbb{R}.$$

Clearly, f is a continuous, and for $\varrho \in [1, e]$ and $w, \varphi, \varsigma, \bar{w}, \bar{\varsigma}, \bar{\varphi} \in \mathbb{R}$, we get

$$|f(\varrho, \varsigma, \varphi, w) - f(\varrho, \bar{\varsigma}, \bar{\varphi}, \bar{w})| \leq L_1 |\varsigma - \bar{\varsigma}| + L_2 |\varphi - \bar{\varphi}| + L_3 |w - \bar{w}|,$$

with $L_1 = \frac{1}{21}, L_2 = \frac{3}{7}, L_3 = 1$ and

$$|K(\varrho, s, v) - K(\varrho, s, y)| \leq \frac{1}{3s} \log\left(\frac{\varrho}{s}\right)^{\kappa-1} |v - y|,$$

$$\beta_k = \sup \int_1^e k(\varrho, s) ds = \frac{1}{3} \int_1^e \log\left(\frac{\varrho}{s}\right)^{\kappa-1} \frac{ds}{s} = \frac{(\log \varrho)^\kappa}{3\kappa} \leq \frac{1}{3\kappa} = \frac{2}{9}.$$

Hence, conditions (H2) and (H3) are satisfied. Moreover,

$$\Lambda = \frac{\alpha (\log \eta)^{q+1}}{\Gamma(q+2)} + \frac{\beta (\log \Psi)^{1-\gamma}}{\Gamma(2-\gamma)} = 3.4887.$$

Thus,

$$\rho := \frac{L_1 + \beta_k L_3}{1 - L_2} \left[\frac{(\log \Psi)^\kappa}{\Gamma(\kappa + 1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa + q + 1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa - \gamma + 1)} \right] = 0.75728 < 1.$$

Thus, the given system has a unique solution $v \in C_\delta^n([1, e], \mathbb{R})$, according to Theorem 3.1,

$$|f(\varrho, \varsigma, \varphi, w)| \leq \frac{\varrho^4}{11 + \varrho^2} + \frac{|\varsigma|}{7\varrho^4(|\varsigma| + 3)} + \frac{3}{7\varrho^2} |\cos \varphi| + \frac{1}{3s} \log\left(\frac{\varrho}{s}\right)^{\kappa-1} |\sin w|,$$

so condition (H4) is satisfied with

$$p(\varrho) = \frac{\varrho^4}{11 + \varrho^2}, \quad \varphi(\varrho) = \frac{1}{21\varrho^4}, \quad \omega(\varrho) = \frac{3}{7\varrho^2}, \quad z(\varrho) = \frac{1}{3s} \log\left(\frac{t}{s}\right)^{\kappa-1}$$

and

$$\varphi^* = \frac{1}{21}, \quad \omega^* = \frac{3}{7}, \quad z^* = 1.$$

We have

$$M := \frac{(\log \Psi)^\kappa}{\Gamma(\alpha + 1)} + \frac{|\alpha| (\log \Psi) (\log \eta)^{\kappa+q}}{|\Lambda| \Gamma(\kappa + q + 1)} + \frac{|\beta| (\log \Psi)^{\kappa-\gamma+1}}{|\Lambda| \Gamma(\kappa - \gamma + 1)} = 1.9712.$$

We will demonstrate that condition (3.5) is true for $\Psi = e$. Indeed,

$$\omega^* + M(\varphi^* + \beta_k z^*) = 0.96048 < 1.$$

Simple calculations demonstrate that conditions of Theorem 3.2 are all satisfied. Example 4.1 must thus have at least one solution specified on $[1, e]$.

REFERENCES

- [1] R. P. Agarwal, M. Benchohra and B. A. Slimani, *Existence results for differential equations with fractional order and impulses*, Mem. Differ. Equ. Math. Phys. **44** (2008), 1–21.
- [2] R. P. Agarwal, M. Meehan and D. O'Regan, *Fixed Point Theory and Applications*, Cambridge University Press, Cambridge, UK, 2001.
- [3] Y. Arioua, B. Basti and N. Benhamidouche, *Initial value system for nonlinear implicit fractional differential equations with Katugampola derivative*, Appl. Math. E-Notes **19** (2019), 397–412.
- [4] M. Alesemi, N. Iqbal and A. A. Hamoud, *The analysis of fractional-order proportional delay physical models via a novel transform*, Complexity **2022** (2022), 1–13.
- [5] K. Balachandran and J. J. Trujillo, *The nonlocal Cauchy system for nonlinear fractional integro-differential equations in Banach spaces*, Nonlinear Anal. **72** (2010), 4587–4593.
- [6] T. D. Benavides, *An existence theorem for implicit differential equations in a Banach space*, Ann. Math. Pura Appl. **4** (1978), 119–130.
- [7] M. Benchohra, S. Bouriah and J. R. Graef, *Boundary value problems for nonlinear implicit Hadamard-Caputo-type fractional differential equations with impulses*, Mediterr. J. Math. **14**(206) (2017), 1–21.
- [8] M. Benchohra, J. Henderson and D. Seba, *Boundary value systems for fractional differential inclusions in Banach Space*, Fractional Differential Calculus **2**(1) (2012), 99–108.
- [9] M. Benchohra and J. E. Lazreg, *Nonlinear fractional implicit differential equations*, Commun. Appl. Anal. **17**(3–4) (2013), 471–482.
- [10] A. Boutiara, K. Guerbati and M. Benbachir, *Hadamard-Caputo fractional differential equation with three-point boundary conditions in Banach spaces*, AIMS Mathematics **5**(1) (2019), 259–272.
- [11] P. L. Butzer, A. A. Kilbas and J. J. Trujillo, *Fractional calculus in the Mellin setting and Hadamard-type fractional integrals*, J. Math. Anal. Appl. **269**(1) (2002), 1–27.
- [12] N. Dardar, *Nonlinear implicit Hadamard-Caputo fractional differential equation with fractional boundary conditions*, Jordan J. Math. Stat. **15**(4B) (2022), 999–1014. <https://doi.org/10.47013/15.4.14>
- [13] K. Diethelm, *The Analysis of Fractional Differential Equations*, Springer, Berlin, 2004.
- [14] G. Emmanuele, *Convergence of successive approximations for implicit ordinary differential equations in Banach spaces*, Funkcialaj Ekvacioj **24** (1981), 325–330.
- [15] G. Emmanuele and B. Ricceri, *On the existence of solutions of ordinary differential equations in implicit form in Banach spaces*, Ann. Mat. Pura Appl. **129** (1981), 367–382.
- [16] A. A. Hamoud, *Existence and uniqueness of solutions for fractional neutral Volterra-Fredholm integro differential equations*, Advances in the Theory of Nonlinear Analysis and its Application **4**(4) (2020), 321–331.
- [17] A. A. Hamoud, *Uniqueness and stability results for Caputo fractional Volterra-Fredholm integro-differential equations*, Journal of Siberian Federal University. Mathematics & Physics **14**(3) (2021), 313–325. <https://doi:10.17516/1997-1397-2021-14-3-313-325>
- [18] A. A. Hamoud and K. P. Ghadle, *The approximate solutions of fractional Volterra-Fredholm integro-differential equations by using analytical techniques*, Probl. Anal. Issues Anal. **7**(25)(1) (2018), 41–58. <https://doi:10.15393/j3.art.2018.4350>
- [19] A. A. Hamoud, M. SH. Bani Issa and K. P. Ghadle, *Existence and uniqueness results for nonlinear Volterra-Fredholm integro-differential equations*, Nonlinear Functional Analysis and Applications **23**(4) (2018), 797–805.
- [20] K. Hilal, L. Ibnelazyz, K. Guida and S. Melliani, *Existence of Mild Solutions for an Impulsive Fractional Integro-differential Equations with Non-local Condition*, Springer Nature Switzerland AG, New York, NY, USA, 2019.
- [21] K. Hilal, K. Guida, L. Ibnelazyz and M. Oukessou, *Existence Results for an Impulsive Fractional Integro-differential Equations with Non-compact Semigroup*, Springer Nature Switzerland AG, New York, NY, USA, 2019.

- [22] K. Hussain, A. Hamoud and N. Mohammed, *Some new uniqueness results for fractional integro-differential equations*, *Nonlinear Functional Analysis and Applications* **24**(4) (2019), 827–836.
- [23] V. M. Hokkanen, *Continuous dependence for an implicit nonlinear equation*, *J. Differ. Equ.* **110** (1994), 67–85.
- [24] F. Jarad, D. Baleanu and T. Abdeljawad, *Caputo-type modification of the Hadamard fractional derivatives*, *Adv. Differ. Equ.* **2012**(1) (2012), 1–8.
- [25] K. Ivaz, I. Alasadi and A. Hamoud, *On the Hilfer fractional Volterra-Fredholm integro differential equations*, *IAENG International Journal of Applied Mathematics* **52**(2) (2022), 426–431.
- [26] A. A. Kilbas, *Hadamard-type fractional calculus*, *J. Korean Math. Soc.* **38**(6) (2001), 1191–1204.
- [27] P. Karthikeyan and R. Arul, *Integral boundary value problems for implicit fractional differential equations involving Hadamard and Hadamard-Caputo fractional derivatives*, *Kragujevac J. Math.* **45**(3) (2021), 331–341. <https://doi.org/10.46793/KgJMat2103.331K>
- [28] K. D. Kucche, J. J. Nieto and V. Venkatesh, *Theory of nonlinear implicit fractional differential equations*, *Differ. Equ. Dyn. Syst.* **28**(1) (2020), 1–17. <https://doi.org/10.1007/s12591-016-0297-7>
- [29] V. Lakshmikantham and M. Rao, *Theory of Integro-Differential Equations*, Gordon & Breach, London, 1995.
- [30] J. J. Nieto, A. Ouahab and V. Venkatesh, *Implicit fractional differential equations via the Liouville-Caputo derivative*, *Mathematics* **3**(2) (2015), 398–411.
- [31] I. Podlubny, *Fractional Differential Equations*, Academic press, New York, USA, 1999.
- [32] S. Unhalely and S. Kendre, *On existence and uniqueness results for iterative fractional integro-differential equation with deviating arguments*, *Appl. Math. E-Notes* **19** (2019), 1–16.
- [33] D. Vivek, E. M. Elsayed and K. Kanagarajan, *Theory of fractional implicit differential equations with complex order*, *Journal of Universal Mathematics* **2**(2) (2019), 154–165.
- [34] D. Vivek, K. Kanagarajan and E. M. Elsayed, *Some existence and stability results for Hilfer-fractional implicit differential equations with nonlocal conditions*, *Mediterr. J. Math.* **15**(1) (2018), 1–14. <https://doi.org/10.1007/s00009-017-1061-0>
- [35] Y. Zhou, *Basic Theory of Fractional Differential Equations*, World Scientific, Singapore, 2014.

¹DEPARTMENT OF MATHEMATICS,
 TAIZ UNIVERSITY,
 TAIZ P.O. BOX 6803, YEMEN.
Email address: ahmed.hamoud@taiz.edu.ye

²DEPARTMENT OF MATHEMATICS AND INFORMATICS,
 UNIVERSITY OF SOUK AHRAS,
 P.O. BOX 1553, SOUK AHRAS, 41000, ALGERIA.
Email address: c.kechar@univ-soukahras.dz

Email address: abd-ardjouni@yahoo.fr

³DEPARTMENT OF MATHEMATICS,
 SAVEETHA SCHOOL OF ENGINEERING,
 SAVEETHA INSTITUTE OF MEDICAL AND TECHNICAL SCIENCES,
 SAVEETHA UNIVERSITY,
 CHENNAI-602 105, TAMIL NADU, INDIA.

⁴MEU RESEARCH UNIT,
 MIDDLE EAST UNIVERSITY,
 AMMAN 11831, JORDAN.

⁵DEPARTMENT OF MATHEMATICS,
HAMEDAN BRANCH, ISLAMIC AZAD UNIVERSITY,
HAMEDAN, IRAN.
Email address: homan_emadi@yahoo.com

⁶DEPARTMENT OF MATHEMATICS,
DAYALBAGH EDUCATIONAL INSTITUTE,
INDIA.
Email address: atul1230123@gmail.com

⁷ARTIFICIAL INTELLIGENCE AND SENSING TECHNOLOGIES (AIST) RESEARCH CENTER,
UNIVERSITY OF TABUK,
TABUK 71491, SAUDI ARABIA

⁸COMPUTER SCIENCE DEPARTMENT,
AL AL-BAYT UNIVERSITY,
MAFRAQ 25113, JORDAN.

⁹HOURLANI CENTER FOR APPLIED SCIENTIFIC RESEARCH,
AL-AHLYYYA AMMAN UNIVERSITY,
AMMAN 19328, JORDAN.

¹⁰DEPARTMENT OF ELECTRICAL AND COMPUTER ENGINEERING,
LEBANESE AMERICAN UNIVERSITY,
BYBLOS 13-5053, LEBANON.

¹¹SCHOOL OF ENGINEERING AND TECHNOLOGY,
SUNWAY UNIVERSITY MALAYSIA,
PETALING JAYA 27500, MALAYSIA.

¹²APPLIED SCIENCE RESEARCH CENTER,
APPLIED SCIENCE PRIVATE UNIVERSITY,
AMMAN 11931, JORDAN.
Email address: aligah.2020@gmail.com