

## An Alternative Moving Frame for Tubular Surface around the Spacelike Curve with a Spacelike Binormal in Minkowski 3-Space

MURAT KEMAL KARACAN AND BAHADDIN BUKCU

ABSTRACT. We describe a robust method for constructing a tubular surface surrounding a spacelike curve with a spacelike binormal in Minkowski 3-Space. Our method is designed to eliminate undesirable twists and wrinkles in the tubular surface's skin at points where the curve experiences high torsion. In our construction the tubular surface's twist is bounded by the spacelike curve's curvature and is independent of the spacelike curve's torsion.

### 1. PRELIMINARIES

Let  $IR^3 = \{(x_1, x_2, x_3) \mid x_1, x_2, x_3 \in IR\}$  be a 3-dimensional vector space, and let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$  be two vectors in  $IR^3$ . The Lorentz scalar product of  $x$  and  $y$  is defined by

$$\langle x, y \rangle_L = x_1y_1 - x_2y_2 + x_3y_3,$$

$IE_1^3 = (IR^3, \langle x, y \rangle_L)$  is called 3-dimensional Lorentzian space, Minkowski 3-Space or 3-dimensional Semi-Euclidean space. The vector  $x$  in  $IE_1^3$  is called a spacelike vector, null vector or a timelike vector if  $\langle x, x \rangle_L > 0$  or  $x = 0$ ,  $\langle x, x \rangle_L = 0$  or  $\langle x, x \rangle_L < 0$ , respectively. For  $x \in IE_1^3$ , the norm of the vector  $x$  defined by  $\|x\|_L = \sqrt{|\langle x, x \rangle_L|}$ , and  $x$  is called a unit vector if  $\|x\|_L = 1$ . For any  $x, y \in IE_1^3$ , Lorentzian vectorial product of  $x$  and  $y$  is defined by

$$x \wedge_L y = (x_2y_3 - x_3y_2, x_1y_3 - x_3y_1, x_1y_2 - x_2y_1).$$

Similarly, an arbitrary curve  $\alpha = \alpha(s)$  in  $IE_1^3$  is locally spacelike, timelike or null (lightlike), if all of its velocity vectors  $\alpha'(s) = T(s)$  are respectively spacelike, timelike or null, for each  $s \in I \subset IR$ . The vectors  $v, w \in IE_1^3$  are orthogonal if and only if  $\langle v, w \rangle_L = 0$ . The Lorentzian sphere of center  $m = (m_1, m_2, m_3)$  and radius  $r \in R^+$  in the space  $IE_1^3$  is defined by [3]

$$S_1^2 = \{a = (a_1, a_2, a_3) \in IE_1^3 \mid \langle a - m, a - m \rangle_L = r^2\}.$$

---

2000 *Mathematics Subject Classification*. Primary: 53A04, 53A05, 53A35.

*Key words and phrases*. Spacelike Curve, Canal Surface, Tubular Surface, Minkowski 3-Space.

## 2. INTRODUCTION

Consider a spacelike curve  $\alpha(s)$  in  $IE_1^3$ , parameterized by its arc length  $s$ . Let  $T(s)$  be its tangent vector, i.e.,  $T(s) = \alpha'(s) = \frac{d\alpha(s)}{ds}$ . The arc length parameterization of the curve makes  $T(s)$  a unit vector, i.e.,  $\|T(s)\|_L = 1$ , therefore its derivative is orthogonal to  $T$ . The principal normal vector  $N$  is defined as  $N = \frac{T'}{\|T'\|_L}$ . The binormal vector  $B$  is defined as the cross product  $B = T \wedge_L N$ . The Frenet-Serret equations, express the rate of change of the moving orthonormal triad  $\{T, N, B\}$  along the spacelike curve with a spacelike binormal  $\alpha(s)$ ,

$$(2.1) \quad \begin{cases} T' = \kappa N \\ N' = \kappa T + \tau B \\ B' = \tau N \end{cases}$$

and

$$\begin{aligned} \langle T, T \rangle_L &= \langle B, B \rangle_L = 1, \langle N, N \rangle_L = -1 \\ \langle T, N \rangle_L &= \langle T, B \rangle_L = \langle B, N \rangle_L = 0. \end{aligned}$$

The coefficients  $\kappa$  and  $\tau$  are the curve's curvature and torsion.

A. Gray [1] has noted that wild gyrations of the Frenet-Serret system can be expected at points where the curvature  $\kappa$  is small and the torsion  $\tau$  is large. An inspection of the Frenet-Serret equations (2.1) shows that at such points  $T'$  is small and  $N'$  and  $B'$  are large, in effect indicating that the Frenet-Serret system is spinning about its  $T$  axis.

The curve's torsion is a function of the curve's third derivative as evidenced by the explicit formulas

$$\kappa = \frac{\|x' \wedge_L x''\|_L}{\|x'\|_L^3}, \quad \tau = \frac{\det(x', x'', x''')}{\|x' \wedge_L x''\|_L^2}.$$

## 3. CHARACTERIZATION OF CANAL AND TUBULAR SURFACES

**Theorem 3.1.** *Suppose the center curve of a canal surface is a unit speed spacelike curve with a spacelike binormal*

$$\alpha : I \rightarrow IE_1^3$$

*with nonzero curvature. Then the canal surface can be parameterized*

$$(3.1) \quad \begin{aligned} C(s, t) &= \alpha(s) + r(s)r'(s)T(s) \mp r(s)\sqrt{1-r'^2(s)}N(s) \sinh t \pm \\ &\quad \pm r(s)\sqrt{1-r'^2(s)}B(s) \cosh t, \end{aligned}$$

*where  $T, N$  and  $B$  denote the tangent, principal normal and binormal of  $\alpha$ .*

*Proof.* Let  $C$  denote a patch that parametrizes the envelope of Lorentzian spheres defining the canal surface. Since the curvature of  $\alpha$  is nonzero, the Frenet-Serret frame  $\{T, N, B\}$  is well-defined, and we can write

$$(3.2) \quad C(s, t) - \alpha(s) = a(s, t)T + b(s, t)N + c(s, t)B$$

where  $a$ ,  $b$  and  $c$  are differentiable on the interval on which  $\alpha$  is defined. We must have

$$(3.3) \quad \|C(s, t) - \alpha(s)\|_L^2 = r^2(s)$$

Equation (3.3) expresses analytically the geometric fact that  $C(s, t)$  lies on a Lorentzian sphere  $S_1^2(s)$  of radius  $r(s)$  centered at  $\alpha(s)$ . Furthermore,  $C(s, t) - \alpha(s)$  is a normal vector to the canal surface; this fact implies that

$$(3.4) \quad \langle C(s, t) - \alpha(s), C_s \rangle_L = 0,$$

$$(3.5) \quad \langle C(s, t) - \alpha(s), C_t \rangle_L = 0,$$

Equations (3.3) and (3.4) say that the vectors  $C_s$  and  $C_t$  are tangent to  $S_1^2(s)$ . From (3.2) and (3.3) we get

$$(3.6) \quad \begin{cases} a^2 - b^2 + c^2 = r^2 \\ aa_s - bb_s + cc_s = rr' \end{cases}$$

When we differentiate (3.2) with respect to  $t$  and use the Frenet-Serret formulas, we obtain

$$(3.7) \quad C_s = (1 + a_s + b\kappa)T + (a\kappa + c\tau + b_s)N + (c_s + b\tau)B.$$

Then (3.6), (3.7), (3.2) and (3.4) imply that

$$(3.8) \quad a + rr' = 0,$$

and from (3.6) and (3.7) we get

$$(3.9) \quad -b^2 + c^2 = r^2(1 - r'^2).$$

We can write

$$\begin{aligned} b &= \mp r \sqrt{1 - r'^2} \sinh t, \\ c &= \pm r \sqrt{1 - r'^2} \cosh t. \end{aligned}$$

Thus (3.2) becomes

$$\begin{aligned} C(s, t) &= \alpha(s) + r(s)r'(s)T(s) \mp r(s)\sqrt{1 - r'^2(s)}N(s) \sinh t \pm \\ &\quad \pm r(s)\sqrt{1 - r'^2(s)}B(s) \cosh t. \end{aligned}$$

With the Frenet-Serret system in hand, we can construct a “tubular surface” of radius  $r = \text{const.}$  about the spacelike curve with a spacelike binormal by defining a surface with parameters  $s$  and  $t$ :

$$(3.10) \quad \text{Tube}(s, t) = \alpha(s) + r(N(s) \sinh t + B(s) \cosh t). \quad \square$$

#### 4. AN ALTERNATIVE MOVING FRAME

We introduce an alternative, more tamely behaved moving triad for spacelike curves with a spacelike binormal. For this, we let  $\alpha(s)$  be a regular spacelike curve in  $IE_1^3$  parameterized by its arc length, and  $T(s) = \alpha'(s)$  be its unit tangent vector the same as before. We will define unit vector fields  $P(s)$  and  $Q(s)$  such that  $\{T, P, Q\}$  is orthonormal at each point along the spacelike curve. We will chose  $P(s)$  and  $Q(s)$  in such a way as to minimize its gyrations as the triad moves along the spacelike curve.

Unlike the Frenet-Serret triad which is defined locally on the spacelike curve, our triad is denned in terms of the solution of a differential equation, hence it depends not only on the spacelike curve's local properties but also on the location and value of the differential equation's initial conditions.

The construction of the new triad is based on the set of solutions of the system of differential equations

$$(4.1) \quad \frac{d}{ds} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 & -\tau(s) \\ -\tau(s) & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$$

where  $a$  and  $b$  are functions of  $s$  and  $\tau(s)$  is the torsion of the spacelike curve  $\alpha(s)$ . Let denote the semi group of solutions of this differential equation, that is,  $(a(s), b(s)) = \phi_{s_0, s}(a(s_0), b(s_0))$  is the solution of the differential equation corresponding to the initial data  $(a(s_0), b(s_0))$  at  $s = s_0$ . If the spacelike curve  $\alpha(s)$  is sufficiently regular, then the standard theory of linear systems, guarantees the existence, uniqueness and maximal continuation of its solutions.

**Definition 4.1.** Let  $U(s)$  denote the normal plane of the curve  $\alpha(s)$  at any  $s$ . That is,  $U(s)$  is perpendicular to  $T(s)$  and contains the principal normal and binormal vectors  $N(s)$  and  $B(s)$ . We equip each  $U(s)$  with the inner product inherited from  $IE_1^3$ . We define the mapping  $\phi_{s_0, s} : U(s_0) \rightarrow U(s)$  with

$$\phi_{s_0, s} : V_0 \in U(s_0) \rightarrow a(s)N(s) + b(s)B(s)$$

where  $(a(s), b(s)) = \phi_{s_0, s}(\langle V_0, N_0(s_0) \rangle_L, \langle V_0, B_0(s_0) \rangle_L)$ .

**Lemma 4.1.** Let  $\phi_{s_0, s}$  be as in Definition 4.1. Then for any  $V_0 \in U(s_0)$  the vector valued function  $V(s)$  defined by  $V(s) = \phi_{s_0, s}(V_0)$  satisfies the differential equation  $V' = -\langle V, T' \rangle_L T$ , where  $T$  is the spacelike curve's unit tangent vector.

*Proof.* According to the definition of  $\phi_{s_0, s}$ , the components  $\langle V(s), N(s) \rangle_L$  and  $\langle V(s), B(s) \rangle_L$  of  $V(s)$  satisfy the differential equation (4.1), therefore

$$\langle V, N \rangle'_L = -\tau \langle V, B \rangle_L$$

and

$$\langle V, B \rangle'_L = -\tau \langle V, N \rangle_L.$$

We now compute  $V'(s)$  while making use of the Frenet-Serret equations:

$$\begin{aligned}
 V'(s) &= [\langle V, N \rangle_L N + \langle V, B \rangle_L B]' \\
 &= \langle V, N \rangle_L' N + \langle V, N \rangle_L N' + \langle V, B \rangle_L' B + \langle V, B \rangle_L B' \\
 &= -\tau \langle V, B \rangle_L N + \langle V, N \rangle_L (\kappa T + \tau B) - \tau \langle V, N \rangle_L B + \tau \langle V, B \rangle_L N \\
 &= \kappa \langle V, N \rangle_L T \\
 &= \langle V, \kappa N \rangle_L T \\
 &= \langle V, T' \rangle_L T
 \end{aligned} \quad \square$$

as asserted.

**Theorem 4.2.** *The mapping  $\phi_{s_0, s} : U(s_0) \rightarrow U(s)$  is an isometry.*

*Proof.* Take any two vectors  $P_0$  and  $Q_0$  in  $U(s_0)$  and let  $P(s) = \phi_{s_0, s}(P_0)$  and  $Q(s) = \phi_{s_0, s}(Q_0)$ . Then applying the result of Lemma 4.1 we get

$$\begin{aligned}
 \langle P(s), Q(s) \rangle_L' &= \langle P', Q \rangle_L + \langle P, Q' \rangle_L \\
 &= \langle \langle P, T' \rangle_L T, Q \rangle_L + \langle \langle Q, T' \rangle_L T, P \rangle_L \\
 &= 0
 \end{aligned}$$

because  $\langle T, P \rangle_L = 0$  and  $\langle T, Q \rangle_L = 0$ . Therefore  $\phi_{s_0, s}$  preserves the inner product hence is an isometry.  $\square$

**Definition 4.2.** Let  $\alpha(s)$  be a regular spacelike curve in  $IE_1^3$  parameterized by its arc length, and let  $T(s)$  denote its unit tangent. Arbitrarily fix a parameter value  $s_0$  and let  $T_0 = T(s_0)$ . Choose any pair of vectors  $P_0$  and  $Q_0$  such that  $\{T_0, P_0, Q_0\}$  forms an orthonormal set. Then let  $P(s) = \phi_{s_0, s}(P_0)$  and  $Q(s) = \phi_{s_0, s}(Q_0)$ . According to Theorem 4.2, the triad  $\{T(s), P(s), Q(s)\}$  is orthonormal for all  $s$ . We call  $\{T(s), P(s), Q(s)\}$  a tubular surface triad for the spacelike curve.

**Remark 4.1.** If the initial triad  $\{T_0, P_0, Q_0\}$  is chosen such that it is positively oriented, i.e.,  $\det \{T_0, P_0, Q_0\} = 1$ , then the tubular surface triad will be positively oriented for all  $s$ . This is an immediate consequence of the continuity of solutions of the differential equations (4.1).

**Theorem 4.3.** *Let  $\{T, P, Q\}$  be the tubular surface triad as in Definition 4.2 and let  $\kappa$  be the curvature of the spacelike curve. We have the following pointwise bounds on the rate of change of the triad [3]:*

$$(4.2) \quad \begin{aligned}
 \|T'(s)\|_L &= \kappa(s), \\
 \|P'(s)\|_L &\leq \kappa(s), \\
 \|Q'(s)\|_L &\leq \kappa(s).
 \end{aligned}$$

**Remark 4.2.** Rather than computing both  $P$  and  $Q$  components of the tubular surface triad with  $P(s) = \phi_{s_0, s}(P_0)$  and  $Q(s) = \phi_{s_0, s}(Q_0)$  as in Definition 4.2, it is more practical to compute one, say  $P(s)$ , then define  $Q(s)$  as  $Q(s) = T(s) \wedge_L P(s)$ ,

which also implies the permutations  $-P = Q \wedge_L T$  and  $T = P \wedge_L Q$ . We can verify directly that  $Q$ , defined this way, satisfies the differential equation (4.1):

$$\begin{aligned}
 Q' &= (T \wedge_L P)' \\
 Q' &= T' \wedge_L P + T \wedge_L P' \\
 &= (k_1 P - k_2 Q) \wedge_L P + T \wedge_L (\langle P, T' \rangle_L T) \\
 &= k_1 (P \wedge_L P) - k_2 (Q \wedge_L P) + (T \wedge_L T) \langle P, T' \rangle_L \\
 &= -k_2 (Q \wedge_L P) \\
 &= -k_2 (-T) \\
 &= k_2 T \\
 &= \langle Q, T' \rangle_L T
 \end{aligned}$$

therefore  $Q(s) = \phi_{s_0, s}(Q_0)$ , as required. In this derivation we have made use of the fact that. We have also used the general vector algebra identities  $a \wedge_L a = 0$  and  $a \wedge_L (b \wedge_L c) = -\langle a, c \rangle_L b + \langle a, b \rangle_L c$ .

**Theorem 4.4.** *Let  $\{T(s), P(s), Q(s)\}$  be a positively oriented tubular surface triad. Then its rate of change is expressed by:*

$$(4.3) \quad \begin{cases} T' = k_1 P - k_2 Q \\ P' = k_1 T \\ Q' = k_2 T \end{cases}$$

where  $k_1(s)$  and  $k_2(s)$  are scalar functions defined along the spacelike curve with a spacelike binormal.

*Proof.* The second and third equations in (4.2) are consequences of Lemma 4.1. To obtain the first equation we note that the positive orientation of the tubular surface triad implies  $T = P \wedge_L Q$  and  $-P = Q \wedge_L T$  and  $Q = T \wedge_L P$ . Therefore we have

$$\begin{aligned}
 T' &= P \wedge_L Q \\
 T' &= P' \wedge_L Q + P \wedge_L Q' \\
 T' &= \langle P, T' \rangle_L T \wedge_L Q + P \wedge_L \langle Q, T' \rangle_L T \\
 T' &= \langle P, T' \rangle_L (T \wedge_L Q) + (P \wedge_L T) \langle Q, T' \rangle_L \\
 T' &= \langle P, T' \rangle_L (P) - \langle Q, T' \rangle_L (-Q) \\
 T' &= \langle P, T' \rangle_L P - \langle Q, T' \rangle_L Q \\
 T' &= k_1 P - k_2 Q
 \end{aligned}$$

as asserted. □

**Example 4.1.** Let  $c^2 = a^2 + b^2 > 0$ . For illustration, we compute the  $P$  and  $Q$  vectors of the tubular surface triad for the unit speed spacelike helix:

$$(4.4) \quad \alpha(s) = \left( a \sinh \frac{s}{c}, a \cosh \frac{s}{c}, \frac{bs}{c} \right)$$

with  $a \neq 0$ . For the corresponding Frenet-Serret triad is:

$$\begin{aligned} T(s) &= \left( \frac{a}{c} \cosh \frac{s}{c}, \frac{a}{c} \sinh \frac{s}{c}, \frac{b}{c} \right) \\ N(s) &= \left( \sinh \frac{s}{c}, \cosh \frac{s}{c}, 0 \right) \\ B(s) &= \left( -\frac{b}{c} \cosh \frac{s}{c}, -\frac{b}{c} \sinh \frac{s}{c}, \frac{a}{c} \right). \end{aligned}$$

The curvature and torsion are constants, independent of  $s$ , and are given by

$$\begin{aligned} \kappa &= \frac{a}{a^2 + b^2} \\ \tau &= \frac{b}{a^2 + b^2}. \end{aligned}$$

Since torsion is constant, the system (4.1) reduces to a differential equation with constant coefficients which can be solved readily once a set of initial conditions is supplied. Suppose that we take  $P(0) = B(0)$ . Note that  $P(0)$  is of unit length and orthogonal to  $T(0)$ , as required. Then

$$a(0) = \langle P(0), N(0) \rangle_L = 1$$

and

$$b(0) = \langle P(0), B(0) \rangle_L = 0.$$

Solving (4.1) with these initial conditions we obtain  $a(s) = \langle P, N \rangle_L = \cosh \tau s$  and  $b(s) = \langle P, B \rangle_L = \sinh \tau s$ , whence, according to Definition 4.1:

$$P(s) = N(s) \cosh \tau s + B(s) \sinh \tau s,$$

where  $N(s), B(s)$  and  $\tau$  are given in terms of  $s$  by the explicit formulas above. The vector  $Q$  can also be computed in the same manner, or simply through  $Q(s) = T(s) \wedge_L P(s)$ . Since  $T \wedge_L N = B$  and  $T \wedge_L B = N$ , we get:

$$Q(s) = N(s) \sinh \tau s + B(s) \cosh \tau s.$$

## REFERENCES

- [1] A. Gray, "Modern Differential Geometry of Curves and Surfaces with Mathematica", CRC Press, 1998.
- [2] M.K. Karacan, "Kinematic Applications of Two Parameter Motions", Ankara University Graduate School, Ph.D Thesis, 2004.

- [3] M.K. Karacan, B. Bukcu, *An Alternative Moving Frame for Tubular Surfaces around Timelike Curves in the Minkowski 3-space*, Balkan Journal of Geometry and its Applications, Vol. **12**, No. **2** (2007), 73–80.
- [4] M. Petrovic and E. Sucurovic, *Some Characterizations of the Spacelike, the Timelike and the Null Curves on the Pseudohyperbolic Space  $H_0^2$  in  $E_1^3$* , Kragujevac J. Math., Vol. **22** (2000), 71–82.
- [5] R.Rostamian, "The Differential Geometry of Tube Plots in Computer Graphics", Technical Report, January 2000.

MURAT KEMAL KARACAN  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCES AND LETTERS  
K. MARAS SUTCU IMAM UNIVERSITY  
AVSAR CAMPUS  
46100, K.MARAS  
TURKEY

*E-mail address:* [mkkaracan@ksu.edu.tr](mailto:mkkaracan@ksu.edu.tr)

BAHADDIN BUKCU  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCES AND LETTERS  
GAZI OSMAN PASA UNIVERSITY  
TOKAT  
TURKEY

*E-mail address:* [bbukcu@yahoo.com](mailto:bbukcu@yahoo.com)