

ON THE SOLUTIONS OF THE FUNCTIONAL EQUATION
 $x(t) + A(t)x(f(t)) = F(t)$ WHEN THE FUNCTION F
SATISFIES SPECIAL CONDITIONS

M. Malenica

Abstract. The results of this paper are concerned with the solution $x(t)$ of the functional equation $x(t) + A(t)x(f(t)) = F(t)$. Using regular summability methods T , we derive some necessary and also some sufficient conditions for the T -sum $x(t)$ of the series $\sum_{i=0}^{\infty} (-1)^i F(f^i(t))$ to be a solution of the above mentioned equation under the specific conditions for $F(t)$.

1. Introduction

We consider here the linear functional equation

$$x(t) + A(t)x(f(t)) = F(t) \quad (1.1)$$

on a given set $S \subset \mathbf{R}$, where $f: S \rightarrow S$, $F: S \rightarrow \mathbf{R}$ and $A: S \rightarrow \mathbf{R}$ are given functions, $x: S \rightarrow \mathbf{R}$ denotes the unknown function and \mathbf{R} is the set of real numbers. Throughout the paper we assume that the functions A and f satisfy the condition $A(t) = A(f(t))$. We define

$$f^0(t) = t, \quad f^{i+1}(t) = f(f^i(t)), \quad i = 0, 1, 2, \dots$$

Equations of this type have been considered by many mathematicians. H. Steinhaus [10] discussed the equation $\varphi(x) + \varphi(x^2) = x$ and G. H. Hardy [4] considered the equation $\varphi(x) + \varphi(x^\alpha) = x$ ($\alpha > 0$), where φ denotes the unknown function. R. Raclis [9] discusses the equation $\varphi(x) + \varphi(f(x)) = F(x)$ for complex x and finds meromorphic solutions. Under certain hypothesis in regard to the function f , N. M. Gersevanoff [2] solves the equation $x(t) + A(t)x(f(t)) = F(t)$ and Ghermanescu [3] solves the equation

$$A_0\varphi + A_1\varphi(f) + A_2\varphi(f(f)) + \dots + A_n\varphi(f(\dots)) = F(x),$$

where φ denotes the unknown function, and f and F are given functions. Equation (1.1) is a direct generalization of the equation $\varphi(x) + \varphi(f(x)) = F(x)$ considered by M. Kuczma in [6]. The basic method used in that paper is the iteration of the function f , and solutions are given in the form of the sums of a convergent series

of functions. M. Bajraktarević [1] has also considered solutions of the equation $\varphi(x) + \varphi(f(x)) = F(x)$ under assumption that the series of functions given by M. Kuczma is divergent, but summable by some regular method of summability. M. Malenica [7], [8], using method of M. Bajraktarević has examined solutions of the same equation. Papers by M. Kuczma [6], M. Bajraktarević [1] and M. Malenica [8] are of special interest to us.

In this paper, considering the series

$$\sum_{i=0}^{\infty} (-1)^i F(f^i(t)) \quad (1.2)$$

and using regular summability methods T we derive some sufficient and also some necessary conditions for T -sum $x(t)$ of the series (1.2) to be a solution of equation (1.1). The function $F(t)$ has a special form. Whenever we say that (1.2) is T -summable with sum $x(t)$ we assume that $T = (a_{kn})$ is a regular sequence matrix transformation, i.e. its elements a_{kn} satisfy the following conditions

- (i) $a_{kn} \rightarrow 0$ ($k \rightarrow \infty, n \geq 0$),
- (ii) $\sum_{i=0}^n |a_{ki}| \leq K$ ($n \geq 0, k \geq 0, K$ fixed),
- (iii) $\sum_{n=0}^{\infty} a_{kn} = A_k \rightarrow 1$ ($k \rightarrow \infty$).

2. Results

It is interesting to consider the equation (1.1) in the case when the form of F causes the series (1.2) to diverge and to look for the conclusions on $T = (a_{kn})$ that assure that the series (1.2) is T -summable to the function $x(t)$ so that the relations

$$Ts_n(t) \rightarrow x(t), \quad x(t) + A(t)x(f(t)) = F(t), \quad (2.1)$$

are valid. The sum $s_n(t)$ is a partial sum of order n of the series (1.2). If we take F to be a function satisfying

$$F(f(t)) = F(t), \quad (2.2)$$

and for $T = (a_{kn})$ regular transformation with the property

$$\sum_{m=0}^{\infty} a_{k,2m} \rightarrow \frac{1}{1+A(t)}, \quad k \rightarrow \infty \quad (A(t) \neq -1),$$

we get the following theorem.

THEOREM 2.1 Suppose $F(t)$ satisfies (2.2) and suppose $Ts_n(t) \rightarrow x(t)$, where T is a regular transformation.

(a) For $x(t)$ to be a solution of the equation (1.1), it is sufficient that T satisfies condition (2.3). In this case the solution has the form $x(t) = \frac{F(t)}{1+A(t)}$.

(b) For $x(t)$ to be a solution of the equation (1.1), it is necessary that T satisfies condition (2.3) with $F(t) \not\equiv 0$.

(c) If $F(t) \equiv 0$, then $x(t) \equiv 0$ is always a solution of the equation (1.1).

Proof. For the function F satisfying (2.2) we have

$$\sum_{i=0}^n (-1)^i F(f^i(t)) = \sum_{i=0}^n (-1)^i F(t) = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ F(t), & \text{if } n \text{ is even;} \end{cases}$$

$$s'_k(t) = \sum_{n=0}^{\infty} a_{kn} s_n(t) = \sum_{m=0}^{\infty} a_{k,2m+1} 0 + \sum_{m=0}^{\infty} a_{k,2m} F(t),$$

i.e.

$$s'_k(t) = F(t) \sum_{m=0}^{\infty} a_{k,2m}. \quad (2.4)$$

(a) Let $T = (a_{kn})$ be the transformation satisfying (2.3). If we let $k \rightarrow \infty$ in (2.4) we get

$$x(t) = \lim_{k \rightarrow \infty} s'_k(t) = \frac{F(t)}{1 + A(t)}. \quad (2.5)$$

It is easy to verify that if $x(t)$ has the form given in (2.5), then it is a solution of the equation (1.1).

(b) Let $x(t)$ be a solution of the equation (1.1). Then by (2.4), the following relations hold.

$$\begin{aligned} G_k(t) &= s'_k(t) + A(t)s'_k(f(t)) - F(t) \\ &= F(t) \sum_{m=0}^{\infty} a_{k,2m} + A(t)F(t) \sum_{m=0}^{\infty} a_{k,2m} - F(t) \rightarrow 0 \quad (k \rightarrow \infty), \\ F(t)(1 + A(t)) \sum_{m=0}^{\infty} a_{k,2m} - F(t) &= G_k(t) \rightarrow 0 \quad (k \rightarrow \infty), \\ F(t)(1 + F(t)) \sum_{m=0}^{\infty} a_{k,2m} &= F(t) + G_k(t). \end{aligned}$$

If $F(t) \not\equiv 0$, when $k \rightarrow \infty$, we get $\lim_{k \rightarrow \infty} \sum_{m=0}^{\infty} a_{k,2m} = \frac{1}{1 + A(t)}$, i.e. $T = (a_{kn})$ is a regular transformation with the property (2.3).

(c) If $F(t) \not\equiv 0$, then $x(t) \equiv 0$ is always a solution of the equation (1.1). ■

It is easy to verify that C_1 -method, E -method and E_r -method, $r \geq 1$, given in [5] are examples of methods satisfying property (2.3).

REMARK 2.1. Obviously, if $A(t) = -1$, the equation (1.1) has the solution $x(t) = \frac{1}{2}F(t)$, under the condition $F(f(t)) = -F(t)$.

At the end, let $F(t)$ has the following properties

$$F(f^p(t)) = 0, \quad F(f^i(t)) \neq 0, \quad i = 1, 2, \dots, p-1. \quad (2.6)$$

We look for the solution $x(t)$ of the equation (1.1) in the form

$$x(t) = a + \sum_{i=0}^{p-1} b_i F(f^i(t)). \quad (2.7)$$

In connection with that, the following theorem is valid.

THEOREM 2.2. *In order that the equation (1.1), in which the function $F(t)$ has properties (2.6), has a solution $x(t)$ of the form (2.7), it is sufficient that*

$$a = 0, \quad b_i = (-1)^i (A(t))^i, \quad i = 0, 1, \dots, p-1, \quad (2.8)$$

i.e. that the function $x(t)$ has the form

$$x(t) = \sum_{i=0}^{p-1} (-1)^i (A(t))^i F(f^i(t)). \quad (2.9)$$

Proof. From (2.9) and (2.6) we have

$$\begin{aligned} x(t) + A(t)x(f(t)) &= \sum_{i=0}^{p-1} (-1)^i (A(t))^i F(f^i(t)) + A(t) \sum_{i=0}^{p-1} (-1)^i (A(f(t)))^i F(f^{i+1}(t)) \\ &= F(t) + \sum_{i=1}^{p-1} (-1)^i (A(t))^i F(f^i(t)) \\ &\quad - \sum_{i=1}^{p-1} (-1)^i (A(t))^i F(f^i(t)) - (-1)^p (A(t))^p F(f^p(t)) = F(t) \quad \blacksquare \end{aligned}$$

REFERENCES

- [1] M. Bajraktarević, *Sur une solution de l'équation fonctionelle $\varphi(x) + \varphi(f(x)) = F(x)$* , Glasnik mat. fiz. i astr., Zagreb **15** (1960), 91–98.
- [2] N. Gercevanoff, *Quelques procédés de la résolution des équations fonctionnelles linéaire par la méthode d'itération*, Compte Rendus (Doklady) de l'Academie de Sciences de l'URSS **29** (1953), 207–209.
- [3] M. Ghermanescu, *Équations fonctionnelles linéaires à argument fonctionnel n-periodique*, Compte Rendue de l'Acad. Sci. Paris **243** (1956), 1593–1596.
- [4] G. H. Hardy, *Divergent Series*, Oxford 1949.
- [5] K. Knopp, *Theorie und Anwendungen der unendlichen Reihen*, Berlin 1931.
- [6] M. Kuczma, *On the functional equation $\varphi(x) + \varphi(f(x)) = F(x)$* , Ann. Pol. Math. **6** (1959).
- [7] M. Malenica, *O jednom rješenju jednačine $\varphi(x) + \varphi(f(x)) = F(x)$ kada funkcija zadovoljava uslov $F(f(x)) = F(x)$* , Akademija nauka i umjetnosti BiH, Odjeljenje prirodnih i matematičkih nauka, **LXIX/20**, Sarajevo, 1982, 17–21.
- [8] M. Malenica, *On some solutions of equation $\varphi(x) + \varphi(f(x)) = F(x)$ under the condition that F satisfies $F(f^p(x)) = F(x)$* , Publ. Inst. Math. (Beograd) **29(43)** (1981).
- [9] F. Raclis, *Sur la solution méromorphe d'une équation fonctionnelle*, Bull. Math. de la Soc. Roum. de Sciences **30** (1927), 101–105.
- [10] H. Steinhaus, *O pewnym szeregu potegowym*, Prace Matematyczne I (1955), 276–284.

(received 03.04.1992)

M. Malenica, Sarajevo, YUGOSLAVIA