

## THE COMPATIBILITY OF THE TANGENCY RELATIONS OF SETS IN GENERALIZED METRIC SPACES

Tadeusz Konik

**Abstract.** In this paper the problem of the compatibility of the tangency relations  $T_{l_i}(a_i, b_i, k, p)$  ( $i = 1, 2$ ) of sets of the classes  $\tilde{M}_{p,k}$  and  $A_{p,k}^*$  in a generalized metric space is considered. Some sufficient conditions for the compatibility of these relations of sets of the above classes are given here.

### Introduction

Let  $E$  be an arbitrary non-empty set and let  $l$  be a non-negative real function defined on the Cartesian product  $E_0 \times E_0$  of the family  $E_0$  of all non-empty subsets of the set  $E$ . Let  $l_0$  be the function defined by the formula

$$l_0(x, y) = l(\{x\}, \{y\}) \quad \text{for } x, y \in E. \quad (1)$$

If we put some conditions on the function  $l$ , then the function  $l_0$  defined by (1) will be a metric on the set  $E$ . By this reason the pair  $(E, l)$  can be treated as a certain generalization of a metric space and we shall call it (see [9]) a generalized metric space. Using (1) we may define in the space  $(E, l)$ , similarly as in a metric space, the following notions: the sphere  $S_l(p, r)$  and the ball  $K_l(p, r)$  with the centre at the point  $p$  and the radius  $r$ .

Let  $S_l(p, r)_u$  denote the so-called  $u$ -neighbourhood of the sphere  $S_l(p, r)$  in the space  $(E, l)$  defined by the formula

$$S_l(p, r)_u = \begin{cases} \bigcup_{q \in S_l(p, r)} K_l(q, u), & \text{for } u > 0, \\ S_l(p, r), & \text{for } u = 0. \end{cases} \quad (2)$$

Let  $a, b$  be arbitrary non-negative real functions defined in a certain right-hand side neighbourhood of 0 such that

$$a(r) \xrightarrow[r \rightarrow 0+]{} 0 \quad \text{and} \quad b(r) \xrightarrow[r \rightarrow 0+]{} 0. \quad (3)$$

We say that the pair  $(A, B)$  of sets  $A, B$  of the family  $E_0$  is  $(a, b)$ -clustered at the point  $p$  of the space  $(E, l)$ , if 0 is the cluster point of the set of all real numbers  $r > 0$  such that the sets of the form  $A \cap S_l(p, r)_{a(r)}$  and  $B \cap S_l(p, r)_{b(r)}$  are non-empty.

---

*AMS Subject Classification:* 53 A 99

*Keywords and phrases:* generalized metric space, compatibility of the tangency of sets

Let (see [9])

$T_l(a, b, k, p) = \{ (A, B) : A, B \in E_0, \text{ the pair } (A, B) \text{ is } (a, b)\text{-clustered at the}$

$$\text{point } p \text{ of the space } (E, l) \text{ and } \frac{1}{r^k} l(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)}) \xrightarrow[r \rightarrow 0+]{} 0 \}. \quad (4)$$

If  $(A, B) \in T_l(a, b, k, p)$  then we say that the set  $A \in E_0$  is  $(a, b)$ -tangent of order  $k$  ( $k > 0$ ) to the set  $B \in E_0$  at the point  $p$  of the space  $(E, l)$ .

We shall call  $T_l(a, b, k, p)$  defined by (4) the relation of  $(a, b)$ -tangency of order  $k$  at the point  $p$ , or shortly: the tangency relation of sets in the generalized metric space  $(E, l)$ .

Two relations of the tangency  $T_{l_1}(a_1, b_1, k, p)$  and  $T_{l_2}(a_2, b_2, k, p)$  are called compatible if  $(A, B) \in T_{l_1}(a_1, b_1, k, p)$  if and only if  $(A, B) \in T_{l_2}(a_2, b_2, k, p)$  for  $A, B \in E_0$ .

We say that the set  $A \in E_0$  has the Darboux property at the point  $p$  of the space  $(E, l)$ , which we write:  $A \in D_p(E, l)$  (see [3]), if there exists a number  $\tau > 0$  such that the set  $A \cap S_l(p, r)$  is non-empty for  $r \in (0, \tau)$ .

Let  $\rho$  be a metric on the set  $E$  and let  $A, B$  be arbitrary sets of the family  $E_0$ . Let us denote

$$\rho(A, B) = \inf\{\rho(x, y) : x \in A, y \in B\}, \quad d_\rho A = \sup\{\rho(x, y) : x, y \in A\}. \quad (5)$$

Let  $f$  be a subadditive increasing real function defined in a certain right-hand side neighbourhood of 0 such that  $f(0) = 0$ . By  $\overline{F}_f$  we shall denote the class of all functions  $l$  fulfilling the conditions:

- 1°  $l: E_0 \times E_0 \rightarrow (0, \infty)$ ,
- 2°  $f(\rho(A, B)) \leq l(A, B) \leq f(d_\rho(A \cup B))$  for  $A, B \in E_0$ .

Since

$$f(\rho(x, y)) = f(\rho(\{x\}, \{y\})) \leq l(\{x\}, \{y\}) \leq f(d_\rho(\{x\} \cup \{y\})) = f(\rho(x, y)),$$

then from here and from (1) it follows that

$$l_0(x, y) = l(\{x\}, \{y\}) = f(\rho(x, y)) \quad \text{for } l \in \overline{F}_f \text{ and } x, y \in E. \quad (6)$$

It is easy to prove that the function  $l_0$  defined by (6) is a metric on the set  $E$ .

In the present paper the problem of the compatibility of the tangency relations of sets of the classes  $\tilde{M}_{p,k}$  and  $A_{p,k}^*$  having the Darboux property at the point  $p$  of the space  $(E, l)$ , for the functions  $l$  belonging to the class  $\overline{F}_f$ , is considered.

### 1. On the compatibility of the tangency relations of sets of the classes $\tilde{M}_{p,k}$

By  $A'$  we shall denote the set of all cluster points of the set  $A \in E_0$ . Let  $k$  be any fixed positive real number and let

$$\rho(x, A) = \inf\{\rho(x, y) : y \in A\}. \quad (1.1)$$

Let us put by definition (see [4])

$$\begin{aligned} \tilde{M}_{p,k} = \{ A \in E_0 : p \in A' \text{ and there exists } \mu > 0 \text{ such that for an arbitrary } \varepsilon > 0 \\ \text{there exists } \delta > 0 \text{ such that for every pair of points } (x, y) \in [A, p; \mu, k] \\ \text{if } \rho(x, y) < \delta \text{ and } \frac{\rho(x, A)}{\rho^k(x, p)} < \delta \text{ then } \frac{\rho(x, y)}{\rho^k(x, p)} < \varepsilon \}, \end{aligned} \quad (1.2)$$

where

$$[A, p; \mu, k] = \{ (x, y) : x \in E, y \in A \text{ and } \mu \rho(x, A) < \rho^k(x, p) = \rho^k(y, p) \} \quad (1.3)$$

In the paper [4] the following lemma was proved.

LEMMA 1.1. If

$$\frac{a(r)}{r^{k+1}} \xrightarrow[r \rightarrow 0+]{\longrightarrow} \alpha, \quad (1.4)$$

where  $\alpha < \infty$ , then for an arbitrary set  $A \in \tilde{M}_{p,k} \cap D_p(E, \rho)$

$$\frac{1}{r^k} d_\rho(A \cap S_\rho(p, r)_{a(r)}) \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0. \quad (1.5)$$

From this lemma and from the fact that every function  $l \in \overline{F}_f$  generates on the set  $E$  the metric defined by the formula (6) it follows that

$$\frac{1}{r^k} d_l(A \cap S_l(p, r)_{a(r)}) \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0, \quad (1.6)$$

for  $l \in \overline{F}_f$  and  $A \in \tilde{M}_{p,k} \cap D_p(E, l)$ , when the function  $a$  fulfills the condition (1.4).

THEOREM 1.1. If  $l_i \in \overline{F}_f$  ( $i = 1, 2$ ),

$$\frac{a(r)}{r^{k+1}} \xrightarrow[r \rightarrow 0+]{\longrightarrow} \alpha \quad \text{and} \quad \frac{b(r)}{r^{k+1}} \xrightarrow[r \rightarrow 0+]{\longrightarrow} \beta, \quad (1.7)$$

where  $\alpha, \beta < \infty$ , then for arbitrary sets of the classes  $\tilde{M}_{p,k} \cap D_p(E, l)$  the tangency relations  $T_{l_1}(a, b, k, p)$  and  $T_{l_2}(a, b, k, p)$  are compatible.

*Proof.* Let us assume that  $(A, B) \in T_{l_1}(a, b, k, p)$  for  $A, B \in \tilde{M}_{p,k}$ . Hence, from (2) and from the fact that (see (6))

$$l_1(\{x\}, \{y\}) = l_2(\{x\}, \{y\}) = l_0(x, y) \quad \text{for } x, y \in E, \quad (1.8)$$

it follows that the pair of sets  $(A, B)$  is  $(a, b)$ -clustered at the point  $p$  of the space  $(E, l_1)$  and

$$\frac{1}{r^k} l_1(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)}) \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0. \quad (1.9)$$

From the inequality

$$d_\rho(A \cup B) \leq d_\rho A + d_\rho B + \rho(A, B) \quad \text{for } A, B \in E_0, \quad (1.10)$$

from the properties of the function  $f$  and from the fact that  $l_1, l_2 \in \overline{F}_f$  we get

$$\begin{aligned} & \left| \frac{1}{r^k} l_2(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)}) - \frac{1}{r^k} l_1(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)}) \right| \leqslant \\ & \frac{1}{r^k} f(d_\rho((A \cap S_l(p, r)_{a(r)}) \cup (B \cap S_l(p, r)_{b(r)}))) - \frac{1}{r^k} f(\rho(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)})) \\ & \leqslant \frac{1}{r^k} f(d_\rho(A \cap S_l(p, r)_{a(r)}) + d_\rho(B \cap S_l(p, r)_{b(r)}) + \rho(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)})) - \\ & \quad - \frac{1}{r^k} f(\rho(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)})) \\ & \leqslant \frac{1}{r^k} f(d_\rho(A \cap S_l(p, r)_{a(r)})) + \frac{1}{r^k} f(d_\rho(B \cap S_l(p, r)_{b(r)})). \end{aligned} \quad (1.11)$$

From the fact that  $f$  is an increasing function we obtain

$$\begin{aligned} f(d_\rho(A \cap S_l(p, r)_{a(r)})) &= f(\sup\{\rho(x, y) : x, y \in (A \cap S_l(p, r)_{a(r)})\}) \\ &= \sup\{f(\rho(x, y)) : x, y \in (A \cap S_l(p, r)_{a(r)})\} \\ &= \sup\{l_0(x, y) : x, y \in (A \cap S_l(p, r)_{a(r)})\} = d_l(A \cap S_l(p, r)_{a(r)}). \end{aligned} \quad (1.12)$$

Hence and from (1.6) it follows that

$$\frac{1}{r^k} f(d_\rho(A \cap S_l(p, r)_{a(r)})) \xrightarrow[r \rightarrow 0+]{ } 0. \quad (1.13)$$

Analogously

$$\frac{1}{r^k} f(d_\rho(B \cap S_l(p, r)_{b(r)})) \xrightarrow[r \rightarrow 0+]{ } 0. \quad (1.14)$$

From (1.9), (1.13), (1.14) and from the inequality (1.11) we get

$$\frac{1}{r^k} l_2(A \cap S_l(p, r)_{a(r)}, B \cap S_l(p, r)_{b(r)}) \xrightarrow[r \rightarrow 0+]{ } 0. \quad (1.15)$$

Since the functions  $l_1, l_2 \in \overline{F}_f$  generate on the set  $E$  the same metric  $l_0$  (see (6)), from the fact that the pair of sets  $(A, B)$  is  $(a, b)$ -clustered at the point  $p$  of the space  $(E, l_1)$ , it follows that it is  $(a, b)$ -clustered at the point  $p$  of the space  $(E, l_2)$ . Hence and from (1.15) it results that  $(A, B) \in T_{l_2}(a, b, k, p)$ .

If  $(A, B) \in T_{l_2}(a, b, k, p)$ , then similarly we prove that  $(A, B) \in T_{l_1}(a, b, k, p)$ . Hence it follows that the tangency relations  $T_{l_1}(a, b, k, p)$  and  $T_{l_2}(a, b, k, p)$  are compatible in the classes of sets  $\tilde{M}_{p,k} \cap D_p(E, l)$ . ■

Let  $a_i, b_i$  ( $i = 1, 2$ ) be non-negative real functions defined in a certain right-hand side neighbourhood of 0 and fulfilling the condition

$$a_i(r) \xrightarrow[r \rightarrow 0+]{ } 0 \quad \text{and} \quad b_i(r) \xrightarrow[r \rightarrow 0+]{ } 0. \quad (1.16)$$

In the paper [7] the following theorem was proved.

**THEOREM 1.2.** *If  $l \in \overline{F}_f$  and*

$$\frac{a_i(r)}{r^{k+1}} \xrightarrow[r \rightarrow 0+]{ } \alpha_i, \quad \frac{b_i(r)}{r^{k+1}} \xrightarrow[r \rightarrow 0+]{ } \beta_i, \quad (1.17)$$

where  $\alpha_i, \beta_i < \infty$  for  $i = 1, 2$ , then for arbitrary sets of the classes  $\tilde{M}_{p,k} \cap D_p(E, l)$  the tangency relations  $T_l(a_1, b_1, k, p)$  and  $T_l(a_2, b_2, k, p)$  are compatible.

From the Theorems 1.1 and 1.2 it follows

**COROLLARY 1.1.** *If  $l_i \in \overline{F}_f$  and the functions  $a_i, b_i$  ( $i = 1, 2$ ) fulfil the condition (1.17), then the tangency relations  $T_{l_1}(a_1, b_1, k, p)$  and  $T_{l_2}(a_2, b_2, k, p)$  are compatible in the classes of sets  $\tilde{M}_{p,k} \cap D_p(E, l)$ .*

## 2. On the compatibility of the tangency relations of sets of the classes $A_{p,k}^*$

Let  $(E, \rho)$  be a metric space. Let us put by definition (see [3])

$$A_{p,k}^* = \{ A \in E_0 : p \in A' \text{ and there exists a number } \lambda > 0 \text{ such that}$$

$$\limsup_{[A,p;k] \ni (x,y) \rightarrow (p,p)} \frac{\rho(x, y) - \lambda \rho(x, A)}{\rho^k(x, p)} \leq 0 \}, \quad (2.1)$$

where

$$[A, p; k] = \{ (x, y) : x \in E, y \in A \text{ and } \rho(x, A) < \rho^k(x, p) = \rho^k(y, p) \}. \quad (2.2)$$

In the paper [4] it was proved that  $A_{p,k}^* \subset \tilde{M}_{p,k}$  for any  $k > 0$  and  $p \in E$ . With this connection the Theorems 1.1, 1.2 mentioned in Section 1 of this paper are fulfilled in the classes of sets  $A_{p,k}^* \cap D_p(E, l)$ . It appears that these theorems will be true for sets of the classes  $A_{p,k}^* \cap D_p(E, l)$  at slightly weaker conditions concerning the functions  $a, b, a_i, b_i$  ( $i = 1, 2$ ) appearing in the assumptions of the above theorems.

In the paper [3] the following lemma was proved:

**LEMMA 2.1.** *If*

$$\frac{a(r)}{r^k} \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0, \quad (2.3)$$

*then for an arbitrary set  $A \in A_{p,k}^* \cap D_p(E, \rho)$*

$$\frac{1}{r^k} d_\rho(A \cap S_\rho(p, r)_{a(r)}) \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0. \quad (2.4)$$

From this lemma and from (6) it follows that

$$\frac{1}{r^k} d_l(A \cap S_l(p, r)_{a(r)}) \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0, \quad (2.5)$$

for  $l \in \overline{F}_f$  and  $A \in A_{p,k}^* \cap D_p(E, l)$ , when the function  $a$  fulfills the condition (2.3).

Similarly as in the case of the classes of sets  $\tilde{M}_{p,k}$ , using (2.5) we can prove the following theorem.

**THEOREM 2.1.** *If  $l_i \in \overline{F}_f$  ( $i = 1, 2$ ),*

$$\frac{a(r)}{r^k} \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0, \quad \text{and} \quad \frac{b(r)}{r^k} \xrightarrow[r \rightarrow 0+]{\longrightarrow} 0, \quad (2.6)$$

then for arbitrary sets of the classes  $A_{p,k}^* \cap D_p(E,l)$  the tangency relations  $T_{l_1}(a,b,k,p)$  and  $T_{l_2}(a,b,k,p)$  are compatible.

Let  $a_i, b_i$  ( $i = 1, 2$ ) be non-negative real functions defined in a certain right-hand side neighbourhood of 0 and fulfilling the condition (1.16). In the paper [8] the following theorem was proved.

**THEOREM 2.2** *If  $l \in \overline{F}_f$  and*

$$\frac{a_i(r)}{r^k} \xrightarrow[r \rightarrow 0+]{ } 0 \quad \text{and} \quad \frac{b_i(r)}{r^k} \xrightarrow[r \rightarrow 0+]{ } 0 \quad \text{for } i = 1, 2, \quad (2.7)$$

then for arbitrary sets of the classes  $A_{p,k}^* \cap D_p(E,l)$  the tangency relations  $T_l(a_1, b_1, k, p)$  and  $T_l(a_2, b_2, k, p)$  are compatible.

From the Theorems 2.1 and 2.2 the following corollary results.

**COROLLARY 2.1.** *If the functions  $a_i, b_i$  ( $i = 1, 2$ ) fulfil the condition (2.7) and  $l_i \in \overline{F}_f$ , then the tangency relations  $T_{l_1}(a_1, b_1, k, p)$  and  $T_{l_2}(a_2, b_2, k, p)$  are compatible in the classes of sets  $A_{p,k}^* \cap D_p(E,l)$ .*

Let  $id$  denotes the identity function defined in a right-hand side neighbourhood of 0. If we put  $f = id$ , then the class  $\overline{F}_{id}$  of the function  $l$  is equal to the class  $F_\rho^*$  (see [3], [4]). From here it results that all theorems about the problem of the compatibility of the tangency relations of sets for the functions of the class  $F_\rho^*$  given in the papers [3] and [4] follow from the theorems of this paper.

#### REFERENCES

- [1] A. Chądzyńska, *On some classes of sets related to the symmetry of the tangency relation in a metric space*, Ann. Soc. Math. Polon., Comm. Math. **16** (1972), 219–228.
- [2] S. Golab et Z. Moszner, *Sur le contact des courbes dans les espaces métriques généraux*, Colloq. Math. **10** (1963), 105–311.
- [3] T. Konik, *On the compatibility of the tangency relations of sets of the classes  $A_{p,k}^*$  in generalized metric spaces*, Demonstratio Math. **19** (1986), 203–220.
- [4] T. Konik, *On the tangency of sets of some class in generalized metric spaces*, Demonstratio Math. **22** (1989), 1093–1107.
- [5] T. Konik, *On the tangency of sets in generalized metric spaces for certain functions of the class  $F_\rho^*$* , Mat. vesnik **43** (1991), 1–10.
- [6] T. Konik, *On the tangency of sets of the class  $\tilde{M}_{p,k}$* , Publ. Math. Debrecen, **43/3–4** (1993), 329–336.
- [7] T. Konik, *Tangency relations for sets in some classes in generalized metric spaces*, Math. Slowaca (in print).
- [8] T. Konik, *On the compatibility and the equivalence of the tangency relations of sets of the classes  $A_{p,k}^*$* , Journal of Geometry (Basel) (in print).
- [9] W. Waliszewski, *On the tangency of sets in generalized metric spaces*, Ann. Polon. Math. **28** (1973), 275–284.

(received 10.09.1997.)

Institute of Mathematics & Computer Science, Technical University, Dąbrowskiego 73, 42-200 Częstochowa, Poland