

## BAIRE'S SPACE OF PERMUTATIONS OF $\mathbf{N}$ AND REARRANGEMENTS OF SERIES

Tibor Šalát

**Abstract.** In the first part of the paper we investigate the structure of the space  $(S, d)$  of all sequences of positive integers with Baire's metric. In the second part we study properties of the space  $(E, d)$  of all permutations of  $\mathbf{N}$  in connection with rearrangements of non-absolutely convergent series.

### 0. Introduction

There are several papers investigating rearrangements of non-absolutely convergent series from the point of view of permutations of the set  $\mathbf{N}$  considered as points of a metric space (cf. [1], [3], [4], [5], [7], [8], [9], [10]). The mentioned metric space is endowed with the Fréchet's metric  $d_1$  in [1], [3], [4], [7], [9], [10] and with Baire's metric  $d$  in [5]. Let us remark that these two metrics are equivalent on the set  $S$  (and also on  $E$ ) of all sequences of positive integers (of all permutations of  $\mathbf{N}$ ) and therefore many properties of  $(S, d_1)$  can be transferred from  $(S, d_1)$  to  $(S, d)$  (or from  $(E, d_1)$  to  $(E, d)$ ), and conversely.

Several of our considerations will be based on the following classical Riemann's theorem on rearrangements of series with real terms (cf. [5]).

**THEOREM A.** *Let  $\sum_{k=1}^{\infty} a_k$  be a non-absolutely convergent series with real terms, let  $-\infty \leq t_1 \leq t_2 \leq +\infty$ . Then there exists a permutation  $x = (x_j)_{j=1}^{\infty} \in E$  such that  $\liminf_{n \rightarrow \infty} S_n(x) = t_1$ ,  $\limsup_{n \rightarrow \infty} S_n(x) = t_2$ , where  $S_n(x) = \sum_{j=1}^n a_{x_j}$  ( $n = 1, 2, \dots$ ).*

*Particularly, for every  $r \in \mathbf{R}$  there exists a permutation  $x = (x_j)_{j=1}^{\infty} \in E$  such that*

$$\sum_{j=1}^{\infty} a_{x_j} = r. \quad (1)$$

---

*AMS Subject Classification:* 40 A05

*Keywords and phrases:* Rearrangements of series, Baire's space, set of first category, residual set,  $\sigma$ -porosity, strong porosity

The set of all permutations (of  $\mathbf{N}$ )  $x = (x_j)_1^\infty$  with (1) will be denoted by  $P_r$  (in agreement with [5]). The set  $P_r$  depends obviously on the sequence  $a_1, a_2, \dots$  and therefore we shall write in detail  $P_r = P_r(a_1, a_2, \dots)$ . Put

$$E_0 = E_0(a_1, a_2, \dots) = \bigcup_{r \in \mathbf{R}} P_r(a_1, a_2, \dots).$$

Although the sets  $E_0(a_1, a_2, \dots)$ ,  $P_r(a_1, a_2, \dots)$  depend on  $a_1, a_2, \dots$ , they have several common properties for all  $a_1, a_2, \dots$  such that  $\sum_{k=1}^\infty a_k$  is a non-absolutely convergent series.

Recall the concept of metrics  $d_1$  and  $d$  (on  $S$  or  $E$ ). Let  $x = (x_j)_1^\infty \in S$ ,  $y = (y_j)_1^\infty \in S$ . Then we put

$$d_1(x, y) = \sum_{k=1}^{\infty} \frac{1}{2^k} \frac{|x_k - y_k|}{1 + |x_k - y_k|}.$$

If  $x = y$ , then  $d(x, y) = 0$  and if  $x \neq y$ , then  $d(x, y) = 1/m$ , where  $m = \min\{j : x_j \neq y_j\}$  (cf. [1], [2], p. 185).

We shall use the concept of porosity of sets (cf. [12], [13]). Let  $(Y, \rho)$  be a metric space,  $y \in Y$  and  $r > 0$ . Then by  $K(y, r)$  denote the ball with centre  $y$  and radius  $r$ , i.e.  $K(y, r) = \{x \in Y : \rho(x, y) < r\}$ . Let  $M \subseteq Y$ . Put

$$\gamma(y, r, M) = \sup\{t > 0 : (\exists z \in Y) [K(z, t) \subseteq K(y, r)] \wedge [K(z, t) \cap M = \emptyset]\},$$

$\bar{p}(y, M) = \limsup_{r \rightarrow 0+} \gamma(y, r, M)/r$ ,  $\underline{p}(y, M) = \liminf_{r \rightarrow 0+} \gamma(y, r, M)/r$  and if  $\bar{p}(y, M) = \underline{p}(y, M)$ , then set

$$p(y, M) = \bar{p}(y, M) = \underline{p}(y, M) = \lim_{r \rightarrow 0+} \frac{\gamma(y, r, M)}{r}.$$

Obviously, each of the numbers  $\bar{p}(y, M)$ ,  $\underline{p}(y, M)$ ,  $p(y, M)$  belongs to the interval  $[0, 1]$ .

A set  $M \subseteq Y$  is said to be porous ( $c$ -porous) at  $y$ , provided that  $\bar{p}(y, M) > 0$  ( $\bar{p}(y, M) \geq c > 0$ ). A set  $M \subseteq Y$  is said to be  $\sigma$ -porous ( $\sigma$ - $c$ -porous) at  $y$  provided that  $M = \bigcup_{n=1}^\infty M_n$ ,  $M_n$  ( $n = 1, 2, \dots$ ) being porous ( $c$ -porous) at  $y$ .

Let  $Y_0 \subseteq Y$ . A set  $M \subseteq Y$  is said to be porous,  $c$ -porous,  $\sigma$ -porous and  $\sigma$ - $c$ -porous in the set  $Y_0$  if  $M$  is porous,  $c$ -porous,  $\sigma$ -porous and  $\sigma$ - $c$ -porous at every point  $y \in Y_0$ , respectively.

If a set  $M$  is  $c$ -porous and  $\sigma$ - $c$ -porous at  $y$ , then obviously it is porous and  $\sigma$ -porous at  $y$ , respectively.

Every porous set  $M$  in  $Y$  is a nowhere dense set in  $Y$  and therefore every  $\sigma$ -porous set  $M$  in  $Y$  is a set of the first Baire category in  $Y$ . The converse is not true already in  $\mathbf{R}$  (cf. [11]).

From the definition of numbers  $\bar{p}(y, M)$ ,  $\underline{p}(y, M)$  we get at once: If  $M_1 \subseteq M_2 \subseteq Y$ , then for each  $y \in Y$  we have  $\bar{p}(y, M_1) \geq \bar{p}(y, M_2)$ ,  $\underline{p}(y, M_1) \geq \underline{p}(y, M_2)$ .

Let  $Y$  be a metric space again. A set  $M \subseteq Y$  is said to be very porous at  $y \in Y$  if  $\underline{p}(y, M) > 0$  and very strongly porous at  $y$  if  $\underline{p}(y, M) = 1$  (cf. [1], p. 327). The set  $\bar{M}$  is said to be very (strongly) porous in  $Y_0 \subseteq Y$  if it is very (strongly) porous at each point  $y \in Y_0$ .

Obviously, if  $M$  is very porous at  $y$ , it is porous at  $y$ , as well. Analogously, if  $M$  is very strongly porous at  $y$ , it is 1-porous at  $y$ .

Further, a set  $M \subseteq Y$  is said to be uniformly very porous (in  $Y_1 \subseteq Y$ ) provided that there is a  $c > 0$  such that for each  $y \in Y_1$  we have  $\underline{p}(y, M) \geq c > 0$  (cf. [13], p. 327).

In agreement with the previous terminology and in analogy with the notion of  $\sigma$ -porosity we introduce the following concept of uniformly very  $\sigma$ -porous sets.

DEFINITION. 1) A set  $M \subseteq Y$  is said to be uniformly very  $\sigma$ -porous (in  $Y_0 \subseteq Y$ ) provided that  $M = \bigcup_{n=1}^{\infty} M_n$  and there is a  $c > 0$  such that for each  $y \in Y_0$  we have  $\underline{p}(y, M_n) \geq c > 0$  ( $n = 1, 2, \dots$ ).

2) A set  $M \subseteq Y$  is said to be uniformly very strongly  $\sigma$ -porous (in  $Y_0 \subseteq Y$ ) provided that  $M = \bigcup_{n=1}^{\infty} M_n$  and for each  $y \in Y_0$  we have  $\underline{p}(y, M_n) = 1$  ( $n = 1, 2, \dots$ ).

If  $A \subseteq Y$ , then by  $CA$  we denote the complement of  $A$ ,  $CA = Y \setminus A$ .

In the first part of the paper we shall investigate the structure of the space  $(S, d)$ . In the second part we shall study properties of the space  $(E, d)$  in connection with rearrangements of non-absolutely convergent series.

## 1. Structure of the space $(S, d)$

We shall study the structure of  $(S, d)$  from the point of view of its subset  $E$  and closure  $\bar{E} = E_1$  of  $E$  in  $S$ .

It is well-known (cf. [1], [5], [9], [10]) that the set  $E$  is not closed in  $S$ . Its closure  $E_1 = \bar{E}$  equals to the set of all  $x = (x_j)_1^{\infty} \in S$  containing every positive integer at most once (equivalently:  $E_1$  consists of all one-to-one sequences of positive integers).

The metric space  $(E, d)$  (subspace of  $(S, d)$ ) is of the second category at each of its points (cf. [1], [4], [5], [9], [10]). We recall the following well-known result (cf. [1], [4], [5], [7], [9], [10]):

THEOREM B. Let  $\sum_{k=1}^{\infty} a_k$  be a non-absolutely convergent series. Denote  $H = H(a_1, a_2, \dots)$  the set of all  $x = (x_j)_1^{\infty} \in E$  such that  $\liminf_{n \rightarrow \infty} S_n(x) = -\infty$ ,  $\limsup_{n \rightarrow \infty} S_n(x) = +\infty$  ( $S_n(x) = \sum_{j=1}^n a_{x_j}$ ,  $n = 1, 2, \dots$ ). Then the set  $H$  is a residual set in  $E$  (i.e.  $E \setminus H$  is a set of the first Baire category in  $E$ ).

This result has been strengthened in [3], where it is proved that the set of all  $x = (x_j)_1^{\infty} \in E$  with  $(S_n(x))'_n = [-\infty, \infty]$  is residual in  $E$  ( $(S_n(x))'_n$  denotes the set of all limit points of the sequence  $(S_n(x))_{n=1}^{\infty}$ ).

It is well-known that  $E$  is a  $G_\delta$ -set in  $S$  (cf. [1]). We shall complete this result in the following proposition and for the completeness we shall give also the proof of the mentioned result from [1] (see (i) in the following proposition).

PROPOSITION 1.1. *The set  $E \subset S$  has the following properties:*

- (i) *The set  $E$  belongs exactly to the first Borel class, it is a  $G_\delta$ -set in  $S$ .*
- (ii) *The set  $E$  is dense in itself.*
- (iii) *The set  $E$  is nowhere dense in  $S$ .*

*Proof.* (i) Put (as in [1])  $G(n, k) = \{x = (x_j)_1^\infty \in S : x_n = k\}$ . If  $x \in G(n, k)$  then it is easy to see that  $K(x, 1/n) \subseteq G(n, k)$ , so the set  $G(n, k)$  is open in  $S$ . But then the set

$$E_2 = \bigcap_{k=1}^{\infty} \bigcup_{n=1}^{\infty} G(n, k) \quad (2)$$

is a  $G_\delta$ -set in  $S$ . Further we have obviously

$$E = E_1 \cap E_2, \quad (2')$$

$E_1$  being closed in  $S$ . Hence by (2') the set  $E$  is a  $G_\delta$ -set in  $S$ .

Further,  $E$  is neither closed in  $S$  (cf. [1], [5]), nor open in  $S$  (see (iii)), so the set  $E$  belongs exactly to the first Borel class.

(ii) Let  $y \in E$ ,  $\varepsilon > 0$ . Choose  $s \in \mathbf{N}$  such that  $1/s < \varepsilon$ . Define  $z = (z_j)_1^\infty$  as follows: Put  $z_j = y_j$  ( $j = 1, 2, \dots, s$ ),  $z_{s+1} = y_{s+1} + 1$  and construct the one-to-one sequence  $z_{s+2}, z_{s+3}, \dots$  containing all positive integers different from  $z_1, \dots, z_s, z_{s+1}$ . Then  $z = (z_j)_1^\infty \in E$ ,  $z \in K(y, \varepsilon)$  and  $z \neq y$ .

(iii) Let  $y \in S$ ,  $\varepsilon > 0$ . We show that there exists a ball  $K_0 \subseteq K(y, \varepsilon)$  such that  $K_0 \cap E = \emptyset$ . Then on account of a well-known criterion of nowhere density (cf. [6], p. 37), the assertion follows.

Choose  $s \in \mathbf{N}$  such that  $1/s < \varepsilon$ . Put  $z_j = y_j$  ( $j = 1, 2, \dots, s$ ),  $z_{s+1} = z_s$  and  $z_{s+k} = 1$  ( $k = 2, 3, \dots$ ). Then  $z = (z_j)_1^\infty \in K(y, \varepsilon)$ ,  $K(z, 1/(s+2)) \subseteq K(y, \varepsilon)$  and  $K(z, 1/(s+2)) \cap E = \emptyset$ . ■

The nowhere density of the set  $E$  in  $S$  follows also from the following result.

THEOREM 1.1. *The set  $E_1 = \overline{E}$  is very strongly porous in  $S$ .*

COROLLARY. a) *The set  $E_1$  is nowhere dense in  $S$ .*

b) *The set  $E$  is nowhere dense in  $S$ .*

*Proof of Theorem 1.1.* Let  $q = (q_j)_1^\infty \in S$ ,  $0 < \varepsilon < 1$ . Choose  $s \in \mathbf{N}$  such that  $1/s \leq \varepsilon < 1/(s-1)$  ( $s \geq 2$ ). Construct  $y = (y_j)_1^\infty \in S$  in this way: Put  $y_i = q_i$  ( $i = 1, 2, \dots, s$ ),  $y_{s+1} = q_s$ ,  $y_{s+j} = 1$  ( $j = 2, 3, \dots$ ). Then  $y \in K(q, 1/s)$  and  $K(y, 1/(s+1)) \subseteq K(q, 1/s) \subseteq K(q, \varepsilon)$ . If  $z = (z_j)_1^\infty \in K(y, 1/(s+1))$  then  $z_s = z_{s+1}$ , so  $K(y, 1/(s+1)) \cap E_1 = \emptyset$ . But then by the definition of  $\gamma(q, \varepsilon, E_1)$  we get

$$\frac{\gamma(q, \varepsilon, E_1)}{\varepsilon} \geq \frac{s-1}{s+1}.$$

If  $\varepsilon \rightarrow 0+$ , the  $s \rightarrow \infty$  and so we get

$$\underline{p}(q, E_1) = \liminf_{\varepsilon \rightarrow 0+} \frac{\gamma(q, \varepsilon, E_1)}{\varepsilon} = 1. \quad \blacksquare$$

In the representation  $E = E_1 \cap E_2$  of the set  $E$  (see (2')) the set  $E_1$  is very strongly porous in  $S$  (hence it is a nowhere dense set). We now show that  $E_2$  is a residual set in  $S$ . This follows from the following result.

**THEOREM 1.2.** *The set  $CE_2 = S \setminus E_2$  is uniformly very strongly  $\sigma$ -porous in  $S$ .*

*Proof.* By (2) we have  $E_2 = \bigcap_{k=1}^{\infty} B(k)$ , where  $B(k) = \bigcup_{n=1}^{\infty} G(n, k)$ . Using de Morgan's rule we get

$$CE_2 = \bigcup_{k=1}^{\infty} CB(k), \quad (3)$$

where  $CB(k) = \bigcap_{n=1}^{\infty} \{x = (x_j)_1^{\infty} \in S : x_n \neq k\}$ .

We show that  $p(y, CB(k)) = 1$  ( $k = 1, 2, \dots$ ) for each  $y \in S$ . Then (3) gives the assertion.

Let  $k$  be fixed,  $y \in S$ ,  $0 < \varepsilon < 1$ . Choose an  $s \in \mathbf{N}$  such that  $1/s \leq \varepsilon < 1/(s-1)$ . Put  $z_j = y_j$  ( $j = 1, 2, \dots, s$ ) and

$$z_{s+1} = k, \quad (4)$$

$z_{s+j} = 1$  ( $j = 2, 3, \dots$ ). Then  $K(z, 1/(s+1)) \subseteq K(y, 1/s) \subseteq K(y, \varepsilon)$  and by (4) we have  $K(z, 1/(s+1)) \cap CB(k) = \emptyset$ . Hence

$$\frac{\gamma(y, \varepsilon, CB(k))}{\varepsilon} \geq \frac{s-1}{s+1}.$$

If  $\varepsilon \rightarrow 0+$ , then  $s \rightarrow \infty$  and so we get  $p(y, CB(k)) = 1$ .  $\blacksquare$

## 2. Structure of the space $(E, d)$ and rearrangements of series

If  $\sum_{k=1}^{\infty} a_k$  is a non-absolutely convergent series then by Theorem B the set  $E_0 = E_0(a_1, a_2, \dots)$  is a set of the first category in  $E$ . We shall complete this result by showing that  $E_0$  is a  $\sigma$ -1-porous set at points of a large subset of  $S$ .

In the first place we shall investigate to which Borel classes the set  $E_0(a_1, a_2, \dots)$  and the sets

$$H^+ = H^+(a_1, a_2, \dots) = \{x \in E : \limsup_{n \rightarrow \infty} S_n(x) = +\infty\},$$

$$H^- = H^-(a_1, a_2, \dots) = \{x \in E : \liminf_{n \rightarrow \infty} S_n(x) = -\infty\}$$

belong.

**THEOREM 2.1.** *For each non-absolutely convergent series  $\sum_{k=1}^{\infty} a_k$  the set  $E_0 = E_0(a_1, a_2, \dots)$  is an  $F_{\sigma\delta}$ -set in  $E$ .*

REMARK 2.1. The foregoing theorem does not state exactly the Borel class of the set  $E_0$ . According to a result from [10] the set  $E_0$  is dense and simultaneously a boundary set in  $E$ . From this it is easy to see that  $E_0$  cannot belong to the zero Borel class. The set  $E_0$  cannot be a  $G_\delta$ -set in  $E$ , since in the opposite case it would be residual in  $S$  (cf. [6], p. 49). But this contradicts Theorem B. One can conjecture that  $E_0$  is an  $F_\sigma$ -set in  $S$ , but I am not able to prove or disprove this conjecture.

*Proof of Theorem 2.1.* By the definition of the set  $E_0$  and Cauchy's test for convergence a permutation  $x = (x_j)_1^\infty \in E$  belongs to  $E_0$  if and only if the following holds:

$$(\forall k)(\exists m)(\forall j) |S_{m+j}(x) - S_m(x)| \leq \frac{1}{k}.$$

From this we get

$$E_0 = \bigcap_{k=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcap_{j=1}^{\infty} \{x \in E : |S_{m+j}(x) - S_m(x)| \leq 1/k\}. \quad (5)$$

Consider that by fixed  $m$  and  $j$  the function  $S_{m+j}(x) - S_m(x) = \sum_{n=m+1}^{m+j} a_{x_n}$  is constant on every ball  $K(y, 1/(m+j))$ ,  $y \in S$ . Hence it is continuous on  $S$ . The assertion follows from (5). ■

It is proved in [9] that the set  $B = B(a_1, a_2, \dots) = E \setminus (H^+ \cup H^-)$  belongs to the first Borel class and is an  $F_\sigma$ -set in  $S$ . We show that this result is exact.

THEOREM 2.2 *If  $\sum_{k=1}^{\infty} a_k$  is a non-absolutely convergent series, then the sets  $H^+(a_1, a_2, \dots)$ ,  $H^-(a_1, a_2, \dots)$  and  $B(a_1, a_2, \dots)$  belong exactly to the first Borel class, the sets  $H^+$ ,  $H^-$  are  $G_\delta$ -sets and  $B$  is an  $F_\sigma$ -set in  $S$ .*

COROLLARY. *The set  $H^* = H^+ \cup H^-$  belongs exactly to the first Borel class and it is a  $G_\delta$ -set in  $S$ .*

*Proof of Theorem 2.2.* By using Theorem A it is easy to see that each of the sets  $H^+$ ,  $H^-$ ,  $B$  is dense and simultaneously boundary in  $S$ . From this we see that none of these sets belongs to zero Borel class. Further

$$H^+ = \bigcap_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{x \in E : S_n(x) > m\}. \quad (6)$$

We show that the set  $A(n, m) = \{x \in E : S_n(x) > m\}$  is open in  $S$ . Indeed, if  $x \in A(n, m)$  then  $K(x, 1/n) \subseteq A(n, m)$ . But then from (6) we get that  $H^+$  belongs exactly to the first Borel class and it is a  $G_\delta$ -set in  $E$ .

Similarly it can be shown that  $H^-$  belongs exactly to the first Borel class and it is a  $G_\delta$ -set in  $E$ . The statement related to  $B$  follows from  $B = E \setminus H^*$ , where  $H^* = H^+ \cup H^-$ . ■

Put  $B_1 = B_1(a_1, a_2, \dots) = \{x \in E : (S_n(x))_1^\infty \text{ is bounded from above}\}$ ,  $B_2 = B_2(a_1, a_2, \dots) = \{x \in E : (S_n(x))_1^\infty \text{ is bounded from below}\}$ . Hence

$B_1 = S \setminus H^+$ ,  $B_2 = S \setminus H^-$ . Then  $B = B_1 \cap B_2$  and  $E_0 \subseteq B$ . We shall investigate the porosity of sets  $B_1$ ,  $B_2$ ,  $B$ ,  $E_0$ .

LEMMA 2.1. *Let  $\sum_{k=1}^{\infty} a_k$  be an arbitrary non-absolutely convergent series. Then the set  $B_1 = B_1(a_1, a_2, \dots)$  is  $\sigma$ -1-porous in the set  $H^+ = H^+(a_1, a_2, \dots)$ .*

COROLLARY. *The sets  $B$ ,  $E_0$  are  $\sigma$ -1-porous in  $H^+$ .*

*Proof of Lemma 2.1.* Observe that

$$B_1 = \bigcup_{m=1}^{\infty} D(m), \quad (7)$$

where  $D(m) = \bigcap_{j=1}^{\infty} \{x \in E : S_j(x) \leq m\}$  ( $m = 1, 2, \dots$ ).

Let  $y = (y_j)_1^{\infty} \in H^+$ . Then by definition of  $H^+$  there exists a sequence  $v_1 < v_2 < \dots < v_k < \dots$  of positive integers such that  $\lim_{k \rightarrow \infty} S_{v_k}(y) = +\infty$ . Construct the ball  $K(y, 1/v_k)$ . Then for all sufficiently large  $k$ 's (say for  $k > k_0$ ) we have  $K(y, 1/v_k) \cap D(m) = \emptyset$ , thus  $\gamma(y, v_k^{-1}, D(m)) = 1/v_k$  (for  $k > k_0$ ). Since  $v_k^{-1} \rightarrow 0$  ( $k \rightarrow \infty$ ), we get  $v_k \gamma(y, v_k^{-1}, D(m)) = 1$ . Thus  $\bar{p}(y, D(m)) = 1$  ( $m = 1, 2, \dots$ ). Lemma 2 follows from (7). ■

Similarly the following lemma can be proved.

LEMMA 2.2. *Let  $\sum_{k=1}^{\infty} a_k$  be a non-absolutely convergent series. Then the set  $B_2(a_1, a_2, \dots)$  is  $\sigma$ -1-porous in the set  $H^-(a_1, a_2, \dots)$ .*

On account of Theorem B, using Lemma 2.1 and Lemma 2.2 we obtain the following result.

THEOREM 2.3. *Let  $\sum_{k=1}^{\infty} a_k$  be an arbitrary non-absolutely convergent series. Then each of the sets  $B(a_1, a_2, \dots)$ ,  $E_0(a_1, a_2, \dots)$  is  $\sigma$ -1-porous in the residual set  $H^+(a_1, a_2, \dots) \cap H^-(a_1, a_2, \dots)$ .*

#### REFERENCES

- [1] R. P. Agnew, *On rearrangements of series*, Bull. Amer. Math. Soc. **46** (1940), 797–799.
- [2] P. S. Alexandrov, *Úvod do obecné theorie množin a funkcí*, NČSAV, Praha, 1954.
- [3] J. Červeňanský, *Rearrangements of series and a topological characterization of the absolute convergence of series*, Acta Fac. Rer. Nat. Univ. Com. **34** (1979), 75–91.
- [4] P. L. Ganguli, B. K. Lahiri, *Some results on certain sets of series*, Czechosl. Math. J. **18** (93) (1968), 589–594.
- [5] I. Hozo, H. I. Miller, *On Riemann's theorem about conditionally convergent series*, Mat. Vesnik **38** (1986), 279–283.
- [6] C. Kuratowski, *Topologie I*, PWN, Warszawa, 1958.
- [7] Gy. L. Pál, *On a problem from the theory of series* (Hungarian), Mat. Lap. **12** (1961), 38–43.
- [8] M. Bhaskara Rao, K. P. S. Bhaskara Rao, B. V. Rao, *Remarks on subsequences, subseries and rearrangements*, Proc. Amer. Math. Soc. **67** (1977), 293–296.
- [9] H. M. Sengupta, *On rearrangements of series*, Proc. Amer. Math. Soc. **1** (1950), 71–75.
- [10] H. M. Sengupta, *Rearrangements of series*, Proc. Amer. Math. Soc. **7** (1956), 347–350.

- [11] J. Tkadlec, *Constructions of some non- $\sigma$ -porous sets of real line*, Real Anal. Exchange **9** (1983–84), 473–482.
- [12] L. Zajíček, *Sets of  $\sigma$ -porosity and sets of  $\sigma$ -porosity ( $q$ )*, Čas. pěst. mat. **101** (1976), 350–359.
- [13] L. Zajíček, *Porosity and  $\sigma$ -porosity*, Real Anal. Exchange **13** (1987–88), 314–350.

(received 11.06.1996.)

Comenius University, Mlynská dolina, 842 15 Bratislava, Slovakia