

EXTREMAL PROPERTIES OF THE CHROMATIC POLYNOMIALS OF CONNECTED 3-CHROMATIC GRAPHS

Ioan Tomescu

Abstract. In this paper the greatest $\lceil n/2 \rceil$ values of $P(G; 3)$ in the class of connected 3-chromatic graphs G of order n are found, where $P(G; \lambda)$ denotes the chromatic polynomial of G .

1. Preliminary definitions and results

Let G be a graph of order n and let $P(G; \lambda)$ be its chromatic polynomial [1]. A k -color partition of G is a partition of the vertex set $V(G)$ into k classes where each class is an independent set of vertices. The number of k -color partitions of G and the chromatic number of G will be denoted by $\text{Col}_k(G)$ and by $\chi(G)$, respectively. It is well known that $P(G; \lambda)$ can be expressed in terms of the number of k -color partitions as follows

$$P(G; \lambda) = \sum_{k=1}^n (\lambda)_k \text{Col}_k(G),$$

where $(\lambda)_k = \lambda(\lambda - 1) \cdots (\lambda - k + 1)$.

It follows that if $\chi(G) = k$, then $\text{Col}_k(G) = P(G; \lambda)/k!$. Let xy be an edge of G . By $G - xy$ we mean the graph obtained from G by deleting edge xy . Also G/xy denotes the graph obtained from G by identifying vertices x and y , i.e., (i) by deleting both x and y and all the edges incident to them, and (ii) by introducing a new vertex z and joining z to both all the neighbors of x different from y and all the neighbors of y different from x in G .

The following lemma describes some properties of $P(G; \lambda)$, which we will use later [2].

LEMMA 1.1. *The following properties hold:*

(i) *Reduction Formula.* Let a and b be two adjacent vertices of G . Then $P(G; \lambda) = P(G - ab; \lambda) - P(G/ab; \lambda)$.

AMS Subject Classification: 05C15

Keywords and phrases: Chromatic polynomial, connected 3-chromatic graph, 3-color partition, skeleton of a graph.

(ii) Let G and H be two graphs that overlap in a complete graph K_r on r vertices. Then the chromatic polynomial of this overlap graph is

$$P(G; \lambda)P(H; \lambda)/P(K_r; \lambda).$$

Let G be a graph and H an induced subgraph of G . The graph obtained from G by the contraction of H is the graph G_1 derived from G by the following operations: suppress all vertices of H and the edges incident with them, and replace them with a new vertex $w \notin V(G)$ and edges wx such that $wx \in E(G_1)$ if and only if there exists $y \in V(G)$ such that $xy \in E(G)$ and $x \in V(G) - V(H)$.

The cycle with n vertices will be denoted by C_n and C_n^1 will denote the graph consisting of C_n and one more vertex adjacent to only one vertex of C_n . The following theorem was proved in [4].

THEOREM 1.2. *The maximum number of 3-color partitions of a connected graph G having n vertices and chromatic number $\chi(G) = 3$ is $(2^{n-1} - 1)/3$ for odd n , and $(2^{n-1} - 2)/3$ for even n . Moreover, if n is odd, the unique connected graph that achieves the maximum number of 3-color partitions is C_n , while if n is even, the unique graph is C_{n-1}^1 .*

By $H(n, 2r+1)$ we denote the class of connected graphs G of order n containing n edges and a unique cycle C_{2r+1} , where $3 \leq 2r+1 \leq n$. It is clear that the graph deduced from $G \in H(n, 2r+1)$ by contracting C_{2r+1} is a tree on $n - 2r$ vertices. By Rényi's formula [3], the number of labeled graphs in $H(n, 2r+1)$ is equal to $(n-1)_{2r} n^{n-2r-1}/2$.

Let D_n ($n \geq 5$) be the graph consisting of a 4-cycle in which two nonadjacent vertices are connected by a newly added path of length $n-3$. Note that $\chi(D_n) = 3$ for even n and $\chi(D_n) = 2$ for odd n . If "nonadjacent" is replaced by "adjacent", the resulting graph is denoted by F_n . Hence, F_n consists of two cycles C_4 and C_{n-2} having a common edge. Also, $\chi(F_n) = 3$ for odd n and $\chi(F_n) = 2$ for even n .

The following two properties were deduced in [5].

LEMMA 1.3. *For every $n \geq 5$, the following equalities hold: $P(D_n; 3) = 2^n - 2^{n-2} + (-1)^{n-1}6$ and $P(F_n; 3) = 2^n - 2^{n-2} + (-1)^n 6$.*

THEOREM 1.4. (a) *If G is a 2-connected graph of order n , $n \geq 5$, such that $P(G; 3)$ is maximum in the class $\mathcal{F}_n \setminus \{C_n, K_{2,n-2}, D_n\}$, where \mathcal{F}_n denotes the class of all 2-connected graphs of order n , then $G \cong F_n$ for odd n .*

(b) *If G is a 2-connected graph of order 6 such that $P(G; 3)$ is maximum in the class $\mathcal{F}_6 \setminus \{C_6, K_{2,4}, F_6, K_{3,3} - e\}$, then $G \cong K_{3,3}$ or D_6 .*

(c) *If G is a 2-connected graph of order n , $n \geq 8$, such that $P(G; 3)$ is maximum in the class $\mathcal{F}_n \setminus \{C_n, K_{2,n-2}, F_n\}$, then $G \cong D_n$ for even n ; for $n = 8$ there exists another extremal graph, $E_{8,3}$.*

Note that the graph $E_{8,3}$, described in [5], has $\chi(E_{8,3}) = 2$; also $\chi(K_{2,n-2}) = \chi(K_{3,3} - e) = \chi(K_{3,3}) = 2$.

LEMMA 1.5. *Let G be a graph of order $n \geq 5$ consisting of two cycles C_{2r+1} and C_{n-2r} having exactly one vertex in common. Then $P(G; 3) < 2^n - 2^{n-2} - 6$.*

Proof. By Lemma 1.1(ii) we get

$$P(G; \lambda) = ((\lambda - 1)^{2r+1} - (\lambda - 1))((\lambda - 1)^{n-2r} + (-1)^{n-2r}(\lambda - 2))/\lambda$$

since $P(C_n; \lambda) = (\lambda - 1)^n + (-1)^n(\lambda - 1)$. It follows that

$$\begin{aligned} P(G; 3) &= (2^{2r+1} - 2)(2^{n-2r} + (-1)^{n-2r} + (-1)^{n-2r}2)/3 \\ &\leq (2^{2r+1} - 2)(2^{n-2r} + 2)/3 = 2(2^n + 2^{2r+1} - 2^{n-2r} - 2)/3. \end{aligned}$$

Since $n - 2r \geq 3$, we shall consider two subcases: Case I. $2r \leq n - 4$, and Case II. $2r = n - 3$.

Case I. If $2r \leq n - 4$ we deduce $2(2^n + 2^{2r+1} - 2^{n-2r} - 2)/3 \leq 2(2^n + 2^{n-3} - 2^4 - 2)/3 = 2^n - 2^{n-2} - 12 < 2^n - 2^{n-2} - 6$.

Case II. In this case $n - 2r = 3$ and $P(G; 3) = (2^{n-2} - 2)(2^3 - 2)/3 < 2^n - 2^{n-2} - 6$. ■

We define the skeleton $S(G)$ of a connected graph G as follows:

(α) If G has no vertex of degree one, then $S(G) = G$.

(β) Otherwise, let x be a vertex of degree one of G ; then G is replaced by $G - x$. Repeat (α).

For example, $S(T)$ consists of a unique vertex if T is a tree, and $S(G) = C_{2r+1}$ for any graph $G \in H(n, 2r + 1)$.

LEMMA 1.6. *Let G be a graph of order n such that its skeleton $S(G)$ has order r . Then $P(G; \lambda) = P(S(G); \lambda)(\lambda - 1)^{n-r}$.*

Proof. One applies Lemma 1.1(ii) since $P(K_2; \lambda) = \lambda(\lambda - 1)$. ■

COROLLARY 1.7. *For every $G \in H(n, 2r + 1)$, where $3 \leq 2r + 1 \leq n$, we have $P(G; \lambda) = (\lambda - 1)^n - (\lambda - 1)^{n-2r}$.*

LEMMA 1.8. *Let G be a connected graph of order n consisting of two vertex disjoint cycles C_r and C_s , joined by a path of length t ($r + s + t = n + 1$). Then*

$$P(G; \lambda) = P(H; \lambda)(\lambda - 1)^t,$$

where H is the graph of order $r + s - 1$ consisting of cycles C_r and C_s having a unique common vertex.

Proof. This equality is a consequence of Lemma 1.1(ii). ■

LEMMA 1.9. *Let G be a graph of order $2r + s + p$ consisting of two cycles—one cycle with $s \geq 3$ vertices and another odd cycle with $2r + 1 \geq 3$ vertices, having in common a path of length $p \geq 1$. Then*

$$P(G; 3) < P(H; 3) = 2^{2r+s-p} - 2^{2r+s-p-2}, \quad (1)$$

where $H \in H(2r + s - p, 3)$.

Proof. Suppose that the common path with $p + 1$ vertices of the two cycles of G has extremities a and b . It follows that $1 \leq p \leq 2r - 1$ and $p \leq s - 2$. If $p \geq 2$ then vertices a and b are not adjacent and by Lemma 1.1 we deduce

$$\begin{aligned} P(G; \lambda) &= P(G_1; \lambda) + P(G_2; \lambda) = \\ &= ((\lambda - 1)^{s-p} + (-1)^{s-p}(\lambda - 1))((\lambda - 1)^p + (-1)^p(\lambda - 1)) \times \\ &\quad \times ((\lambda - 1)^{2r-p+1} + (-1)^{2r-p+1}(\lambda - 1)) / \lambda^2 + \\ &+ ((\lambda - 1)^{s-p+1} + (-1)^{s-p+1}(\lambda - 1))((\lambda - 1)^{p+1} + (-1)^{p+1}(\lambda - 1)) \times \\ &\quad \times ((\lambda - 1)^{2r-p+2} + (-1)^{2r-p}(\lambda - 1)) / (\lambda^2(\lambda - 1)^2), \end{aligned}$$

where G_1 consists of three cycles with p , $s - p$ and $2r - p + 1$ vertices having a common vertex and G_2 of three cycles with $p + 1$, $s - p + 1$ and $2r - p + 2$ vertices having a common edge. Hence (1) is equivalent to

$$2^{2r+s-p} > (-1)^s 2^{2r-p+4} - 2^{s-p+3} + (-1)^{s+1} 2^{p+3} + (-1)^{s-p+1} 8. \quad (2)$$

For $s = 3$ we deduce $p = 1$ which contradicts our hypothesis. If $s \geq 4$ we can write $2^{2r+s-p} + (-1)^{s+1} 2^{2r-p+4} \geq 2^{2r+s-p} - 2^{2r-p+4} = 2^{2r-p+4}(2^{s-4} - 1) \geq 2^5(2^{s-4} - 1) = 2^{s+1} - 2^5$ since $p \leq 2r - 1$. Since $p \leq s - 2$, $2^{s-p+3} + (-1)^s 2^{p+3} \geq 2^{s-p+3} - 2^{p+3} = 2^5 - 2^{s+1}$ for $p = s - 2$ and $2^{s-p+3} - 2^{p+3} \geq 2^6 - 2^s$ for $p \leq s - 3$, and (2) is verified.

If $p = 1$ then cycles C_s and C_{2r+1} have an edge in common and $P(G; \lambda) = P(C_{2r+s-1}; \lambda) - P(G_3; \lambda)$, where G_3 consists of two cycles with $s - 1$ and $2r$ vertices having a common vertex. It follows that

$$P(G; 3) = 2^{2r+s-1} + (-1)^{s-1} 2 - (2^{s-1} + (-1)^{s-1} 2)(2^{2r} + 2)/3$$

and (1) is equivalent to $2^{2r+s-3} > (-1)^s 2^{2r+1} - 2^s + (-1)^{s-1} 2$. But this inequality can be deduced from (2) for $p = 1$ and it is also true for $s = 3$. ■

2. Main result

We shall denote by $\mathcal{C}_{n,3}$ the class of connected 3-chromatic graphs of order n . The following theorem is an extension of Theorem 1.2.

THEOREM 2.1. *Let $n \geq 5$. Then:*

(a) *For every $r = \lceil n/2 \rceil - 1$, $r = \lceil n/2 \rceil - 2, \dots, 1$, if G is a connected 3-chromatic graph of order n , such that $P(G; 3)$ is maximum in the class of graphs*

$$\mathcal{C}_{n,3} \setminus \bigcup_{s \geq r+1} H(n, 2s + 1),$$

then $G \in H(n, 2r + 1)$ and $P(G; 3) = 2^n - 2^{n-2r}$.

(b) *If $P(G; 3)$ is maximum in the class of graphs*

$$\mathcal{C}_{n,3} \setminus \bigcup_{s \geq 1} H(n, 2s + 1),$$

then $G \cong F_n$ for odd n , $G \cong D_n$ for even n and in this case $P(G; 3) = 2^n - 2^{n-2} - 6$.

Proof. (a) Let $G \in \mathcal{C}_{n,3}$. It follows that G contains an odd cycle C_{2r+1} . If for every edge $e \in E(G) \setminus E(C_{2r+1})$ the graph $G - e$ is not connected then $G \in H(n, 2r+1)$. Otherwise, by Lemma 1.1(ii) we have

$$P(G - e; 3) = P(G; 3) + P(G/e; 3). \quad (3)$$

But $\chi(G/e) = 3$ since G/e contains an odd cycle even if e is a chord of C_{2r+1} . It follows that $P(G/e; 3) > 0$ and (3) implies that $P(G - e; 3) > P(G; 3)$. By applying several times this operation of deleting edges not belonging to C_{2r+1} without disconnecting the resulting graph, one obtains a graph $H \in H(n, 2r+1)$ such that $P(H; 3) > P(G; 3)$. By Corollary 1.7 if $3 \leq 2j+1 < 2i+1 \leq n$ then $G_1 \in H(n, 2i+1)$ and $G_2 \in H(n, 2j+1)$ imply

$$P(G_1; 3) = 2^n - 2^{n-2i} > 2^n - 2^{n-2j} = P(G_2; 3)$$

and (a) is proved for $r = \lceil n/2 \rceil - 1$ (this is the property expressed by Theorem 1.2).

Let $G \in \bigcup_{s \geq 2} H(n, 2s+1)$ and a, b be two nonadjacent vertices of G . We shall prove that if $e = ab$ then

$$P(G + e; 3) < 2^n - 2^{n-2} = P(H; 3), \quad (4)$$

where $H \in H(n, 3)$.

It is clear that the skeleton $S(G + e)$ consists of: I. Two vertex disjoint cycles joined by a path of length $t \geq 1$; II. Two cycles having exactly one common vertex; III. Two cycles having in common a path of length $p \geq 1$. In all cases at least one cycle is odd. Suppose that $|S(G + e)| = m$.

Case I. In this case by Lemmas 1.6 and 1.8 one deduces

$$P(G + e; \lambda) = P(S(G + e); \lambda)(\lambda - 1)^{n-m} = P(H; \lambda)(\lambda - 1)^{n-m+t},$$

where H has order $m - t$ and consists of two cycles (one is odd) having one vertex in common. By Lemma 1.5 we get

$$P(G + e; 3) = P(H; 3)2^{n-m+t} < (2^{m-t} - 2^{m-t-2} - 6)2^{n-m+t} < 2^n - 2^{n-2}.$$

Cases II, III. We have $P(G + e; 3) < (2^m - 2^{m-2})2^{n-m} = 2^n - 2^{n-2}$ by Lemmas 1.5, 1.6 and 1.9. Let now r be such that $1 \leq r \leq \lceil n/2 \rceil - 2$ and G be such that $P(G; 3)$ is maximum in the class $\mathcal{C}_{n,3} \setminus \bigcup_{s \geq r+1} H(n, 2s+1)$. If $G \in \bigcup_{s=1}^r H(n, 2s+1)$ it follows that $G \in H(n, 2r+1)$ and the property is proved. Otherwise, there exists an edge $e \in E(G)$ such that $G - e \in \mathcal{C}_{n,3}$. Since $P(G; 3)$ is maximum in the class $\mathcal{C}_{n,3} \setminus \bigcup_{s \geq r+1} H(n, 2s+1)$, it follows that $G - e \in \bigcup_{s \geq r+1} H(n, 2s+1)$, i.e., there exists a graph H in $\bigcup_{s \geq r+1} H(n, 2s+1)$ such that $G \cong H + e$. By (4) this leads to a contradiction.

(b) Let $G \in \mathcal{C}_{n,3} \setminus \bigcup_{s \geq 1} H(n, 2s+1)$ be such that $P(G; 3)$ is maximum. We have seen that the greatest values of $P(G; 3)$ in the class $\mathcal{C}_{n,3}$ are obtained for graphs in $\bigcup_{s \geq 1} H(n, 2s+1)$, and for graphs not belonging to this class the greatest values of $P(G; 3)$ are obtained for graphs of the form $H + e$, where $H \in \bigcup_{s \geq 1} H(n, 2s+1)$ and

$e \notin E(H)$. It follows that $G \cong H + e$, where $H \in \bigcup_{s \geq 1} H(n, 2s+1)$ and $e \notin E(H)$. Suppose that $|S(H + e)| = m$. As for the case (a) we may distinguish cases I–III concerning the structure of $S(H + e)$. Using the same notation, in the case I one obtains $P(H + e; 3) < (2^{m-t} - 2^{m-t-2} - 6)2^{n-m+t} < 2^n - 2^{n-2} - 6$ since $n - m + t \geq 1$. In the case II by Lemma 1.5, $P(H + e; 3) < (2^m - 2^{m-2} - 6)2^{n-m} \leq 2^n - 2^{n-2} - 6$.

In the case III the skeleton $S(H + e)$ is 2-connected and by Lemmas 1.3, 1.6 and Theorem 1.4 one deduces

$$P(H + e; 3) \leq (2^m - 2^{m-2} - 6)2^{n-m} \leq 2^n - 2^{n-2} - 6$$

and equality holds if and only if $m = n$ and $G \cong F_n$ for odd n and $G \cong D_n$ for even n . ■

Note that $\text{Col}_3(F_n)$ for odd n , resp. $\text{Col}_3(D_n)$ for even n is equal to $\text{Col}_3(H) - 1 = 2^{n-3} - 1$ for any $H \in H(n, 3)$.

REFERENCES

- [1] G.D. Birkhoff, *A determinantal formula for the number of ways of coloring a map*, Ann. Math. (2) **14** (1912), 42–46.
- [2] R.C. Read, *An introduction to chromatic polynomials*, J. Combinatorial Theory **4** (1968), 52–71.
- [3] A. Rényi, *On connected graphs*, Magyar Tud. Akad. Mat. Kutató Int. Közl **4** (1959), 385–388.
- [4] I. Tomescu, *Le nombre maximal de 3-coloration d'un graphe connexe*, Discrete Math. **1** (1972), 351–356.
- [5] I. Tomescu, *Maximum chromatic polynomial of 3-chromatic blocks*, Discrete Math. **172** (1997), 131–139.

(received 25.01.2001)

University of Bucharest, Faculty of Mathematics, Str. Academiei 14, 70109 Bucuresti, Romania
E-mail: ioan@math.math.unibuc.ro