

FRÉCHET FRAMES FOR SHIFT INVARIANT WEIGHTED SPACES¹

Suzana Simić²

Abstract. In the present paper we analyze Fréchet frame of the form $\{\varphi(\cdot - j) \mid j \in \mathbb{Z}^d\}$. With a known condition on φ , we show that the given sequence constitutes a frame for a test space isomorphic to the space of periodic smooth functions so that its dual is the multiple of the space of periodic distributions by $\hat{\varphi}$.

AMS Mathematics Subject Classification (2000): 53C25

Key words and phrases: Banach frame, Fréchet frame, pre- F -frame, F -Bessel sequence, weighted L_s^p -spaces, weighted sequence spaces.

1. Introduction

Frame theory was introduced in [9] and up to now it has developed very much in connection to wavelet theory, time frequency analysis, and sampling theory (see [1], [2], [5], [7], [11], [14], [15], [16], ...). Shift invariant spaces are generated by frames of the form $\{\varphi(x - na)\}_{n \in \mathbb{Z}^d}$ and this field is, in Banach spaces, especially L^p spaces, very much dealt with by Aldroubi, Sun and Tang [4], who studied the frames of the form $\{\varphi_i(\cdot - j) \mid j \in \mathbb{Z}^d, 1 \leq i \leq r\}$ in L^p spaces. On the other hand, in [17] and [18] authors introduced Fréchet frames and thus enabled the analysis of various test function spaces and their duals spaces of distributions.

In Section 2 we recall from [17] and [18] definitions concerning Fréchet frames. Section 3 contains preliminary results on shift invariant weighted spaces, extensions of the corresponding results given in [4]. Our main result is given in Section 4. We prove that $\{\varphi(\cdot - j) \mid j \in \mathbb{R}^d\}$ is a frame for weighted shift invariant spaces through several equivalent conditions. In the end we conclude that $\{\varphi(\cdot - j) \mid j \in \mathbb{R}^d\}$ forms a Fréchet frame for a space of test functions $X_F = \mathcal{F}^{-1}(\hat{\varphi} \cdot \mathcal{P}(-\pi, \pi))$, where \mathcal{P} is the space of periodic test functions.

2. Notation and notions

We will recall basic notions following [6], [12], [17].

We denote by $(X, \|\cdot\|)$ a Banach space, by $(X^*, \|\cdot\|^*)$ its dual space, $(\Theta, \|\cdot\|)$ is a Banach sequence space. If the coordinate functionals on Θ are continuous,

¹Supported by the Project No. 144016.

²Department of Mathematics and Informatics, Faculty of Science, University of Kragujevac, Radoja Domanovića 12, Kragujevac 34000, Serbia, e-mail: suzanasimic@kg.ac.rs

or, equivalently, if the convergence in Θ implies the convergence of the corresponding coordinates, then Θ is called a *BK-space*.

We refer to [12] for the basic definitions for frames. p -frames in shift-invariant spaces of L^p were considered in [4], while p -frames in general Banach spaces were studied in [8].

Let $\{(Y_s, |\cdot|_s)\}_{s \in \mathbb{N}_0}$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, be a family of separable Banach spaces such that

$$(1) \quad \{\mathbf{0}\} \neq \bigcap_{s \in \mathbb{N}_0} Y_s \subseteq \cdots \subseteq Y_2 \subseteq Y_1 \subseteq Y_0,$$

$$(2) \quad |\cdot|_0 \leq |\cdot|_1 \leq |\cdot|_2 \leq \cdots,$$

$$(3) \quad Y_F := \bigcap_{s \in \mathbb{N}_0} Y_s \text{ is dense in } Y_s, \quad s \in \mathbb{N}_0.$$

Then Y_F is a Fréchet space with the sequence of norms $|\cdot|_s$, $s \in \mathbb{N}_0$.

We will always assume that $\{(X_s, \|\cdot\|_s)\}_{s \in \mathbb{N}_0}$ and $\{(\Theta_s, |||\cdot|||_s)\}_{s \in \mathbb{N}_0}$ are the sequences of Banach spaces which satisfy (1)-(3). For a fixed $s \in \mathbb{N}_0$, an operator $V : \Theta_F \rightarrow X_F$ will be called s -bounded if there exists a constant $K_s > 0$ such that $\|V(\{c_i\}_{i \in \mathbb{N}})\|_s \leq K_s |||\{c_i\}_{i \in \mathbb{N}}|||_s$ for all $\{c_i\}_{i \in \mathbb{N}} \in \Theta_F$. If V is s -bounded for every $s \in \mathbb{N}_0$, then V will be called F -bounded.

Let $\{(\Theta_s, |||\cdot|||_s)\}_{s \in \mathbb{N}_0}$ be a sequence of *BK-spaces*, as well. Then a sequence $\{g_i\}_{i \in \mathbb{N}} \in (X_F^*)^{\mathbb{N}}$ is called a *pre- F -frame* for X_F with respect to Θ_F , if for every $s \in \mathbb{N}_0$, there exist constants $0 < A_s \leq B_s < +\infty$ such that

$$(4) \quad \{g_i(f)\}_{i \in \mathbb{N}} \in \Theta_F, \quad f \in X_F,$$

$$(5) \quad A_s \|f\|_s \leq |||\{g_i(f)\}_{i \in \mathbb{N}}|||_s \leq B_s \|f\|_s, \quad f \in X_F.$$

The constants B_s and A_s , $s \in \mathbb{N}_0$, are called resp. upper and lower bounds for $\{g_i\}_{i \in \mathbb{N}}$. If $A_s = B_s$, $s \in \mathbb{N}_0$, then the pre- F -frame is called *tight*. If there exists an F -bounded operator $V : \Theta_F \rightarrow X_F$ such that $V(\{g_i(f)\}_{i \in \mathbb{N}}) = f$ for all $f \in X_F$, then a pre- F -frame $\{g_i\}_{i \in \mathbb{N}}$ is called an F -frame (Fréchet frame) for X_F with respect to Θ_F and V is called an F -frame operator for $\{g_i\}_{i \in \mathbb{N}}$. When (4) holds and at least the upper inequality in (5) holds, then $\{g_i\}_{i \in \mathbb{N}}$ is called an F -Bessel sequence for X_F with respect to Θ_F with bounds B_s , $s \in \mathbb{N}_0$.

When $X = X_F = X_s$ and $\Theta = \Theta_F = \Theta_s$, then one obtains the definitions of Θ -frame, Banach frame and Θ -Bessel sequence, respectively.

If $\{g_i\}_{i \in \mathbb{N}}$ is a pre- F -frame for X_F with respect to Θ_F with lower bounds A_s and upper bounds B_s , $s \in \mathbb{N}_0$, then for every $s \in \mathbb{N}_0$ we have

$$A_s \|f\|_s \leq |||\{g_i^s(f)\}_{i \in \mathbb{N}}|||_s \leq \lambda_s B_s \|f\|_s, \quad f \in X_s,$$

where g_i^s is the continuous extension of g_i on X_s . We will consider the following operators

$$(6) \quad U_s : X_s \rightarrow \Theta_s, \quad U_s f = \{g_i^s(f)\}_{i \in \mathbb{N}}, \quad s \in \mathbb{N}_0,$$

$$(7) \quad U : X_F \rightarrow \Theta_F, \quad Uf = \{g_i(f)\}_{i \in \mathbb{N}},$$

and

$$(8) \quad U_s^{-1} : \mathcal{R}(U_s) \rightarrow X_s, \quad U^{-1} : \mathcal{R}(U) \rightarrow X_F.$$

The shift invariant spaces of the form

$$V(\varphi) = \left\{ \sum_{j \in \mathbb{Z}^d} c_j \varphi(\cdot - j) \right\},$$

where $c = \{c_j\}_{j \in \mathbb{Z}^d}$ is taken from some sequence space, are considered in [4]. φ is called generator of $V(\varphi)$. The space $V_p(\varphi)$ is the shift invariant space of the form $V_p(\varphi) = \left\{ \sum_{k \in \mathbb{Z}^d} c_k \varphi(\cdot - k) \mid c = \{c_k\}_{k \in \mathbb{Z}^d} \in \ell^p \right\}$. Let $V_0(\varphi)$ be the space of finite linear combination of integer translates of φ and $V_{0,p}(\varphi)$ be the L^p closure of $V_0(\varphi)$. Obviously, we have $V_0(\varphi) \subset V_p(\varphi) \subset V_{0,p}(\varphi)$. A function in $V_{0,p}(\varphi)$ is not necessarily generated by ℓ^p coefficients. If $V_p(\varphi)$ itself is closed, i.e. a Banach space, then $V_p(\varphi) = V_{0,p}(\varphi)$.

3. Preliminary result

Considering p -frames for shift invariant subspaces of L^p space, Aldroubi, Sun and Tung [4] proved that when a sequence of translations of a finite set of appropriate functions $\varphi_1, \dots, \varphi_r$ forms an ℓ^p -frame for the shift-invariant space $V_p(\varphi_1, \dots, \varphi_r) \subseteq L^p$, for some $p > 1$, then this sequence is also an ℓ^r -frame for $V_r(\varphi_1, \dots, \varphi_r)$ for all values of $r > 1$.

In this paper we will consider weighted L_s^p , $s \geq 0$, spaces. A function f belongs to L_s^p with weight function $\omega_s(x) = (1 + |x|)^s$, $x \in \mathbb{R}^d$, $s \geq 0$, if $\omega_s f$ belongs to L^p . Equipped with the norm $\|f\|_{L_s^p} = \|\omega_s f\|_{L^p}$, the space L_s^p is a Banach space. Let $s \geq 0$, $1 \leq p < +\infty$ and

$$\mathcal{L}_s^p := \left\{ f \mid \|f\|_{\mathcal{L}_s^p} := \left(\int_{[0,1]^d} \left(\sum_{j \in \mathbb{Z}^d} |f(x+j)|(1+|x+j|^s)^p dx \right)^{1/p} < +\infty \right\},$$

$$\mathcal{L}_s^\infty := \left\{ f \mid \|f\|_{\mathcal{L}_s^\infty} := \sup_{x \in [0,1]^d} \sum_{j \in \mathbb{Z}^d} |f(x+j)|(1+|x+j|^s) < +\infty \right\};$$

$$W_s^p := \left\{ f \mid \|f\|_{W_s^p} := \left(\sum_{j \in \mathbb{Z}^d} \sup_{x \in [0,1]^d} |f(x+j)|^p (1+|j|)^{ps} \right)^{1/p} < +\infty \right\};$$

$$\ell_s^p := \left\{ c = \{c_i\}_{i \in \mathbb{N}} \mid \|c\|_{\ell_s^p} = \left(\sum_{i \in \mathbb{N}} |c_i|^p (1+|i|)^{sp} \right)^{1/p} < +\infty \right\}.$$

Obviously, we have $W_s^p \subset W_s^q \subset \mathcal{L}_s^\infty \subset \mathcal{L}_s^q \subset \mathcal{L}_s^p \subset L_s^p$, where $1 \leq p \leq q \leq +\infty$. For $p = 1$ and $s = 0$ we also have $\mathcal{L}^1 = L^1$.

Next we recall the inequalities from [3].

Lemma 1. a) Let $f \in L_s^p$, $g \in L_s^1$ and $1 \leq p \leq +\infty$. Then

$$(9) \quad \|f * g\|_{L_s^p} \leq \|f\|_{L_s^p} \|g\|_{L_s^1}.$$

b) If $f \in L_s^p$, $1 \leq p \leq +\infty$, and $g \in W_s^1$, then $f * g \in W_s^p$ and

$$(10) \quad \|f * g\|_{W_s^p} \leq \|f\|_{L_s^p} \|g\|_{W_s^1}.$$

c) If $c \in \ell_s^p$ and $d \in \ell_s^1$, then $c * d \in \ell_s^p$ and

$$(11) \quad \|c * d\|_{\ell_s^p} \leq \|c\|_{\ell_s^p} \|d\|_{\ell_s^1}.$$

For any sequence $c = \{c_i\}_{i \in \mathbb{N}} \in \ell_s^p$ and $f \in \mathcal{L}_s^p$, $1 \leq p \leq +\infty$, define, as in [4], their semi-convolution $f *' c$ by

$$(f *' c)(x) = \sum_{j \in \mathbb{Z}^d} c_j f(x - j), \quad x \in \mathbb{R}^d.$$

Lemma 2. a) If $f \in W_s^p$, $1 \leq p \leq +\infty$, and $c \in \ell_s^1$, then the function $f *' c$ belongs to W_s^p and

$$(12) \quad \|f *' c\|_{W_s^p} \leq \|c\|_{\ell_s^1} \|f\|_{W_s^p},$$

and also if $f \in W_s^1$ and $c \in \ell_s^p$, $1 \leq p \leq +\infty$, then the function $f *' c$ belongs to W_s^p and

$$(13) \quad \|f *' c\|_{W_s^p} \leq \|c\|_{\ell_s^p} \|f\|_{W_s^1}.$$

b) If $f \in \mathcal{L}_s^p$ and $c \in \ell_s^1$, then $f *' c$ belongs to $f \in \mathcal{L}_s^p$ and

$$(14) \quad \|f *' c\|_{\mathcal{L}_s^p} \leq \|c\|_{\ell_s^1} \|f\|_{\mathcal{L}_s^p}.$$

c) $f *' \cdot$ is a continuous map from ℓ_s^p to L_s^p , and also from ℓ_s^1 to \mathcal{L}_s^p if $f \in \mathcal{L}_s^p$, $1 \leq p \leq +\infty$.

We will give the proof of the next lemma since it is differently posed in [4].

Lemma 3. Let $f \in L_s^p$ and $g \in W_s^1$, $1 \leq p \leq +\infty$, $s \geq 0$. Then the sequence $\left\{ \int_{\mathbb{R}^d} f(x)g(x-j)dx \right\}_{j \in \mathbb{Z}^d}$ belongs to ℓ_s^p and we have

$$(15) \quad \left\| \left\{ \int_{\mathbb{R}^d} f(x)\overline{g(x-j)}dx \right\}_{j \in \mathbb{Z}^d} \right\|_{\ell_s^p} \leq \|f\|_{L_s^p} \|g\|_{W_s^1}.$$

Proof. Using inequality (11) for fixed $x \in \mathbb{R}^d$, we obtain

$$\begin{aligned}
& \left\| \left\{ \int_{\mathbb{R}^d} f(x) \overline{g(x-j)} dx \right\}_{j \in \mathbb{Z}^d} \right\|_{\ell_s^p} = \left(\sum_{j \in \mathbb{Z}^d} \left| \int_{\mathbb{R}^d} f(x) \overline{g(x-j)} dx \right|^p (1+|j|)^{sp} \right)^{1/p} \\
& \leq \left(\sum_{j \in \mathbb{Z}^d} \left(\int_{\mathbb{R}^d} |f(x)| |g(x-j)| dx \right)^p (1+|j|)^{sp} \right)^{1/p} \\
& = \left(\sum_{j \in \mathbb{Z}^d} \left(\int_{[0,1]^d} \sum_{k \in \mathbb{Z}^d} |f(x+k)| |g(x+k-j)| dx \right)^p (1+|j|)^{sp} \right)^{1/p} \\
& \leq \left(\sum_{j \in \mathbb{Z}^d} \int_{[0,1]^d} \left(\sum_{k \in \mathbb{Z}^d} |f(x+k)| |g(x+k-j)| \right)^p (1+|j|)^{sp} dx \right)^{1/p} \\
& = \left(\sum_{j \in \mathbb{Z}^d} \int_{[0,1]^d} \left(\sum_{k \in \mathbb{Z}^d} |f(x+k)| |g(x+k-j)| (1+|k|)^s \right)^p dx \right)^{1/p} \\
& \leq \left(\int_{[0,1]^d} \sum_{j \in \mathbb{Z}^d} \left(\sum_{k \in \mathbb{Z}^d} |f(x+k)| |g(x+k-j)| (1+|k|)^s \right)^p dx \right)^{1/p} \\
& \leq \left(\int_{[0,1]^d} \sum_{j \in \mathbb{Z}^d} |f(x+j)|^p (1+|j|)^{sp} \left(\sum_{k \in \mathbb{Z}^d} |g(x-k)| (1+|k|)^s \right)^p dx \right)^{1/p} \\
& \leq \|f\|_{L_s^p} \left(\sup_{x \in [0,1]^d} \left(\sum_{k \in \mathbb{Z}^d} |g(x-k)| (1+|k|)^s \right)^p \right)^{1/p} \leq \|f\|_{L_s^p} \|g\|_{W_s^1}.
\end{aligned}$$

□

4. Main result

Our main result is related to Theorem 1 in [4].

Let $\varphi \in \mathcal{L}_s^p$, $1 \leq p \leq \infty$. We consider shift-invariant spaces of the form

$$(16) \quad V_s^p(\varphi) = \left\{ \sum_{j \in \mathbb{Z}^d} c_j \varphi(\cdot - j) \mid c \in \ell_s^p \right\}.$$

Note, if $s = 0$, then we have the space $V^p(\varphi)$ considered in [4].

Theorem 1. *Let $\varphi \in \bigcap_{s \geq 0} W_s^1$. Then the following statements are equivalent to each other.*

- i) $V_s^p(\varphi)$ is closed in L_s^p for all $s \geq 0$ and for all $1 \leq p \leq +\infty$.
- ii) For all $s \geq 0$ and $1 \leq p \leq +\infty$, the family $\{\varphi(\cdot - j) \mid j \in \mathbb{Z}^d\}$ is a p -frame for $V_s^p(\varphi)$, i.e. there exist positive constants A_s, B_s (depending on φ and

s) such that

$$(17) \quad A_s \|f\|_{L_s^p} \leq \left\| \left\{ \int_{\mathbb{R}^d} f(x) \overline{\varphi(x-j)} dx \right\}_{j \in \mathbb{Z}^d} \right\|_{\ell_s^p} \leq B_s \|f\|_{L_s^p}, \quad \forall f \in V_s^p(\varphi).$$

iii) There exist positive constants C_1 and C_2 such that

$$(18) \quad 0 < C_1 \leq \sum_{j \in \mathbb{Z}^d} |\widehat{\varphi}(x+j)|^2 \leq C_2 < +\infty, \quad \text{a.e. } x \in \mathbb{R}^d.$$

iv) There exist positive constants K_s^1 and K_s^2 (depending on φ and s) such that for all $1 \leq p \leq +\infty$ we have

$$(19) \quad K_s^1 \|f\|_{L_s^p} \leq \inf_{c \in M} \|c\|_{\ell_s^p} \leq K_s^2 \|f\|_{L_s^p}, \quad \forall f \in V_s^p(\varphi), \quad s \geq 0,$$

where

$$(20) \quad M = \left\{ c = \{c_k\}_{k \in \mathbb{Z}^d} \in \ell_s^p \mid f(\cdot) = \sum_{k \in \mathbb{Z}^d} c_k \varphi(\cdot - k) \right\}.$$

v) There exists $\psi \in \bigcap_{s \geq 0} W_s^1$ such that

$$(21) \quad f = \sum_{j \in \mathbb{Z}^d} \langle f, \psi(\cdot - j) \rangle \varphi(\cdot - j) = \sum_{j \in \mathbb{Z}^d} \langle f, \varphi(\cdot - j) \rangle \psi(\cdot - j), \quad \forall f \in V_s^p(\varphi).$$

Proof.

$v) \Rightarrow iw)$

Let $f = \sum_{j \in \mathbb{Z}^d} \langle f, \psi(\cdot - j) \rangle \varphi(\cdot - j)$ and let M be given by (20). Using (15) we have

$$\inf_{c \in M} \|c\|_{\ell_s^p} \leq \left\| \left\{ \int_{\mathbb{R}^d} f(x) \overline{\psi(x-j)} dx \right\}_{j \in \mathbb{Z}^d} \right\|_{\ell_s^p} \leq \|f\|_{L_s^p} \|\psi\|_{W_s^1}.$$

For $K_s^2 = \|\psi\|_{W_s^1}$ we have the right-hand side of the inequality.

Using (13), we have

$$\|f\|_{L_s^p} \leq \|f\|_{W_s^p} = \|\varphi *' c\|_{W_s^p} \leq \|\varphi\|_{W_s^1} \|c\|_{\ell_s^p},$$

and for $K_s^1 = \frac{1}{\|\varphi\|_{W_s^1}}$ we prove the left-hand side of inequality (19).

Assertions $v) \Rightarrow ii)$, $ii) \Leftrightarrow iv)$, and $iv) \Rightarrow i)$ are simple and their proofs will be omitted.

$iii) \Rightarrow iv)$

We have already seen that for $\varphi \in W_s^1$ and $c \in \ell_s^p$, $1 \leq p \leq +\infty$, the inequality

$$\|\varphi *' c\|_{W_s^p} \leq \|c\|_{\ell_s^p} \|\varphi\|_{W_s^1},$$

holds. With $\|\varphi *' c\|_{L_s^p} \leq \|\varphi *' c\|_{W_s^p}$ for all $1 \leq p \leq +\infty$, and $K_s^1 = \|\varphi\|_{W_s^1}^{-1}$, we have that the left-hand side of the inequality (17).

The family $\{\varphi(\cdot - k) \mid k \in \mathbb{Z}^d\}$ with the condition (18) is a Riesz basis of $V^2(\varphi)$ (see [3]), so there exists a unique function $\psi \in V^2(\varphi)$ such that $\{\psi(\cdot - k) \mid k \in \mathbb{Z}^d\}$ is also a Riesz basis for $V^2(\varphi)$, and such that it satisfies the biorthogonality relations

$$\langle \psi(x), \varphi(x) \rangle = 1, \quad \langle \psi(x), \varphi(x - k) \rangle = 0, \quad k \neq 0.$$

Theorem 2.3 in [3] says that if $\varphi \in W_s^1$ and the family $\{\varphi(\cdot - k) \mid k \in \mathbb{Z}^d\}$ is a Riesz basis for $V^2(\varphi)$, then the dual generator ψ is in W_s^1 . Since we have that $\varphi \in W_s^1$ for all $s \geq 0$, then we have that $\psi \in \bigcap_{s \geq 0} W_s^1$. Since

$$(\varphi *' c)(x) = \sum_{k \in \mathbb{Z}^d} c_k \varphi(x - k) \in V_s^p(\varphi),$$

then $c_k, k \in \mathbb{Z}^d$, can be expressed in the form

$$c_k = \int_{\mathbb{R}^d} (\varphi *' c)(x) \overline{\psi(x - k)} dx.$$

For $1 \leq p \leq +\infty$ (with usual changes for $p = \infty$), we have

$$\begin{aligned} |c_k(1 + |k|)^s|^p &= \left| \int_{\mathbb{R}^d} (\varphi *' c)(x) \overline{\psi(x - k)} (1 + |k|)^s dx \right|^p \\ &\leq \left(\int_{[0,1]^d} \sum_{j \in \mathbb{Z}^d} |\varphi *' c|(x + j) |\psi(x + j - k)| (1 + |k|)^s dx \right)^p \\ &\leq \int_{[0,1]^d} \left(\sum_{j \in \mathbb{Z}^d} |\varphi *' c|(|\psi(x + j - k)| (1 + |k|)^s) \right)^p dx. \end{aligned}$$

We sum over $k \in \mathbb{Z}^d$ and obtain

$$\begin{aligned} &\sum_{k \in \mathbb{Z}^d} |c_k|^p (1 + |k|)^{sp} \\ &\leq \int_{[0,1]^d} \sum_{j \in \mathbb{Z}^d} \left(\sum_{k \in \mathbb{Z}^d} |\varphi *' c|(x + j) |\psi(x + j - k)| (1 + |k|)^s \right)^p dx \\ &\leq \int_{[0,1]^d} \sum_{k \in \mathbb{Z}^d} |\varphi *' c|^p(x + k) (1 + |k|)^{sp} \left(\sum_{k \in \mathbb{Z}^d} |\psi(x + k)| (1 + |k|)^s \right)^p dx \\ &\leq \|\psi\|_{W_s^1}^p \|\varphi *' c\|_{L_s^p}^p. \end{aligned}$$

It follows

$$\|c\|_{\ell_s^p} \leq \|\psi\|_{W_s^1} \|\varphi *' c\|_{L_s^p}.$$

For the lower bound in the inequality (19) one may choose $K_s^2 = \|\psi\|_{W_s^1}$. Finally,

$$\|c\|_{\ell_s^p} \leq K_s^2 \|f\|_{L_s^p}.$$

i) \Rightarrow iii)

Since $V_s^p(\varphi)$ is closed in L_s^p for all $1 \leq p \leq +\infty$, $s \geq 0$, then for $p = 2$ and $s = 1$ we have the standard assumption on the generator φ , i.e. there exist two constants C_1 and C_2 such that

$$0 < C_1 \leq \sum_{j \in \mathbb{Z}^d} |\widehat{\varphi}(x+j)|^2 \leq C_2 < +\infty, \quad \text{a.e. } x \in \mathbb{R}^d.$$

□

Corollary 1. *Let $\varphi \in \bigcap_{s \geq 0} W_s^1$. Then $V_s^p(\varphi) \subset V_s^q(\varphi)$, for all $1 \leq p \leq q \leq +\infty$ and $s \geq 0$.*

Proof. Let $f(x) = \sum_{k \in \mathbb{Z}^d} c_k \varphi(x-k)$, for some $c = \{c_k\}_{k \in \mathbb{Z}^d} \in \ell_s^p$, $1 \leq p \leq +\infty$. Since $\ell_s^p \subset \ell_s^q$, $1 \leq p \leq q \leq +\infty$, Theorem 1 implies the inequalities

$$\|f\|_{L_s^q} \leq B_s \|c\|_{\ell_s^q} \leq B'_s \|c\|_{\ell_s^p} \leq \|f\|_{L_s^p}, \quad \forall s \geq 0, 1 \leq p \leq q \leq +\infty.$$

□

Remark 1. *From the inequalities (19) and (17) we can conclude that ℓ_s^p and V_s^p are isomorphic Banach spaces for all $s \geq 0$ and $1 \leq p \leq +\infty$, and for $f \in V_s^p(\varphi)$ we have the equivalence between $\inf_{c \in M} \{\|c\|_{\ell_s^p}\}$ and the L_s^p -norm of f .*

As a consequence of Theorem 1 and from [3, Theorem 1], and since $\ell_{s_1}^p \subset \ell_{s_2}^p$, for $0 \leq s_2 \leq s_1$, we have the following corollary.

Corollary 2. *Let $\varphi \in \bigcap_{s \geq 0} W_s^1$. Then $V_{s_1}^p(\varphi) \subset V_{s_2}^p(\varphi)$ for $0 \leq s_2 \leq s_1$ and every $1 \leq p \leq +\infty$.*

We construct Fréchet spaces $X_{F,p}$, $p \geq 1$, as the intersection of translator invariant spaces $V_s^p(\varphi)$, $s \in \mathbb{N}$. Note that, for $1 \leq p \leq +\infty$,

$$\{\mathbf{0}\} \neq \bigcap_{s \in \mathbb{N}_0} V_s^p(\varphi) \subseteq \cdots \subseteq V_2^p(\varphi) \subseteq V_1^p(\varphi) \subseteq V_0^p(\varphi) = V^p(\varphi).$$

Also, we have that $X_{F,p} = \bigcap_{s \in \mathbb{N}_0} V_s^p(\varphi)$ is dense in $V_s^p(\varphi)$ for all $s \in \mathbb{N}_0$. The corresponding sequence space $Q_{F,p}$, $p \geq 1$, is the intersection of the weighted sequence space ℓ_s^p , $s \in \mathbb{N}_0$. Note that $\bigcap_{s \in \mathbb{N}_0} \ell_s^p$, for every $p \geq 1$, is actually the space of rapidly decreasing sequences s . We proved that if $\varphi \in W_s^1$, then a sequence $\{\varphi(\cdot - k) \mid k \in \mathbb{Z}^d\}$ is a p -frame for $V_s^p(\varphi)$ as well as $\{\varphi(\cdot - k) \mid k \in \mathbb{Z}^d\}$

is an r -frame for $V_s^r(\varphi)$, for all $1 \leq r \leq +\infty$. So we have that the definition of $X_{F,p}$ does not depend on $p \geq 1$, so $\{\varphi(\cdot - k) \mid k \in \mathbb{Z}^d\}$ is a pre- F -frame for $X_{F,p}$ as well as that $\{\varphi(\cdot - k) \mid k \in \mathbb{Z}^d\}$ is a pre- F -frame for $X_{F,r}$, for all $1 \leq r \leq +\infty$.

Since the corresponding function space for s is the space of rapidly increasing functions

$$\mathcal{S} = \{f \mid \|f\|_m = \sup_{n \leq m} (1 + |x|^2)^{m/2} |f^{(n)}(x)| < +\infty\},$$

and its dual is \mathcal{S}' - the space of slowly decreasing distributions, we obtain that dual space of Fréchet space $X_F = X_{F,p}$, for any p , is isomorphic to (a complemented subspace of) the space \mathcal{S}' .

Denote by $\mathcal{P}(-\pi, \pi)$ the space of smooth 2π -periodic functions with the family of norms $|\theta|_k = \sup\{|\theta^{(k)}(t)|; t \in (-\pi, \pi)\}$, $k \in \mathbb{N}_0$. It is a Fréchet space and its dual is the space of 2π -periodic tempered distributions. Denote by \mathcal{F} and \mathcal{F}^{-1} the Fourier transformation and its inverse transformation, respectively. We have

Theorem 2. Let $\varphi \in \bigcap_{s \geq 0} W_s^1$ and $X_F = \bigcap_{s \in \mathbb{N}_0} V_s^p(\varphi)$ for some $1 \leq p \leq +\infty$.

Then

$$X_F = \mathcal{F}^{-1}(\widehat{\varphi} \cdot \mathcal{P}(-\pi, \pi)),$$

in the topological sense.

Proof. For $f \in X_F$ we have $f = \sum_{j \in \mathbb{Z}^d} c_j \varphi(\cdot - j)$, for some sequence $c = \{c_j\}_{j \in \mathbb{Z}^d} \in s$. Then

$$\widehat{f} = \sum_{j \in \mathbb{Z}^d} \widehat{c_j \varphi(\cdot - j)} = \left(\sum_{j \in \mathbb{Z}^d} c_j e^{ij \cdot} \right) \widehat{\varphi}.$$

This implies the assertion. \square

References

- [1] Aldroubi A., Oblique projections in atomic spaces. Proc. Am. Math. Soc., 124 (1996), 2051–2060.
- [2] Aldroubi A., Exact iterative reconstruction algorithm for multivariate irregularly sampled functions in spline-like space: the L^p theory. Proc. Am. Math. Soc., 126 (1998), 2677–2686.
- [3] Aldroubi A., Gröchenig K., Non-uniform sampling and reconstruction in shift-invariant spaces. SIAM Review, 43 (2001), 585–620.
- [4] Aldroubi A., Sun Q., Tang W., p -frames and shift invariant subspaces of L^p . J. Fourier Anal. Appl., 7 (2001), 1–21.

- [5] Benedetto J. J., Frame decomposition, sampling and uncertainty principle inequalities. In: *Wavelets: Mathematics and Applications* (Benedetto J. J., Frazier M. W.) pp. 247–304. Boca Raton, FL: CRC Press, 1994.
- [6] Benedetto J. J., Frazier M. W., *Wavelets: Mathematics and Applications*. Boca Raton, FL: CRC Press, 1994.
- [7] Christensen O., Heil C., Perturbations of Banach frames and atomic decompositions. *Math. Nach.*, 185 (1997), 33–47.
- [8] Christensen O., Stoeva D. T., p -frames in separable Banach spaces . *Adv. Comp. Math.* 18 (2-4) (2003), 117-126.
- [9] Duffin R. J., Schaeffer A. C., A class of nonharmonic Fourier series. *Trans. Am. Math. Soc.*, 72 (1952), 341-3666.
- [10] Feichtinger G. H., Gewichtsfunktionen auf lokalkompakten Gruppen. *Sitzungsber.d.osterr. Akad.Wiss.*, Vol.188 (1979), 451–471.
- [11] Gröchenig K., Describing functions: atomic decompositions versus frames. *Monatsh. Math.*, 112 (1991), 1–42.
- [12] Gröchenig K., *Foundations of Time-Frequency Analysis*. Boston: Birkhäuser, 2001.
- [13] Heil C., *A Basis Theory Primer* . Manuscript, 1997.
- [14] Heil C. E., Walnut D. F., Continuous and discrete wavelet transforms. *SIAM Review*, 31 (1989), 628–666.
- [15] Mallat S., *A Wavelet Tour of Signal Processing*. New York: Academic Press, 1998.
- [16] Meyer Y., *Ondelettes*. Paris: Hermann, Editeurs des Science et des Arts, 1990.
- [17] Pilipović S., Stoeva D. T., Series expansions in Fréchet spaces and construction of Fréchet frames. preprint, 2009.
- [18] Pilipović S., Stoeva D. T., Teofanov N.: Frames for Fréchet spaces . *Bull. Cl. Sci. Math. Nat. Sci. Math.*, 32 (2007), 69-84.
- [19] Strichartz R. S., *A Guide to Distribution Theory and Fourier Transforms*. New Jersey: World Scientific, 1994.

Received by the editors June 5, 2009