

A NOTE ON MINORS DETERMINED BY CLONES OF SEMILATTICES

Erkko Lehtonen¹

Abstract. The \mathcal{C} -minor partial orders determined by the clones generated by a semilattice operation (and possibly the constant operations corresponding to its identity or zero elements) are shown to satisfy the descending chain condition.

AMS Mathematics Subject Classification (2010): 08A40, 06A06

Key words and phrases: clones, semilattices, \mathcal{C} -minors, descending chain condition

1. Introduction

This paper is a study of substitution instances of functions of several arguments when the inner functions are taken from a prescribed set of functions. Such an idea has been studied by several authors. Henno [6] generalized Green's relations to Menger algebras (essentially, abstract clones) and described Green's relations on the set of all operations on A for each set A . Harrison [5] considered two Boolean functions to be equivalent if they are substitution instances of each other with respect to the general linear group $\text{GL}(n, \mathbb{F}_2)$ or the affine linear group $\text{AGL}(n, \mathbb{F}_2)$, where \mathbb{F}_2 denotes the two-element field. In [15, 16], a Boolean function f is defined to be a minor of another Boolean function g , if and only if f can be obtained from g by substituting for each variable of g a variable, a negated variable, or one of the constants 0 or 1. Further variants of the notion of minor can be found in [1, 3, 4, 13, 17].

These ideas are unified and generalized by the notions of \mathcal{C} -minor and \mathcal{C} -equivalence, which first appeared in print in [8]. More precisely, let A be a nonempty set, and let $f: A^n \rightarrow A$ and $g: A^m \rightarrow A$ be operations on A . Let \mathcal{C} be a set of operations on A . We say that f is a \mathcal{C} -minor of g , if $f = g(h_1, \dots, h_m)$ for some $h_1, \dots, h_m \in \mathcal{C}$, and we say that f and g are \mathcal{C} -equivalent if f and g are \mathcal{C} -minors of each other. If \mathcal{C} is a clone, then the \mathcal{C} -minor relation is a preorder and it induces a partial order on the \mathcal{C} -equivalence classes. For background and basic results on \mathcal{C} -minors and \mathcal{C} -equivalences, see [8, 10, 11, 12].

In this paper, we study the \mathcal{C} -minors and \mathcal{C} -equivalences induced by the clones generated by semilattice operations (and possibly some constants). Our

¹Computer Science and Communications Research Unit, University of Luxembourg, 6, rue Richard Coudenhove-Kalergi, L-1359 Luxembourg, Luxembourg,
email: erkko.lehtonen@uni.lu

main result (Theorem 3.1) asserts that if $(A; \wedge)$ is a semilattice, then for the clone $\mathcal{C} := \langle \wedge \rangle$ generated by \wedge , the induced \mathcal{C} -minor partial order satisfies the descending chain condition. Furthermore, if $(A; \wedge)$ has an identity element 1 and a zero element 0, then this property is enjoyed by all clones \mathcal{C} such that $\langle \wedge \rangle \subseteq \mathcal{C} \subseteq \langle \wedge, 0, 1 \rangle$. These results find an application in [9], in which the clones of Boolean functions are classified according to certain order-theoretical properties of their induced \mathcal{C} -minor partial orders.

2. Clones, \mathcal{C} -minors and \mathcal{C} -decompositions

2.1. Operations and clones

Throughout this paper, for an integer $n \geq 1$, we denote $[n] := \{1, \dots, n\}$. Let A be a fixed nonempty base set. An *operation* on A is a map $f: A^n \rightarrow A$ for some integer $n \geq 1$, called the *arity* of f . We denote the set of all n -ary operations on A by $\mathcal{O}_A^{(n)}$, and we denote by $\mathcal{O}_A := \bigcup_{n \geq 1} \mathcal{O}_A^{(n)}$ the set of all operations on A . The i -th n -ary *projection* ($1 \leq i \leq n$) is the operation $(a_1, \dots, a_n) \mapsto a_i$, and it is denoted by $x_i^{(n)}$, or simply by x_i when the arity is clear from the context.

If $f \in \mathcal{O}_A^{(n)}$ and $g_1, \dots, g_n \in \mathcal{O}_A^{(m)}$, then the *composition* of f with g_1, \dots, g_n , denoted $f(g_1, \dots, g_n)$ is the m -ary operation defined by

$$f(g_1, \dots, g_n)(\mathbf{a}) = f(g_1(\mathbf{a}), \dots, g_n(\mathbf{a}))$$

for all $\mathbf{a} \in A^m$.

Let $\mathcal{C} \subseteq \mathcal{O}_A$. The n -ary *part of \mathcal{C}* is the set $\mathcal{C}^{(n)} := \mathcal{C} \cap \mathcal{O}_A^{(n)}$ of n -ary members of \mathcal{C} . A *clone* on A is a subset $\mathcal{C} \subseteq \mathcal{O}_A$ that contains all projections and is closed under composition, i.e., $f(g_1, \dots, g_n) \in \mathcal{C}$ whenever $f, g_1, \dots, g_n \in \mathcal{C}$ and the composition is defined.

The clones on A constitute a complete lattice under inclusion. Therefore, for each set $F \subseteq \mathcal{O}_A$ of operations there exists a smallest clone that contains F , which will be denoted by $\langle F \rangle$ and called the *clone generated by F* . See [2, 7, 14] for general background on clones.

2.2. \mathcal{C} -minors

Let $\mathcal{C} \subseteq \mathcal{O}_A$, and let $f, g \in \mathcal{O}_A$. We say that f is a \mathcal{C} -minor of g , if $f = g(h_1, \dots, h_m)$ for some $h_1, \dots, h_m \in \mathcal{C}$, and we denote this fact by $f \leq_{\mathcal{C}} g$. We say that f and g are \mathcal{C} -equivalent, denoted $f \equiv_{\mathcal{C}} g$, if f and g are \mathcal{C} -minors of each other.

The \mathcal{C} -minor relation $\leq_{\mathcal{C}}$ is a preorder (i.e., a reflexive and transitive relation) on \mathcal{O}_A if and only if \mathcal{C} is a clone. If \mathcal{C} is a clone, then the \mathcal{C} -equivalence relation $\equiv_{\mathcal{C}}$ is an equivalence relation on \mathcal{O}_A , and, as for preorders, $\leq_{\mathcal{C}}$ induces a partial order $\preceq_{\mathcal{C}}$ on the quotient $\mathcal{O}_A / \equiv_{\mathcal{C}}$. It follows from the definition of \mathcal{C} -minor, that if \mathcal{C} and \mathcal{K} are clones such that $\mathcal{C} \subseteq \mathcal{K}$, then $\leq_{\mathcal{C}} \subseteq \leq_{\mathcal{K}}$ and $\equiv_{\mathcal{C}} \subseteq \equiv_{\mathcal{K}}$. For further background and properties of \mathcal{C} -minor relations, see [8, 10, 11, 12].

2.3. \mathcal{C} -decompositions

Let \mathcal{C} be a clone on A , and let $f \in \mathcal{O}_A^{(n)}$. If $f = g(\phi_1, \dots, \phi_m)$ for some $g \in \mathcal{O}_A^{(m)}$ and $\phi_1, \dots, \phi_m \in \mathcal{C}$, then we say that the $(m+1)$ -tuple $(g, \phi_1, \dots, \phi_m)$ is a \mathcal{C} -decomposition of f . We often avoid referring explicitly to the tuple and we simply say that $f = g(\phi_1, \dots, \phi_m)$ is a \mathcal{C} -decomposition. Clearly, there always exists a \mathcal{C} -decomposition of every operation f for every clone \mathcal{C} , because $f = f(x_1^{(n)}, \dots, x_n^{(n)})$ and projections are members of every clone. A \mathcal{C} -decomposition of a nonconstant function f is *minimal* if the arity m of g is the smallest possible among all \mathcal{C} -decompositions of f . This smallest possible m is called the \mathcal{C} -degree of f , denoted $\deg_{\mathcal{C}} f$. We agree that the \mathcal{C} -degree of every constant function is 0.

Lemma 2.1. *If $f \leq_{\mathcal{C}} g$, then $\deg_{\mathcal{C}} f \leq \deg_{\mathcal{C}} g$.*

Proof. Let $\deg_{\mathcal{C}} g = m$, and let $g = h(\gamma_1, \dots, \gamma_m)$ be a minimal \mathcal{C} -decomposition of g . Since $f \leq_{\mathcal{C}} g$, there exist $\phi_1, \dots, \phi_n \in \mathcal{C}$ such that $f = g(\phi_1, \dots, \phi_n)$. Then

$$f = h(\gamma_1, \dots, \gamma_m)(\phi_1, \dots, \phi_n) = h(\gamma_1(\phi_1, \dots, \phi_n), \dots, \gamma_m(\phi_1, \dots, \phi_n)),$$

and since $\gamma_i(\phi_1, \dots, \phi_n) \in \mathcal{C}$ for $1 \leq i \leq m$, we have that $(h, \gamma_1(\phi_1, \dots, \phi_n), \dots, \gamma_m(\phi_1, \dots, \phi_n))$ is a \mathcal{C} -decomposition of f , not necessarily minimal, so $\deg_{\mathcal{C}} f \leq m$. \square

An immediate consequence of Lemma 2.1 is that \mathcal{C} -equivalent functions have the same \mathcal{C} -degree.

Let (ϕ_1, \dots, ϕ_m) be an m -tuple ($m \geq 2$) of n -ary operations on A . If there is an $i \in \{1, 2, \dots, m\}$ and $g: A^{m-1} \rightarrow A$ such that

$$\phi_i = g(\phi_1, \dots, \phi_{i-1}, \phi_{i+1}, \dots, \phi_m),$$

then we say that the m -tuple (ϕ_1, \dots, ϕ_m) is *functionally dependent*. Otherwise we say that (ϕ_1, \dots, ϕ_m) is *functionally independent*. We often omit the m -tuple notation and simply say that ϕ_1, \dots, ϕ_m are functionally dependent or independent.

Remark 2.2. Every m -tuple containing a constant function is functionally dependent. Also if $f_i = f_j$ for some $i \neq j$, then f_1, \dots, f_n are functionally dependent.

Lemma 2.3. *If $(g, \phi_1, \dots, \phi_m)$ is a minimal \mathcal{C} -decomposition of f , then ϕ_1, \dots, ϕ_m are functionally independent.*

Proof. Suppose, on the contrary, that ϕ_1, \dots, ϕ_m are functionally dependent. Then there is an i and an $h: A^{m-1} \rightarrow A$ such that

$$\phi_i = h(\phi_1, \dots, \phi_{i-1}, \phi_{i+1}, \dots, \phi_m).$$

Then

$$\begin{aligned} f &= g(\phi_1, \dots, \phi_{i-1}, h(\phi_1, \dots, \phi_{i-1}, \phi_{i+1}, \dots, \phi_m), \phi_{i+1}, \dots, \phi_m) \\ &= g(x_1^{(m-1)}, \dots, x_{i-1}^{(m-1)}, h, x_i^{(m-1)}, \dots, x_{m-1}^{(m-1)})(\phi_1, \dots, \phi_{i-1}, \phi_{i+1}, \dots, \phi_m), \end{aligned}$$

which shows that $(g(x_1, \dots, x_{i-1}, h, x_i, \dots, x_{m-1}), \phi_1, \dots, \phi_{i-1}, \phi_{i+1}, \dots, \phi_m)$ is a \mathcal{C} -decomposition of f , contradicting the minimality of $(g, \phi_1, \dots, \phi_m)$. \square

3. \mathcal{C} -minors determined by clones of semilattices

In this section we will prove our main result, namely Theorem 3.1. It will find an application in [9] where the clones of Boolean functions are classified according to certain order-theoretical properties that their induced \mathcal{C} -minor partial orders enjoy.

A binary operation \wedge on A is called a *semilattice operation*, if for all $x, y, z \in A$, the following identities hold:

$$x \wedge (y \wedge z) = (x \wedge y) \wedge z, \quad x \wedge y = y \wedge x, \quad x \wedge x = x,$$

i.e., \wedge is associative, commutative and idempotent.

A partial order $(P; \leq)$ is said to satisfy the *descending chain condition*, or it is called *well-founded*, if it contains no infinite descending chains, i.e., given any sequence of elements of P

$$\dots \leq a_3 \leq a_2 \leq a_1,$$

there exists a positive integer n such that

$$a_n = a_{n+1} = a_{n+2} = \dots.$$

Theorem 3.1. *Let \mathcal{S} be the clone generated by a semilattice operation \wedge on A . Then the \mathcal{S} -minor partial order $\preceq_{\mathcal{S}}$ satisfies the descending chain condition.*

Proof. Let $(\phi_1, \dots, \phi_m) \in (\mathcal{S}^{(n)})^m$. Then, for $1 \leq j \leq m$, ϕ_j is of the form

$$(1) \quad \phi_j = \bigwedge_{i \in \Phi_j} x_i^{(n)}$$

for some $\emptyset \neq \Phi_j \subseteq [n]$. For $1 \leq i \leq n$, denote

$$(2) \quad X_i := \{j \in [m] : i \in \Phi_j\},$$

and let $X(\phi_1, \dots, \phi_m) := \{X_1, \dots, X_n\} \subseteq \mathcal{P}([m])$. It follows from the definitions of Φ_j and X_i that

$$(3) \quad j \in X_i \iff i \in \Phi_j.$$

Correspondingly, for any $\emptyset \neq E \subseteq \mathcal{P}([m])$, denote $\Psi_E := (\psi_1, \dots, \psi_m)$, where $\psi_j \in \mathcal{S}^{(|E|)}$ is given by

$$\psi_j = \bigwedge_{s \in E} x_{\sigma_E(s)},$$

where $\sigma_E: E \rightarrow [|E|]$ is any fixed bijection.

Let $(g, \phi_1, \dots, \phi_m)$ be an \mathcal{S} -decomposition of $f: A^n \rightarrow A$. Then each ϕ_j is of the form (1) for some $\emptyset \neq \Phi_j \subseteq [n]$. Let $E := X(\phi_1, \dots, \phi_m)$, $(\psi_1, \dots, \psi_m) := \Psi_E$, and let $f' = g(\psi_1, \dots, \psi_m)$. We will show that $f \equiv_{\mathcal{S}} f'$.

As in (2), for $1 \leq i \leq n$, let $X_i = \{j \in [m] : i \in \Phi_j\}$. Let $\pi: [n] \rightarrow [|E|]$ be defined as $\pi(i) := \sigma_E(X_i)$. Then

$$\begin{aligned} f(x_{\pi(1)}, \dots, x_{\pi(n)}) &= g(\phi_1, \dots, \phi_m)(x_{\pi(1)}, \dots, x_{\pi(n)}) \\ &= g(\phi_1(x_{\pi(1)}, \dots, x_{\pi(n)}), \dots, \phi_m(x_{\pi(1)}, \dots, x_{\pi(n)})) = g(\psi_1, \dots, \psi_m) = f', \end{aligned}$$

where the second last equality holds because for $1 \leq j \leq m$,

$$\phi_j(x_{\pi(1)}, \dots, x_{\pi(n)}) = \bigwedge_{i \in \Phi_j} x_{\pi(i)} = \bigwedge_{i \in \Phi_j} x_{\sigma(X_i)} = \bigwedge_{j \in S \in E} x_{\sigma(S)} = \psi_j.$$

Since all projections are members of \mathcal{S} , we have that $f' \leq_C f$. On the other hand, for $1 \leq j \leq |E|$, let $\Xi_j := \{i \in [n] : X_i = \sigma_E^{-1}(j)\}$, and let

$$\xi_j := \bigwedge_{i \in \Xi_j} x_i.$$

It is easy to see that $\Xi_j \neq \emptyset$; hence $\xi_j \in \mathcal{S}$. Then

$$\begin{aligned} f'(\xi_1, \dots, \xi_{|E|}) &= g(\psi_1, \dots, \psi_m)(\xi_1, \dots, \xi_{|E|}) \\ &= g(\psi_1(\xi_1, \dots, \xi_{|E|}), \dots, \psi_m(\xi_1, \dots, \xi_{|E|})) = g(\phi_1, \dots, \phi_m) = f, \end{aligned}$$

where the second last equality holds because for $j = 1, \dots, m$,

$$\begin{aligned} \psi_j(\xi_1, \dots, \xi_{|E|}) &= \left(\bigwedge_{j \in S \in E} x_{\sigma_E(S)} \right) (\xi_1, \dots, \xi_{|E|}) = \bigwedge_{j \in S \in E} \xi_{\sigma_E(S)} \\ &= \bigwedge_{j \in S \in E} \left(\bigwedge_{i \in \Xi_{\sigma_E(S)}} x_i \right) = \bigwedge_{j \in S \in E} \left(\bigwedge_{\substack{i \in [n] \\ X_i = S}} x_i \right) = \bigwedge_{i \in \Phi_j} x_i = \phi_j. \end{aligned}$$

Here, the third last equality holds, because $\Xi_{\sigma_E(S)} = \{i \in [n] : X_i = S\}$, and the second last equality holds by (3) and the associativity, commutativity and idempotency of \wedge . Since $\xi_j \in \mathcal{S}$, we have that $f \leq_C f'$. We conclude that $f \equiv_C f'$, as desired.

Claim. If $f_1 = g(\phi_1, \dots, \phi_m)$ and $f_2 = g(\varphi_1, \dots, \varphi_m)$ are \mathcal{S} -decompositions and $X(\phi_1, \dots, \phi_m) = X(\varphi_1, \dots, \varphi_m)$, then $f_1 \equiv_{\mathcal{S}} f_2$.

Proof of the claim. Let $(\psi_1, \dots, \psi_m) := \Psi_{X(\phi_1, \dots, \phi_m)}$ ($= \Psi_{X(\varphi_1, \dots, \varphi_m)}$), and let $f' = g(\psi_1, \dots, \psi_m)$. It follows from what was shown above that $f_1 \equiv_{\mathcal{S}} f' \equiv_{\mathcal{S}} f_2$. The claim follows by the transitivity of $\equiv_{\mathcal{S}}$. \diamond

To finish the proof that $\preceq_{\mathcal{S}}$ satisfies the descending chain condition, assume that $f_1 \prec_{\mathcal{S}} f_2$, $f_2 = g(\phi_1, \dots, \phi_m)$ is a minimal \mathcal{S} -decomposition, and $f_1 = f_2(h_1, \dots, h_n)$ for some $h_1, \dots, h_n \in \mathcal{S}$. For $i = 1, \dots, m$, denote $\phi'_i = \phi_i(h_1, \dots, h_n)$, so that $f_1 = g(\phi'_1, \dots, \phi'_m)$. By Lemma 2.1, either $\deg_{\mathcal{S}} f_1 <$

$\deg_{\mathcal{S}} f_2$, or $\deg_{\mathcal{S}} f_1 = \deg_{\mathcal{S}} f_2$ and $X(\phi_1, \dots, \phi_m) \neq X(\phi'_1, \dots, \phi'_m)$. Since \mathcal{S} -degrees are nonnegative integers and $\mathcal{P}([m])$ is a finite set, there are only a finite number of $\equiv_{\mathcal{S}}$ -classes preceding the $\equiv_{\mathcal{S}}$ -class of f_2 in the \mathcal{S} -minor partial order $\preceq_{\mathcal{S}}$. This completes the proof of the theorem. \square

Corollary 3.2. *Assume that a semilattice $(A; \wedge)$ has identity and zero elements 1 and 0, respectively. Let \mathcal{C} be a clone on A such that $\langle \wedge \rangle \subseteq \mathcal{C} \subseteq \langle \wedge, 0, 1 \rangle$. Then the \mathcal{C} -minor partial order $\preceq_{\mathcal{C}}$ satisfies the descending chain condition.*

Proof. The proof of Theorem 3.1 in fact shows that $\preceq_{\mathcal{C}}$ satisfies the descending chain condition. For, in this case $\mathcal{C} \setminus \mathcal{S}$ contains only constant operations. Remark 2.2 and Lemma 2.3 guarantee that $f = g(h_1, \dots, h_m)$ is a minimal \mathcal{S} -decomposition if and only if it is a minimal \mathcal{C} -decomposition, and since $\mathcal{S} \subseteq \mathcal{C}$, \mathcal{S} -equivalence implies \mathcal{C} -equivalence. \square

References

- [1] Couceiro, M., Pouzet, M., On a quasi-ordering on Boolean functions. *Theoret. Comput. Sci.* 396 (2008), 71–87.
- [2] Denecke, K. Wismath, S.L., *Universal Algebra and Applications in Theoretical Computer Science*. Chapman & Hall/CRC, Boca Raton, 2002.
- [3] Ekin, O., Foldes, S., Hammer, P.L., Hellerstein, L., Equational characterizations of Boolean function classes. *Discrete Math.* 211 (2000), 27–51.
- [4] Feigelson, A., Hellerstein, L., The forbidden projections of unate functions. *Discrete Appl. Math.* 77 (1997), 221–236.
- [5] Harrison, M.A., On the classification of Boolean functions by the general linear and affine groups. *J. Soc. Indust. Appl. Math.* 12(2) (1964), 285–299.
- [6] Henno, J., Green’s equivalences in Menger systems. *Tartu Riikl. Üli. Toimetised* 277 (1971), 37–46 [Russian].
- [7] Lau, D., *Function Algebras on Finite Sets – A Basic Course on Many-Valued Logic and Clone Theory*. Springer-Verlag, Berlin, Heidelberg, 2006.
- [8] Lehtonen, E., Descending chains and antichains of the unary, linear, and monotone subfunction relations. *Order* 23 (2006), 129–142.
- [9] Lehtonen, E., Nešetřil, J., Minors of Boolean functions with respect to clique functions and hypergraph homomorphisms. *Eur. J. Combin.* 31 (2010), 1981–1995.
- [10] Lehtonen, E., Szendrei, Á., Equivalence of operations with respect to discriminator clones. *Discrete Math.* 309 (2009), 673–685.
- [11] Lehtonen, E., Szendrei, Á., Clones with finitely many relative \mathcal{R} -classes. *Algebra Univers.*, to appear, arXiv:0905.1611.
- [12] Lehtonen, E., Szendrei, Á., The submaximal clones on the three-element set with finitely many relative \mathcal{R} -classes. *Discuss. Math. Gen. Algebra Appl.*, to appear, arXiv:0905.1614.
- [13] Pippenger, N., Galois theory for minors of finite functions. *Discrete Math.* 254 (2002), 405–419.

- [14] Szendrei, Á., Clones in Universal Algebra. Séminaire de mathématiques supérieures, vol. 99. Les Presses de l'Université de Montréal, Montréal, 1986.
- [15] Wang, C., Boolean minors. *Discrete Math.* 141 (1995), 237–258.
- [16] Wang, C., Williams, A.C., The threshold order of a Boolean function. *Discrete Appl. Math.* 31 (1991), 51–69.
- [17] Zverovich, I.E., Characterizations of closed classes of Boolean functions in terms of forbidden subfunctions and Post classes. *Discrete Appl. Math.* 149 (2005), 200–218.

Received by the editors March 31, 2010. Revised August 13, 2010.