

## New decomposition forms of bioperation-continuity

Carlos Carpintero<sup>1,2</sup>, Neelamegarajan Rajesh<sup>3</sup>,  
Ennis Rosas<sup>4</sup> and Jorge Vielma<sup>5</sup>

**Abstract.** In this paper, we introduce some new types of sets via bioperation and obtain new decomposition forms of bioperation-continuity using these sets and finally using the notions of a bioperation some well known concepts of continuity are generalized.

*AMS Mathematics Subject Classification* (2010): 54A05; 54A10; 54C05; 54D10

*Key words and phrases:* topological spaces;  $(\gamma, \gamma')$ -open set

### 1. Introduction

Generalized open sets play a very important role in General Topology and they are now the research topics of many topologist worldwide. Indeed a significant theme in General Topology and Real analysis concerns the various modified forms of continuity, separation axioms etc. By utilizing generalized open sets. Kasahara [5] introduced the concept of an operation on topological spaces and the notion of  $\gamma$ -open sets, the collection of all  $\gamma$ -open sets is denoted by  $\tau_\gamma$ . Ogata and Maki [11] introduced the notion of  $\tau_{\gamma \vee \gamma'}$  which is the collection of all  $\gamma \vee \gamma'$ -open sets in a topological space  $(X, \tau)$  and Umehara et al. [13] introduced the notion of  $\tau_{(\gamma, \gamma')}$  which is the collection of all  $(\gamma, \gamma')$ -open sets in a topological space  $(X, \tau)$  that generalized the notions of  $\gamma \vee \gamma'$ -open sets in a topological space  $(X, \tau)$ . In this paper, using the bioperation  $(\gamma, \gamma')$ , we introduce new types of sets and find the relationships between them and obtain a new forms of decomposition of bioperation-continuity. Finally we can see that these new concepts of continuity using these bioperations, generalizes well-known concepts of continuity.

### 2. Preliminaries

The closure and the interior of a subset  $A$  of  $(X, \tau)$  are denoted by  $\text{Cl}(A)$  and  $\text{Int}(A)$ , respectively.

---

<sup>1</sup>Departamento de Ciencias Básicas, Corporación Universitaria del Caribe-CECAR, Sincelejo, Colombia, e-mail: [carpintero.carlos@gmail.com](mailto:carpintero.carlos@gmail.com)

<sup>2</sup>Corresponding author

<sup>3</sup>Department of Mathematics, Rajah Serfoji Govt. College, Thanjavur-613005, Tamilnadu, India, e-mail: [nrajesh\\_topology@yahoo.co.in](mailto:nrajesh_topology@yahoo.co.in)

<sup>4</sup>Universidad de la Costa, Department of Natural Sciences and Exact, Barranquilla, Colombia, e-mail: [ennisrafael@gmail.com](mailto:ennisrafael@gmail.com)

<sup>5</sup>Escuela Superior Politécnica del Litoral, Facultad de Ciencias Naturales y Matemáticas, Campus Gustavo Galindo, Km 30.5, Vía perimetral, P.O Box 09-01-5863, Guayaquil, Ecuador, e-mail: [jvielma@espol.edu.ec](mailto:jvielma@espol.edu.ec)

**Definition 2.1.** [5] Let  $(X, \tau)$  be a topological space. An operation  $\gamma$  on the topology  $\tau$  is a function from  $\tau$  on to power set  $\mathcal{P}(X)$  of  $X$  such that  $V \subset V^\gamma$  for each  $V \in \tau$ , where  $V^\gamma$  denotes the value of  $\tau$  at  $V$ . It is denoted by  $\gamma : \tau \rightarrow \mathcal{P}(X)$ .

**Definition 2.2.** [11] A topological space  $(X, \tau)$  equipped with two operations, say,  $\gamma$  and  $\gamma'$  defined on  $\tau$  is called a bioperation-topological space, it is denoted by  $(X, \tau, \gamma, \gamma')$ .

**Definition 2.3.** [11] A subset  $A$  of a topological space  $(X, \tau)$  is said to be  $\gamma \vee \gamma'$ -open set if for each  $x \in A$  there exists an open neighborhood  $U$  of  $x$  such that  $U^\gamma \cup U^{\gamma'} \subset A$ . The complement of  $\gamma \vee \gamma'$ -open set is called  $\gamma \vee \gamma'$ -closed.  $\tau_{\gamma \vee \gamma'}$  denotes set of all  $\gamma \vee \gamma'$ -open sets in  $(X, \tau)$ .

**Definition 2.4.** [13] A subset  $A$  of a topological space  $(X, \tau)$  is said to be  $(\gamma, \gamma')$ -open set if for each  $x \in A$  there exist open neighborhoods  $U$  and  $V$  of  $x$  such that  $U^\gamma \cup V^{\gamma'} \subset A$ . The complement of  $(\gamma, \gamma')$ -open set is called  $(\gamma, \gamma')$ -closed.  $\tau_{(\gamma, \gamma')}$  denotes set of all  $(\gamma, \gamma')$ -open sets in  $(X, \tau)$ .

**Remark 2.5.** Observe that from Definitions 2.3 and 2.4, each  $\gamma \vee \gamma'$ -open set is a  $(\gamma, \gamma')$ -open set, but the converse is not necessarily true as we can see in the following example.

**Example 2.6.** Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ . We define the operations  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  as follows:  $\gamma(\{a\}) = \{a\}$ ,  $\gamma(\{b\}) = \{b, c\}$ ,  $\gamma(\{a, b\}) = \{a, b\}$   
 $\gamma'(\{a\}) = \{a, b\}$ ,  $\gamma'(\{b\}) = \{b\}$ ,  $\gamma'(\{a, b\}) = X$   
 Observe that:  $\tau_{\gamma \vee \gamma'} = \{\emptyset, X\}$  and  $\tau_{(\gamma, \gamma')} = \{\emptyset, X, \{a, b\}\}$

**Example 2.7.** Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ . We define the operators  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  by  $\gamma(A) = \text{Cl}(A)$  and  $\gamma'(A) = \text{Int}(\text{Cl}(A))$  for all  $A \in \tau$ . Then  $\tau_{(\gamma, \gamma')} = \tau_{\gamma \vee \gamma'} = \{\emptyset, X\}$ .

**Definition 2.8.** [13] For a subset  $A$  of  $(X, \tau)$ ,  $\text{Cl}_{(\gamma, \gamma')}(A)$  denotes the intersection of all  $(\gamma, \gamma')$ -closed sets containing  $A$ , that is,  $\text{Cl}_{(\gamma, \gamma')}(A) = \cap \{F : A \subset F, X \setminus F \in \tau_{(\gamma, \gamma')}\}$ .

**Definition 2.9.** Let  $A$  be any subset of  $X$ . The  $\text{Int}_{(\gamma, \gamma')}(A)$  is defined as  $\text{Int}_{(\gamma, \gamma')}(A) = \cup \{U : U \text{ is a } (\gamma, \gamma')\text{-open set and } U \subset A\}$ .

**Definition 2.10.** Let  $(X, \tau)$  be a topological space and  $A$  be a subset of  $X$  and  $\gamma$  and  $\gamma'$  be operations on  $\tau$ . Then  $A$  is said to be

1.  $(\gamma, \gamma')$ - $\alpha$ -open if  $A \subset \text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A)))$
2.  $(\gamma, \gamma')$ -preopen [3] if  $A \subset \text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A))$
3.  $(\gamma, \gamma')$ -semiopen [10] if  $A \subset \text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A))$
4.  $(\gamma, \gamma')$ -semipreopen (or  $(\gamma, \gamma')$ - $\beta$ -open) if  $A \subset \text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A)))$

5.  $(\gamma, \gamma')$ -regular open [9] if  $A = \text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A))$ .

**Remark 2.11.** The union of all  $(\gamma, \gamma')$ -semipreopen sets contained in  $A$  is called the  $(\gamma, \gamma')$ -semipreinterior of  $A$  and denoted by  $\text{spInt}_{(\gamma, \gamma')}(A)$ . The complement of a  $(\gamma, \gamma')$ -semipreopen set is called a  $(\gamma, \gamma')$ -semipreclosed set. It is clear that  $\text{spInt}_{(\gamma, \gamma')}(A) = A \cap \text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A)))$ .

**Remark 2.12.** Observe that if in Definition 2.10, the operations  $\gamma$  and  $\gamma'$  are the identity operations, we obtain well-known concepts studied in general topology such as:  $\alpha$ -open set [7], [12], preopen set [8], semiopen set [2], semipreopen set [2].

**Definition 2.13.** Let  $(X, \tau)$  and  $(Y, \sigma)$  be two topological spaces and let  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  be operations on  $\tau$ . A mapping  $f : (X, \tau) \rightarrow (Y, \sigma)$  is said to be  $(\gamma, \gamma')$ -continuous (resp.  $(\gamma, \gamma')$ - $\alpha$ -continuous,  $(\gamma, \gamma')$ -precontinuous,  $(\gamma, \gamma')$ -semicontinuous,  $(\gamma, \gamma')$ -semiprecontinuous) if for each  $x \in X$  and each open set  $V$  of  $Y$  containing  $f(x)$  there exists a  $(\gamma, \gamma')$ -open set  $U$  containing  $x$  (resp.  $(\gamma, \gamma')$ - $\alpha$ -open set,  $(\gamma, \gamma')$ -preopen set,  $(\gamma, \gamma')$ -semiopen set,  $(\gamma, \gamma')$ -semipreopen set) such that  $f(U) \subset V$ .

### 3. Some subsets in topological spaces

Through this section, let  $(X, \tau)$  and  $(Y, \sigma)$  be two topological spaces, and let  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  be operations on  $\tau$ .

**Definition 3.1.** A subset  $A$  of a topological space  $(X, \tau)$  with the operations  $\gamma, \gamma'$  is called:

1.  $\alpha^*_{(\gamma, \gamma')}$ -set if  $\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A))) = \text{Int}_{(\gamma, \gamma')}(A)$ ,
2.  $t_{(\gamma, \gamma')}$ -set if  $\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A)) = \text{Int}_{(\gamma, \gamma')}(A)$ ,
3.  $s_{(\gamma, \gamma')}$ -set if  $\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A)) = \text{Int}_{(\gamma, \gamma')}(A)$ ,
4.  $\beta^*_{(\gamma, \gamma')}$ -set if  $\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A))) = \text{Int}_{(\gamma, \gamma')}(A)$ .

**Remark 3.2.** Observe that if in Definition 3.1, the operations  $\gamma$  and  $\gamma'$  are the identity operations, we obtain well-known concepts studied in general topology such as  $\beta$ -set [1],  $t$ -set and  $\alpha^*$ -set.

**Example 3.3.** Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}, \{a, b\}\}$ . We define the operations  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  as follows

$$A^\gamma = \begin{cases} A & \text{if } A = \{a\} \text{ or } \{c\}, \\ A \cup \{a, c\} & \text{if } A \neq \{a\} \text{ and } \{c\} \end{cases}$$

and  $A^{\gamma'} = \text{int}(\text{Cl}(A))$ .

1.  $\tau_{(\gamma, \gamma')} = \{\emptyset, X, \{c\}\}$
2.  $\alpha^*_{(\gamma, \gamma')} \text{-set} = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ .

3.  $t_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ .
4.  $s_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ .
5.  $\beta_{(\gamma, \gamma')}^*$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ .

**Proposition 3.4.** *The following are equivalent for a subset  $A$  of a space  $(X, \tau)$  with the operations  $\gamma, \gamma'$*

1.  $A$  is a  $\alpha_{(\gamma, \gamma')}^*$ -set,
2.  $A$  is a  $(\gamma, \gamma')$ -semipreclosed set,
3.  $\text{Int}_{(\gamma, \gamma')}(A)$  is a  $(\gamma, \gamma')$ -regular open set.

*Proof.* Straightforward. □

**Proposition 3.5.** *Let  $A$  be a subset of a space  $(X, \tau)$  with the operations  $\gamma, \gamma'$*

1.  $A$   $(\gamma, \gamma')$ -semiopen set  $A$  is a  $t_{(\gamma, \gamma')}$ -set if and only if it is an  $\alpha_{(\gamma, \gamma')}^*$ -set.
2.  $A$  is  $(\gamma, \gamma')$ - $\alpha$ -open and  $\alpha_{(\gamma, \gamma')}^*$ -set if and only if it is  $(\gamma, \gamma')$ -regular open.

*Proof.* 1. Let  $A$  be a  $(\gamma, \gamma')$ -semiopen and  $A$  an  $\alpha_{(\gamma, \gamma')}^*$ -set. Since  $A$  is  $(\gamma, \gamma')$ -semiopen,

$$\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A)) = \text{Cl}_{(\gamma, \gamma')}(A)$$

and

$$\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A)) = \text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A))) = \text{Int}_{(\gamma, \gamma')}(A).$$

Therefore,  $A$  is a  $t_{(\gamma, \gamma')}$ -set.

2. Let  $A$  be a  $(\gamma, \gamma')$ - $\alpha$ -open set and an  $\alpha_{(\gamma, \gamma')}^*$ -set. Then  $\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A)) = A$  and hence  $\text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(A)) = \text{Int}_{(\gamma, \gamma')}(\text{Cl}_{(\gamma, \gamma')}(\text{Int}_{(\gamma, \gamma')}(A))) = A$ . The converse is obvious. □

**Definition 3.6.** *A subset  $A$  of a topological space  $(X, \tau)$  with the operations  $\gamma, \gamma'$  is called a*

1.  $C_{(\gamma, \gamma')}$ -set if  $A = U \cap V$ , where  $U \in \tau_{(\gamma, \gamma')}$  and  $V$  is an  $\alpha_{(\gamma, \gamma')}^*$ -set,
2.  $B_{(\gamma, \gamma')}$ -set if  $A = U \cap V$ , where  $U \in \tau_{\gamma, \gamma'}$  and  $V$  is a  $t_{(\gamma, \gamma')}$ -set,
3.  $S_{(\gamma, \gamma')}$ -set if  $A = U \cap V$ , where  $U \in \tau_{(\gamma, \gamma')}$  and  $V$  is a  $s_{(\gamma, \gamma')}$ -set,
4.  $\beta_{(\gamma, \gamma')}$ -set if  $A = U \cap V$ , where  $U \in \tau_{(\gamma, \gamma')}$  and  $V$  is a  $\beta_{(\gamma, \gamma')}^*$ -set,
5.  $\beta^{**}$ -open set if  $\text{sp Int}_{(\gamma, \gamma')}(A) = \text{Int}_{(\gamma, \gamma')}(A)$ .

**Example 3.7.** *Observe that in Example 2.6,*

1.  $\tau_{(\gamma, \gamma')} = \{\emptyset, X, \{a, b\}\}$
2.  $\alpha_{(\gamma, \gamma')}^*$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}$ .
3.  $t_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{c\}\}$ .

4.  $s_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}$ .
5.  $\beta^*_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{c\}\}$ .
6.  $C_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}\}$ .
7.  $B_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{c\}, \{a, b\}\}$ .
8.  $S_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}\}$ .
9.  $\beta_{(\gamma, \gamma')}$ -set =  $\{\emptyset, X, \{c\}, \{a, b\}\}$ .
10.  $\beta^{**}$ -open set =  $\{\emptyset, X, \{a, b\}\}$ .

**Proposition 3.8.** *Let  $(X, \tau)$  be a topological space with the operations  $\gamma, \gamma'$  and  $A$  a subset of  $X$ . Then the following hold:*

1. *If  $A$  is a  $t_{(\gamma, \gamma')}$ -set, then  $A$  is an  $\alpha^*_{(\gamma, \gamma')}$ -set,*
2. *If  $A$  is a  $s_{(\gamma, \gamma')}$ -set, then  $A$  is an  $\alpha^*_{(\gamma, \gamma')}$ -set,*
3. *If  $A$  is a  $\beta^*_{(\gamma, \gamma')}$ -set, then  $A$  is both a  $t_{(\gamma, \gamma')}$ -set and a  $s_{(\gamma, \gamma')}$ -set.*
4.  *$t_{(\gamma, \gamma')}$ -set and  $s_{(\gamma, \gamma')}$ -set are independent notions.*

*Proof.* (1). Let  $A$  be a  $t_{(\gamma, \gamma')}$ -set. Then  $\tau_{(\gamma, \gamma')} \text{-Int}(\tau_{(\gamma, \gamma')} \text{-Cl}(A)) = \tau_{(\gamma, \gamma')} \text{-Int}(A) \supset \tau_{(\gamma, \gamma')} \text{-Int}(\tau_{(\gamma, \gamma')} \text{-Cl}(\tau_{\gamma} \text{-Int}(A))) \supset \tau_{(\gamma, \gamma')} \text{-Int}(A)$  and hence  $\tau_{\gamma \vee \gamma'} \text{-Int}(\tau_{\gamma \vee \gamma'} \text{-Cl}(\tau_{\gamma \vee \gamma'} \text{-Int}(A))) = \tau_{\gamma \vee \gamma'} \text{-Int}(A)$ . Therefore,  $A$  is an  $\alpha^*_{\gamma \vee \gamma'}$ -set. (2) and (3) are proved in a similar form as (1). (4) The following examples shows that the notions of a  $t_{(\gamma, \gamma')}$ -set and a  $s_{(\gamma, \gamma')}$ -set are independent. □

**Remark 3.9.** *The converse of the statement in Proposition 3.8 are not true as seen in the following examples.*

**Example 3.10.** *Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}, \{a, b\}\}$ . We define the operations  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  as follows*

$$A^\gamma = A^{\gamma'} = \begin{cases} A & \text{if } A = \{a\} \text{ or } \{c\}, \\ A \cup \{a, c\} & \text{if } A \neq \{a\} \text{ and } \{c\}. \end{cases}$$

*Then  $\tau_{(\gamma, \gamma')} = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}\}$ . If we take  $A = \{a\}$ , then  $A$  is an  $\alpha^*_{(\gamma, \gamma')}$ -set and a  $t_{(\gamma, \gamma')}$ -set, but it is neither a  $s_{(\gamma, \gamma')}$ -set and nor a  $\beta^*_{(\gamma, \gamma')}$ -set.*

**Example 3.11.** *If in Example 2.6 and Example 3.7, we take  $A = \{b\}$ , then it is an  $\alpha^*_{(\gamma, \gamma')}$ -set and a  $s_{(\gamma, \gamma')}$ -set, but it is neither a  $t_{(\gamma, \gamma')}$ -set and nor a  $\beta^*_{(\gamma, \gamma')}$ -set.*

**Proposition 3.12.** *Let  $(X, \tau)$  be a topological space with the operations  $\gamma, \gamma'$  and  $A$  a subset of  $X$ . Then the following hold:*

1. *If  $A$  is an  $\alpha^*_{(\gamma, \gamma')}$ -set, then it is a  $C_{(\gamma, \gamma')}$ -set,*

2. If  $A$  is a  $t_{(\gamma, \gamma')}$ -set, then it is a  $B_{(\gamma, \gamma')}$ -set,
3. If  $A$  is a  $s_{(\gamma, \gamma')}$ -set, then it is a  $S_{(\gamma, \gamma')}$ -set,
4. If  $A$  is a  $\beta_{(\gamma, \gamma')}^*$ -set, then it is a  $\beta_{(\gamma, \gamma')}$ -set.

*Proof.* 1. Let  $A$  be an  $\alpha_{(\gamma, \gamma')}^*$ -set. If we take  $U = X \in \tau_{(\gamma, \gamma')}$ , then  $A = U \cap A$  and hence  $A$  is a  $C_{(\gamma, \gamma')}$ -set.

2. Let  $A$  be a  $t_{(\gamma, \gamma')}$ -set. If we take  $U = X \in \tau_{(\gamma, \gamma')}$ , then  $A = U \cap A$  and hence  $A$  is a  $B_{(\gamma, \gamma')}$ -set.

3. Let  $A$  be a  $s_{(\gamma, \gamma')}$ -set. If we take  $U = X \in \tau_{(\gamma, \gamma')}$ , then  $A = U \cap A$  and hence  $A$  is a  $S_{(\gamma, \gamma')}$ -set.

4. Let  $A$  be a  $\beta_{(\gamma, \gamma')}^*$ -set. If we take  $U = X \in \tau_{(\gamma, \gamma')}$ , then  $A = U \cap A$  and hence  $A$  is a  $\beta_{(\gamma, \gamma')}$ -set. □

**Remark 3.13.** *The converse of the statements in Proposition 3.12 are not true. In Example 3.7,  $\{a, b\}$  is a  $C_{(\gamma, \gamma')}$ -set (resp.  $B_{(\gamma, \gamma')}$ -set,  $S_{(\gamma, \gamma')}$ -set,  $\beta_{(\gamma, \gamma')}$ -set), but it is not an  $\alpha_{(\gamma, \gamma')}^*$ -set (resp.  $t_{(\gamma, \gamma')}$ -set,  $s_{(\gamma, \gamma')}$ -set,  $\beta_{(\gamma, \gamma')}^*$ -set).*

**Proposition 3.14.** *Let  $(X, \tau)$  be a topological space with the operations  $\gamma, \gamma'$ .*


1. Every  $B_{(\gamma, \gamma')}$ -set is a  $C_{(\gamma, \gamma')}$ -set,
2. Every  $S_{(\gamma, \gamma')}$ -set is a  $C_{(\gamma, \gamma')}$ -set,
3. Every  $\beta_{(\gamma, \gamma')}$ -set is both a  $B_{(\gamma, \gamma')}$ -set and a  $S_{(\gamma, \gamma')}$ -set.

*Proof.* The proof follows from Proposition 3.12 and Definition 3.6. □

**Remark 3.15.** *The converse of the statements in Proposition 3.14 are not true and  $B_{(\gamma, \gamma')}$ -set and  $S_{(\gamma, \gamma')}$ -set are independent notions. In Example 3.10,  $\{a, b\}$  is a  $B_{(\gamma, \gamma')}$ -set but it is not a  $S_{(\gamma, \gamma')}$ -set and not a  $\beta_{(\gamma, \gamma')}$ -set. In Example 2.7,  $\{b\}$  is a  $C_{(\gamma, \gamma')}$ -set and a  $S_{(\gamma, \gamma')}$ -set but it is neither a  $B_{(\gamma, \gamma')}$ -set nor a  $\beta_{(\gamma, \gamma')}$ -set.*

**Remark 3.16.** *Observe that if  $(X, \tau)$  is a topological space with the operations  $\gamma, \gamma'$ . Then  $\beta^{**}$ -open and  $\beta^*$ -set are independent notions. See Example 3.7.*

**Remark 3.17.** *We have the following implication diagram.*





**Proposition 4.3.** *Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function and  $\gamma, \gamma'$  operations on  $\tau$ . Then*

1. *Every  $B_{(\gamma, \gamma')}$ -continuous function is  $C_{(\gamma, \gamma')}$ -continuous.*
2. *Every  $S_{(\gamma, \gamma')}$ -continuous function is  $C_{(\gamma, \gamma')}$ -continuous.*
3. *Every  $\beta_{(\gamma, \gamma')}$ -continuous is both  $B_{(\gamma, \gamma')}$ -continuous and  $S_{(\gamma, \gamma')}$ -continuous.*

*Proof.* The proof follows from Proposition 3.14. □

**Theorem 4.4.** *Let  $(X, \tau)$  and  $(Y, \sigma)$  be two topological spaces and let  $\gamma, \gamma'$  operations on  $\tau$ . For a function  $f : (X, \tau) \rightarrow (Y, \sigma)$ , the following properties are equivalent:*

1.  *$f$  is  $(\gamma, \gamma')$ -continuous.*
2.  *$f$  is  $(\gamma, \gamma')$ - $\alpha$ -continuous and  $C_{(\gamma, \gamma')}$ -continuous.*
3.  *$f$  is  $(\gamma, \gamma')$ -precontinuous and  $B_{(\gamma, \gamma')}$ -continuous.*
4.  *$f$  is  $(\gamma, \gamma')$ -semicontinuous and  $S_{(\gamma, \gamma')}$ -continuous.*
5.  *$f$  is  $(\gamma, \gamma')$ -semiprecontinuous and  $\beta_{(\gamma, \gamma')}$ -continuous.*

*Proof.* The proof follows from Theorem 3.18. □

**Remark 4.5.** *The notions of  $(\gamma, \gamma')$ - $\alpha$ -continuity,  $C_{(\gamma, \gamma')}$ -continuity,  $(\gamma, \gamma')$ -continuity,  $B_{(\gamma, \gamma')}$ -continuity,  $(\gamma, \gamma')$ -semicontinuity,  $S_{(\gamma, \gamma')}$ -continuity,  $(\gamma, \gamma')$ -semiprecontinuity and  $(\gamma, \gamma')$ -continuity are independent of each other as seen in the following examples.*

**Example 4.6.** *Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}, \{a, b\}\}$  and  $\sigma = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}\}$ . We define the operators  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  by*

$$A^\gamma = A^{\gamma'} = \begin{cases} A & \text{if } A = \{a\}, \\ A \cup \{a, c\} & \text{if } A \neq \{a\}. \end{cases}$$

*Then  $\tau_{(\gamma, \gamma')} = \{\emptyset, X, \{a\}, \{a, c\}\}$ . Define a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  as  $f(a) = f(b) = a, f(c) = c$ . Then  $f$  is  $C_{(\gamma, \gamma')}$ -continuous (resp.  $B_{(\gamma, \gamma')}$ -continuous,  $(\gamma, \gamma')$ -semicontinuous and  $(\gamma, \gamma')$ -semiprecontinuous), but it is not  $(\gamma, \gamma')$ - $\alpha$ -continuous (resp.  $\gamma \vee \gamma'$ -precontinuous,  $S_{\gamma \vee \gamma'}$ -continuous and  $\beta_{(\gamma, \gamma')}$ -continuous).*

**Example 4.7.** *Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{a, b\}\}$  and  $\sigma = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ . We define the operators  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  by  $\gamma(A) = \text{Cl}(A)$  and  $\gamma'(A) = \text{Int}(\text{Cl}(A))$  for all  $A \in \tau$ . Then  $\tau_{(\gamma, \gamma')} = \{\emptyset, X\}$ . Define a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  as  $f(a) = f(c) = a, f(b) = b$ . Then  $f$  is both  $S_{(\gamma, \gamma')}$ -continuous and  $(\gamma, \gamma')$ -precontinuous, but it is neither  $(\gamma, \gamma')$ -semicontinuous nor  $B_{(\gamma, \gamma')}$ -continuous.*



**Example 4.8.** Let  $X = \{a, b, c, d\}$  and  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, \{a, b, d\}\}$  and  $\sigma = \{\emptyset, Y, \{a\}, \{b\}, \{a, b\}\}$ . We define the operations  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  by

$$A^\gamma = A^{\gamma'} = \begin{cases} \text{Int}(\text{Cl}(A)) & \text{if } A = \{a\}, \\ \text{Cl}(A) & \text{if } A \neq \{a\}. \end{cases}$$

Then  $\tau_{(\gamma, \gamma')} = \{\emptyset, \{a\}, \{c\}, \{a, c\}, \{a, b, d\}, X\}$ . Define a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  as  $f(a) = f(c) = a$ ,  $f(b) = f(d) = b$ . Then  $f$  is  $\beta_{(\gamma, \gamma')}$ -continuous, but it is not  $(\gamma, \gamma')$ -semiprecontinuous.

**Example 4.9.** Let  $X = \{a, b, c\}$  and  $\tau = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}, \{b, c\}\}$  and  $\sigma = \{\emptyset, X, \{a\}\}$ . We define the operations  $\gamma, \gamma' : \tau \rightarrow \mathcal{P}(X)$  by

$$A^\gamma = A^{\gamma'} = \begin{cases} \text{Int}(\text{Cl}(A)) & \text{if } A = \{a\}, \\ X & \text{if } A \neq \{a\}. \end{cases}$$

Then  $\tau_{(\gamma, \gamma')} = \{\emptyset, \{a\}, X\}$ . Define a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  as  $f(a) = f(c) = a$ ,  $f(b) = b$ . Then  $f$  is  $(\gamma, \gamma')$ - $\alpha$ -continuous but it is not  $C_{(\gamma, \gamma')}$ -continuous.

## Acknowledgement

Thanks to the referees for the valuable observations made that improve the content of the article.

## References

- [1] ABD EL-MONSEF, M. E., EL-DEEB, S. N., AND MAHMOUD, R. A.  $\beta$ -open sets and  $\beta$ -continuous mapping. *Bull. Fac. Sci. Assiut Univ. A* 12, 1 (1983), 77–90.
- [2] ANDRIJEVIĆ, D. Semipreopen sets. *Mat. Vesnik* 38, 1 (1986), 24–32.
- [3] CARPINTERO, C., RAJESH, N., AND ROSAS, E. On a class of  $(\gamma, \gamma')$ -preopen sets in a topological space. *Fasc. Math.*, 46 (2011), 25–35.
- [4] CARPINTERO, C., RAJESH, N., AND ROSAS, E. On decomposition of bioperation-continuity. *C. R. Acad. Bulgare Sci.* 73, 4 (2020), 443–450.
- [5] KASAHARA, S. Operation-compact spaces. *Math. Japon.* 24, 1 (1979/80), 97–105.
- [6] LEVINE, N. Semi-open sets and semi-continuity in topological spaces. *Amer. Math. Monthly* 70 (1963), 36–41.
- [7] MASHBOUR, A., ABD-EL-MONSEF, M., AND EL-DEEP, S.  $\alpha$ -continuous and  $\alpha$ -open mapping. 213–218.
- [8] MASHHOUR, A. S., ABD EL-MONSEF, M. E., AND EL-DEEP, S. N. On precontinuous and weak precontinuous mappings. *Proc. Math. Phys. Soc. Egypt*, 53 (1982), 47–53 (1983).
- [9] NIRMALA, R., AND RAJESH, N. Bioperation-regular open sets. *Aryabhata J. Math. Infor.* 8(1) (2016), 53–62.

- [10] NIRMALA, R., AND RAJESH, N. Generalization of semiopen sets via bioperation.
- [11] OGATA, H., AND MAKI, H. Bioperations on topological spaces. *Math. Japon.* 38, 5 (1993), 981–985.
- [12] REILLY, I., AND M. VAMANAMUNTY, M. On  $\alpha$ -continuity in topological spaces. *J. Indian Acad. Math.* 18 (1996), 89–99.
- [13] UMEHARA, J.-I., MAKI, H., AND NOIRI, T. Bioperations on topological spaces and some separation axioms. *Mem. Fac. Sci. Kochi Univ. Ser. A Math.* 13 (1992), 45–59.

*Received by the editors June 13, 2021*

*First published online January 27, 2022*