

ON ALMOST ω_1 - n -SIMPLY PRESENTED ABELIAN p -GROUPS

Peter Danchev

ABSTRACT. We define and investigate the class of almost ω_1 - n -simply presented p -torsion abelian groups, which class properly contains the subclasses of almost n -simply presented groups and ω_1 - n -simply presented groups, respectively. The obtained results generalize those obtained by us in Korean J. Math. (2014) and J. Algebra Appl. (2015).

1. Introduction and background

In what follows, all considered groups are additive p -primary abelian, where p is a fixed prime integer. Also, let $n \geq 0$ be a nonnegative integer. As usual, for any ordinal α , the symbol $p^\alpha G$ denotes the p^α -th power subgroup of G , that is, the subgroup of G consisting of all elements with heights $\geq \alpha$. In the case when $\alpha = \omega$, $p^\omega G$ is just called the *first Ulm subgroup* of G . We say that the group G is *separable* if $p^\omega G = \{0\}$.

All unexplained notions and notations are mainly standard and follow essentially those from [5] and [6]. For the specific terminology, we provide the reader with the following:

- [9] A reduced group G is said to be *almost totally projective* if it has a collection \mathcal{C} consisting of nice subgroups of G satisfying the following three conditions:

(1) $\{0\} \in \mathcal{C}$;

(2) \mathcal{C} is closed with respect to ascending unions, i.e., if $H_i \in \mathcal{C}$ with $H_i \subseteq H_j$ whenever $i \leq j$ ($i, j \in I$), then $\bigcup_{i \in I} H_i \in \mathcal{C}$;

(3) If K is a countable subgroup of G , then there is $L \in \mathcal{C}$ (that is, a nice subgroup L of G) such that $K \subseteq L$ and L is countable.

If G is separable, it is known as an *almost direct sum of cyclic groups*.

- A group G is said to be *almost simply presented* if it is the direct sum of a divisible group and an almost totally projective group.

The last concept can be generalized for any natural number n as follows:

2010 *Mathematics Subject Classification*: Primary 20K10; Secondary 20K15.

Key words and phrases: Abelian groups, almost n -simply presented groups, almost ω_1 - n -simply presented groups.

Communicated by Žarko Mijajlović.

- [2] A group G is said to be *almost n -simply presented* if there is $L \leq G[p^n]$ with G/L almost simply presented.

If L is nice in G , the latter group will be called *nice almost n -simply presented*.

- [3] A group G is said to be *almost $p^{\omega+n}$ -projective* if there is $P \leq G[p^n]$ with G/P an almost direct sum of cyclic groups.

- [3] A group G is said to be *almost ω_1 - $p^{\omega+n}$ -projective* if there is a countable subgroup $C \leq G$ such that G/C is almost $p^{\omega+n}$ -projective.

These two concepts can be extended in the sense of [4] like this:

DEFINITION 1.1. A group G is called *almost ω_1 - n -simply presented* if there exists a countable subgroup K of G such that G/K is almost n -simply presented. In addition, if K is finite, G is said to be *almost ω - n -simply presented*.

When K is a priori chosen to be nice in G , one may state the following:

DEFINITION 1.2. The group G is called *nice almost ω_1 - n -simply presented* if there exists a countable nice subgroup N of G such that G/N is almost n -simply presented. In addition, if K is finite, G is said to be *nice almost ω - n -simply presented*.

Apparently, almost ω_1 - $p^{\omega+n}$ -projective groups and almost ω - n -simply presented groups are themselves nice almost ω_1 - n -simply presented and nice almost ω - n -simply presented, respectively. Besides, owing to Theorem 2.25 of [2], nice almost ω_1 - n -simply presented groups with countable first Ulm subgroup are almost n -simply presented (compare with Corollary 3.1 listed below).

On the other vein, Hill and Megibben gave in [7] the definition of a *c.c. group* as a group G such that $p^\omega(G/C)$ is countable whenever $C \leq G$ is a countable subgroup. Since $p^\omega G/(p^\omega G \cap C) \cong (p^\omega G + C)/C \subseteq p^\omega(G/C)$ is countable, it easily follows that $p^\omega G$ must be countable as well.

Our purpose here is to give a systematic study of the defined above two group classes as thereby we somewhat settle Problem 3 in [2]. Our work is organized as follows: In the next section we state and prove some preliminary technical claims and a background material. After that, in the third section, we proceed by proving the most of our basic results as we leave some specific of them in the fourth section. We close in the final section with some left-open problems of interest and importance.

2. Preliminaries and backgrounds

The following two technicalities are pivotal for the "niceness" property (cf. [2] too).

LEMMA 2.1. *Suppose that α is an ordinal, and that G and F are groups where F is finite. Then the following formula is fulfilled:*

$$p^\alpha(G + F) = p^\alpha G + [F \cap p^\alpha(G + F)] \subseteq p^\alpha G + F.$$

PROOF. We will use a transfinite induction on α . First, if $\alpha - 1$ exists, we have

$$\begin{aligned}
p^\alpha(G + F) &= p(p^{\alpha-1}(G + F)) = p(p^{\alpha-1}G + [F \cap p^{\alpha-1}(G + F)]) \\
&= p(p^{\alpha-1}G) + p(F \cap p^{\alpha-1}(G + F)) \\
&\subseteq p^\alpha G + [F \cap p(p^{\alpha-1}(G + F))] = p^\alpha G + [F \cap p^\alpha(G + F)].
\end{aligned}$$

Since the reverse inclusion " \supseteq " is obvious, we obtain the desired equality.

If now $\alpha - 1$ does not exist, we have that

$$p^\alpha(G + F) = \bigcap_{\beta < \alpha} (p^\beta(G + F)) \subseteq \bigcap_{\beta < \alpha} (p^\beta G + F) = \bigcap_{\beta < \alpha} p^\beta G + F = p^\alpha G + F.$$

In fact, the second sign " $=$ " follows like this: Given $x \in \bigcap_{\beta < \alpha} (p^\beta G + F)$, we write that $x = g_{\beta_1} + f_1 = \dots = g_{\beta_s} + f_s = \dots$ where $f_1, \dots, f_s \in F$ are the all elements of F ; $g_{\beta_1} \in p^{\beta_1}G, \dots, g_{\beta_s} \in p^{\beta_s}G$ with $\beta_1 < \dots < \beta_s < \dots$.

Since F is finite, while the number of equalities is infinite due to the infinite cardinality of α , we infer that $g_{\beta_s} \in p^\beta G$ for any ordinal $\beta < \alpha$ which means that $g_{\beta_s} \in \bigcap_{\beta < \alpha} p^\beta G = p^\alpha G$. Thus $x \in \bigcap_{\beta < \alpha} p^\beta G + F = p^\alpha G + F$, as claimed. Furthermore, $p^\alpha(G + F) \subseteq (p^\alpha G + F) \cap p^\alpha(G + F) = p^\alpha G + [F \cap p^\alpha(G + F)]$ which is obviously equivalent to an equality. \square

LEMMA 2.2. *Let N be a nice subgroup of a group G . Then*

- (i) $N + R$ is nice in G for every finite subgroup $R \leq G$;
- (ii) N is nice in $G + F$ for each finite group F .

PROOF. (i) For any limit ordinal γ , we deduce that

$$\bigcap_{\delta < \gamma} (N + R + p^\delta G) \subseteq R + \bigcap_{\delta < \gamma} (N + p^\delta G) = R + N + p^\gamma G$$

as required. Indeed, the relation " \subseteq " follows like this: Given $x \in \bigcap_{\delta < \gamma} (N + R + p^\delta G)$, we write $x = a_1 + r_1 + g_1 = \dots = a_s + r_s + g_s = \dots = a_k + r_1 + g_k = \dots$, where $a_1, \dots, a_k \in N$; $r_1, \dots, r_k \in R$; $g_1 \in p^{\delta_1}G, \dots, g_k \in p^{\delta_k}G$ with $\delta_1 < \dots < \delta_k$. So $a_1 + g_1 = \dots = a_k + g_k = \dots \in \bigcap_{\delta < \gamma} (N + p^\delta G)$ and hence $x \in R + \bigcap_{\delta < \gamma} (N + p^\delta G)$, as required.

(ii) Since N is nice in G , we may write $\bigcap_{\delta < \gamma} [N + p^\delta G] = N + p^\gamma G$ for every limit ordinal γ . Furthermore, with Lemma 2.1 at hand, we subsequently deduce that

$$\begin{aligned}
\bigcap_{\delta < \gamma} [N + p^\delta(G + F)] &= \bigcap_{\delta < \gamma} [N + p^\delta G + (F \cap p^\delta(G + F))] \\
&\subseteq \bigcap_{\delta < \gamma} (N + p^\delta G) + [F \cap p^\gamma(G + F)] \\
&= N + p^\gamma G + [F \cap p^\gamma(G + F)] = N + p^\gamma(G + F).
\end{aligned}$$

The inclusion " \subseteq " follows thus: Given $x \in \bigcap_{\delta < \gamma} [N + p^\delta G + (F \cap p^\delta(G + F))]$, we write $x = a_1 + g_1 + f_1 = \dots = a_s + g_s + f_s = \dots = a_k + g_k + f_1 = \dots$, where $a_1, \dots, a_k \in N$; $g_1 \in p^{\delta_1}G, \dots, g_k \in p^{\delta_k}G$; $f_1 \in F \cap p^{\delta_1}(G + F), \dots, f_k \in F \cap p^{\delta_k}(G + F)$ with $\delta_1 < \dots < \delta_k$. Hence $a_1 + g_1 = \dots = a_k + g_k = \dots \in \bigcap_{\delta < \gamma} (N + p^\delta G)$ and

because the number of the f_i 's ($1 \leq i \leq k$) is finite whereas the number of equalities is not, we can deduce that $f_1 \in \bigcap_{\delta < \gamma} (F \cap p^\delta(G+F)) = F \cap p^\gamma(G+F)$, as needed. \square

The following can be seen as Proposition 2.20 from [2]. It is listed here only for the sake of completeness and the readers' convenience.

LEMMA 2.3. *If T is almost n -simply presented and G/T is countable, then G is almost n -simply presented.*

PROOF. Write $G = T + K$, where $K \leq G$ is countable. With bullet three listed above at hand, there exists $P \leq T[p^n]$ such that T/P is almost simply presented. Furthermore, $G/P = (T/P) + (K + P)/P$, where $(K + P)/P \cong K/(K \cap P)$ is countable. Thus Theorem 1 of [1] (see [8] too) can be employed to show that G/P is almost simply presented, as required. \square

The next statement appears as Proposition 2.23 (b) in [2] in the case of almost n -simply presented groups. We here extend it even in the situation of nicely almost n -simply presented groups.

LEMMA 2.4. *If S is a subgroup of a group G such that G/S is finite, then G is (nicely) almost n -simply presented if and only if S is (nicely) almost n -simply presented.*

PROOF. Write $G = S + F$, where $F \leq G$ is finite. Suppose first that S is nicely almost n -simply presented. With bullet three quoted above in hand, there is $Z \leq S[p^n]$ which is nice in S such that S/Z is almost simply presented. We therefore have that $G/Z = [S/Z] + [(F + Z)/Z]$, where $(F + Z)/Z \cong F/(F \cap Z)$ is finite. Again by virtue of Theorem 1 in [1], G/Z should be almost simply presented. But Z is nice in G utilizing Lemma 2.2 (ii), as required.

Reciprocally, let G be nicely almost n -simply presented. Since $p^t G = p^t S$ for some $t \in \mathbb{N}$, and in Lemma 2.3 (iii) of [2] it was established that any group A is nicely almost n -simply presented if and only if so is $p^t A$, one may derive that S is nicely almost n -simply presented. Actually, this idea also provides a new verification of the sufficiency, considered above.

The same method works for almost n -simply presented groups as well. \square

So, we are now coming to the following (see also Proposition 2.23 (a) from [2]).

LEMMA 2.5. *A group G is almost n -simply presented if and only if G/F is almost n -simply presented for some finite subgroup F of G .*

PROOF. The "and only if" direction was proved as Corollary 2.19 in [2].

To treat the "if" one, write $(G/F)/(A/F) \cong G/A$ is almost simply presented for some $A \leq G$ such that $p^n A \subseteq F \subseteq A$. Since $p^n A$ is finite, it is a routine technical exercise to check that $A = L + A[p^n]$ for some finite $L \leq A$. Furthermore, $G/A \cong (G/A[p^n])/(A/A[p^n])$ being almost simply presented with finite $A/A[p^n] \cong L/L[p^n]$ implies with the help of [1] (see [8] or [9] as well) that $G/A[p^n]$ is almost simply presented, as required. \square

With the last assertion at hand, one observes that almost ω - n -simply presented groups are exactly the almost n -simply presented ones.

3. Main results

We begin this section with some different characterizations of almost ω_1 - n -simply presented groups. The first major result is the following:

THEOREM 3.1. *The following points are equivalent:*

- (i) G is almost ω_1 - n -simply presented;
- (ii) $G/(C \oplus L)$ is almost simply presented, where C is a countable subgroup of G and L is a p^n -bounded subgroup of G ;
- (iii) G/L is almost ω_1 -simply presented for some $L \leq G[p^n]$.

PROOF. (i) \Leftrightarrow (ii). Foremost, letting point (i) be fulfilled, given G/K is almost n -simply presented for some countable subgroup $K \leq G$. Thus there is A/K with $A \leq G$ and $p^n A \subseteq K$ such that G/A is almost simply presented. But it is well known that $A = C \oplus L$, and hence (ii) holds.

Conversely, assume that clause (ii) is true. Thus $G/(C \oplus L) \cong [G/C]/[(C \oplus L)/C]$ is almost simply presented, where $(C \oplus L)/C \cong L$ is p^n -bounded. Therefore G/C is almost n -simply presented, as required.

(ii) \Leftrightarrow (iii). First, assuming that point (ii) is valid, we see that $G/(C \oplus L) \cong [G/L]/[(C \oplus L)/L]$ is almost simply presented, where $(C \oplus L)/L \cong C$ is countable. So, G/L is almost ω_1 -simply presented.

Reciprocally, let clause (iii) be true, so given G/L is almost ω_1 -simply presented for some p^n -bounded subgroup L . Hence there is a countable subgroup B/L with $B \leq G$ such that $(G/L)/(B/L) \cong G/B$ is almost simply presented. Besides, $B = L + K$ for some countable $K \leq B$. Since $p^n L = \{0\}$, we write $L = L_1 \oplus L_2$, where L_2 is countable and $L \cap K \subseteq L_2$. Observe that $B = L_1 + (K + L_2)$, where L_1 is p^n -bounded and $K + L_2$ is countable. Moreover, $L_1 \cap (K + L_2) = \{0\}$; indeed take $a = b + c$, where $a \in L_1$, $b \in K$ and $c \in L_2$. Furthermore, $a - c \in L \cap K \subseteq L_2$, whence $a \in L_1 \cap L_2 = \{0\}$ and so $a = 0$. Finally, $B = L_1 \oplus (K + L_2)$ and thus $G/(C \oplus M)$ is almost simply presented for the countable $C = K + L_2$ and the p^n -bounded $M = L_1$, as stated. \square

We continue with some other structural affirmations.

PROPOSITION 3.1. *If G is nicely almost ω_1 - n -simply presented, then $G/p^\omega G$ is almost $p^{\omega+n}$ -projective. In particular, separable nicely almost ω_1 - n -simply presented groups are always almost $p^{\omega+n}$ -projective.*

PROOF. According to Corollary 2.8 of [3], one observes that the quotient $(G/N)/p^\omega(G/N) \cong G/(p^\omega G + N)$ is almost $p^{\omega+n}$ -projective, where N is a countable nice subgroup of G . But $G/(p^\omega G + N) \cong [G/p^\omega G]/[(p^\omega G + N)/p^\omega G]$, where it is obvious that $(p^\omega G + N)/p^\omega G \cong N/(N \cap p^\omega G)$ is countable. Henceforth, we apply Proposition 2.10 from [3] to get the first claim. The second part is its trivial consequence. \square

COROLLARY 3.1. *Suppose G is a group for which $p^\omega G$ is countable. Then G is nicely almost ω_1 - n -simply presented if and only if G is almost ω_1 - $p^{\omega+n}$ -projective.*

PROOF. In view of Proposition 3.1, the factor-group $G/p^\omega G$ is almost $p^{\omega+n}$ -projective. Hence, in virtue of [3], G is almost ω_1 - $p^{\omega+n}$ -projective, as expected.

The reverse implication is obvious. \square

PROPOSITION 3.2. *If G is both an almost ω_1 - n -simply presented group and a c.c. group, then $G/p^\omega G$ is almost $p^{\omega+n}$ -projective.*

PROOF. Let G/K be an almost n -simply presented group, where K is a countable subgroup of G . Therefore, we apply Corollary 2.8 from [3] to show that

$$(G/K)/p^\omega(G/K) \cong G/\left[\bigcap_{i<\omega}(p^i G + K)\right] \cong (G/p^\omega G)/\left[\bigcap_{i<\omega}(p^i G + K)/p^\omega G\right]$$

is almost $p^{\omega+n}$ -projective. Since $[\bigcap_{i<\omega}(p^i G + K)]/p^\omega G$ is countable, again Proposition 2.10 in [3] applies to get that $G/p^\omega G$ remains almost $p^{\omega+n}$ -projective, as desired. \square

As two immediate consequences, we deduce:

COROLLARY 3.2. *Suppose G is a c.c. group. Then G is almost ω_1 - n -simply presented if and only if G is almost ω_1 - $p^{\omega+n}$ -projective.*

PROOF. The sufficiency being elementary, we deal with the necessity. Since c.c. groups are obviously with countable first Ulm subgroup, Proposition 3.2 allows us to conclude with the help of [3] that G is almost ω_1 - $p^{\omega+n}$ -projective, as stated. \square

We recollect that a group is termed *weakly ω_1 -separable* if it is a separable c.c. group. So, we directly obtain:

COROLLARY 3.3. *Suppose G is a weakly ω_1 -separable group. Then G is almost ω_1 - n -simply presented if and only if G is almost $p^{\omega+n}$ -projective.*

Furthermore, we come to the following.

PROPOSITION 3.3. *Suppose that A is a group with a countable subgroup L . Then A is almost ω_1 - n -simply presented if and only if A/L is almost ω_1 - n -simply presented.*

PROOF. First, let us assume that A be almost ω_1 - n -simply presented, hence A/K is almost n -simply presented for some countable $K \leq A$. But

$$[A/L]/[(L+K)/L] \cong A/(L+K) \cong [A/K]/[(L+K)/K],$$

where the last factor-group $[A/K]/[(L+K)/K]$ is almost n -simply presented by Proposition 2.18 (or Corollary 2.19) of [2] since $(L+K)/K$ is countable. Therefore, $[A/L]/[(L+K)/L]$ is almost n -simply presented with countable $(L+K)/L \cong K/(K \cap L)$, as wanted.

Reciprocally, let us now A/L be almost ω_1 - n -simply presented, and so let C/L be a countable subgroup of A/L for some $C \leq A$ such that $(A/L)/(C/L) \cong A/C$ is almost n -simply presented. Observing that C is of necessity countable, we deduce via Definition 1.1 that A is almost ω_1 - n -simply presented, as formulated. \square

As an easy consequence, we deduce:

COROLLARY 3.4. *Suppose A is a group such that $p^\alpha A$ is countable for some ordinal α . Then A is almost ω_1 - n -simply presented if and only if $A/p^\alpha A$ is almost ω_1 - n -simply presented.*

PROPOSITION 3.4. *Let A be a group with a subgroup G such that A/G is countable. Then A is almost ω_1 - n -simply presented if and only if G is almost ω_1 - n -simply presented.*

PROOF. Write $A = G + C$ where C is countable and assume that G is almost ω_1 - n -simply presented. Now, Definition 1.1 insures that there is a countable subgroup K such that G/K is almost n -simply presented. Consequently, $A/K = (G/K) + (C + K)/K$. Employing Lemma 2.3, A/K is almost n -simply presented since $(C + K)/K$ is obviously countable. This gives that A is almost ω_1 - n -simply presented, as desired.

Conversely, let us assume that A is almost ω_1 - n -simply presented. Now, Proposition 3.3 guarantees that $(G + C)/C \cong G/(G \cap C)$ is almost ω_1 - n -simply presented. But $G \cap C$ is countable and again Proposition 3.3 will work to get that G is almost ω_1 - n -simply presented, as wanted. \square

We are now ready to prove the following central result:

THEOREM 3.2. *The class of almost ω_1 - n -simply presented groups is closed under the formation of ω_1 -bijections, and is the smallest class containing almost n -simply presented groups with this property.*

In other words, if $f : G \rightarrow A$ is an ω_1 -bijective homomorphism and G is an almost ω_1 - n -simply presented group, then A is an almost ω_1 - n -simply presented group, and thus almost ω_1 - n -simply presented groups form the minimal class of groups possessing that property.

PROOF. The first part follows by [10, Lemma 1.9] accomplished with Propositions 3.3 and 3.4.

For the second one, that it is the minimal class possessing that property, we making use Proposition 1.10 of [10] and Theorem 3.1. \square

PROPOSITION 3.5. *Suppose A is a group with a finite subgroup F . Then A is nicely almost ω_1 - n -simply presented if and only if A/F is nicely almost ω_1 - n -simply presented.*

PROOF. Assume first that A is nicely almost ω_1 - n -simply presented, i.e., there is a countable nice subgroup N such that A/N is almost n -simply presented. Observing as above that

$$[A/F]/[(F + N)/F] \cong A/(F + N) \cong [A/N]/[(F + N)/N],$$

and that $[A/N]/[(F + N)/N]$ is almost n -simply presented, it follows that A/F is nicely almost ω_1 - n -simply presented, because $(F + N)/F$ is countable and nice in A/F in accordance with Lemma 2.2 (i).

Conversely, given that A/F is nicely almost ω_1 - n -simply presented, so there exists a countable nice subgroup C/F of A/F with $C \leq A$ such that the factor-group $(A/F)/(C/F) \cong A/C$ is almost n -simply presented. Since F is nice in A , one can see that C is countable and nice in A (see [5]), whence Definition 1.2 gives the claim. \square

PROPOSITION 3.6. *Let A be a group with a subgroup G such that A/G is finite. Then A is almost ω_1 - $p^{\omega+n}$ -projective if and only if G is almost ω_1 - $p^{\omega+n}$ -projective.*

PROOF. Repeating the same method as in Proposition 3.3 combined with Proposition 3.5, we complete the arguments. \square

We are now ready to establish the following main result:

THEOREM 3.3. *The class of nicely almost ω_1 - n -simply presented groups is closed under taking ω -bijections.*

PROOF. Follows by our discussion in Section 2 (see again [10, Lemma 1.9]) along with Propositions 3.5 and 3.6. \square

PROPOSITION 3.7. *Let A be a group with a countable nice subgroup N . If A/N is nicely almost ω_1 - n -simply presented, then so is A .*

4. Nunke-like theorems

We will now prove some versions of Nunke-esque results for the new group classes defined in the introductory section. Generalizing the given above concept of a c.c. group, introduced in [7], one may define the following notion:

DEFINITION 4.1. Let λ be an ordinal. A group G is said to be λ -countably if for any countable subgroup $K \leq G$ the quotient $p^\lambda(G/K)/(p^\lambda G + K)/K = p^\lambda((G/K)/(p^\lambda G + K)/K)$ is countable.

THEOREM 4.1. *Suppose G is a $\lambda + n$ -countably group for some ordinal λ such that $p^\lambda G$ is almost n -simply presented. Then G is almost ω_1 - n -simply presented if and only if $G/p^{\lambda+n}G$ is almost ω_1 - n -simply presented.*

PROOF. (\Rightarrow) Given G/K is an almost n -simply presented group for some countable $K \leq G$. Consequently, Theorem 2.7 (a) in [2] forces that

$$\begin{aligned} (G/K)/p^{\lambda+n}(G/K) &\cong [(G/K)/(p^{\lambda+n}G + K)/K]/[p^{\lambda+n}(G/K)/(p^{\lambda+n}G + K)/K] \\ &= [(G/K)/(p^{\lambda+n}G + K)/K]/p^{\lambda+n}((G/K)/(p^{\lambda+n}G + K)/K) \end{aligned}$$

is also almost n -simply presented. Because of the countability of the quotient $p^{\lambda+n}(G/K)/[(p^{\lambda+n}G + K)/K] = p^{\lambda+n}((G/K)/(p^{\lambda+n}G + K)/K)$, a simple appeal to Theorem 2.10 of [2] leads to almost n -simply presentness of

$$[G/K]/[(p^{\lambda+n}G + K)/K] \cong G/(p^{\lambda+n}G + K) \cong [G/p^{\lambda+n}G]/[(p^{\lambda+n}G + K)/p^{\lambda+n}G].$$

And since $(p^{\lambda+n}G + K)/p^{\lambda+n}G \cong K/(K \cap p^{\lambda+n}G)$ is countable, we are done.

(\Leftarrow) Let $(G/p^{\lambda+n}G)/(C/p^{\lambda+n}G) \cong G/C$ be almost n -simply presented for some countable $C/p^{\lambda+n}G$ with $C \leq G$. Write $C = p^{\lambda+n}G + K$ for some countable subgroup K . So, $G/(p^{\lambda+n}G + K) \cong [G/K]/[(p^{\lambda+n}G + K)/K]$ is almost n -simply presented; note that this gives with the aid of Theorem 2.7 (a) from [2] that $p^{\lambda+n}((G/K)/(p^{\lambda+n}G + K)/K) = p^{\lambda+n}(G/K)/(p^{\lambda+n}G + K)/K$ is almost n -simply presented – however we have by assumption the more restrictive condition that this quotient is countable. Moreover, again Theorem 2.7(a) or Corollary 2.19 in [2] applies to conclude that

$$\begin{aligned} & [(G/K)/(p^{\lambda+n}G + K)/K]/p^{\lambda+n}((G/K)/(p^{\lambda+n}G + K)/K) \\ & \cong [(G/K)/(p^{\lambda+n}G + K)/K]/[p^{\lambda+n}(G/K)/(p^{\lambda+n}G + K)/K] \\ & \cong (G/K)/p^{\lambda+n}(G/K) \end{aligned}$$

is almost n -simply presented. But, on the other hand, by Theorem 2.7 (a) in [2] we deduce that $p^{\lambda+n}G$ is almost n -simply presented. Moreover, $(p^{\lambda+n}G + K)/K \cong p^{\lambda+n}G/(p^{\lambda+n}G \cap K)$, which means by Proposition 2.18 of [2] that the second term, and hence the first one, are almost n -simply presented because $p^{\lambda+n}G \cap K$ is countable. We therefore may apply Lemma 2.3 to derive that $p^{\lambda+n}(G/K)$ is almost n -simply presented. Finally, utilizing Theorem 2.10 in [2] to get after all that G/K is almost n -simply presented, as expected. \square

Notice that we have not used in the necessity the condition that $p^\lambda G$ is almost n -simply presented, so that what immediately arises is whether or not this limitation can be dropped off in the formulation of the theorem.

PROPOSITION 4.1. *Let G be a group and α an ordinal. If G is nicely almost ω_1 - n -simply presented, then $p^\alpha G$ and $G/p^\alpha G$ are nicely almost ω_1 - n -simply presented.*

PROOF. Let G/N be almost n -simply presented for some countable nice subgroup N of G . Hence, using [2], $p^\alpha(G/N) = (p^\alpha G + N)/N \cong p^\alpha G/(p^\alpha G \cap N)$ is almost n -simply presented, where $p^\alpha G \cap N$ is countable and nice in $p^\alpha G$ (cf. [5]).

Moreover, $(G/N)/p^\alpha(G/N) \cong G/(p^\alpha G + N) \cong [G/p^\alpha G]/[(p^\alpha G + N)/p^\alpha G]$ is almost n -simply presented, where $(p^\alpha G + N)/p^\alpha G \cong N/(p^\alpha G \cap N)$ is countable and $(p^\alpha G + N)/p^\alpha G$ is nice in $G/p^\alpha G$, because $N + p^\alpha G$ is so in G . \square

PROPOSITION 4.2. *Suppose that $G/p^{\lambda+n}G$ is almost n -simply presented for some ordinal λ . Then G is nicely almost ω_1 - n -simply presented if and only if $p^\lambda G$ is nicely almost ω_1 - n -simply presented.*

PROOF. The “only if” part follows by a direct application of Proposition 4.1.

As for the “if” part, let $p^\lambda G/Y = p^\lambda(G/Y)$ be almost n -simply presented for some nice countable subgroup Y . Hence Y is also nice in G (see, e.g., [5]), and besides $p^{\lambda+n}(G/Y)$ remains almost n -simply presented in conjunction with Theorem 2.7 (a) from [2]. But $G/p^{\lambda+n}G \cong (G/Y)/p^{\lambda+n}(G/Y)$ is almost n -simply presented by assumption. Now, the application of Theorem 2.10 of [2] leads us to G/Y is almost n -simply presented, as wanted. \square

5. Open problems

In closing, we shall state two left-open questions which remain unanswered.

PROBLEM 1. Does it follow that if G is an almost ω_1 - n -simply presented group with $p^\omega G = \{0\}$, then it is almost $p^{\omega+n}$ -projective?

PROBLEM 2. Let α be an ordinal. Does it follow that G is (nicely) almost ω_1 - n -simply presented if and only if both $p^\alpha G$ and $G/p^\alpha G$ are (nicely) almost ω_1 - n -simply presented?

Acknowledgment. The author is very grateful to the referee for the careful reading of the manuscript.

References

1. P. V. Danchev, *On extensions of primary almost totally projective groups*, Math. Bohemica **133** (2) (2008), 149–155.
2. ———, *On almost n -simply presented abelian p -groups*, Korean J. Math. **21**(4) (2013), 401–419.
3. ———, *On almost ω_1 - $p^{\omega+n}$ -projective abelian p -groups*, Korean J. Math. **22**(3) (2014), 501–516.
4. ———, *On ω_1 - n -simply presented abelian p -groups*, J. Algebra Appl. **14**(3) (2015).
5. L. Fuchs, *Infinite Abelian Groups*, volumes I and II, Acad. Press, New York and London, 1970 and 1973.
6. P. Griffith, *Infinite Abelian Group Theory*, The University of Chicago Press, Chicago–London, 1970.
7. P. D. Hill, C. K. Megibben, *Primary abelian groups whose countable subgroups have countable closure*; in: A. Facchini, C. Menini (eds.), *Abelian Groups and Modules, Proc. Padova conf., 1994*, Math. Appl., Dordr. **343**, Kluwer, Dordrecht, 1995, 283–290.
8. P. D. Hill, W. D. Ullery, *Isotype separable subgroups of totally projective subgroups*; in: A. Facchini, C. Menini (eds.), *Abelian Groups and Modules, Proc. Padova conf., 1994*, Math. Appl., Dordr. **343**, Kluwer, Dordrecht, 1995, 291–300.
9. ———, *Almost totally projective groups*, Czech. Math. J. **46**(2) (1996), 249–258.
10. P. W. Keef, *On ω_1 - $p^{\omega+n}$ -projective primary abelian groups*, J. Algebra Numb. Th. Acad. **1**(1) (2010), 41–75.

Department of Mathematics
Plovdiv State University
4000 Plovdiv
Bulgaria
pvdanchev@yahoo.com

(Received 26 02 2014)