

## THE INDUCED CONNECTIONS ON TOTAL SPACES OF FIBRED MANIFOLDS

Włodzimierz M. Mikulski

**ABSTRACT.** Let  $Y \rightarrow M$  be a fibred manifold with  $m$ -dimensional base and  $n$ -dimensional fibres. If  $m \geq 2$  and  $n \geq 3$ , we classify all linear connections  $A(\Gamma, \Lambda, \Theta) : TY \rightarrow J^1(TY \rightarrow Y)$  in  $TY \rightarrow Y$  (i.e., classical linear connections on  $Y$ ) depending canonically on a system  $(\Gamma, \Lambda, \Theta)$  consisting of a general connection  $\Gamma : Y \rightarrow J^1Y$  in  $Y \rightarrow M$ , a torsion free classical linear connection  $\Lambda : TM \rightarrow J^1(TM \rightarrow M)$  on  $M$  and a linear connection  $\Theta : VY \rightarrow J^1(VY \rightarrow Y)$  in the vertical bundle  $VY \rightarrow Y$ .

### Introduction

All manifolds considered in the paper are assumed to be Hausdorff, second countable, without boundary, finite dimensional and smooth (of class  $C^\infty$ ). Maps between manifolds are assumed to be smooth (infinitely differentiable).

Let  $Y \rightarrow M$  be a fibred manifold with  $m$ -dimensional base  $M$  and  $n$ -dimensional fibres. Let  $\Gamma : Y \rightarrow J^1Y$  be a general connection in a fibred manifold  $Y \rightarrow M$  (i.e., a section of the first jet prolongation  $\pi_0^1 : J^1Y \rightarrow Y$  of  $Y \rightarrow M$ ),  $\Lambda : TM \rightarrow J^1(TM \rightarrow M)$  be a torsion free linear connection in the tangent bundle  $TM \rightarrow M$  of  $M$  (i.e., a torsion free classical linear connection on  $M$ ) and  $\Theta : VY \rightarrow J^1(VY \rightarrow Y)$  be a linear connection in the vertical bundle  $VY \rightarrow Y$  of  $Y \rightarrow M$  (i.e., a vertical classical linear connection on  $Y \rightarrow M$ ). More on connections can be found in [6].

Here we study how to construct canonically a linear connection  $A(\Gamma, \Lambda, \Theta) : TY \rightarrow J^1(TY \rightarrow Y)$  in  $TY \rightarrow Y$  (i.e., a classical linear connection on the total space  $Y$ ) from the system  $(\Gamma, \Lambda, \Theta)$  as above.

For example, one can construct a linear connection  $\Psi = \Psi(\Gamma, \Lambda, \Theta) : TY \rightarrow J^1(TY \rightarrow Y)$  in  $TY \rightarrow Y$  as follows. We decompose  $Z \in T_yY$  into the horizontal part  $h(Z) = \Gamma(y, Z_0)$ ,  $Z_0 \in T_xM$ ,  $x = p(y)$  and the vertical part  $vZ$ . We take a vector field  $X$  on  $M$  such that  $j_x^1X = \Lambda(Z_0)$  and construct its  $\Gamma$ -lift  $\Gamma X : Y \rightarrow TY$ ,

---

2010 *Mathematics Subject Classification*: Primary 53C05; Secondary 58A32.

*Key words and phrases*: general connection, linear connection, natural operator.

Communicated by Stevan Pilipović.

and we take a vertical vector field  $\theta Z : Y \rightarrow VY$  such that  $j_y^1(\theta Z) = \Theta(vZ)$ . For every  $Z \in T_y Y$  we define

$$\Psi(Z) = j_y^1(\Gamma X + \theta Z).$$

The coordinate expression of  $\Psi$  can be found in Section 1.

In Section 2, using the torsion of  $\Psi$ , we produce 12 tensor fields  $\tau_i = \tau_i(\Gamma, \Lambda, \Theta)$  ( $i = 1, \dots, 12$ ) of type  $T^* \otimes T^* \otimes T$  on  $Y$ .

The main result of the paper is the following one. If  $m \geq 2$  and  $n \geq 3$ , then the canonical constructions in question form the 12-parameter family  $\Psi + \sum_i \lambda^i \tau_i$  for real numbers  $\lambda^i$ ,  $i = 1, \dots, 12$ .

### 1. The coordinate expression

Let  $x^1, \dots, x^m$  be the usual coordinates on  $\mathbf{R}^m$ . Let  $\mathbf{R}^{m,n}$  be the trivial bundle over  $\mathbf{R}^m$  with the standard fiber  $\mathbf{R}^n$  and  $x^1, \dots, x^m, y^1, \dots, y^n$  be the usual fiber coordinates on  $\mathbf{R}^{m,n}$ . Let  $\eta^1, \dots, \eta^n$  be the additional coordinates on  $V\mathbf{R}^{m,n}$  and  $\xi^i$  be the additional coordinates in  $T\mathbf{R}^m$ . Let  $(\Gamma, \Lambda, \Theta)$  and  $\Psi = \Psi(\Gamma, \Lambda, \Theta)$  be as in Introduction. Let  $dy^p = F_i^p(x, y)dx^i$  be the coordinate expression of  $\Gamma$ ,

$$(1.1) \quad d\xi^i = \Lambda_{jk}^i(x)\xi^j dx^k$$

be the coordinate expression of  $\Lambda$ , and  $d\eta^p = \Theta_{qi}^p(x, y)\eta^q dx^i + \Theta_{qs}^p(x, y)\eta^q dy^s$  be the coordinate expression of  $\Theta$ . Then we have the following lemma.

LEMMA 1.1. *The coordinate expression of  $\Psi = \Psi(\Gamma, \Lambda, \Theta)$  is (1.1) and*

$$(1.2) \quad d\eta^p = \left( \frac{\partial F_i^p}{\partial x^j} \xi^i + F_i^p \Lambda_{kj}^i \xi^k + \Theta_{qj}^p (\eta^q - F_i^q \xi^i) \right) dx^j + \left( \frac{\partial F_i^p}{\partial y^s} \xi^i + \Theta_{qs}^p (\eta^q - F_i^q \xi^i) \right) dy^s.$$

where (the same letters)  $\xi^1, \dots, \xi^m, \eta^1, \dots, \eta^n$  denote the usual additional coordinates on  $T\mathbf{R}^{m,n}$ .

PROOF. Let  $\xi^i = X^i(x)$  and  $\eta^p = (\theta Z)^p(x, y)$  be the coordinate expression of  $X$  or  $\theta Z$ , respectively. Hence

$$\frac{\partial X^i}{\partial x^j} = \Lambda_{kj}^i X^k, \quad \frac{\partial (\theta Z)^p}{\partial x^j} = \Theta_{qj}^p (\theta Z)^q, \quad \frac{\partial (\theta Z)^p}{\partial y^s} = \Theta_{qs}^p (\theta Z)^q.$$

Then the coordinate expression of  $\Gamma X + \theta Z$  is

$$\xi^i = X^i(x) \text{ and } \eta^p = F_i^p(x, y)X^i(x) + (\theta Z)^p(x, y).$$

Differentiating this relation, we obtain (1.2). □

### 2. Main examples

Let  $(\Gamma, \Lambda, \Theta)$  be the triple as in the introduction. According to the usual  $\Gamma$ -decomposition  $TY = VY \oplus_Y H^\Gamma Y$  we have the decomposition

$$\begin{aligned} T^*Y \otimes TY &= (V^*Y \otimes VY) \oplus_Y (V^*Y \otimes H^\Gamma Y) \\ &\oplus_Y ((H^\Gamma Y)^* \otimes VY) \oplus_Y ((H^\Gamma Y)^* \otimes H^\Gamma Y). \end{aligned}$$

Let  $\text{id}_{HY}$  be the tensor field of type  $T^* \otimes T$  on  $Y$  being the  $(H^\Gamma Y)^* \otimes H^\Gamma Y$ -component of the identity tensor field  $\text{id}_{TY}$  on  $Y$  (the other 3 component of  $\text{id}_{HY}$  are zero). Let  $\text{id}_{VY}$  be the tensor field of type  $T^* \otimes T$  on  $Y$  being the  $V^* Y \otimes VY$ -component of  $\text{id}_{TY}$  (the other 3 components of  $\text{id}_{VY}$  are zero).

Quite similarly, we have the decomposition

$$\begin{aligned} T^* Y \otimes T^* Y \otimes TY &= (V^* Y \otimes V^* Y \otimes VY) \oplus_Y (V^* Y \otimes V^* Y \otimes H^\Gamma Y) \\ &\oplus_Y (V^* Y \otimes (H^\Gamma Y)^* \otimes VY) \oplus_Y (V^* Y \otimes (H^\Gamma Y)^* \otimes H^\Gamma Y) \\ &\oplus_Y ((H^\Gamma Y)^* \otimes V^* Y \otimes VY) \oplus_Y ((H^\Gamma Y)^* \otimes V^* Y \otimes H^\Gamma Y) \\ &\oplus_Y ((H^\Gamma Y)^* \otimes H^\Gamma Y)^* \otimes VY \oplus_Y ((H^\Gamma Y)^* \otimes (H^\Gamma Y)^* \otimes H^\Gamma Y). \end{aligned}$$

Let  $\text{Tor}^{H^* \otimes V^* \otimes V}(\Psi)$  be the  $(H^\Gamma Y)^* \otimes V^* Y \otimes VY$ -component of the torsion tensor field  $\text{Tor}(\Psi)$  of the classical linear connection  $\Psi = \Psi(\Gamma, \Lambda, \Theta)$  (from Introduction). This components can be treated as the tensor field of type  $T^* \otimes T^* \otimes T$  on  $Y$  (the other 7 components of it are zero). Taking contraction  $C_2^1 : T^* Y \otimes T^* Y \otimes TY \rightarrow T^* Y$ ,  $C_2^1(\omega_1 \otimes \omega_2 \otimes v_1) = \langle \omega_2, v_1 \rangle \omega_1$ , we produce tensor field  $C_2^1 \text{Tor}^{H^* \otimes V^* \otimes V}(\Psi)$  of type  $T^*$  on  $Y$  (horizontal vector field). Similarly, let  $\text{Tor}^{H^* \otimes H^* \otimes V}(\Psi)$  or  $\text{Tor}^{V^* \otimes H^* \otimes V}(\Psi)$  or  $\text{Tor}^{V^* \otimes V^* \otimes V}(\Psi)$  be the (treated as the tensor field of type  $T^* \otimes T^* \otimes T$  on  $Y$ )  $(H^\Gamma Y)^* \otimes (H^\Gamma Y)^* \otimes VY$ - or  $V^* Y \otimes (H^\Gamma Y)^* \otimes VY$ - or  $V^* Y \otimes V^* Y \otimes VY$ -component of  $\text{Tor}(\Psi)$ , respectively. Thus we have the following tensor fields of type  $T^* \otimes T^* \otimes T$  on  $Y$  canonically depending on  $(\Gamma, \Lambda, \Theta)$ .

$$\text{EXAMPLE 2.1. } \tau_1(\Gamma, \Lambda, \Theta) := \text{Tor}^{H^* \otimes H^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.2. } \tau_2(\Gamma, \Lambda, \Theta) := \text{Tor}^{H^* \otimes V^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.3. } \tau_3(\Gamma, \Lambda, \Theta) := \text{id}_{HY} \otimes C_2^1 \text{Tor}^{H^* \otimes V^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.4. } \tau_4(\Gamma, \Lambda, \Theta) := C_2^1 \text{Tor}^{H^* \otimes V^* \otimes V}(\Psi) \otimes \text{id}_{HY} .$$

$$\text{EXAMPLE 2.5. } \tau_5(\Gamma, \Lambda, \Theta) := C_2^1 \text{Tor}^{H^* \otimes V^* \otimes V}(\Psi) \otimes \text{id}_{VY} .$$

$$\text{EXAMPLE 2.6. } \tau_6(\Gamma, \Lambda, \Theta) := \text{id}_{VY} \otimes C_2^1 \text{Tor}^{H^* \otimes V^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.7. } \tau_7(\Gamma, \Lambda, \Theta) = \text{Tor}^{V^* \otimes V^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.8. } \tau_8(\Gamma, \Lambda, \Theta) := \text{id}_{HY} \otimes C_2^1 \text{Tor}^{V^* \otimes V^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.9. } \tau_9(\Gamma, \Lambda, \Theta) := C_2^1 \text{Tor}^{V^* \otimes V^* \otimes V}(\Psi) \otimes \text{id}_{HY} .$$

$$\text{EXAMPLE 2.10. } \tau_{10}(\Gamma, \Lambda, \Theta) := \text{id}_{VY} \otimes C_2^1 \text{Tor}^{V^* \otimes V^* \otimes V}(\Psi) .$$

$$\text{EXAMPLE 2.11. } \tau_{11}(\Gamma, \Lambda, \Theta) := C_2^1 \text{Tor}^{V^* \otimes V^* \otimes V}(\Psi) \otimes \text{id}_{VY} .$$

$$\text{EXAMPLE 2.12. } \tau_{12}(\Gamma, \Lambda, \Theta) := \text{Tor}^{V^* \otimes H^* \otimes V}(\Psi) .$$

### 3. Natural operators

Let  $\mathcal{FM}_{m,n}$  be the category of fibred manifolds with  $m$ -dimensional bases and  $n$ -dimensional fibres and their fibred (local) diffeomorphisms. The general concept of natural operators can be found in [6]. We need the following particular cases of natural operators, only.

**DEFINITION 3.1.** An  $\mathcal{FM}_{m,n}$ -natural operator  $A$  sending systems  $(\Gamma, \Lambda, \Theta)$  as in Introduction on fibred manifolds  $Y \rightarrow M$  into classical linear connections  $A_Y(\Gamma, \Lambda, \Theta)$  on  $Y$  is an  $\mathcal{FM}_{m,n}$ -invariant system of regular operators

$$A_Y : \text{Con}(Y) \times \text{Con}_{\text{clas}}^0(M) \times \text{Con}_{\text{vert-clas}}(Y) \rightarrow \text{Con}_{\text{clas}}(Y)$$

for any  $\mathcal{FM}_{m,n}$ -object  $Y = (Y \rightarrow M)$ , where  $\text{Con}(Y)$  is the set of general connections  $\Gamma$  in  $Y \rightarrow M$ ,  $\text{Con}_{\text{clas}}^0(M)$  is the set of torsion free classical linear connections  $\Lambda$  on  $M$ ,  $\text{Con}_{\text{vert-clas}}(Y)$  is the set of vertical classical linear connections  $\Theta$  on  $Y \rightarrow M$  and  $\text{Con}_{\text{clas}}(Y)$  is the set of classical linear connections on  $Y$ . The  $\mathcal{FM}_{m,n}$ -invariance of  $A$  means that if  $(\Gamma, \Lambda, \Theta) \in \text{Con}(Y) \times \text{Con}_{\text{clas}}^0(M) \times \text{Con}_{\text{vert-clas}}(Y)$  is  $f$ -related to  $(\Gamma_1, \Lambda_1, \Theta_1) \in \text{Con}(Y_1) \times \text{Con}_{\text{clas}}^0(M_1) \times \text{Con}_{\text{vert-clas}}(Y_1)$  for a  $\mathcal{FM}_{m,n}$ -map  $f : Y \rightarrow Y_1$  with the base map  $\underline{f} : M \rightarrow M_1$ , then  $A_Y(\Gamma, \Lambda, \Theta)$  and  $A_{Y_1}(\Gamma_1, \Lambda_1, \Theta_1)$  are  $f$ -related. The regularity of  $A$  means that  $A_Y$  transforms smoothly parametrized families into smoothly parametrized families.

Clearly, the construction of classical linear connection  $\Psi(\Gamma, \Lambda, \Theta)$  (from Introduction) determines a natural operator in the above sense.

To classify all natural operators in the sense of Definition 3.1, it suffices to classify all natural operators in the following sense.

**DEFINITION 3.2.** An  $\mathcal{FM}_{m,n}$ -natural operator  $A$  sending systems  $(\Gamma, \Lambda, \Theta)$  as in Introduction on fibred manifolds  $Y \rightarrow M$  into tensor fields  $A_Y(\Gamma, \Lambda, \Theta)$  of type  $T^* \otimes T^* \otimes T$  on  $Y$  is an  $\mathcal{FM}_{m,n}$ -invariant system of regular operators

$$A_Y : \text{Con}(Y) \times \text{Con}_{\text{clas}}^0(M) \times \text{Con}_{\text{vert-clas}}(Y) \rightarrow \text{Ten}^{(1,2)}(Y)$$

for any  $\mathcal{FM}_{m,n}$ -object  $Y \rightarrow M$ , where  $\text{Ten}^{(1,2)}(Y)$  is the space of tensor fields of type  $T^* \otimes T^* \otimes T$  on  $Y$ .

Of course, the constructions of tensor fields  $\tau_i(\Gamma, \Lambda, \Theta)$  from Examples 2.1–2.12 determine natural operators in the sense of Definition 3.2.

### 4. Estimation of dimension

We denote the trivial general connection on  $\mathbf{R}^{m,n}$  by  $\Gamma^0$  (i.e.,  $\Gamma^0 = \sum_{i=1}^m dx^i \otimes \frac{\partial}{\partial x^i}$ ), the torsion free flat classical linear connection on  $\mathbf{R}^m$  by  $\Lambda^0$  (i.e.,  $\Lambda^0 = (0)$ ) and the trivial vertical classical linear connection on  $\mathbf{R}^{m,n}$  by  $\Theta^0$  (i.e.,  $\Theta^0 = \sum_{i=1}^m dx^i \otimes \frac{\partial}{\partial x^i} + \sum_{p=1}^n dy^p \otimes \frac{\partial}{\partial y^p}$ ).

We study a natural operators  $A$  in the sense of Definition 3.2. For simplicity, we will omit the subscripts  $Y$  on  $A_Y$ . It is clear that  $A$  is determined by the values  $A(\Gamma, \Lambda, \Theta)(y) \in T_y^* Y \otimes T_y^* Y \otimes T_y Y$  for fibred manifolds  $p_Y : Y \rightarrow M$  with  $m$ -dimensional bases and  $n$ -dimensional fibres, general connections  $\Gamma$  on  $Y \rightarrow M$ ,

torsion free classical linear connections  $\Lambda$  on  $M$  and vertical classical linear connections  $\Theta$  on  $Y \rightarrow M$  and  $y \in Y_x$ ,  $x \in M$ . More, using the invariance of  $A$  with respect to (respective) fibred manifold charts, we can assume  $Y = \mathbf{R}^{m,n}$ ,  $y = (0, 0)$ . Further, using Corollary 19.8 in [6], we may assume

$$(4.1) \quad \Gamma = \Gamma^0 + \sum F_{j;\alpha\beta}^p x^\alpha y^\beta dx^j \otimes \frac{\partial}{\partial y^p},$$

where the sum is over all  $m$ -tuples  $\alpha$  and all  $n$ -tuples  $\beta$  of non-negative integers and  $j = 1, \dots, m$  and  $p = 1, \dots, n$  with  $1 \leq |\alpha| + |\beta| \leq K$  (we can assume  $F_{j;(0)(0)}^p = 0$  by the existence of respective "adapted" (for  $\Gamma$ ) fibred coordinates),

$$(4.2) \quad \Lambda = \left( \sum \Lambda_{jk;\gamma}^i x^\gamma \right)_{i,j,k=1,\dots,m}, \quad \Lambda_{jk;\gamma}^i = \Lambda_{kj;\gamma}^i,$$

where the sums are over all  $m$ -tuples  $\gamma$  of non-negative integers with  $1 \leq |\gamma| \leq K$  (we can assume  $\Lambda_{jk;(0)}^i = 0$  by the existence of  $\Lambda$ -normal coordinates on the base and the fact that torsion free classical linear connection has vanishing symbols in the center of normal coordinates),

$$(4.3) \quad \Theta = \Theta^0 + \sum \Theta_{ip;\delta\sigma}^r x^\delta y^\sigma \eta^p dx^i \otimes \frac{\partial}{\partial \eta^r} + \sum \Theta_{sp;\delta\sigma}^r x^\delta y^\sigma \eta^p dy^s \otimes \frac{\partial}{\partial \eta^r},$$

where the first sum is over all  $m$ -tuples  $\delta$  and all  $n$ -tuples  $\sigma$  of non-negative integers and  $i = 1, \dots, m$  and  $r, p = 1, \dots, n$  with  $0 \leq |\delta| + |\sigma| \leq K$  and the second sum is over all  $m$ -tuples  $\delta$  and  $n$ -tuples  $\sigma$  of non-negative integers and  $r, s, p = 1, \dots, n$  with  $0 \leq |\delta| + |\sigma| \leq K$ , where  $K$  is an arbitrary positive integer.

(More precisely, from Corollary 19.8 in [6], given  $(\Gamma, \Lambda, \Theta) \in \text{Con}(\mathbf{R}^{m,n}) \times \text{Con}_{\text{clas}}^0(\mathbf{R}^m) \times \text{Con}_{\text{vert-clas}}(\mathbf{R}^{m,n})$ , there exists a finite number  $r = r(\Gamma, \Lambda, \Theta)$  such that for any  $\Gamma_1 \in \text{Con}(\mathbf{R}^{m,n})$  we have the following implication

$$j_{(0,0)}^r \Gamma_1 = j_{(0,0)}^r \Gamma \Rightarrow A(\Gamma_1, \Lambda, \Theta)(0, 0) = A(\Gamma, \Lambda, \Theta)(0, 0),$$

i.e., we may replace  $\Gamma$  by  $\Gamma^1$  being polynomial. Next, by the quite similar argument, we can replace  $\Lambda$  by  $\Lambda^1$  being polynomial. Next, by the quite similar argument, we can replace  $\Theta$  by  $\Theta^1$  being polynomial.)

So,  $A$  is determined by the collection of smooth maps  $A_K : \mathbf{R}^{n(K)} \rightarrow \mathbf{R}^q = T_{(0,0)}^* \mathbf{R}^{m,n} \otimes T_{(0,0)}^* \mathbf{R}^{m,n} \otimes T_{(0,0)} \mathbf{R}^{m,n}$  ( $K = 1, 2, \dots$ ) given by

$$A_K((F_{j;\alpha\beta}^p), (\Lambda_{jk;\gamma}^i), (\Theta_{ip;\delta\sigma}^r), (\Theta_{sp;\delta\sigma}^r)) := A(\Gamma, \Lambda, \Theta)(0, 0),$$

where  $\Gamma, \Lambda, \Theta$  are as in (4.1), (4.2) and (4.3).

Using the invariance of  $A$  with respect to  $\varphi_t \times \phi_t$ ,  $\varphi_t = t \text{id}_{\mathbf{R}^m}$ ,  $\phi_t = t \text{id}_{\mathbf{R}^n}$ ,  $t > 0$ , we get the homogeneous condition

$$\begin{aligned} t A_K((F_{j;\alpha\beta}^p), (\Lambda_{jk;\gamma}^i), (\Theta_{ip;\delta\sigma}^r), (\Theta_{sp;\delta\sigma}^r)) \\ = A_K((t^{|\alpha|+|\beta|} F_{j;\alpha\beta}^p), (t^{|\gamma|+1} \Lambda_{jk;\gamma}^i), (t^{|\delta|+|\sigma|+1} \Theta_{ip;\delta\sigma}^r), (t^{|\delta|+|\sigma|+1} \Theta_{sp;\delta\sigma}^r)). \end{aligned}$$

By the homogeneous function theorem [6], from this homogeneity condition we obtain.

LEMMA 4.1.  $A_K$  is independent of  $F_{j;\alpha\beta}^p$  with  $|\alpha| + |\beta| \geq 2$ ,  $A$  is independent of  $\Lambda_{jk;\gamma}^i$  with  $|\gamma| \geq 1$ ,  $A_K$  is independent of  $\Theta_{ip;\delta\sigma}^r$  with  $|\delta| + |\sigma| \geq 1$  and  $A_K$  is independent of  $\Theta_{sp;\delta\sigma}^r$  with  $|\delta| + |\sigma| \geq 1$ . Even,  $A_K$  is a linear combination with real coefficients of  $\Theta_{ip;(0)(0)}^r$ ,  $\Theta_{sp;(0)(0)}^r$  and  $F_{j;\alpha\beta}^p$  with  $|\alpha| + |\beta| = 1$ ,  $i, j = 1, \dots, m$ ,  $p, r, s = 1, \dots, n$ . In particular,  $A_K(\Gamma^0, \Lambda^0, \Theta^0)(0, 0) = 0$ .

Now, we prove the following two lemmas.

LEMMA 4.2. Let  $m \geq 2$  and  $n \geq 2$ . Any natural operator  $A$  in the sense of Definition 3.2 is fully determined by the collection of values

$$\begin{aligned} A^1 &:= A\left(\Gamma^0 + x^2 dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0\right)(0, 0), \\ A^2 &:= A\left(\Gamma^0 + y^1 dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0\right)(0, 0), \\ A^3 &:= A\left(\Gamma^0, \Lambda^0, \Theta^0 + \eta^1 dy^2 \otimes \frac{\partial}{\partial \eta^1}\right)(0, 0), \end{aligned}$$

where  $\Gamma^0, \Lambda^0, \Theta^0$  are the trivial connections.

PROOF. We know that the collection of maps  $A_K$  for  $K = 1, 2, \dots$  determines  $A$ . Then, using Lemma 4.1, it remains to prove that:

- (a)  $A(\Gamma^0 + x^{j_0} dx^{i_0} \otimes \frac{\partial}{\partial y^{p_0}}, \Lambda^0, \Theta^0)(0, 0)$  is determined by  $A^1$ ,
- (b)  $A(\Gamma^0 + y^{q_0} dx^{i_0} \otimes \frac{\partial}{\partial y^{p_0}}, \Lambda^0, \Theta^0)(0, 0)$  is determined by  $A^2$ ,
- (c)  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^{q_0} dx^{i_0} \otimes \frac{\partial}{\partial \eta^{p_0}})(0, 0)$  is determined by  $A^2$ , and
- (d)  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^{q_0} dy^{r_0} \otimes \frac{\partial}{\partial \eta^{p_0}})(0, 0)$  is determined by  $A^3$

We start with the proof of (a). By the invariance of  $A$  with respect to the (local)  $\mathcal{FM}_{m,n}$ -map  $k = (x^1, \dots, x^m, y^1, \dots, y^{p_0-1}, y^{p_0} + \frac{1}{2}(x^{i_0})^2, y^{p_0+1}, \dots, y^n)$ , from  $A(\Gamma^0, \Lambda^0, \Theta^0)(0, 0) = 0$ , we get  $A(\Gamma^0 + x^{i_0} dx^{i_0} \otimes \frac{\partial}{\partial y^{p_0}}, \Lambda^0, \Theta^0)(0, 0) = 0$  as  $k$  preserves  $\Lambda^0$  and  $\Theta^0$  and sends  $\Gamma^0$  into  $\Gamma^0 + x^{i_0} dx^{i_0} \otimes \frac{\partial}{\partial y^{p_0}}$ . If  $i_0 \neq j_0$ , there exists a permutation of coordinates sending  $A^1$  into  $A(\Gamma^0 + x^{i_0} dx^{j_0} \otimes \frac{\partial}{\partial y^{p_0}}, \Lambda^0, \Theta^0)(0, 0)$ . So, (a) is complete.

Now, we prove (b). By the invariance of  $A$  with respect to the  $\mathcal{FM}_{m,n}$ -map  $f := (x^1, \dots, x^m, y^1 + y^2, y^2, \dots, y^n)$  we see that  $A^0 = A(\Gamma^0 + (y^1 - y^2) dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0)(0, 0)$  is the image of  $A^2$  by  $f$ . Therefore  $A(\Gamma^0 + y^2 dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0)(0, 0) = A^2 - A^0$  is determined by  $A^2$ . So, using the invariance of  $A$  with respect to respective permutations of coordinates, we complete (b).

Next we prove (c). By the invariance of  $A$  with respect to the (local)  $\mathcal{FM}_{m,n}$ -map  $g := (x^1, \dots, x^m, y^1 + x^1 y^1, y^2, \dots, y^n)$  (then  $Vg = (g, \eta^1 + x^1 \eta^1, \eta^2, \dots, \eta^n)$ ) from  $A(\Gamma^0, \Lambda^0, \Theta^0) = 0$  we get that  $A(\Gamma^0 + y^1 dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0 + \eta^1 dx^1 \otimes \frac{\partial}{\partial \eta^1})(0, 0) = 0$ . More precisely,  $g$  preserves  $\Lambda^0$  and transforms  $\Gamma^0$  and  $\Theta^0$  into  $\Gamma^0 + \frac{y^1}{1+x^1} dx^1 \otimes \frac{\partial}{\partial y^1} = \Gamma^0 + y^1 dx^1 \otimes \frac{\partial}{\partial y^1} + \dots$  and  $\Theta^0 + \frac{\eta^1}{1+x^1} dx^1 \otimes \frac{\partial}{\partial \eta^1} = \Theta^0 + \eta^1 dx^1 \otimes \frac{\partial}{\partial \eta^1} + \dots$  (where the dots have the 1-jets at  $(0, 0)$  equal to 0) which can be replaced by  $\Gamma^0 + y^1 dx^1 \otimes \frac{\partial}{\partial y^1}$  and  $\Theta^0 + \eta^1 dx^1 \otimes \frac{\partial}{\partial \eta^1}$  in  $A$  in account of Lemma 4.1. Then

$A' = A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^1 dx^1 \otimes \frac{\partial}{\partial \eta^1})(0, 0)$  is determined by  $A^2$  (it is  $-A^2$ ). Then using the invariance of  $A$  with respect to  $f$  (the one of case (b)) we see that  $A'' = A(\Gamma^0, \Lambda^0, \Theta^0 + (\eta^1 - \eta^2) dx^1 \otimes \frac{\partial}{\partial \eta^1})(0, 0)$  is determined by  $A^2$  (it is the image of  $A'$  by  $f$ ). Then  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^2 dx^1 \otimes \frac{\partial}{\partial \eta^1})(0, 0)$  is determined by  $A^2$  (it is  $A' - A''$ ). Now, using the invariance of  $A$  with respect to respective permutations of coordinates, we complete (c).

Finally, we prove (d). By the invariance of  $A$  with respect to the (local)  $\mathcal{FM}_{m,n}$ -map  $H = (x^1, \dots, x^m, y^1 + \frac{1}{2}(y^1)^2, y^2, \dots, y^n)$  (then  $VH = (H, \eta^1 + y^1 \eta^1, \eta^2, \dots, \eta^n)$ ), from  $A(\Gamma^0, \Lambda^0, \Theta^0)(0, 0) = 0$  and Lemma 4.1 we get (using similar arguments as in (c)) that  $A''' = A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^1 dy^1 \otimes \frac{\partial}{\partial \eta^1})(0, 0) = 0$ . If  $q_0 \geq 2$ , by the invariance of  $A$  with respect to the  $\mathcal{FM}_{m,n}$ -map  $h = (x^1, \dots, x^m, y^1 + y^{q_0}, y^2, \dots, y^n)$ , we see that  $A'''' = A(\Gamma^0, \Lambda^0, \Theta^0 + (\eta^1 - \eta^{q_0}) dy^2 \otimes \frac{\partial}{\partial \eta^1})(0, 0)$  is determined by  $A^3$  (it is the image of  $A^3$  by  $h$ ). Then  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^{q_0} dy^2 \otimes \frac{\partial}{\partial \eta^1})(0, 0)$  is determined by  $A^3$  (it is  $A^3 - A''''$ ). In particular, for  $q_0 = 2$ ,  $A'''' = A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^2 dy^2 \otimes \frac{\partial}{\partial \eta^1})(0, 0)$  is determined by  $A^3$ . Then using the invariance of  $A$  with respect to  $f$  (the one of case (b)) from  $A''' = 0$  we get

$$A\left(\Gamma^0, \Lambda^0, \Theta^0 + (\eta^1 - \eta^2)(dy^1 - dy^2) \otimes \frac{\partial}{\partial \eta^1}\right)(0, 0) = 0,$$

and consequently  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^2 dy^1 \otimes \frac{\partial}{\partial \eta^1})(0, 0) = A'''' - A^3 + A''' = A'''' - A^3$  is determined by  $A^3$ . So, using the invariance of  $A$  with respect to respective permutations of coordinates, we complete (d).  $\square$

LEMMA 4.3. *Let  $m \geq 2$  and  $n \geq 3$ . Let  $A^1, A^2, A^3$  be the values from the last lemma. There are real numbers  $a_1, \dots, a_{12}$  such that*

$$\begin{aligned} A^1 &= a_1 \left( d_{(0,0)} x^2 \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} - d_{(0,0)} x^1 \otimes d_{(0,0)} x^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} \right), \\ A^2 &= a_2 \sum_{p=1}^n d_{(0,0)} x^1 \otimes d_{(0,0)} y^p \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} + a_3 \sum_{p=1}^n d_{(0,0)} y^p \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} \\ &\quad + a_4 d_{(0,0)} x^1 \otimes d_{(0,0)} y^1 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} + a_5 d_{(0,0)} y^1 \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} \\ &\quad + a_6 \sum_{i=1}^m d_{(0,0)} x^1 \otimes d_{(0,0)} x^i \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)} + a_7 \sum_{i=1}^m d_{(0,0)} x^i \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)}, \\ A^3 &= a_8 \sum_{p=1}^n d_{(0,0)} y^p \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} + a_9 \sum_{p=1}^n d_{(0,0)} y^2 \otimes d_{(0,0)} y^p \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} \\ &\quad + a_{10} \sum_{i=1}^m d_{(0,0)} x^i \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)} + a_{11} \sum_{i=1}^m d_{(0,0)} y^2 \otimes d_{(0,0)} x^i \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)} \\ &\quad + a_{12} \left( d_{(0,0)} y^2 \otimes d_{(0,0)} y^1 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} - d_{(0,0)} y^1 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} \right). \end{aligned}$$

PROOF. We start with the proof of the first formula. By the invariance of  $A$  with respect to the  $\mathcal{FM}_{m,n}$ -maps  $a_{t,\tau} := (t^1x^1, \dots, t^mx^m, \tau^1y^1, \dots, \tau^ny^n)$  for  $t^1 > 0, \dots, t^m > 0$  and  $\tau^1 > 0, \dots, \tau^n > 0$  we get easily

$$A^1 = b_1 d_{(0,0)} x^2 \otimes d_{(0,0)} x^1 \frac{\partial}{\partial y^1} \Big|_{(0,0)} + b_2 d_{(0,0)} x^1 \otimes d_{(0,0)} x^2 \frac{\partial}{\partial y^1} \Big|_{(0,0)}$$

for some real numbers  $b_1, b_2$ . Then (by the invariance of  $A$  with respect to permuting  $x^1$  and  $x^2$ )

$$\begin{aligned} A(\Gamma^0 + x^1 dx^2 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0)(0,0) \\ = b_1 d_{(0,0)} x^1 \otimes d_{(0,0)} x^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} + b_2 d_{(0,0)} x^2 \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)}. \end{aligned}$$

But by the invariance of  $A$  with respect to the (local)  $\mathcal{FM}_{m,n}$ -map  $(x^1, \dots, x^m, y^1 + x^1 x^2, y^3, \dots, y^n)$  from  $A(\Gamma^0, \Lambda^0, \Theta^0)(0,0) = 0$ , we get (using the similar arguments as in (a) of the proof of Lemma 4.2) that  $A(\Gamma^0 + x^2 dx^1 \otimes \frac{\partial}{\partial y^1} + x^1 dx^2 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0)(0,0) = 0$ . Therefore  $b_1 = -b_2$ . That is why, the first formula of the lemma is complete.

Now, we prove the second formula of the lemma. By the invariance of  $A$  with respect to  $a_{t,\tau}$  (the same as above), we get immediately

$$\begin{aligned} A^2 = \sum_{p=1}^n b_p d_{(0,0)} y^p \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} + \sum_{p=1}^n c_p d_{(0,0)} x^1 \otimes d_{(0,0)} y^p \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} \\ + \sum_{i=1}^m d_i d_{(0,0)} x^i \otimes d_{(0,0)} x^1 \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)} + \sum_{i=1}^m e_i d_{(0,0)} x^1 \otimes d_{(0,0)} x^i \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)}. \end{aligned}$$

Then by the invariance of  $A$  with respect to respective permutation of coordinates, we deduce  $b_2 = \dots = b_n$ ,  $c_2 = \dots = c_n$ ,  $d_2 = \dots = d_m$ ,  $e_2 = \dots = e_m$ . Then  $A^2 =$  the right-hand side of the second formula of the lemma  $+ b dx^1 \otimes dx^1 \otimes \frac{\partial}{\partial x^1}$ . Now, by the invariance of  $A$  with respect to  $(x^1, x^2 + x^1, x^3, \dots, x^m, y^1, \dots, y^n)$  one can obtain that  $b = 0$ . That is why, the second formula of the lemma is true.

Finally we prove the last formula of the lemma. By the invariance of  $A$  with respect to  $a_{t,\tau}$  (the same as above) we get immediately

$$\begin{aligned} A^3 = \sum_{p=1}^n b_p d_{(0,0)} y^2 \otimes d_{(0,0)} y^p \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} + \sum_{p=1}^n c_p d_{(0,0)} y^p \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} \\ + \sum_{i=1}^m d_i d_{(0,0)} y^2 \otimes d_{(0,0)} x^i \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)} + \sum_{i=1}^m e_i d_{(0,0)} x^i \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)}. \end{aligned}$$

Then by the invariance of  $A$  with respect to respective permutation of coordinates, we deduce  $b_3 = \dots = b_n$ ,  $c_3 = \dots = c_n$ ,  $d_1 = \dots = d_m$  and  $e_1 = \dots = e_m$ . Then

$$(*) \quad A^3 = \lambda_1 d_{(0,0)} y^2 \otimes d_{(0,0)} y^1 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)}$$



$$\begin{aligned}
& + \lambda_2 d_{(0,0)} y^1 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} + \lambda_3 d_{(0,0)} y^2 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^2} \Big|_{(0,0)} \\
& + \lambda_4 \sum_{p=1}^n d_{(0,0)} y^p \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} + \lambda_5 \sum_{p=1}^n d_{(0,0)} y^2 \otimes d_{(0,0)} y^p \otimes \frac{\partial}{\partial y^p} \Big|_{(0,0)} \\
& + \lambda_6 \sum_{i=1}^m d_{(0,0)} x^i \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)} + \lambda_7 \sum_{i=1}^m d_{(0,0)} y^2 \otimes d_{(0,0)} x^i \otimes \frac{\partial}{\partial x^i} \Big|_{(0,0)}.
\end{aligned}$$

Then by invariance of  $A$  with respect to  $\mathcal{FM}_{m,n}$ -map  $(x^1, \dots, x^m, y^1 - y^2, y^2, \dots, y^n)$ , from (\*) we deduce

$$\begin{aligned}
A^3 + A \left( \Gamma^0, \Lambda^0, \Theta^0 + \eta^2 dy^2 \otimes \frac{\partial}{\partial \eta^1} \right) (0, 0) &= A^3 + \lambda_1 d_{(0,0)} y^2 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} \\
&+ \lambda_2 d_{(0,0)} y^2 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} - \lambda_3 d_{(0,0)} y^2 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)}.
\end{aligned}$$

On the other hand, from the invariance of  $A$  with respect to the (local)  $\mathcal{FM}_{m,n}$ -map  $G = (x^1, \dots, x^m, y^1 + \frac{1}{2}(y^2)^2, y^2, \dots, y^n)$  (then  $VG = (G, \eta^1 + y^2 \eta^2, \eta^2, \dots, \eta^n)$ ) from  $A(\Gamma^0, \Lambda^0, \Theta^0)(0, 0) = 0$  and Lemma 4.1 (using similar arguments as in (c) of the proof of Lemma 4.2), we get  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^2 dy^2 \otimes \frac{\partial}{\partial \eta^1})(0, 0) = 0$ . So,  $\lambda_1 + \lambda_2 - \lambda_3 = 0$ . Further, from the invariance of  $A$  with respect to  $(x^1, \dots, x^m, y^1 - y^3, y^2, \dots, y^n)$  (we assume  $n \geq 3$ ) from (\*) we get (after cancelling  $A^3$ )

$$\begin{aligned}
A \left( \Gamma^0, \Lambda^0, \Theta^0 + \eta^3 dy^2 \otimes \frac{\partial}{\partial \eta^1} \right) (0, 0) &= \lambda_1 d_{(0,0)} y^2 \otimes d_{(0,0)} y^3 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} \\
&+ \lambda_2 d_{(0,0)} y^3 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)}.
\end{aligned}$$

Then from the invariance of  $A$  with respect to the switching  $y^2$  and  $y^3$  we get

$$\begin{aligned}
A \left( \Gamma^0, \Lambda^0, \Theta^0 + \eta^2 dy^3 \otimes \frac{\partial}{\partial \eta^1} \right) (0, 0) &= \lambda_1 d_{(0,0)} y^3 \otimes d_{(0,0)} y^2 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)} \\
&+ \lambda_2 d_{(0,0)} y^2 \otimes d_{(0,0)} y^3 \otimes \frac{\partial}{\partial y^1} \Big|_{(0,0)}.
\end{aligned}$$

On the other hand from the invariance of  $A$  with respect to  $K = (x^1, \dots, x^m, y^1 + y^2 y^3, y^2, \dots, y^n)$  (then  $VK = (K, \eta^1 + y^2 \eta^3 + y^3 \eta^2, \eta^2, \dots, \eta^n)$ ) from Lemma 4.1 and  $A(\Gamma^0, \Lambda^0, \Theta^0)(0, 0) = 0$ , we get (using similar arguments as in (c) of the proof of Lemma 4.2) that  $A(\Gamma^0, \Lambda^0, \Theta^0 + \eta^3 dy^2 \otimes \frac{\partial}{\partial \eta^1} + \eta^2 dy^3 \otimes \frac{\partial}{\partial \eta^1})(0, 0) = 0$ . So,  $\lambda_1 = -\lambda_2$  (and then  $\lambda_3 = 0$ ). That is why, the last formula of the lemma holds.  $\square$

From Lemmas 4.2 and 4.3 it follows immediately the following proposition.

**PROPOSITION 4.1.** *If  $m \geq 2$  and  $n \geq 3$ , the vector space of all natural operators in the sense of Definition 3.2 is of dimension not more than 12.*

## 5. Linear independence

We prove the following proposition.

**PROPOSITION 5.1.** *Let  $m \geq 2$  and  $n \geq 3$ . The natural operators  $\tau_i$  ( $i = 1, \dots, 12$ ) in the sense of Definition 3.2 from Examples 2.1–2.12 are linearly independent.*

**PROOF.** By Lemma 4.2, it is sufficient to study the values  $A^1, A^2, A^3$  from Lemma 4.2 for  $A = \tau_i$ ,  $i = 1, \dots, 12$ . To compute these values, we use Lemma 1.1.

*The case  $\Sigma = (\Gamma^0 + x^2 dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0)$ .* In this case, we have (in the notation of Lemma 1.1)  $F_1^1(x, y) = x^2$  and other  $F_i^p(x, y) = 0$ ,  $\Lambda_{ij}^k = 0$ ,  $\Theta_{sj}^p = 0$ ,  $\Theta_{qs}^p = 0$ . Then (by Lemma 1.1)  $d\eta^1 = \xi^1 dx^2$  and other  $d\eta^p = 0$ , and  $d\xi^i = 0$ . Then (modulo signum)

$$\text{Tor}(\Psi(\Sigma))(0, 0) = d_{(0,0)}x^1 \otimes d_{(0,0)}x^2 \otimes \frac{\partial}{\partial y^1}|_{(0,0)} - d_{(0,0)}x^2 \otimes d_{(0,0)}x^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)}.$$

Hence (modulo signum)

$$\tau_1(\Sigma)(0, 0) = d_{(0,0)}x^1 \otimes d_{(0,0)}x^2 \otimes \frac{\partial}{\partial y^1}|_{(0,0)} - d_{(0,0)}x^2 \otimes d_{(0,0)}x^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)}$$

and  $\tau_i(\Sigma)(0, 0) = 0$  for  $i = 2, \dots, 12$ .

*The case  $\Sigma = (\Gamma^0 + y^1 dx^1 \otimes \frac{\partial}{\partial y^1}, \Lambda^0, \Theta^0)$ .* In this case, by Lemma 1.1,  $d\eta^1 = \xi^1 dy^1$  and other  $d\eta^p = 0$ , and  $d\xi^i = 0$ . Then (modulo signum)

$$\text{Tor}(\Psi(\Sigma))(0, 0) = d_{(0,0)}x^1 \otimes d_{(0,0)}y^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)} - d_{(0,0)}y^1 \otimes d_{(0,0)}x^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)}.$$

Then  $\tau_1(\Sigma)(0, 0) = 0$  and (modulo signum)

$$\tau_2(\Sigma)(0, 0) = d_{(0,0)}x^1 \otimes d_{(0,0)}y^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)},$$

$$\tau_3(\Sigma)(0, 0) = \sum_{i=1}^m d_{(0,0)}x^i \otimes d_{(0,0)}x^1 \otimes \frac{\partial}{\partial x^i}|_{(0,0)},$$

$$\tau_4(\Sigma)(0, 0) = \sum_{i=1}^m d_{(0,0)}x^1 \otimes d_{(0,0)}x^i \otimes \frac{\partial}{\partial x^i}|_{(0,0)},$$

$$\tau_5(\Sigma)(0, 0) = \sum_{p=1}^n d_{(0,0)}x^1 \otimes d_{(0,0)}y^p \otimes \frac{\partial}{\partial y^p}|_{(0,0)},$$

$$\tau_6(\Sigma)(0, 0) = \sum_{p=1}^n d_{(0,0)}y^p \otimes d_{(0,0)}x^1 \otimes \frac{\partial}{\partial y^p}|_{(0,0)},$$

$\tau_i(\Sigma)(0, 0) = 0$  for  $i = 7, \dots, 11$  and (modulo signum)  $\tau_{12}(\Sigma)(0, 0) = d_{(0,0)}y^1 \otimes d_{(0,0)}x^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)}$ .

The case  $\Sigma = (\Gamma^0, \Lambda^0, \Theta^0 + \eta^1 dy^2 \otimes \frac{\partial}{\partial \eta^1})$ . In this case, by Lemma 1.1,  $d\eta^1 = \eta^1 dy^2$  and  $d\eta^p = 0$  for other  $p$ , and  $d\xi^i = 0$ . Then (modulo signum)

$$\text{Tor}(\Psi(\Sigma))(0, 0) = d_{(0,0)}y^1 \otimes d_{(0,0)}y^2 \otimes \frac{\partial}{\partial y^1}|_{(0,0)} - d_{(0,0)}y^2 \otimes d_{(0,0)}y^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)}$$

. Then  $\tau_i(\Sigma)(0, 0) = 0$  for  $i = 1, \dots, 6$ , and (modulo signum)

$$\tau_7(\Sigma)(0, 0) = d_{(0,0)}y^2 \otimes d_{(0,0)}y^1 \otimes \frac{\partial}{\partial y^1}|_{(0,0)} - d_{(0,0)}y^1 \otimes d_{(0,0)}y^2 \otimes \frac{\partial}{\partial y^1}|_{(0,0)},$$

$$\tau_8(\Sigma)(0, 0) = \sum_{i=1}^m d_{(0,0)}x^i \otimes d_{(0,0)}y^2 \otimes \frac{\partial}{\partial x^i}|_{(0,0)},$$

$$\tau_9(\Sigma)(0, 0) = \sum_{i=1}^m d_{(0,0)}y^2 \otimes d_{(0,0)}x^i \otimes \frac{\partial}{\partial x^i}|_{(0,0)},$$

$$\tau_{10}(\Sigma)(0, 0) = \sum_{p=1}^n d_{(0,0)}y^p \otimes d_{(0,0)}y^2 \otimes \frac{\partial}{\partial y^p}|_{(0,0)},$$

$$\tau_{11}(\Sigma)(0, 0) = \sum_{p=1}^n d_{(0,0)}y^2 \otimes d_{(0,0)}y^p \otimes \frac{\partial}{\partial y^p}|_{(0,0)}$$

and  $\tau_{12}(\Sigma)(0, 0) = 0$ .

Now, it is easily seen that the natural operators  $\tau_1, \dots, \tau_{12}$  are linearly independent. Proposition 5.1 is complete.  $\square$

## 6. The main result

From Propositions 4.1 and 5.1 the main theorem of the paper follows immediately.

**THEOREM 6.1.** *Let  $m \geq 2$  and  $n \geq 3$ . Any natural operator  $A$  in the sense of Definition 3.1 is of the form  $A(\Gamma, \Lambda, \Theta) = \Psi(\Gamma, \Lambda, \Theta) + \sum_i \lambda^i \tau_i(\Gamma, \Lambda, \Theta)$  for some (uniquely determined by  $A$ ) real numbers  $\lambda^i$ ,  $i = 1, \dots, 12$ , where  $\tau_i$  are the natural operators described in Examples 2.1–2.12 and  $\Psi(\Gamma, \Lambda, \Theta)$  is the connection from Introduction.*

## 7. Final remarks

Let  $Y \rightarrow M$  be a fibred manifold with  $m$ -dimensional basis and  $n$ -dimensional fibres. Let  $E \rightarrow M$  be a vector bundle with the same base  $M$  and  $n$ -dimensional fibres (the same  $n$ ). A vertical parallelism on  $Y \rightarrow M$  is a vector bundle isomorphism  $\Phi : Y \times_M E \rightarrow VY$  covering the identity map of  $Y$ , i.e., a system of parallelism  $\Phi_x : Y_x \times E_x \rightarrow TY_x$ ,  $x \in M$ . Let  $\Gamma$  be a general connection on  $Y \rightarrow M$ ,  $\Lambda$  be a classical linear connection on  $M$ ,  $\Phi : Y \times_M E \rightarrow VY$  be a vertical parallelism on  $Y \rightarrow M$  and  $\Delta$  be a linear connection on  $E \rightarrow M$ .

In [5], Kolář constructed the classical linear connection  $(\Gamma, \Lambda, \Phi, \Delta)$  on  $Y$  depending canonically on  $(\Gamma, \Lambda, \Phi, \Delta)$ . Using our connection  $\Psi(\Gamma, \Lambda, \Theta)$  from Introduction, we can reobtain the connection  $(\Gamma, \Lambda, \Phi, \Delta)$  by Kolář as follows. The system  $(\Delta, \Phi)$  determines a vertical classical linear connection  $\Theta = \Theta(\Delta, \Phi) : VY \rightarrow$

$J^1(VY \rightarrow Y)$  on  $Y \rightarrow M$ . Indeed, for any point  $v = \Phi(y, v_0) \in V_y Y$ ,  $y \in Y_x$ ,  $v_0 \in E_x$ ,  $x = p(y)$ , we take a section  $\sigma : M \rightarrow E$  such that  $j_x^1 \sigma = \Delta(v_0)$  and we define  $\Theta(v) = j_y^1(\varphi(\sigma))$ , where  $\varphi(\sigma) : Y \rightarrow VY$  is a vertical vector field given by  $\varphi(\sigma)(y) = \Phi(y, \sigma(p(y)))$ . Now, if we additionally use the connections  $\Gamma$  and  $\Lambda$  we can produce the classical linear connection  $\Psi(\Gamma, \Lambda, \Theta(\Delta, \Phi))$ , which (as easily to see) coincides with the Kolář connection  $(\Gamma, \Lambda, \Phi, \Delta)$ . So, the construction of  $\Psi(\Gamma, \Lambda, \Theta)$  from Introduction is a generalization of the construction of  $(\Gamma, \Lambda, \Phi, \Delta)$  by Kolář [5].

If  $Y = E \rightarrow M$ ,  $\Gamma = \Delta$  and  $\Phi$  is the canonical vertical parallelism, the connection by Kolář (and then the connection  $\Psi$  from Introduction) is a generalization of the classical linear connection on  $E$  from  $\Delta$  by means of  $\Lambda$  presented by Gancarzewicz [3]. If  $P = Y \rightarrow M$  is a principal  $G$ -bundle,  $\Gamma$  is principal (i.e., right invariant),  $E \rightarrow M$  is the usual  $\mathcal{L}(G)$ -algebra bundle of  $P$  and  $\Phi$  is the canonical vertical parallelism, the connection by Kolář (and then the connection  $\Psi$  from Introduction) is a generalization of the classical linear connection  $N(\Gamma, \Lambda)$  on  $P$  considered in [6, p. 415], see [5].

In [7], we described all classical linear connections  $A(\Gamma, \Lambda, \Phi, \Delta)$  on  $Y$  canonically depending on the system  $(\Gamma, \Lambda, \Phi, \Delta)$ . Thus the present paper can be treated as the generalization of [7]. In [7], we showed that all  $A(\Gamma, \Lambda, \Phi, \Delta)$  form the 12-parameter family, too.

Let us also remark, why we must use an auxiliary object (in the paper we use  $\Theta$ ) to construct from  $\Gamma$  and  $\Lambda$  a classical linear connection  $A(\Gamma, \Lambda)$  on  $Y$ . In the other case we would have a classical linear connection  $A(\Gamma^0, \Lambda^0)$  on  $\mathbf{R}^m \times \mathbf{R}^n$  which would be  $GL(m) \times \text{Diff}(\mathbf{R}^n, \mathbf{R}^n)$  invariant. But this is impossible as the group of affine transformations of a classical linear connection is a (finite dimensional) Lie group.

Classifications of constructions on connections has been studied in many papers, e.g., [1, 2, 4, 6, 7], e.t.c.

## References

1. J. Dębecki, *Affine liftings of torsion-free connections to Weil bundles*, Colloq. Math. **114**(1) (2009), 1–8.
2. M. Doupovec, W.M. Mikulski, *Reduction theorems for principal and classical connections*, Acta Math. Sinica (E-S) **26**(1)(2010), 169–184.
3. J. Gancarzewicz, *Horizontal lifts of linear connections to the natural vector bundles*, Research Notes in Math. (Pitman) **121** 1985, 318–341.
4. J. Janyška, J. Vondra, *Natural principal connections on the principal gauge prolongation of a principal bundle*, Rep. Math. Phys. **64**(3) (2009), 395–415.
5. I. Kolář, *Induced connections on total spaces of fibred bundles*, Int. J. Geom. Methods Mod. Phys. **7**(4) (2010), 705–711.
6. I. Kolář, P.W. Michor, J. Slovák, *Natural Operations in Differential Geometry*, Springer-Verlag, 1993.
7. W.M. Mikulski, *On the Kolář connection*, Arch. Math. (Brno) **49** (2013), 223–240.

Institute of Mathematics, Jagiellonian University  
 Łojasiewicza 6, Cracow, Poland  
 Wlodzimierz.Mikulski@im.uj.edu.pl

(Received 26 05 2014)