

## CONVERGENCE THEOREMS OF A SCHEME FOR $I$ -ASYMPTOTICALLY QUASI-NONEXPANSIVE TYPE MAPPING IN BANACH SPACE

Seyit Temir

ABSTRACT. Let  $X$  be a Banach space. Let  $K$  be a nonempty subset of  $X$ . Let  $T : K \rightarrow K$  be an  $I$ -asymptotically quasi-nonexpansive type mapping and  $I : K \rightarrow K$  be an asymptotically quasi-nonexpansive type mappings in the Banach space. Our aim is to establish the necessary and sufficient conditions for the convergence of the Ishikawa iterative sequences with errors of an  $I$ -asymptotically quasi-nonexpansive type mapping in Banach spaces to a common fixed point of  $T$  and  $I$ . Also, we study the convergence of the Ishikawa iterative sequences to common fixed point for nonself  $I$ -asymptotically quasi-nonexpansive type mapping in Banach spaces.

The results presented in this paper extend and generalize some recent work of Chang and Zhou [1], Wang [19], Yao and Wang [20] and many others.

### 1. Introduction

Let  $X$  be a real Banach space,  $K$  be a nonempty subset of Banach space and  $T, I : K \rightarrow K$ . Let  $F(T) = \{x \in K : Tx = x\}$  and  $F(I) = \{x \in K : Ix = x\}$  denote the set of fixed points of mappings  $T$  and  $I$ , respectively. Recall some definitions and notations.  $T$  is called nonexpansive if  $\|Tx - Ty\| \leq \|x - y\|$  for all  $x, y \in K$ . The quasi-nonexpansive mappings defined as the following were studied by Diaz and Metcalf [4] and Dotson [5] in Banach spaces.  $T$  is called a quasi-nonexpansive mapping if  $F(T) \neq \emptyset$  and  $\|Tx - p\| \leq \|x - p\|$  for all  $x \in K$  and  $p \in F(T)$ . The concept of asymptotically nonexpansiveness defined as the following was introduced by Goebel and Kirk [7].  $T$  is called asymptotically quasi-nonexpansive mapping if  $F(T) \neq \emptyset$  and there exists a sequence  $\{k_n\} \subset [1, \infty)$  with  $\lim_{n \rightarrow \infty} k_n = 1$  such that  $\|T^n x - p\| \leq k_n \|x - p\|$  for all  $x \in K$  and  $p \in F(T)$  and  $n \geq 1$ . Let  $X$  be a Banach space and  $K$  be a nonempty subset of the Banach space. Let  $T, I : K \rightarrow K$  be two mappings.  $T$  is called  $I$ -nonexpansive if  $\|Tx - Ty\| \leq \|Ix - Iy\|$  for all  $x, y \in K$ .  $T$  is called  $I$ -quasi-nonexpansive if  $F(T) \cap F(I) \neq \emptyset$  and  $\|Tx - p\| \leq \|Ix - p\|$  for all  $x \in K$  and  $p \in F(T) \cap F(I)$ .

---

2010 *Mathematics Subject Classification*: 47H09; 47H10.

*Key words and phrases*:  $I$ -asymptotically quasi-nonexpansive type mapping, nonself  $I$ -asymptotically quasi-nonexpansive type mapping, Ishikawa iterative schemes.

Communicated by Stevan Pilipović.

From the above definitions, it follows that if  $F(T) \cap F(I)$  is nonempty, an  $I$ -nonexpansive mapping must be  $I$  quasi-nonexpansive, and linear  $I$  quasi-nonexpansive mappings are  $I$ -nonexpansive mappings. But it is easily seen that there exist nonlinear continuous  $I$  quasi-nonexpansive mappings which are not  $I$ -nonexpansive.  $T$  is called  $I$ -asymptotically quasi-nonexpansive if there exists a sequence  $\{k_n\} \subset [1, \infty)$  with  $\lim_{n \rightarrow \infty} k_n = 1$  such that  $\|T^n x - p\| \leq k_n \|I^n x - p\|$  for all  $x \in K$  and  $p \in F(T) \cap F(I)$  and  $n \geq 1$ .  $T$  is called  $I$ -asymptotically nonexpansive type mapping if  $\limsup_{n \rightarrow \infty} \{\sup\{\|T^n x - T^n y\| - \|I^n x - I^n y\|\}\} \leq 0$  for all  $x, y \in K$ .

$T$  is called  $I$ -asymptotically quasi-nonexpansive type if  $F(T) \cap F(I) \neq \emptyset$  and

$$(1.1) \quad \limsup_{n \rightarrow \infty} \{\sup\{\|T^n x - p\| - \|I^n x - p\|\}\} \leq 0$$

for all  $x \in K$  and  $p \in F(T) \cap F(I)$ .

$I$  is called asymptotically quasi-nonexpansive type if  $F(I) \neq \emptyset$  and

$$(1.2) \quad \limsup_{n \rightarrow \infty} \{\sup\{\|I^n x - p\| - \|x - p\|\}\} \leq 0$$

for all  $x \in K$  and  $p \in F(I)$ .

From the above definitions, it follows that if  $F(I)$  is nonempty, quasi-nonexpansive mappings, asymptotically nonexpansive mappings, asymptotically quasi-nonexpansive mappings and asymptotically nonexpansive type mappings all are special cases of asymptotically quasi-nonexpansive type mappings.

Let  $\{x_n\}$  be of the Ishikawa iterative scheme [8] associated with  $T$ ,  $x_0 \in K$ ,

$$\begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_n T x_n \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n \end{aligned}$$

for every  $n \in \mathbb{N}$ , where  $0 \leq \alpha_n, \beta_n \leq 1$ .

Let  $S, T : K \rightarrow K$  be two mappings. In 2006, Lan [9] introduced the following iterative scheme with errors. The sequence  $x_n$  in  $K$  defined by

$$(1.3) \quad \begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_n T^n x_n + \psi_n \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n S^n y_n + \varphi_n \end{aligned}$$

for every  $n \in \mathbb{N}$ , where  $0 \leq \{\alpha_n\}, \{\beta_n\} \leq 1$  and  $\{\varphi_n\}, \{\psi_n\}$  are two sequences in  $K$ .

The iterative approximation problems for nonexpansive mapping, asymptotically nonexpansive mapping and asymptotically quasi-nonexpansive mapping were studied Ghosh and Debnath [6], Goebel and Kirk [7], Liu [10, 11], Petryshyn and Williamson [13] in the settings of Hilbert spaces and uniformly convex Banach spaces. The strong and weak convergences of the sequence of Mann iterates to a fixed point of quasi-nonexpansive maps were studied by Petryshyn and Williamson [13]. Subsequently, the convergence of Ishikawa iterates of quasi-nonexpansive mappings in Banach spaces were discussed by Ghosh and Debnath [6]. The above results and some necessary and sufficient conditions for Ishikawa iterative sequences obtained to converge to a fixed point for asymptotically quasi-nonexpansive mappings were extended by Liu [10]. In [11], the results of Liu [10] were extended and some sufficient and necessary conditions for Ishikawa iterative sequences of

asymptotically quasi-nonexpansive mappings with error member to converge to fixed points were proved. Recently, Temir and Gul [17] obtained the weakly almost convergence theorems for I-asymptotically quasi-nonexpansive mapping in a Hilbert space. In [20], Yao and Wang established the strong convergence of an iterative scheme with errors involving I-asymptotically quasi-nonexpansive mappings in a uniformly convex Banach space. Temir [18], studied the convergence to common fixed point of Ishikawa iterative process of generalized I-asymptotically quasi-nonexpansive mappings to common fixed point in Banach space. In [1], the convergence theorems for Ishikawa iterative sequences with mixed errors of asymptotically quasi-nonexpansive type mappings in Banach spaces were studied.

**2. Preliminaries and notations**

We first recall the following definitions. A Banach space  $X$  is said to satisfy Opial's condition [12] if, for each sequence  $\{x_n\}$  in  $X$ , the condition  $x_n \rightarrow x$  implies

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|$$

for all  $y \in X$  with  $y \neq x$ . It is well known from [12] that all  $l_r$  spaces for  $1 < r < \infty$  have this property. However, the  $L_r$  space do not have unless  $r = 2$ .

In order to prove the main results of this paper, we need the following lemmas.

LEMMA 2.1. [16] *Let  $\{a_n\}$ ,  $\{b_n\}$  be sequences of nonnegative real numbers satisfying the following conditions:  $\forall n \geq 1, a_{n+1} \leq a_n + b_n$ , where  $\sum_{n=1}^{\infty} b_n < \infty$ . Then  $\lim_{n \rightarrow \infty} a_n$  exists.*

LEMMA 2.2. [15] *Let  $K$  be a nonempty closed bounded convex subset of a uniformly convex Banach space  $X$  and  $\{\alpha_n\} \subseteq [\epsilon, 1 - \epsilon] \subset (0, 1)$ . Let  $\{x_n\}$  and  $\{y_n\}$  be two sequences in  $K$  such that  $\limsup_{n \rightarrow \infty} \|x_n\| \leq c$ ,  $\limsup_{n \rightarrow \infty} \|y_n\| \leq c$ , and  $\limsup_{n \rightarrow \infty} \|\alpha_n x_n + (1 - \alpha_n)y_n\| = c$  for some  $c \geq 0$ . Then  $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$ .*

LEMMA 2.3. [2] *Let  $X$  be a uniformly convex Banach space,  $K$  a nonempty closed convex subset of  $X$  and  $T : K \rightarrow K$  an asymptotically nonexpansive mapping with a sequence  $\{k_n\} \subset [1, \infty)$  and  $\lim_{n \rightarrow \infty} k_n = 1$ . Then  $E - T$  is semi-closed (demi-closed) at zero, i.e., for each sequence  $\{x_n\}$  in  $K$ , if  $\{x_n\}$  converges weakly to  $q \in K$  and  $(E - T)\{x_n\}$  converges strongly to 0, then  $(E - T)q = 0$ .*

**3. Convergence theorems for I-asymptotically quasi-nonexpansive type mapping**

In this section,  $X$  is a Banach space and  $K$  is its nonempty subset. Let  $T, I : K \rightarrow K$  be two mappings, where  $T$  is an I-asymptotically quasi-nonexpansive type mapping and  $I : K \rightarrow K$  is an asymptotically quasi-nonexpansive type mapping. We study the strong and weak convergences of the sequence of Ishikawa iterates with mixed errors to a common fixed point of  $T$  and  $I$ .

THEOREM 3.1. *Let  $X$  be a Banach space,  $K$  its nonempty subset, and  $T, I : K \rightarrow K$  two mappings. Let  $T$  be an I-asymptotically quasi-nonexpansive type and  $I$  be an asymptotically quasi-nonexpansive type in the Banach space satisfying*

(3.1) 
$$\|Tx - p\| \leq L\|Ix - p\|$$

for all  $x \in K$  and  $p \in F(T) \cap F(I)$ , where  $L > 0$  is a constant and

$$(3.2) \quad \|Ix - p\| \leq \Gamma \|x - p\|$$

for all  $x \in K$  and  $p \in F(I)$ , where  $\Gamma > 0$  is a constant. Write  $I : K \rightarrow K$  instead of  $S : K \rightarrow K$  in (1.3) and get

$$(3.3) \quad \begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_n T^n x_n + \psi_n \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n I^n y_n + \varphi_n \end{aligned}$$

for every  $n \in \mathbb{N}$ , where  $0 \leq \{\alpha_n\}, \{\beta_n\} \leq 1$  and  $\{\varphi_n\}, \{\psi_n\}$  be two sequences in  $K$  satisfying: (i)  $\sum_{n=1}^{\infty} \alpha_n < \infty$ ; (ii)  $\{\psi_n\}$  is bounded,  $\varphi_n = \varphi'_n + \varphi''_n$ ,  $n \in \mathbb{N}$  and  $\sum_{n=1}^{\infty} \|\varphi'_n\| < \infty$ ,  $\|\varphi''_n\| = o(\alpha_n)$ .

Then  $\{x_n\}$  converges strongly to a common fixed point of  $T$  and  $I$  in  $K$  iff

$$(3.4) \quad \liminf_{n \rightarrow \infty} d(x_n, F(T) \cap F(I)) = 0.$$

LEMMA 3.1. Suppose all conditions in Theorem 3.1 are satisfied; then for  $\varepsilon > 0$ , there exists a positive integer  $n_0$  and  $M > 0$  such that

$$\|x_{n+1} - p\| \leq \|x_n - p\| + \alpha_n M + \|\varphi'_n\|$$

for all  $p \in F(T) \cap F(I)$ ,  $n \geq n_0$  and

$$\|x_{n+m} - p\| \leq \|x_n - p\| + M \sum_{i=n}^{n+m-1} \alpha_i + \sum_{i=n}^{n+m-1} \|\varphi'_i\|,$$

for all  $p \in F(T) \cap F(I)$ ,  $n \geq n_0$ ,  $\forall m \geq 1$ , where  $M = \sup_{n \geq 0} \{\varepsilon_n + \|\psi_n\|\} + 3\varepsilon \leq \infty$ , and  $\varepsilon_n$  is a sequence with  $\varepsilon_n > 0$  and  $\varepsilon_n \rightarrow 0$  such that  $\|\varphi''_n\| = \varepsilon_n \alpha_n$ .

PROOF. For  $p \in F(T) \cap F(I)$ , from (3.3), we have

$$(3.5) \quad \begin{aligned} \|x_{n+1} - p\| &= \|(1 - \alpha_n)(x_n - p) + \alpha_n(I^n y_n - p) + \varphi_n\| \\ &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n\|I^n y_n - p\| + \|\varphi_n\| \\ &= (1 - \alpha_n)\|x_n - p\| + \alpha_n\{\|I^n y_n - p\| - \|y_n - p\|\} \\ &\quad + \alpha_n\|y_n - p\| + \|\varphi_n\| \end{aligned}$$

Now we consider the second term on the right-hand side of (3.5). From (1.1) and (1.2), for any given  $\varepsilon > 0$ , there exists a positive integer  $n_0$  such that  $n \geq n_0$ , so we have

$$\begin{aligned} \sup_{x \in K, p \in F(T) \cap F(I)} \{\|T^n x - p\| - \|I^n x - p\|\} &< \varepsilon, \\ \sup_{x \in K, p \in F(I)} \{\|I^n x - p\| - \|x - p\|\} &< \varepsilon. \end{aligned}$$

Therefore, in particular, we have

$$(3.6) \quad \{\|T^n x_n - p\| - \|I^n x_n - p\|\} < \varepsilon,$$

for all  $p \in F(T) \cap F(I)$  and  $\forall n \geq n_0$ .

$$(3.7) \quad \{\|I^n y_n - p\| - \|y_n - p\|\} < \varepsilon,$$

for all  $p \in F(I)$  and  $\forall n \geq n_0$ . From (3.7), we have

$$(3.8) \quad \|x_{n+1} - p\| \leq (1 - \alpha_n)\|x_n - p\| + \alpha_n\varepsilon + \alpha_n\|y_n - p\| + \|\varphi_n\|$$

Consider the third term on the right-hand side of (3.8). From (3.6) and (3.7), we get

$$(3.9) \quad \begin{aligned} \|y_n - p\| &= \|(1 - \beta_n)(x_n - p) + \beta_n(T^n x_n - p) + \psi_n\| \\ &\leq (1 - \beta_n)\|x_n - p\| + \beta_n\{\|T^n x_n - p\| - \|I^n x_n - p\|\} \\ &\quad + \beta_n\{\|I^n x_n - p\| - \|x_n - p\|\} + \beta_n\|x_n - p\| + \|\psi_n\| \\ &\leq (1 - \beta_n)\|x_n - p\| + 2\beta_n\varepsilon + \beta_n\|x_n - p\| + \|\psi_n\| \\ &= \|x_n - p\| + 2\beta_n\varepsilon + \|\psi_n\| \end{aligned}$$

Now consider the fourth term on the right side of (3.5); we have  $\|\varphi_n\| \leq \|\varphi'_n\| + \|\varphi''_n\|$ ,  $\forall n \geq 0$ . Substituting (3.9) into (3.8), we have

$$\begin{aligned} \|x_{n+1} - p\| &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n\varepsilon + \alpha_n\{\|x_n - p\| + 2\beta_n\varepsilon + \|\psi_n\|\} + \|\varphi_n\| \\ &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n\varepsilon + \alpha_n\|x_n - p\| \\ &\quad + 2\alpha_n\beta_n\varepsilon + \alpha_n\|\psi_n\| + \|\varphi'_n\| + \|\varphi''_n\| \\ &= \|x_n - p\| + \alpha_n\varepsilon(1 + 2\beta_n) + \alpha_n\varepsilon_n + \alpha_n\|\psi_n\| + \|\varphi'_n\| \end{aligned}$$

Taking  $M = \sup_{n \geq 0} \{\varepsilon_n + \|\psi_n\|\} + 3\varepsilon$  we obtain

$$(3.10) \quad \|x_{n+1} - p\| \leq \|x_n - p\| + \alpha_n M + \|\varphi'_n\|$$

for all  $p \in F(T) \cap F(I)$ ,  $n \geq n_0$ . Writing  $n + m - 1$  instead of  $n$  in inequality (3.10), for  $m \geq 1$ , we get

$$\begin{aligned} \|x_{n+m} - p\| &\leq \|x_{n+m-1} - p\| + \alpha_{n+m-1}M + \|\varphi'_{n+m-1}\| \\ &\leq \|x_{n+m-2} - p\| + (\alpha_{n+m-1} + \alpha_{n+m-2})M + \|\varphi'_{n+m-2}\| + \|\varphi'_{n+m-1}\| \\ &\quad \vdots \\ &\leq \|x_n - p\| + M \sum_{i=n}^{n+m-1} \alpha_i + \sum_{i=n}^{n+m-1} \|\varphi'_i\| \end{aligned}$$

for all  $p \in F(T) \cap F(I)$ ,  $n \geq n_0$ . Thus Lemma 3.1 is proved. □

Since  $\{\psi_n\}$  is bounded,  $\varphi_n = \varphi'_n + \varphi''_n$ ,  $n \in \mathbb{N}$  and  $\sum_{n=0}^{\infty} \|\varphi'_n\| < \infty$ ,  $\|\varphi''_n\| = o(\alpha_n)$ , then we have  $\sum_{n=0}^{\infty} (M\alpha_n + \|\varphi'_n\|) < \infty$ . From Lemma 2.1, we take  $\{a_n\} = \{x_n - p\}$  and  $\{b_n\} = M\alpha_n + \|\varphi'_n\|$ . This implies that  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists.

**PROOF OF THEOREM 3.1.** We only prove the sufficiency of Theorem 3.1. Suppose that (3.4) is satisfied; then  $\lim_{n \rightarrow \infty} d(x_n, F(T) \cap F(I)) = 0$ .

First we show that  $\{x_n\}$  is a Cauchy sequence in  $K$ . For  $\varepsilon > 0$  and  $n \geq n_1$  there exists  $n_1 \geq n_0$  such that  $d(x_n, F(T) \cap F(I)) < \varepsilon$ ,  $\sum_{n=n_1}^{\infty} \alpha_n < \frac{\varepsilon}{M}$ ,  $\sum_{n=n_1}^{\infty} \|\varphi'_n\| < \varepsilon$ . By the definition of infimum and  $d(x_n, F(T) \cap F(I)) < \varepsilon$  there exists  $p_0 \in F(T) \cap$

$F(I)$  such that  $d(x_{n_1}, p) < 2\varepsilon$ . Furthermore, for  $n \geq n_1 \geq n_0$  and  $\forall m \geq 1$

$$\begin{aligned} \|x_{n+m} - x_n\| &\leq \|x_{n+m} - p_0\| + \|x_n - p_0\| \\ &\leq \|x_{n_1} - p_0\| + M \sum_{i=n_1}^{n+m-1} \alpha_i + \sum_{i=n_1}^{n+m-1} \|\varphi'_i\| \\ &\quad + \|x_{n_1} - p_0\| + M \sum_{i=n_1}^{n-1} \alpha_i + \sum_{i=n_1}^{n-1} \|\varphi'_i\|. \end{aligned}$$

Then for  $n \geq n_1 \geq n_0$  and  $\forall m \geq 1$  we have  $\|x_{n+m} - x_n\| \leq 8\varepsilon$ . Since  $\varepsilon$  is arbitrary, then  $\{x_n\}$  is a Cauchy sequence in  $K$ . Since  $X$  is a Banach space, let  $\{x_n\} \rightarrow p^*$  as  $n \rightarrow \infty$ . We prove that  $p^* \in F(T) \cap F(I)$ . We have  $\{x_n\} \rightarrow p^*$  as  $n \rightarrow \infty$  and  $\lim_{n \rightarrow \infty} d(x_n, F(T) \cap F(I)) = 0$ , for  $\varepsilon > 0$ , there exists a positive integer  $n_2 \geq n_1 \geq n_0$  and  $n \geq n_2$  we have  $\|x_n - p^*\| < \varepsilon$ ,  $d(x_n, F(T) \cap F(I)) < \varepsilon$ . Then there exists  $q \in F(T) \cap F(I)$  such that  $d(x_{n_2}, q) < 2\varepsilon$ . Furthermore, for  $n \geq n_2$

$$\begin{aligned} \|T^n p^* - p^*\| &\leq \{\|T^n p^* - q\| - \|p^* - q\|\} + 2\|p^* - q\| \\ &\leq \{\|T^n p^* - q\| - \|I^n p^* - q\|\} + \{\|I^n p^* - q\| - \|p^* - q\|\} + 3\|p^* - q\| \\ &< 2\varepsilon + 3\{3\varepsilon\} = 11\varepsilon \end{aligned}$$

Since  $T$  is  $I$ -asymptotically quasi nonexpansive type and  $I$  is asymptotically quasi nonexpansive type, this implies that  $\{T^n p^*\} \rightarrow p^*$  as  $n \rightarrow \infty$ . Furthermore,

$$\|T^n p^* - T p^*\| \leq \{\|T^n p^* - q\| - \|p^* - q\|\} + \|p^* - q\| + \|T p^* - q\|.$$

Then for  $n \geq n_2$  by (3.1), (3.2), (3.6) and (3.7) we have

$$\begin{aligned} \|T^n p^* - T p^*\| &\leq \{\|T^n p^* - q\| - \|I^n p^* - q\|\} + \{\|I^n p^* - q\| - \|p^* - q\|\} \\ &\quad + 2\|p^* - q\| + L\|I p^* - q\| \\ &\leq 2\varepsilon + 2\|p^* - q\| + L\Gamma\|p^* - q\| \\ &\leq 2\varepsilon + (2 + L\Gamma)\{\|x_{n_2} - p^*\| + \|x_{n_2} - q\|\} \\ &< 2\varepsilon + (2 + L\Gamma)3\varepsilon < \varepsilon(8 + 3L\Gamma) \end{aligned}$$

Since  $\varepsilon$  is arbitrary,  $\{T^n p^*\} \rightarrow T p^*$  as  $n \rightarrow \infty$ , implying  $T p^* = p^* \in F(T) \cap F(I)$ .

Further we apply for  $I : K \rightarrow K$  asymptotically quasi nonexpansive type mapping. Then for  $n \geq n_2$  we have

$$\begin{aligned} \|I^n p^* - p^*\| &\leq \{\|I^n p^* - q\| - \|q - p^*\|\} + 2\|p^* - q\| \\ &\leq \varepsilon + 2\{\|x_{n_2} - p^*\| + \|x_{n_2} - q\|\} < \varepsilon + 2\{\varepsilon + 2\varepsilon\} = 7\varepsilon \end{aligned}$$

This implies that  $\{I^n p^*\} \rightarrow p^*$  as  $n \rightarrow \infty$ . Furthermore,

$$\|I^n p^* - I p^*\| \leq \{\|I^n p^* - q\| - \|p^* - q\|\} + \|p^* - q\| + \|I p^* - q\|.$$

Then for  $n \geq n_2$  by (3.2) and (3.7) we have

$$\begin{aligned} \|I^n p^* - Ip^*\| &\leq \{\|I^n p^* - q\| - \|p^* - q\|\} + \|p^* - q\| + \Gamma \|Ip^* - q\| \\ &\leq \varepsilon + \|p^* - q\| + \Gamma \|p^* - q\| \\ &\leq \varepsilon + (1 + \Gamma)\{\|x_{n_2} - p^*\| + \|x_{n_2} - q\|\} \\ &< \varepsilon + (1 + \Gamma)3\varepsilon < \varepsilon(4 + 3\Gamma). \end{aligned}$$

Since  $\varepsilon$  is arbitrary,  $\{I^n p^*\} \rightarrow p^*$  as  $n \rightarrow \infty$ . Also

$$\|I^n p^* - Ip^*\| \leq \|I^n p^* - q\| + \|Ip^* - q\| < 2\varepsilon$$

Since  $\varepsilon$  is arbitrary,  $\{I^n p^*\} \rightarrow Ip^*$  as  $n \rightarrow \infty$ . This shows that  $Ip^* = p^* \in F(I)$ . From this we obtain  $p^* \in F(T) \cap F(I)$ .

Thus  $\{x_n\}$  converges strongly to a common fixed point of  $T$  and  $I$  in  $K$ , subset of  $X$  Banach space.  $\square$

Now we establish the weak convergence theorem for Ishikawa iterates of  $I$ -asymptotically quasi-nonexpansive type mappings in Banach spaces. First, we prove the following lemma.

LEMMA 3.2. *Let  $X$  be a uniformly convex Banach space and  $K$  be a nonempty closed convex subset of  $X$ . Let  $T, I$  and  $\{x_n\}$  be the same as in Lemma 3.1. If  $F = F(T) \cap F(I) \neq \emptyset$ , then  $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = \lim_{n \rightarrow \infty} \|Ix_n - x_n\| = 0$ .*

PROOF. By Lemma 3.1, for any  $p \in F(T) \cap F(I)$ ,  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists. Let  $\lim_{n \rightarrow \infty} \|x_n - p\| = c$ . If  $c = 0$ , then the proof is completed.

Now suppose  $c > 0$ . From (3.9), we have  $\|y_n - p\| \leq \|x_n - p\| + 2\beta_n \varepsilon + \|\psi_n\|$ . Taking  $\limsup$  on both sides in the above inequality,

$$(3.11) \quad \limsup_{n \rightarrow \infty} \|y_n - p\| \leq c.$$

Since  $I$  is asymptotically nonexpansive type self-mappings on  $K$ , from (3.7), which is on taking  $\limsup_{n \rightarrow \infty}$  and using (3.11), then we get  $\limsup_{n \rightarrow \infty} \|I^n y_n - p\| \leq c$ . Further,  $\lim_{n \rightarrow \infty} \|x_{n+1} - p\| = c$  means that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|\alpha_n I^n y_n + (1 - \alpha_n)x_n - p\| &= c \\ \lim_{n \rightarrow \infty} (1 - \alpha_n)\|x_n - p\| + \alpha_n \|I^n y_n - p\| &= c. \end{aligned}$$

It follows from Lemma 2.2

$$(3.12) \quad \lim_{n \rightarrow \infty} \|I^n y_n - x_n\| = 0.$$

Further,

$$\lim_{n \rightarrow \infty} \|\alpha_n(T^n x_n - p) + (1 - \alpha_n)(x_n - p)\| = \lim_{n \rightarrow \infty} \|y_n - p\| = c.$$

By Lemma 2.2, we have

$$(3.13) \quad \lim_{n \rightarrow \infty} \|T^n x_n - x_n\| = 0.$$

From (3.12) and (3.13), we have

$$(3.14) \quad \lim_{n \rightarrow \infty} \|I^n x_n - x_n\| = 0.$$

Using (3.1), (3.2), (3.3), (3.13) and (3.14), it is easy to show that

$$(3.15) \quad \lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$$

$$(3.16) \quad \lim_{n \rightarrow \infty} \|Ix_n - x_n\| = 0.$$

Then the proof is completed.  $\square$

**THEOREM 3.2.** *Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition,  $K$  be a nonempty closed convex subset of  $X$ . Let  $T, I$  and  $\{x_n\}$  be the same as in Lemma 3.1. If  $F(T) \cap F(I) \neq \emptyset$ , the mappings  $E - T$  and  $E - I$  are semi-closed at zero, then  $\{x_n\}$  converges weakly to a common fixed point of  $T$  and  $I$ .*

**PROOF.** By assumption,  $F(T) \cap F(I)$  is nonempty. Take  $p \in F(T) \cap F(I)$ . It follows from Lemma 3.1 that the limit  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists. Therefore,  $\{x_n - p\}$  is a bounded sequence in  $X$ . Since  $X$  is a uniformly convex Banach space and  $K$  is a nonempty closed convex subset of  $X$ , then  $K$  is weakly compact. This implies that there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that  $\{x_{n_k}\}$  converges to a point  $p \in w(\{x_n\})$ , where  $w(\{x_n\})$  denotes the weak limit set of  $\{x_n\}$ , which shows that  $w(\{x_n\})$  is nonempty. For any  $p \in w(\{x_n\})$ , there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that  $\{x_{n_k}\} \rightarrow p$  weakly. Hence, it follows from (3.15) and (3.16) in Lemma 3.2 that  $Tp = p$  and  $Ip = p$ . By Opial's condition,  $\{x_n\}$  has only one weak limit point, i.e.,  $\{x_n\}$  converges weakly to a common fixed point of  $T$  and  $I$ .  $\square$

#### 4. Convergence for nonself $I$ -asymptotically quasi-nonexpansive type mappings

In this section, the convergence of the Ishikawa iterative sequences to common fixed point for nonself  $I$ -asymptotically quasi-nonexpansive type mappings is obtained in Banach spaces.

A subset  $K$  of  $X$  is called a *retract* of  $X$  if there exists a continuous map  $P : X \rightarrow K$  such that  $Px = x$  for all  $x \in K$ . A map  $P : X \rightarrow K$  is called a *retraction* if  $P^2 = P$ . In particular, a subset  $K$  is called a *nonexpansive retract* of  $X$  if there exists a *nonexpansive retraction*  $P : X \rightarrow K$  such that  $Px = x$  for all  $x \in K$ .

Next, we introduce the following concepts for nonself mappings. Let  $X$  be a real Banach space. A subset  $K$  of  $X$  be nonempty nonexpansive retraction of  $X$  and  $P$  be nonexpansive retraction from  $X$  onto  $K$ . A nonself mapping  $T : K \rightarrow X$  is called *asymptotically nonexpansive* if there exists a sequence  $\{v_n\} \subset [1, \infty)$  with  $\lim_{n \rightarrow \infty} v_n = 1$  such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq v_n \|x - y\|$$

for all  $x, y \in K$  and  $n \geq 1$ .  $T$  is called *uniformly  $L$ -Lipschitzian* if there exists a constant  $L > 0$  such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq L \|x - y\|$$

for all  $x, y \in K$  and  $n \geq 1$ . From the above definition, it is obvious that nonself asymptotically nonexpansive mappings is uniformly  $L$ -Lipschitzian.



Let  $I : K \rightarrow X$  be a nonself asymptotically quasi-nonexpansive type mappings and  $T : K \rightarrow X$  be a nonself  $I$ -asymptotically quasi-nonexpansive type mappings with  $F(T) \cap F(I) = \{x \in K : Tx = x = Ix\} \neq \emptyset$ . A mapping  $T : K \rightarrow X$  is called  $\Lambda$ -Lipschitzian if there exists constant  $\Lambda > 0$  such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq \Lambda \|I(PI)^{n-1}x - I(PI)^{n-1}y\|$$

for all  $x, y \in K$  and  $n \geq 1$ .

Iterative techniques for converging fixed points of nonexpansive non-self mappings have been studied by many authors (see, for example, [3, 19, 14]). The concept of nonself asymptotically nonexpansive mappings was introduced in [3] as a generalization of asymptotically nonexpansive self-mappings and some strong and weak convergence theorems for such mappings were obtained. The sequence  $\{x_n\}_{n \geq 1}$  generated as follows:  $x_1 \in K$ ,

$$\begin{aligned} y_n &= P(\alpha_n T(PT)^{n-1}x_n + \beta_n x_n), \\ x_{n+1} &= P(\alpha'_n T(PT)^{n-1}y_n + \beta'_n x_n), \quad \forall n \geq 1, \end{aligned}$$

where  $\{\alpha_n\}, \{\beta_n\}, \{\alpha'_n\}, \{\beta'_n\} \in (0, 1)$ .

Let  $T : K \rightarrow X$  be a nonself  $I$ -asymptotically quasi-nonexpansive type mapping and  $I : K \rightarrow X$  be a nonself asymptotically quasi-nonexpansive type mapping.

Now we define an  $\{x_n\}_{n \geq 1}$  sequence as follows:

$$(4.1) \quad \begin{aligned} y_n &= P(\alpha_n T(PT)^{n-1}x_n + \beta_n x_n + \gamma_n \psi_n), \\ x_{n+1} &= P(\alpha'_n I(PI)^{n-1}y_n + \beta'_n x_n + \gamma'_n \varphi_n), \quad \forall n \geq 1, \end{aligned}$$

where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\alpha'_n\}, \{\beta'_n\}, \{\gamma'_n\}$  are sequences in  $(0, 1)$  with  $\alpha_n + \beta_n + \gamma_n = 1 = \alpha'_n + \beta'_n + \gamma'_n$  and  $\{\psi_n\}, \{\varphi_n\}$  are bounded sequences in  $K$ .

$$\limsup_{n \rightarrow \infty} \left( \sup_{x \in X, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\| - \|I(PI)^{n-1}x - p\| \} \right) \leq 0.$$

Observe that

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \left( \sup_{x \in X, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\| - \|I(PI)^{n-1}x - p\| \} \right) \\ &\quad \times \limsup_{n \rightarrow \infty} \left( \sup_{x \in X, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\| + \|I(PI)^{n-1}x - p\| \} \right) \\ &= \limsup_{n \rightarrow \infty} \left( \sup_{x \in X, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\|^2 - \|I(PI)^{n-1}x - p\|^2 \} \right) \leq 0. \end{aligned}$$

Therefore we have

$$\limsup_{n \rightarrow \infty} \left( \sup_{x \in X, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\| - \|I(PI)^{n-1}x - p\| \} \right) \leq 0.$$

This implies that for any given  $\varepsilon > 0$ , there exists a positive integer  $n_0$  such that for  $n \geq n_0$  we have

$$\left( \sup_{x \in X, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\| - \|I(PI)^{n-1}x - p\| \} \right) \leq 0.$$

**THEOREM 4.1.** *Let  $X$  be a Banach space and  $K$  be a nonempty subset of the Banach space. Let  $T, I : K \rightarrow X$  be two nonself mappings. Let  $T$  be a nonself  $I$ -asymptotically quasi-nonexpansive type and  $I$  be a nonself asymptotically quasi-nonexpansive type in Banach space with  $F(T) \cap F(I) \neq \emptyset$ . Let the sequence  $\{x_n\}$  be defined by (4.1) and for every  $n \in \mathbb{N}$ , where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\alpha'_n\}, \{\beta'_n\}, \{\gamma'_n\}$  are sequences in  $(0, 1)$  with  $\alpha_n + \beta_n + \gamma_n = 1 = \alpha'_n + \beta'_n + \gamma'_n$ ,  $\sum_{i=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{i=1}^{\infty} \gamma'_n < \infty$ , and  $\{\psi_n\}, \{\varphi_n\}$  are bounded sequences in  $K$ .*

*Then  $\{x_n\}$  converges strongly to a common fixed point of  $T$  and  $I$  in  $K$  iff*

$$(4.2) \quad \liminf_{n \rightarrow \infty} d(x_n, F(T) \cap F(I)) = 0.$$

**PROOF.** The necessity of condition (4.2) is obvious. Next we prove the sufficiency of condition (4.2). Let the sequence  $\{x_n\}$  be defined by (4.1). Let  $p \in F(T) \cap F(I)$ , by boundedness of the sequences  $\{\psi_n\}, \{\varphi_n\}$ , so we can put

$$M = \max\{\sup_{n \geq 1} \|\psi_n - p\|, \sup_{n \geq 1} \|\varphi_n - p\|\}.$$

For any given  $\varepsilon > 0$ , there exists a positive integer  $n_0$  such that  $n \geq n_0$

$$\begin{aligned} \sup_{x \in K, p \in F(T) \cap F(I)} \{ \|T(PT)^{n-1}x - p\| - \|I(PI)^{n-1}x - p\| \} &< \varepsilon. \\ \sup_{x \in K, p \in F(I)} \{ \|I(PI)^{n-1}x - p\| - \|x - p\| \} &< \varepsilon. \end{aligned}$$

Therefore, in particular, we have

$$(4.3) \quad \{ \|T(PT)^{n-1}x_n - p\| - \|I(PI)^{n-1}x_n - p\| \} < \varepsilon,$$

for all  $p \in F(T) \cap F(I)$  and  $\forall n \geq n_0$ .

$$(4.4) \quad \{ \|I(PI)^{n-1}y_n - p\| - \|y_n - p\| \} < \varepsilon,$$

for all  $p \in F(I)$  and  $\forall n \geq n_0$ . Thus for each  $n \geq 1$  and for any  $p \in F(T) \cap F(I)$ , using (4.1), (4.3) and (4.4), we have

$$\begin{aligned} (4.5) \quad \|x_{n+1} - p\| &= \|P(\alpha'_n x_n + \beta'_n I(PI)^{n-1}y_n + \gamma'_n \varphi_n - p)\| \\ &\leq \alpha'_n \|x_n - p\| + \beta'_n \|I(PI)^{n-1}y_n - p\| + \gamma'_n \|\varphi_n - p\| \\ &= \alpha'_n \|x_n - p\| + \beta'_n \{ \|I(PI)^{n-1}y_n - p\| - \|y_n - p\| \} \\ &\quad + \beta'_n \|y_n - p\| + \gamma'_n \|\varphi_n - p\| \\ &\leq \alpha'_n \|x_n - p\| + \beta'_n \{\varepsilon\} + \beta'_n \|y_n - p\| + \gamma'_n M \end{aligned}$$

and

$$\begin{aligned}
 (4.6) \quad \|y_n - p\| &= \|P(\alpha_n x_n + \beta_n T(PT)^{n-1} x_n + \gamma_n \psi_n - p)\| \\
 &\leq \alpha_n \|x_n - p\| + \beta_n \|T(PT)^{n-1} x_n - p\| + \gamma_n \|\psi_n - p\| \\
 &\leq \alpha_n \|x_n - p\| + \beta_n \{\|T(PT)^{n-1} x_n - p\| - \|I(PI)^{n-1} x_n - p\|\} \\
 &\quad + \beta_n \{\|I(PI)^{n-1} x_n - p\| - \|x_n - p\|\} + \beta_n \|x_n - p\| + \gamma_n M \\
 &\leq \alpha_n \|x_n - p\| + 2\beta_n \{\varepsilon\} + \beta_n \|x_n - p\| + \gamma_n M \\
 &\leq (\alpha_n + \beta_n) \|x_n - p\| + 2\beta_n \varepsilon + \gamma_n M \\
 &\leq (1 - \gamma_n) \|x_n - p\| + 2\beta_n \varepsilon + \gamma_n M \\
 &\leq \|x_n - p\| + D_n
 \end{aligned}$$

where  $D_n = 2\beta_n \varepsilon + \gamma_n M$ . Then  $\sum_{n=1}^{\infty} D_n < \infty$  since  $\sum_{n=1}^{\infty} \gamma_n < \infty$ .

Substituting (4.6) into (4.5), we have

$$\begin{aligned}
 (4.7) \quad \|x_{n+1} - p\| &\leq \alpha'_n \|x_n - p\| + \beta'_n \varepsilon + \beta'_n (\|x_n - p\| + D_n) + \gamma'_n M \\
 &\leq (\alpha'_n + \beta'_n) \|x_n - p\| + \beta'_n (\varepsilon + D_n) + \gamma'_n M \\
 &\leq (1 - \gamma'_n) \|x_n - p\| + G_n \\
 &\leq \|x_n - p\| + G_n
 \end{aligned}$$

where  $G_n = \beta'_n (\varepsilon + D_n) + \gamma'_n M$ . Then  $\sum_{n=1}^{\infty} G_n < \infty$  since  $\sum_{n=1}^{\infty} \gamma'_n < \infty$  and  $\sum_{n=1}^{\infty} D_n < \infty$ .

It follows from (4.7) that  $d(x_{n+1}, F(T) \cap F(I)) \leq d(x_n, F(T) \cap F(I)) + G_n$ .

By Lemma 2.1, we can get that  $\lim_{n \rightarrow \infty} d(x_n, F(T) \cap F(I))$  exists. By condition  $\liminf_{n \rightarrow \infty} d(x_n, F(T) \cap F(I)) = 0$ , we have

$$(4.8) \quad \lim_{n \rightarrow \infty} d(x_n, F(T) \cap F(I)) = 0.$$

Next we prove that  $\{x_n\}$  is a Cauchy sequence in  $X$ . In fact, for any  $n \geq n_0$ , any  $m \geq n_1$  and any  $p \in F(T) \cap F(I)$  we have

$$\begin{aligned}
 (4.9) \quad \|x_{n+m} - p\| &\leq \|x_{n+m-1} - p\| + G_{n+m-1} \\
 &\leq \|x_{n+m-2} - p\| + G_{n+m-1} + G_{n+m-2} \\
 &\leq \dots \leq \|x_n - p\| + \sum_{k=n}^{\infty} G_k.
 \end{aligned}$$

So by (4.9), we have

$$(4.10) \quad \|x_{n+m} - x_n\| \leq \|x_{n+m} - p\| + \|x_n - p\| \leq 2\|x_n - p\| + \sum_{k=n}^{\infty} G_k.$$

By the arbitrariness of  $p \in F(T) \cap F(I)$  and (4.10), we have

$$\|x_{n+m} - p\| \leq 2d(x_n, F(T) \cap F(I)) + \sum_{k=n}^{\infty} G_k \quad \forall n \geq n_0.$$

For any given  $\varepsilon > 0$ , there exists a positive integer  $n_1 \geq n_0$ , such that for any  $n \geq n_1$ ,  $d(x_n, F(T) \cap F(I)) < \frac{\varepsilon}{4}$  and  $\sum_{k=n}^{\infty} G_k < \frac{\varepsilon}{2}$ , we have  $\|x_{n+m} - x_n\| < \varepsilon$ , and so for any  $m \geq 1$

$$\lim_{n \rightarrow \infty} \|x_{n+m} - x_n\| = 0.$$

This shows that  $\{x_n\}$  is a Cauchy sequence in  $X$ . Since  $X$  is complete, there exists a  $p^* \in X$  such that  $x_n \rightarrow p^*$  as  $n \rightarrow \infty$ .

Finally, by the routine method, we have to prove that  $p^* \in F(T) \cap F(I)$ . By contradiction, we assume that  $p^*$  is not in  $F(T) \cap F(I)$ . Since  $F(T) \cap F(I)$  is a closed set,  $d(p^*, F(T) \cap F(I)) > 0$ . Hence for all  $p \in F(T) \cap F(I)$ , we have

$$\|p^* - p\| \leq \|x_n - p^*\| + \|x_n - p\|.$$

This implies that

$$(4.11) \quad d(p^*, F(T) \cap F(I)) \leq \|x_n - p^*\| + d(x_n, F(T) \cap F(I)).$$

Letting  $n \rightarrow \infty$  in (4.11) and noting (4.8), we have  $d(p^*, F(T) \cap F(I)) \leq 0$ . This is a contradiction. Hence  $p^* \in F(T) \cap F(I)$ . This completes the proof of Theorem 4.1.  $\square$

## References

1. S. S. Chang, Y. Y. Zhou, *Some convergence theorems for mappings of asymptotically quasi-nonexpansive type in Banach spaces*, J. Appl. Math. Comput. **12**(1-2) (2003), 119–127.
2. S. S. Chang, Y. J. Cho, H. Zhou, *Demi-closed principle and weak convergence problems for asymptotically nonexpansive mappings*, J. Korean Math. Soc. **38**(6) (2001), 1245–1260.
3. C. E. Chidume, E. U. Ofoedu, H. Zegeye, *Strong and weak convergence theorems for asymptotically nonexpansive mappings*, J. Math. Anal. Appl. **280** (2003), 364–374.
4. J. B. Diaz, F. T. Metcalf, *On the set of sequential limit points of successive approximations*, Trans. Am. Math. Soc. **135** (1969), 459–485.
5. W. G. Dotson Jr., *On the Mann iterative process*, Trans. Am. Math. Soc. **149** (1970), 65–73.
6. M. K. Ghosh, L. Debnath, *Convergence of Ishikawa iterates of quasi-nonexpansive mappings*, J. Math. Anal. Appl. **207** (1997), 96–103.
7. K. Goebel, W. A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Am. Math. Soc. **35** (1972), 171–174.
8. S. Ishikawa, *Fixed points by a new iteration method*, Proc. Am. Math. Soc. **44** (1974), 147–150.
9. H. Y. Lan, *Common fixed point iterative processes with errors for generalized asymptotically quasi-nonexpansive mappings*, Comput. Math. Appl. **52** (2006) 1403–1412.
10. Q. H. Liu, *Iterative sequences for asymptotically quasi-nonexpansive mappings*, J. Math. Anal. Appl. **259** (2001), 1–7.
11. ———, *Iterative sequences for asymptotically quasi-nonexpansive mappings with error member*, J. Math. Anal. Appl. **259** (2001), 18–24.
12. Z. Opial, *Weak convergence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
13. W. V. Petryshyn, T. E. Williamson, *Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings*, J. Math. Anal. Appl. **43** (1973), 459–497.
14. N. Shahzad, *Approximating fixed points of non-self nonexpansive mappings in Banach spaces*, Nonlinear Anal. **61** (2005), 1031–1039.
15. J. Schu, *Weak and strong convergence to fixed points of asymptotically nonexpansive mappings*, Bull. Aust. Math. Soc. **43**(1) (1991), 153–159.
16. K. K. Tan, H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iterative process*, J. Math. Anal. Appl. **178** (1993), 301–308.

17. S. Temir, O. Gul, *Convergence theorem for I-asymptotically quasi-nonexpansive mapping in Hilbert space*, J. Math. Anal. Appl. **329** (2007) 759–765.
18. S. Temir, *Convergence of iterative process for generalized I-asymptotically quasi-nonexpansive mappings*, Thai J. Math. **7**(2) (2009), 367–379.
19. L. Wang, *Strong and weak convergence theorems for common fixed points of nonself asymptotically nonexpansive mappings*, J. Math. Anal. Appl. **323** (2006), 550–557.
20. S.S. Yang, L. Wang, *Convergence of the Ishikawa iteration scheme with errors for I-asymptotically quasi-nonexpansive mappings*, Thai J. Math. **5**(2) (2007), 199–207.

Department of Mathematics  
Art and Science Faculty  
Harran University  
Sanliurfa  
Turkey  
temirsevit@harran.edu.tr

(Received 14 04 2012)  
(Revised 16 11 2012)