

## ALEXANDER $r$ -TUPLES AND BIER COMPLEXES

Duško Jojić, Ilya Nekrasov,  
Gaiane Panina, and Rade Živaljević

**ABSTRACT.** We introduce and study *Alexander  $r$ -tuples*  $\mathcal{K} = \langle K_i \rangle_{i=1}^r$  of simplicial complexes, as a common generalization of pairs of Alexander dual complexes (Alexander 2-tuples) and  $r$ -unavoidable complexes of [3] and [11]. In the same vein, the *Bier complexes*, defined as the deleted joins  $\mathcal{K}_\Delta^*$  of Alexander  $r$ -tuples, include both standard *Bier spheres* and *optimal multiple chessboard complexes* (Section 2.2) as interesting, special cases.

Our main results are Theorem 4.1 saying that (1) the  $r$ -fold deleted join of Alexander  $r$ -tuple is a pure complex homotopy equivalent to a wedge of spheres, and (2) the  $r$ -fold deleted join of a collective unavoidable  $r$ -tuple is  $(n - r - 1)$ -connected, and a classification theorem (Theorem 5.1 and Corollary 5.1) for Alexander  $r$ -tuples and Bier complexes.

### 1. Introduction

Topological combinatorics utilizes methods from algebraic (combinatorial) topology to solve problems in combinatorics and discrete geometry. Among the highlights (that strongly influenced the subsequent developments), and early achievements of topological combinatorics are the solution of Kneser conjecture (L. Lovász, 1978), topological Tverberg theorem (I. Bárány, S. B. Shlosman, A. Szűcs, 1981), N. Alon's 'Splitting necklace theorem' (1987), and many others, see [1, 12, 14] for an overview and introduction.

Simplicial complexes are among the central objects of study in topological combinatorics. Their role in this subject can be compared to the role of manifolds in differential geometry and topology, for illustration R. Forman's 'Discrete Morse theory' (Section 2.4) exemplifies a fruitful interplay of ideas and techniques from these areas.

In this paper we introduce "Alexander  $r$ -tuples of simplicial complexes" and closely related "collective  $r$ -unavoidable complexes" (Section 3), as unifying concepts that bring together *Alexander pairs* of mutually dual complexes, and  *$r$ -unavoidable complexes* [3, 10, 11].

---

2010 *Mathematics Subject Classification*: Primary 05E45; Secondary 05D15; 52A35; 55M05.

*Key words and phrases*: Bier spheres, Alexander duality, chessboard complexes, unavoidable complexes, discrete Morse theory.

Communicated by Žarko Mijajlović.

The deleted join operation, applied to an Alexander pair  $(K, K^\circ)$ , yields a combinatorial sphere  $\text{Bier}(K) = K *_\Delta K^\circ$ , known as the Bier sphere associated to  $K$ , see [12, Section 5.6]. The special case of a self-dual complex  $K = K^\circ \subset 2^{[n]}$  is of particular importance. In this case the Bier sphere  $\text{Bier}(K) = K *_\Delta K$  is a  $\mathbb{Z}_2$ -complex and its equivariant  $\mathbb{Z}_2$ -index is  $\text{Ind}_{\mathbb{Z}_2}(\text{Bier}(K)) = \text{Ind}_{\mathbb{Z}_2}(S^{n-2}) = n - 2$ . This fact alone has many interesting consequences, including the Van Kampen-Flores theorem [12, Theorem 5.1.1] which says that the  $d$ -skeleton  $(\sigma^{2d+2})^{\leq d}$  of a  $(2d + 2)$ -dimensional simplex is nonembeddable in  $\mathbb{R}^{2d}$ .

The 2-unavoidable complexes are easily identified as superdual complexes  $K \supseteq K^\circ$ . From here it easily follows that self-dual complexes are precisely (inclusion) minimal 2-unavoidable complexes.

The outline above leads to the conclusion that  $r$ -unavoidable complexes can be interpreted as  $r$ -fold analogues (relatives) of Alexander self-dual complexes. It may be tempting to extend this analogy further, to include  $r$ -fold generalization of (not necessarily symmetric) Alexander dual pairs. The following research problem summarizes the desirable properties of such an extension.

**PROBLEM 1.1.** Find a property  $\mathcal{P}_r$  of a collection  $\mathcal{K} = \langle K_i \rangle_{i=1}^r = \langle K_1, \dots, K_r \rangle$  of simplicial complexes on the same vertex set,  $K_i \subset 2^{[n]}$ , such that:

- (1) If  $r = 2$  then a pair of complexes  $\langle K_1, K_2 \rangle$  satisfies  $\mathcal{P}_2$  if and only if  $\langle K_1, K_2 \rangle$  is an Alexander superdual pair in the sense that  $K_1 \supseteq K_2^\circ$  (equivalently  $K_2 \supseteq K_1^\circ$ );
- (2) If  $K_1 = \dots = K_r = K$  then  $\mathcal{K}$  satisfies  $\mathcal{P}_r$  if and only if  $K$  is an  $r$ -unavoidable complex;
- (3) If  $\mathcal{K} \in \mathcal{P}_r$  then the deleted join  $\mathcal{K}_\Delta^* = K_1 *_\Delta \dots *_\Delta K_r$  is an  $(n - r - 1)$ -connected complex.

Moreover, it is desirable to describe a stronger property  $\mathcal{P}_r^\sharp \subset \mathcal{P}_r$  such that:

- (1<sup>♯</sup>)  $\langle K_1, K_2 \rangle \in \mathcal{P}_2^\sharp$  if and only if  $K_1 = K_2^\circ$ ;
- (2<sup>♯</sup>) If  $K_1 = \dots = K_r = K$  and  $\mathcal{K} \in \mathcal{P}_r^\sharp$ , then  $K$  is an (inclusion) minimal  $r$ -unavoidable complex;
- (3<sup>♯</sup>) If  $\mathcal{K} \in \mathcal{P}_r^\sharp$  then the deleted join  $\mathcal{K}_\Delta^* = K_1 *_\Delta \dots *_\Delta K_r$  has the homotopy type of a wedge of  $(n - r)$ -dimensional spheres.

Motivated by Problem 1.1, we describe (Definition 3.1) the class  $CU_r$  of “collective  $r$ -unavoidable complexes”, as our primary candidate for the class  $\mathcal{P}_r$ . Individual  $r$ -unavoidable complexes often arise from the ‘pigeonhole principle’ (see [3, Lemma 4.2]). For this reason we may occasionally say that an ordered collection  $\mathcal{K} = \langle K_i \rangle_{i=1}^r \in CU_r$  has the *pigeonhole property*, or that  $\mathcal{K}$  itself is a *pigeonhole  $r$ -tuple*.

We introduce the class  $\mathcal{A}_r$  of “Alexander  $r$ -tuples of simplicial complexes” (Definition 3.2), as the most regular class of “collective  $r$ -unavoidable complexes”, and as our primary candidate for the class  $\mathcal{P}_r^\sharp$ .

Finally, *Bier complexes* (Section 4) arise as the deleted joins of Alexander  $r$ -tuples, in perfect analogy with the case of standard Bier spheres, which arise as deleted joins of Alexander pairs of complexes.

**1.1. Summary of the main results.** The core of the paper are the results showing that the collective  $r$ -unavoidable complexes (and their deleted joins) as well as the Alexander  $r$ -tuples (and the associated Bier complexes) indeed satisfy the properties listed in Problem 1.1. Perhaps the most interesting among them are the following (see Sections 3 and 4).

If  $\mathcal{K} = \langle K_i \rangle_{i=1}^r = \langle K_1, \dots, K_r \rangle$  is a collective  $r$ -unavoidable collection of subcomplexes of  $2^{[n]}$  (Definition 3.1), then (by Problem 1.1 (3)) the associated deleted join  $\mathcal{K}_\Delta^* = K_1 *_\Delta \cdots *_\Delta K_r$  is expected to be  $(n - r - 1)$ -connected. This is indeed the case, as shown in the first part of Theorem 4.1. In particular we recover the result that  $K *_\Delta K^\circ$  is an  $(n - 2)$ -dimensional homotopy sphere, whenever  $K \neq 2^{[n]}$  is *superdual* in the sense that  $K \supseteq K^\circ$ .

In the special case when  $\mathcal{K} = \langle K_i \rangle_{i=1}^r$  is an Alexander  $r$ -tuple (Definition 3.2), we have a stronger result (see the second half of Theorem 4.1), that the associated Bier complex  $\mathcal{K}_\Delta^*$  is a wedge of spheres of the same dimension  $n - r$  (Property (3<sup>#</sup>) in Problem 1.1). We describe an algorithm how the number of these spheres can be explicitly calculated (Corollary 4.1) and illustrate the calculation in the case of ‘optimal chessboard complexes’ (Section 8).

A classification theorem for Alexander  $r$ -tuples (Theorem 5.1) is proved in Section 5. It turns out, somewhat unexpectedly and as a pleasant surprise, that the ‘optimal chessboard complexes’ (introduced in Section 2.2) are the central examples of Bier complexes (Section 4) for  $r \geq 3$ .

Among the corollaries of our results are exact connectivity bounds for some classes of generalized chessboard complexes (including the main case of [8, Theorem 3.2]).

The rest of the paper is organized as follows. Section 2 is an overview of basic notions and facts, including a brief exposition of the discrete Morse theory [6, 7] (which is our central tool in this paper). We develop a version of this method which appears to be particularly well adapted for the analysis of Bier spheres (Section 6). We show in Section 7 how the method can be extended to the case of deleted joins of collective  $r$ -unavoidable complexes and general Bier complexes (introduced in Section 4). The highlights include the construction of a perfect discrete Morse function in the case of ‘optimal chessboard complexes’ (Section 8) and their relatives ‘long chessboard complexes’ (Section 9).

## 2. An overview of basic definitions and facts

In this section we collect some standard definitions and facts, as a reminder for the reader. This is also an opportunity to introduce some less standard notation and concepts, used in the rest of the paper. For other standard facts and definitions the reader is referred to [12].

**2.1. Simplicial complexes.** A simplicial complex on a set  $V$  of vertices is a subset  $K \subset 2^V$  such that (1)  $\emptyset \in K$  and (2) if  $A \subset B \in K$  then  $A \in K$ . By definition it is possible that  $\{v\} \notin K$  for some  $v \in V$ , however  $K \neq \emptyset$  (since  $\emptyset \in K$  by property (1)).

The complex  $2^V$  is often referred to as the simplex spanned by  $V$ , and denoted by  $\Delta(V)$ . We use, side by side, topological and combinatorial language (and notation). For example,

$$\binom{[n]}{\leq k},$$

is the  $(k-1)$ -skeleton of the  $(n-1)$ -dimensional simplex  $\Delta([n])$ .

The *deleted join* [12, Section 6] of a family  $\mathcal{K} = \langle K_i \rangle_{i=1}^r = \langle K_1, \dots, K_r \rangle$  of subcomplexes of  $2^{[n]}$  is the complex  $\mathcal{K}_\Delta^* = K_1 *_\Delta \dots *_\Delta K_r \subset (2^{[n]})^{*r}$  where  $A = A_1 \uplus \dots \uplus A_r \in \mathcal{K}_\Delta^*$  if and only if  $A_j$  are pairwise disjoint and  $A_i \in K_i$  for each  $i = 1, \dots, r$ .

**2.2. Multiple chessboard complexes.** A ‘chessboard complex’, in a very broad sense, is any subcomplex  $K \subset 2^{([n] \times [r])}$  of the simplex  $\Delta([n] \times [r])$  spanned by elementary squares of an  $(n \times r)$ -chessboard. Following [8, Section 2.1], the multiple chessboard complex

$$\Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}} = \Delta_{n,r}^{m_1, \dots, m_r; 1, \dots, 1}$$

is described by the condition that  $S \in \Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}}$  if and only if the cardinality of the set  $S \cap ([n] \times \{i\})$  is at most  $m_i$  for each  $i = 1, \dots, r$ , and the cardinality of the set  $S \cap (\{j\} \times [r])$  is at most 1 for each  $j = 1, \dots, n$ .

A moment’s reflections reveal that there is a relation,

$$\Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}} \cong \binom{[n]}{\leq m_1} *_\Delta \dots *_\Delta \binom{[n]}{\leq m_r},$$

which says that the multiple chessboard complex can be always expressed as the deleted join of skeletons of the simplex  $\Delta([n]) \cong \Delta^{n-1}$ .

One of the central results of [8, Theorem 3.2] says that  $\Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}}$  is  $(\nu-2)$ -connected where  $\nu = m_1 + \dots + m_r$ , provided  $n \geq m_1 + \dots + m_r + r - 1$ . For this reason the chessboard complex  $\Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}}$  is often called *optimal*, if  $n = m_1 + \dots + m_r + r - 1$ . Similarly we say that a multiple chessboard complex is *long* if  $n > m_1 + \dots + m_r + r - 1$ .

**2.3. Alexander duality and Bier spheres.** The Alexander dual [12, Section 5.6] of  $K \subset 2^V$  is the set  $K^\circ$  of all complements of nonsimplices in  $K$ ,

$$K^\circ = \{F \subset V \mid V \setminus F \notin K\}.$$

In order to rule out the possibility  $K^\circ = \emptyset$ , we tacitly assume throughout the paper that  $K \neq 2^V$ , whenever we are dealing with Alexander pairs  $(K, K^\circ)$  of complexes.

For a given simplicial complex  $K \subset 2^{[n]}$ , the associated Bier sphere  $\text{Bier}(K) = K *_\Delta K^\circ$  is described as the deleted join of  $K$  with its Alexander dual  $K^\circ$ . The simplices of the deleted join  $K *_\Delta K^\circ$  are by definition disjoint unions  $A_1 \uplus A_2 \subset [n] \uplus [n] \cong [n] \times [2]$ , where  $A_1 \in K, A_2 \in K^\circ$  and  $A_1 \cup A_2 \neq \emptyset$ . They can be also described as ordered partitions of the set  $[n]$  into three parts  $(A_1, A_2; B)$  (where  $B := [n] \setminus (A_1 \cup A_2)$ ).

Note that a partition  $(A_1, A_2; B)$  corresponds to a simplex in the deleted join  $K *_\Delta K^\circ$  if and only if:

- (1)  $A_1 \in K$ ,
- (2)  $A_2 \in K^\circ$  (or equivalently  $[n] \setminus A_2 \notin K$ );
- (3)  $\emptyset \neq B \neq [n]$  (equivalently  $\emptyset \neq A_1 \cup A_2 \neq [n]$ ).

The incidence relation of the simplices is described by the rule:  $(A_1, A_2; B) \subseteq (A'_1, A'_2; B')$  iff  $A_1 \subseteq A'_1$ , and  $A_2 \subseteq A'_2$ .

**2.4. Discrete Morse theory.** Robin Forman's discrete Morse theory [6, 7] is, as a tool, as powerful as the smooth Morse theory. It has been used in computations of the homology, the cup-product, Novikov homology, and other topological and combinatorial computations and applications. Major advantage of discrete Morse theory (compared to smooth Morse theory) is its applicability to a considerably larger class of objects which include simplicial and cellular complexes (and not only smooth manifolds).

In our paper we make use of a relatively small and quite reduced piece of the general theory. For our purposes it suffices to think of a 'Morse function' as a special kind of matching on the set of simplices. Here is a brief overview of some of the central definitions and results of the discrete Morse theory.

Let  $K$  be a simplicial complex. Its  $p$ -dimensional simplices ( $p$ -simplices for short) are denoted by  $\alpha^p, \beta^p, \sigma^p$ , etc. A *discrete vector field*  $D$  is a set of pairs  $(\alpha^p, \beta^{p+1})$  (called a matching) such that:

- (1) each simplex of the complex participates in at most one pair, and
- (2) in each pair, the simplex  $\alpha^p$  is a facet of  $\beta^{p+1}$ .

The pair  $(\alpha^p, \beta^{p+1})$  can be informally thought of as a vector in the vector field  $D$ . For this reason it is occasionally denoted by  $\alpha^p \rightarrow \beta^{p+1}$  (and in this case  $\beta^{p+1}$  is referred to as *the end* of the arrow  $\alpha^p \rightarrow \beta^{p+1}$ ).

Given a discrete vector field  $D$ , a *gradient path* in  $D$  is a sequence of simplices

$$\alpha_0^p, \beta_0^{p+1}, \alpha_1^p, \beta_1^{p+1}, \alpha_2^p, \beta_2^{p+1}, \dots, \alpha_m^p, \beta_m^{p+1}, \alpha_{m+1}^p,$$

which satisfies the following conditions:

- (1)  $p \geq 0$ , that is, the empty set  $\emptyset \in K$  is never matched,
- (2)  $(\alpha_i^p, \beta_i^{p+1})$  is a pair in  $D$  for each  $i$ ,
- (3) for each  $i = 0, \dots, m$  the simplex  $\alpha_{i+1}^p$  is a facet of  $\beta_i^{p+1}$ .
- (4)  $\alpha_i \neq \alpha_{i+1}$ .

A path is *closed* if  $\alpha_{m+1}^p = \alpha_0^p$ . A *discrete Morse function* (DMF for short) is a discrete vector field without closed paths.

Assuming that a discrete Morse function is fixed, the *critical simplices* are those simplices of the complex that are not matched. The Morse inequality [7] states that critical simplices cannot be completely avoided.

A discrete Morse function is a *perfect Morse function* whenever the number of critical  $k$ -simplices equals the  $k$ -th Betty number of the complex. It is equivalent to the condition that the number of all critical simplices equals the sum of Betty numbers.

Perhaps the main idea of discrete Morse theory, as summarized in the following theorem of R. Forman, is to contract all matched pairs of simplices and to reduce

the simplicial complex  $K$  to a cell complex (where critical simplices correspond to the cells).

**THEOREM 2.1.** [6, 7] *Assume that a discrete Morse function on a simplicial complex  $K$  has a single zero-dimensional critical simplex  $\sigma^0$  and that all other critical simplices have the same dimension  $N > 1$ . Then  $K$  is homotopy equivalent to a wedge of  $N$ -dimensional spheres.*

*More generally, if all critical simplices, aside from  $\sigma^0$ , have dimension  $\geq N$ , then the complex  $K$  is  $(N - 1)$ -connected.*

### 3. Collectively unavoidable $r$ -tuples and Alexander $r$ -tuples of complexes

In this section we introduce the central objects of our paper. Our tacit assumption is that all complexes  $K$  are *proper subcomplexes* of  $2^{[n]}$  in the sense that  $K \subsetneq 2^{[n]}$ .

**DEFINITION 3.1.** An ordered  $r$ -tuple  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  of subcomplexes of  $2^{[n]}$  is *collective  $r$ -unavoidable* (we also say that  $\mathcal{K}$  is a *pigeonhole  $r$ -tuple* on  $[n]$ ), if for each ordered collection  $(A_1, \dots, A_r)$  of disjoint sets in  $[n]$  there exists  $i$  such that  $A_i \in K_i$ . The class of collective  $r$ -unavoidable complexes is denoted by  $CU_{r,n}$ , or by  $CU_r$  if  $n$  is fixed or clear from the context.

On closer inspection, the definition can be usefully rephrased as follows. For the ordered  $r$ -tuple  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  and for an ordered disjoint collection  $(A_1, \dots, A_r)$  of subsets of  $[n]$ , we construct a bipartite graph  $\Gamma \subset K_{r,r}$ , where by definition there is an edge  $(i, j) \in \Gamma$  if and only if  $A_i \notin K_j$ . Then the collective  $r$ -unavoidability of  $\mathcal{K}$  is equivalent to the condition that the graph  $\Gamma$  does not contain a complete matching (does not satisfy the *marriage condition* of Hall's classical theorem). We therefore conclude that the pigeonhole property does not depend on the ordering of simplicial complexes.

**REMARK 3.1.** The bipartite graph  $\Gamma = \{(i, j) \in [r]^2 \mid A_i \notin K_j\}$  interpretation naturally leads to an extension of Definition 3.1 to the case of collections  $\mathcal{K} = \langle K_1, \dots, K_s \rangle$  where  $s$  is not necessarily equal to  $r$ . Note however that the symmetric case  $s = r$  is somewhat exceptional. For example the classical 'Hilfsatz' of Frobenius [13] implies that  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  is collective  $r$ -unavoidable if and only if for each ordered collection  $(A_1, \dots, A_r)$  of disjoint sets in  $[n]$  there exists a pair  $(S, T)$  of subsets of  $[r]$ , such that  $|S| + |T| = r + 1$ , and  $A_i \in K_j$  for each  $i \in S$  and  $j \in T$ .

It is easy to characterize all pigeonhole 2-tuples:  $(K_1, K_2)$  is collective unavoidable if and only if  $K_1^\circ \subset K_2$  (or equivalently  $K_2^\circ \subset K_1$ ).

For an  $r$ -tuple of complexes  $\langle K_1, \dots, K_r \rangle$  we shall use a natural partial ordering on the set of all set of pairwise disjoint  $r$ -tuples  $(A_1, \dots, A_r)$  with  $A_i \in K_i$ : say that  $(A_1, \dots, A_r) \leq (A'_1, \dots, A'_r)$  whenever  $\forall i : A_i \subseteq A'_i$ .

We also put a partial ordering on the set of all  $r$ -tuples of complexes by the same rule. So we automatically have the notion of *emphminimal unavoidable  $r$ -tuple* of complexes  $\langle K_1, \dots, K_r \rangle$ .

LEMMA 3.1. *Suppose that an  $r$ -tuple  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  of simplicial complexes is collective  $r$ -unavoidable. Then for each maximal disjoint collection  $(A_1, \dots, A_r)$  with  $A_i \in K_i$ , the set  $[n] \setminus \bigcup_{i=1}^r A_i$  contains at most  $r - 1$  elements.*

PROOF. Let  $\langle K_1, \dots, K_r \rangle$  be collective  $r$ -unavoidable, and let  $(A_1, \dots, A_r)$  be a maximal disjoint collection satisfying the condition  $A_i \in K_i$  for each  $i = 1, \dots, r$ . Suppose (for contradiction) that  $\{a_1, \dots, a_r\} \subset [n] \setminus \bigcup_{i=1}^r A_i$ . Then  $A'_i = A_i \cup \{a_i\} \notin K_i$  (by the maximality of the collection  $(A_1, \dots, A_r)$ ) and the collection  $(A'_1, \dots, A'_r)$  clearly violates the collective  $r$ -unavoidability condition for  $\langle K_1, \dots, K_r \rangle$ .  $\square$

DEFINITION 3.2. An  $r$ -tuple of complexes  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  on one and the same set of vertices  $[n]$  is an *Alexander  $r$ -tuple* if,

- (1) it is collective  $r$ -unavoidable, and
- (2) for each  $r$ -tuple of sets  $A_1, \dots, A_r$  with  $A_i \in K_i$  the set  $[n] \setminus \bigcup_{i=1}^r A_i$  has at least  $r - 1$  elements.

The class of Alexander  $r$ -tuples of subcomplexes of  $2^{[n]}$  is denoted by  $\mathcal{A}_r$  (or by  $\mathcal{A}_{r,n}$  if the set  $[n]$  of vertices should be emphasized).

PROPOSITION 3.1. *Given an Alexander  $r$ -tuple on  $[n]$ , for each maximal  $r$ -tuple of disjoint sets  $(A_1, \dots, A_r)$  with  $A_i \in K_i$  the set  $[n] \setminus \bigcup_{i=1}^r A_i$  has exactly  $r - 1$  elements.*

PROOF. This follows from Lemma 3.1 and the property (2) from the definition of the Alexander  $r$ -tuple (Definition 3.2).  $\square$

PROPOSITION 3.2. *An Alexander  $r$ -tuple of complexes is always a minimal pigeonhole  $r$ -tuple of complexes.*

PROOF. Assume  $\langle K_1, \dots, K_r \rangle$  is an Alexander  $r$ -tuple which is not a minimal collective  $r$ -unavoidable collection of complexes. This means that (possibly after a re-enumeration) the collection  $\langle K_1 \setminus \{A_1\}, K_2, \dots, K_r \rangle$  is also collective  $r$ -unavoidable for some maximal simplex  $A_1 \in K_1$ . As a consequence the restrictions  $\langle K_2|_{[n] \setminus A_1}, \dots, K_r|_{[n] \setminus A_1} \rangle$  form a collective  $(r-1)$ -unavoidable family of complexes. Lemma 3.1 implies that for any maximal disjoint collection  $(A_2, \dots, A_r)$  such that  $A_j \in K_j|_{[n] \setminus A_1}$  for each  $j = 2, \dots, r$ , the set  $[n] \setminus \bigcup_{i=1}^r A_i$  contains strictly less than  $r - 1$  elements. Then  $(A_1, \dots, A_r)$  is a maximal family satisfying  $A_i \in K_i$  for each  $i = 1, \dots, r$ , which is in contradiction with the condition (2) from Definition 3.2.  $\square$

The converse of Proposition 3.2 is in general not true.

EXAMPLE 3.1.

$$K_1 = K_2 = K_3 = \binom{[10]}{\leq 2} \sqcup \binom{[9]}{\leq 3}$$

is a minimal collective unavoidable 3-tuple which is not an Alexander 3-tuple.

EXAMPLE 3.2. A 2-tuple of complexes is an Alexander 2-tuple iff it is a pair of mutually dual complexes  $(K, K^\circ)$ .

The following examples describes the Alexander  $r$ -tuples  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  where each of the complexes  $K_i$  is a skeleton of the simplex  $2^{[n]}$ .

EXAMPLE 3.3. The collection of subcomplexes of  $2^{[n]}$ ,

$$\left( \binom{[n]}{\leq m_1}, \dots, \binom{[n]}{\leq m_r} \right)$$

is always an Alexander  $r$ -tuple, provided  $n = \sum_{i=1}^r m_i + r - 1$ .

EXAMPLE 3.4. Define a simplicial complex  $K \subset 2^{[6]}$  as the cone with apex 1 over the five-element set  $\{2, 3, 4, 5, 6\}$ . The complex  $K$  is essentially a graph with five edges  $\{1, 2\}, \{1, 3\}, \{1, 4\}, \{1, 5\}, \{1, 6\}$ . It is not difficult to see that  $\langle K, K, K \rangle$  is indeed an Alexander 3-tuple.

This example is the simplest case of a more general construction. For given integers  $m_1, m_2, \dots, m_r$ , let  $n = m_1 + m_2 + \dots + m_r + r - 1$ . Choose a simplex  $\Delta(C)$  spanned by  $C \neq \emptyset$  (where  $[n] \cap C = \emptyset$ ) and define the complexes,

$$K_i = \binom{[n]}{\leq m_i} * \Delta(C), \text{ for all } i = 1, 2, \dots, r.$$

It can be easily seen that  $\langle K_1, K_2, \dots, K_r \rangle$  is an Alexander  $r$ -tuple.

**3.1. Operations generating collective  $r$ -unavoidable complexes.** As demonstrated by the classification theorem (Theorem 5.1), Alexander  $r$ -tuples are scarce, and a very special class of simplicial complexes. The situation with the collective  $r$ -unavoidable complexes is quite the opposite, as illustrated by the following construction.

Let  $r \geq 2$  and let  $K_i \subset 2^{[n]}$  be a collection of not necessarily distinct simplicial complexes. Assume that the  $(r-1)$ -tuple  $\mathcal{K} = \langle K_1, K_2, \dots, K_{r-1} \rangle$  is NOT collective  $(r-1)$ -unavoidable on  $[n]$ .

Define  $R(\mathcal{K}) = R_r(\mathcal{K}) = R_r(K_1, K_2, \dots, K_{r-1})$  as the subcomplex of  $2^{[n]}$  where  $F \in R(\mathcal{K})$  if and only if there exists an ordered partition  $F_1 \uplus \dots \uplus F_{r-1} = F^c$  of the complement of  $F$  such that  $F_i \notin K_i$  for each  $i = 1, \dots, r-1$ .

Observe that  $R(\mathcal{K})$  is generated by the sets  $(F_1 \cup \dots \cup F_{r-1})^c$  where  $F_j$  are pairwise disjoint and  $F_j$  is a minimal nonface in  $K_j$  for each  $j = 1, \dots, r-1$ .

Note that  $R_r(\mathcal{K})$  can be described as the unique minimal simplicial complex  $Z$  such that  $\langle K_1, \dots, K_{r-1}, Z \rangle$  is a collective unavoidable  $r$ -tuple on  $[n]$ . Observe that  $\emptyset \in R_r(\mathcal{K})$  follows from the assumption that  $\mathcal{K}$  is not  $(r-1)$ -unavoidable.

DEFINITION 3.3. The complex  $R_r(\mathcal{K})$  is referred to as the *residual complex* of the  $(r-1)$ -tuple  $\mathcal{K} = \langle K_1, K_2, \dots, K_{r-1} \rangle$ .

Observe that in the case  $r = 2$  the residual complex of  $K \subset 2^{[n]}$  is precisely the Alexander dual,  $R(K) = K^\circ$ . More generally, for a complex  $K \subseteq 2^{[n]}$  we define the associated  $r^{\text{th}}$  residual complex  $R_r(K) = R(K_1, \dots, K_{r-1})$  where  $K_1 = \dots = K_{r-1} = K$ . Note that  $K$  is a minimal  $r$ -unavoidable complex if and only if  $R_r(K) = K$ .



PROBLEM 3.1. Find interesting examples of ordered collections of complexes  $\mathcal{K} = \langle K_1, K_2, \dots, K_{r-1} \rangle$  such that  $\langle K_1, K_2, \dots, K_{r-1}, R(\mathcal{K}) \rangle$  satisfies the condition (3<sup>#</sup>) (in Problem 1.1).

#### 4. Bier complexes

For each Alexander 2-tuple  $\langle K_1, K_2 \rangle = \langle K_1, K_1^\circ \rangle = \langle K_2^\circ, K_2 \rangle$ , the associated deleted join  $K_1 *_{\Delta} K_2$  is the standard Bier sphere  $\text{Bier}(K_1) \cong \text{Bier}(K_2)$  (Example 3.2). This observation motivates the following definition.

DEFINITION 4.1. Suppose that  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  is an Alexander  $r$ -tuple of complexes  $K_i \subset 2^{[n]}$ . Then the associated *Bier complex* is defined as the deleted join,

$$\text{Bier}(\mathcal{K}) := \mathcal{K}_{\Delta}^{*r} = K_1 *_{\Delta} K_2 *_{\Delta} \dots *_{\Delta} K_r.$$

It is well known that the ‘join’ and the ‘deleted join’ operations commute (see for example [12, Lemma 6.4.3.]). The following lemma is a natural generalization.

LEMMA 4.1. Let  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  and  $\mathcal{L} = \langle L_1, \dots, L_r \rangle$  be two collections of simplicial complexes where  $K_i \subset 2^{[m]}$  and  $L_i \subset 2^{[n]}$  for each  $i = 1, \dots, r$ . Then,

$$(\mathcal{K} * \mathcal{L})_{\Delta}^{*r} \cong \mathcal{K}_{\Delta}^{*r} * \mathcal{L}_{\Delta}^{*r}$$

where by definition  $\mathcal{K} * \mathcal{L} := \langle K_1 * L_1, \dots, K_r * L_r \rangle$ .

PROOF. If  $A = A_1 \uplus \dots \uplus A_r \in \mathcal{K}_{\Delta}^{*r}$  and  $B = B_1 \uplus \dots \uplus B_r \in \mathcal{L}_{\Delta}^{*r}$  then  $A * B \in \mathcal{K}_{\Delta}^{*r} * \mathcal{L}_{\Delta}^{*r}$  corresponds to the simplex  $C_1 \uplus \dots \uplus C_r \in (\mathcal{K} * \mathcal{L})_{\Delta}^{*r}$  where  $C_i := A_i \uplus B_i$  for each  $i = 1, \dots, r$ .  $\square$

The following theorem is one of the main results of our paper. It says that the classes  $CU_r$  and  $\mathcal{A}_r$  respectively, satisfy the central properties (3) and (3<sup>#</sup>), listed in Problem 1.1.

THEOREM 4.1. Let  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  be a collection of subcomplexes of  $2^{[n]}$ .

- (1) The deleted join  $\mathcal{K}_{\Delta}^{*r} = K_1 *_{\Delta} K_2 *_{\Delta} \dots *_{\Delta} K_r$  of a collective  $r$ -unavoidable collection  $\mathcal{K}$  of complexes is always  $(n - r - 1)$ -connected.
- (2) The Bier complex  $\text{Bier}(\mathcal{K}) = K_1 *_{\Delta} K_2 *_{\Delta} \dots *_{\Delta} K_r$ , associated to an Alexander  $r$ -tuple  $\mathcal{K}$ , is a pure complex of dimension  $n - r$ , homotopy equivalent to a wedge of  $(n - r)$ -dimensional spheres.

The following corollary of the proof of Theorem 4.1 emphasizes the computational efficiency of the approach based on the discrete Morse function described in Section 7.

COROLLARY 4.1. For an Alexander  $r$ -tuple  $\mathcal{K}$  the number of spheres in the wedge  $\text{Bier}(\mathcal{K})$  can be efficiently calculated as the number of critical simplices of the discrete Morse function  $D$  constructed in Section 7.

The efficiency of the method is illustrated in Section 8 by the calculation of the number of spheres in the important particular case of the optimal multiple chessboard complex (Section 2.2).

Recall that the number of spheres in a wedge decomposition can be in principle calculated as the reduced Euler characteristic of the complex. This calculation is typically very slow and inefficient, as it is based on an ‘inclusion-exclusion’ type formula which involves enumeration of all simplices in  $\text{Bier}(\mathcal{K})$ .

**4.1. Bier complexes and discrete Morse theory.** The proof of Theorem 4.1 (Section 7) and the proofs of other connectivity results in this paper rely on Discrete Morse theory (Theorem 2.1). All our discrete Morse functions (DMF) are defined on deleted joins  $\mathcal{K}_\Delta^{*r} = K_1 *_\Delta \cdots *_\Delta K_r$  of complexes  $K_i \subset 2^{[n]}$  and they all have some common features.

A simplex  $\beta \in \mathcal{K}_\Delta^{*r}$  is usually recorded as a disjoint sum  $\beta = A_1 \uplus \cdots \uplus A_r$ , see [12, Sections 5 and 6]. We find it convenient (for bookkeeping purposes) to use an alternative ‘partition notation’  $\beta = (A_1, \dots, A_r; B)$  where  $B = [n] \setminus \bigcup_{i=1}^r A_i$ . To match a  $p$ -simplex  $\alpha^p = (A'_1, \dots, A'_r; B')$  with a  $(p+1)$ -simplex  $\beta^{p+1} = (A_1, \dots, A_r; B)$  is the same as to ‘migrate’ an element  $i \in B'$  to one of the sets  $A'_j$ . This is possible if  $\alpha^p$  is a facet of  $\beta^{p+1}$  i.e. if  $B' = B \uplus \{i\}$  for some  $i \in B'$ .

CAVEAT. In the paper, we simplify the notation by omitting the braces and by writing simply  $B \cup i$  instead of  $B \cup \{i\}$  (with the tacit assumption that  $i \notin B$ ). We also write  $j < B$  ( $j > B$ ) if  $j < i$  for each  $i \in B$  (respectively if  $j > i$  for each  $i \in B$ ).

## 5. Classification theorem for Alexander $r$ -tuples

THEOREM 5.1. *If  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  is an Alexander  $r$ -tuple, then*

- (1)  $r = 2$  and  $(K_1, K_2) = (K, K^\circ)$  is an Alexander pair of dual complexes (Example 3.2), or
- (2)  $r \geq 3$  and  $K_i = \binom{[n]}{\leq m_i}$  (Example 3.3) where  $n = m_1 + \cdots + m_r + r - 1$ , or
- (3)  $r \geq 3$  and  $K_i = \binom{[n]}{\leq m_i} * \Delta(C)$  (Example 3.4) where  $n = m_1 + \cdots + m_r + r - 1$  and  $\Delta(C) = 2^C$  is the simplex spanned by a nonempty set  $C$  such that  $C \cap [n] = \emptyset$ .

PROOF. Suppose that  $r \geq 3$ . A minimal nonsimplex of a simplicial complex  $K \subset 2^{[n]}$  is called a  $K$ -blocker. Equivalently,  $A \subset [n]$  is a  $K$ -blocker if  $A \notin K$  and  $\partial(A) \subset K$ .

Let  $\mathcal{A} = (A_1, \dots, A_r)$  be a maximal, pairwise disjoint  $r$ -tuple of sets in  $[n]$  such that  $A_i \in K_i$  for each  $i = 1, \dots, r$ . ( $\mathcal{A}$  is referred to as a maximal  $r$ -tuple in  $\mathcal{K}$  or simply a maximal  $r$ -tuple).

We say that a maximal  $r$ -tuple  $\mathcal{A} = (A_1, \dots, A_r)$  is  $A_r$ -maximal (similarly  $A_i$ -maximal for  $i \neq r$ ) if for each maximal  $r$ -tuple  $\mathcal{A}' = (A'_1, \dots, A'_r)$  in  $\mathcal{K}$ ,

$$(5.1) \quad A_r \subseteq A'_r \quad \Rightarrow \quad A_r = A'_r.$$

From here on we assume that  $\mathcal{A}$  is  $A_r$ -maximal so in particular if  $x \in A_\nu$  for  $\nu \neq r$  then  $A_r \cup \{x\} \notin K_r$ .

Since  $\mathcal{K}$  is an Alexander  $r$ -tuple we know (Proposition 3.1) that if  $\mathcal{A}$  is a maximal  $r$ -tuple then the set  $[n] \setminus \bigcup_{i=1}^r A_i = \{t_1, \dots, t_{r-1}\}$  has exactly  $(r-1)$  elements.

Let  $X_{\mathcal{A}} = X_1 \uplus \cdots \uplus X_{r-1} \uplus X_r$  be the associated ‘blocker partition’ where  $X_i := A_i \cup \{t_i\}$  for each  $i = 1, \dots, r-1$  and  $X_r := A_r = [n] \setminus \bigcup_{i=1}^{r-1} X_i$ . The name is justified by the fact that  $X_i$  is a  $K_i$ -blocker for each  $i = 1, \dots, r-1$ .

Indeed, suppose that  $X_\nu$  is not a  $K_\nu$ -blocker for some  $\nu = 1, \dots, r-1$ , which means that there exists  $x \in A_\nu$  such that  $X_\nu \setminus \{x\} \notin K_\nu$ . The maximality of  $A_r$  (implication (5.1)) guarantees that  $A_r \cup \{x\} \notin K_r$ . This is a contradiction since the partition  $\mathcal{Z} = \langle Z_1, \dots, Z_r \rangle$  where  $Z_\nu := X_\nu \setminus \{x\}$ ,  $Z_r := A_r \cup \{x\}$  and  $Z_j = X_j$  for  $j \notin \{\nu, r\}$  clearly violates the condition that  $\mathcal{K}$  is collective  $r$ -unavoidable.

Let  $V \subset [n]$ . We say that a simplicial complex  $K \subset 2^{[n]}$  is  $V$ -homogeneous if  $S \in K \Leftrightarrow \phi(S) \in K$  for each permutation  $\phi: V \rightarrow V$  and each  $S \subset V$ .

CLAIM 1. Each of the complexes  $\{K_j\}_{j=1}^{r-1}$  is  $X$ -homogeneous where  $X = \bigcup_{j=1}^{r-1} X_j = [n] \setminus A_r$ .

PROOF OF THE CLAIM 1. Let us show for illustration that  $K_1$  is  $X$ -homogeneous. This is deduced from the observation that for each bijection  $\phi: X \rightarrow X$ ,

$$(5.2) \quad (a) \quad \phi(X_1) \notin K_1 \quad \text{and} \quad (b) \quad \phi(\partial(X_1)) \subset K_1.$$

This is obvious if  $\phi(X_1) = X_1$ . Moreover, it is sufficient to establish (5.2) in the case when  $\phi$  is a transposition, say  $\phi(x_1) = x_2$  where  $x_1 \in X_1$  and  $x_2 \in X_2$  (the case  $x_2 \in X_j$  for  $j > 2$  is treated similarly).

(a) is equivalent to  $(X_1 \setminus \{x_1\}) \cup \{x_2\} \notin K_1$ . This is true since otherwise,

$(X_1 \setminus \{x_1\}) \cup \{x_2\} \in K_1$ ,  $X_2 \setminus \{x_2\} \in K_2$ ,  $A_3 \in K_3, \dots, A_{r-1} \in K_{r-1}$ ,  $A_r \in K_r$  would be a disjoint family of sets covering all but  $(r-2)$  elements of  $[n]$  (contradicting (2) in Definition 3.4).

In order to prove (b), let  $X_1 \setminus \{y\}$  be a facet of  $\partial(X_1)$  (the interesting case is  $y \neq x_1$ ). Then,  $X'_1 := \phi(X_1 \setminus \{y\}) = (X_1 \setminus \{x_1, y\}) \cup \{x_2\} \in K_1$ . Otherwise the disjoint collection,

$X'_1 \notin K_1$ ,  $(X_2 \setminus \{x_2\}) \cup \{x_1\} \notin K_2$ ,  $X_3 \notin K_3, \dots, X_{r-1} \notin K_{r-1}$ ,  $X_r \cup \{y\} \notin K_r$  would violate the collective  $r$ -unavoidability of  $\mathcal{K}$ . (Note that  $(X_2 \setminus \{x_2\}) \cup \{x_1\} = \phi(X_2) \notin K_2$  follows from (a)).  $\square$

Summarizing, we have so far established that for each  $j = 1, \dots, r-1$  the restriction of  $K_j$  on  $X$  is the complex  $\binom{X}{\leq m_j}$  where  $m_i$  is the cardinality of the set  $A_i$ . In particular the sets  $A_1, A_2, \dots, A_{r-1}$  can be replaced by any disjoint family  $A'_1, A'_2, \dots, A'_{r-1}$  of subsets of  $X$  such that  $|A'_i| = m_i$  for each  $i$ .

CLAIM 2. If  $x \in X$  then  $X_r \cup \{x\} = A_r \cup \{x\} \notin K_r$ .

PROOF OF THE CLAIM 2. By Claim 1 we can assume that  $x \in X \setminus \bigcup_{i=1}^{r-1} A_i$ . Then the assumption  $A'_r := A_r \cup \{x\} \in K_r$  would contradict the fact that  $\mathcal{A} = (A_1, \dots, A_r)$  is a maximal disjoint  $r$ -tuple of sets in  $[n]$  such that  $A_i \in K_i$  for each  $i = 1, \dots, r$ .  $\square$

It follows from Claim 2 that either  $X_r \cup \{x\} = A_r \cup \{x\}$  is a  $K_r$ -blocker (this corresponds to the case (2) of the theorem) or there exists a proper subset  $S \subset X_r$  such that  $S \cup \{x\}$  is a  $K_r$ -blocker. The following claim makes this observation more

precise by showing (eventually) that  $S \subset X_r$  is unique with this property (and in particular independent of  $x$ ).

CLAIM 3. Choose  $x \in X$ . Let  $X_r = S \uplus C$  be a partition of  $X_r$  such that  $\{x\} \cup S$  is a  $K_r$ -blocker, i.e., such that  $\{x\} \cup S \notin K_r$  and  $\partial(\{x\} \cup S) \subset K_r$ . Then  $T \cup C \in K_i$  for each  $T \subset X \setminus \{x\}$  of cardinality  $m_i$  where  $i = 1, \dots, r-1$ . Moreover,  $T \cup C$  is a facet (maximal simplex) of  $K_i$ .

REMARK 5.1. The case  $C = \emptyset$  is NOT ruled out. As it will turn out from the proof if  $X_r = S' \uplus C'$  is another decomposition such that  $\{x\} \cup S'$  is a  $K_r$ -blocker then  $S' = S$  and  $C' = C$ .

PROOF OF THE CLAIM 3. Assume that  $i = 1$  (the proof in other cases is analogous). Since the sets  $A_1, \dots, A_{r-1}$  can be replaced by any disjoint family  $A'_1, \dots, A'_{r-1}$  of subsets of  $X$  such that  $|A'_i| = m_i$  for each  $i$ , we assume that  $T = A_1$ . For a similar reason we can assume that  $x \notin \bigcup_{i=1}^{r-1} A_i$ .

Then  $A_1 \cup C \in K_1$  since otherwise,

$$A_1 \cup C \notin K_1, X_2 \notin K_2, \dots, X_{r-1} \notin K_{r-1}, \{x\} \cup S \notin K_r$$

would violate the collective  $r$ -unavoidability of  $\mathcal{K}$ .

Suppose that  $A_1 \cup C$  is not a facet of  $K_1$ . It follows that  $A_1 \cup C \cup \{z\} \in K_1$  for some  $z \in S$ , hence  $(S \cup \{x\}) \setminus \{z\} \in K_r$ . This is a contradiction since the family,

$$A_1 \cup C \cup \{z\} \in K_1, A_2 \in K_2, A_3 \in K_3, \dots, A_{r-1} \in K_{r-1}, (S \cup \{x\}) \setminus \{z\} \in K_r$$

is a disjoint family of sets covering all but  $(r-2)$  elements of  $[n]$  (contradicting (2) in Definition 3.4).  $\square$

It follows from Claim 3 that  $K_1$  and  $K_r$  can interchange roles. More explicitly  $B_r := S$  can be included in a disjoint family  $\mathcal{B} := \{B_j\}_{j=1}^r$  (replacing the family  $\{A_i\}_{i=1}^r$ ) where  $B_j = A_j$  for each  $j = 2, \dots, r-1$  and  $B_1 := A_1 \cup C$ .

Since (by Claim 3)  $B_1$  is a facet of  $K_1$ , we observe that the maximal  $r$ -tuple  $\mathcal{B}$  is  $B_1$ -maximal (in the sense of definition preceding implication (5.1)). As a consequence Claims 1, 2, and 3 are all valid for the  $r$ -tuple  $\mathcal{B}$ , with  $B_1$  playing the role of  $A_r$ .

In light of Claim 1 each of the complexes  $\{K_j\}_{j=2}^r$  is  $Y$ -homogeneous where  $Y = [n] \setminus (\{x\} \cup B_1)$  and  $x$  is an arbitrary element in  $[n] \setminus B_1$ . Moreover the decomposition  $B_1 = A_1 \uplus C$  corresponds to the decomposition  $X_r = A_r = S \uplus C$  in Claim 3.

From here it is not difficult to conclude that for each  $i = 1, \dots, r$  there is a decomposition  $K_i \cong W_i * F$  where  $W_i \cong \binom{[n]}{\leq m_i}$  and  $F$  is either empty or  $F = \Delta(C)$  is the simplex spanned by a finite, nonempty set  $C$ .  $\square$

COROLLARY 5.1. *If  $K = K_1 *_{\Delta} \dots *_{\Delta} K_r$  is a Bier complex (Definition 4.1) then either,*

- (1)  $r = 2$  and  $K = K_1 *_{\Delta} K_2 = K *_{\Delta} K^{\circ}$  is a Bier sphere, or
- (2)  $r \geq 3$  and  $K = \Delta_{n,r}^{m_1, \dots, m_r; 1}$  is an optimal chessboard complex where  $n = m_1 + \dots + m_r + r - 1$  (Section 2.2), or

- (3)  $r \geq 3$  and  $K = \Delta * [r]^{*k}$  where  $\Delta = \Delta_{n,r}^{m_1, \dots, m_r; 1}$  is an optimal chessboard complex and  $[r]^{*k} = [r] * \dots * [r]$  is the join of  $k \geq 1$  copies of the 0-dimensional complex  $[r]$ .

(Note that (2) is a formal ‘consequence’ of (3) if we allow  $k = 0$ .)

PROOF. Assume  $r \geq 3$ . It follows from Theorem 5.1 and Lemma 4.1 that  $K \cong E * F$  where (Section 2.2)  $E = \Delta_{n,r}^{m_1, \dots, m_r; 1}$  is an optimal chessboard complex and  $F$  is either empty (the case (2)) or  $F = (\Delta(C))_{\Delta}^{*r}$  for a nonempty set  $C$  of cardinality  $|C| = k$ . The proof is completed by the observation that,

$$(\Delta(C))_{\Delta}^{*r} \cong (\{p\}^{*k})_{\Delta}^{*r} \cong (\{p\}_{\Delta}^{*r})^{*k} \cong [r]^{*k}. \quad \square$$

## 6. Two perfect discrete Morse functions on the Bier sphere

We illustrate the method of constructing DMF on deleted joins (by the method of ‘migrating elements’) first in the case of classical Bier spheres.

It is known that Bier spheres are always shellable, see [2] and [5]. A method of Chari [4] can be used to turn this shelling into a perfect DMF on a Bier sphere. The construction of the ‘first perfect DMF’ on a Bier sphere (Section 6.1) essentially follows this path.

The ‘second perfect DMF’ (Section 6.2) differs from the first DMF, although the ‘migration rules’ look very similar. The advantage of the second DMF is that it can be generalized to Alexander  $r$ -tuples and the associated Bier complexes.

**6.1. First perfect DMF.** We construct a discrete vector field  $D_1$  on the Bier sphere  $\text{Bier}(K)$  in two steps:

- (1) We match the simplices

$$\alpha = (A_1, A_2; B \cup i) \quad \text{and} \quad \beta = (A_1, A_2 \cup i; B)$$

iff the following holds:

- (a)  $i < B$ ,  $i < A_2$  (that is,  $i$  is smaller than all the entries of  $B$  and  $A_2$ ).
- (b)  $A_2 \cup i \in K^\circ$ .

Before we pass to step 2, let us observe that the nonmatched simplices are labelled by  $(A_1, A_2; B \cup i)$  such that  $A_2 \in K^\circ$ , but  $A_2 \cup i \notin K^\circ$ . As a consequence, for nonmatched simplices  $A_1 \cup B \in K$ .

- (2) In the second step we match together the simplices

$$\alpha = (A_1, A_2; B \cup j) \quad \text{and} \quad \beta = (A_1 \cup j, A_2; B)$$

iff the following holds:

- (a) None of the simplices  $\alpha$  and  $\beta$  is matched in the first step.
- (b)  $j > B$ ,  $j > A_1$ .
- (c)  $A_1 \cup j \in K$ .

Observe that the condition (c) always holds (provided that the condition (a) is satisfied), except for the case  $B = \emptyset$ .

LEMMA 6.1. *The discrete vector field  $D_1$  is a discrete Morse function on the Bier sphere  $\text{Bier}(K)$ .*

PROOF. Since  $D_1$  is (by construction) a discrete vector field, it remains to check that there are no closed gradient paths. Observe that in each pair of simplices in the discrete vector field  $D_1$  there is exactly one *migrating element*. More precisely, in the case (1) the element  $i$  migrates to  $A_2$ , and in the case (2) the element  $j$  migrates to  $A_1$ .

The lemma follows from the observation that (along a gradient path) the values of the migrating element that move to  $A_2$  strictly decrease. Similarly, the values of migrating elements that move to  $A_1$  can only increase. This is certified through the following simple case analysis: (1) After a first step pairing comes a splitting of  $A_2$ . Then the gradient path terminates. (2) After a first step pairing (with migrating element  $i$ ) comes a splitting of  $A_1$ . The gradient path proceeds only if the splitted element is smaller than  $i$ . (3) After a second step pairing comes a splitting of  $A_1$ . Then the gradient path terminates. (4) After a second step pairing (with migrating element  $i$ ) comes a splitting of  $A_2$ . The gradient path proceeds only if the splitted element is bigger than  $i$ .

Let us illustrate this observation by an example which captures the above case analysis. Assume we have a fragment of a gradient path that contains two matchings of type 1. We have:

$$(A_1 \cup k, A_2; B \cup i) \rightarrow (A_1 \cup k, A_2 \cup i; B) \rightarrow (A_1, A_2 \cup i; B \cup k) \rightarrow (A_1, A_2 \cup k \cup i; B)$$

The migrating elements here are  $i$  and  $k$ . The definition of the matching  $D_1$  implies  $k < i$ . Otherwise  $(A_1, A_2 \cup i; B \cup k)$  is matched with  $(A_1, A_2; B \cup k \cup i)$ , and the path would terminate after its second term.  $\square$

It is not difficult to see that there are precisely two critical simplices in  $D_1$ :

- (1) An  $(n - 2)$ -dimensional simplex,  $(A_1, A_2; i)$  where  $A_1 < i < A_2$ , (this condition describes this simplex uniquely, in light of the fact that  $A_1 \in K$  and  $A_2 \in K^\circ$ ),
- (2) and the 0-dimensional simplex,  $(\emptyset, \{1\}; \{2, 3, 4, \dots, n\})$ .

(Here we make a simplifying assumption that  $\{1\} \in K^\circ$ , which can be always achieved by a re-enumeration, except in the trivial case  $K^\circ = \{\emptyset\}$ .)

**6.2. Second perfect DMF.** The construction of the second discrete vector field  $D_2$  is also in two steps:

The first step remains the same:

- (1) We match the simplices

$$\alpha = (A_1, A_2; B \cup i) \text{ and } \beta = (A_1, A_2 \cup i; B)$$

iff the following holds:

- (a)  $i < B$ ,  $i < A_2$  (that is,  $i$  is smaller than all elements in  $B$  and  $A_2$ ).
- (b)  $A_2 \cup i \in K^\circ$ .

Before we pass to the second step, let us remind ourselves that the nonmatched simplices are labelled by  $(A_1, A_2; B \cup i)$  such that  $A_2 \in K^\circ$ , but  $A_2 \cup i \notin K^\circ$ . As a consequence, for nonmatched simplices  $A_1 \cup B \in K$ .

(2) We match together the simplices

$$\alpha = (A_1, A_2, B \cup i \cup j) \text{ and } \beta = (A_1 \cup j, A_2, B \cup i)$$

iff the following holds:

- (a) None of the simplices  $\alpha$  and  $\beta$  was matched in the first step, i.e.,  
 $i < j$ ,  $i < B$ ,  $i < A_2$ , and  $i \cup A_2 \notin K^\circ$ .
- (b)  $j < B$ ,  $j < A_1 \setminus [1, i]$ .
- (c)  $A_1 \cup j \in K^\circ$ .

Note that condition (c) is always satisfied (provided that condition (a) above holds).

We omit the proof that  $D_2$  is indeed a discrete Morse function since a more general fact will be established in the proof of Theorem 4.1 (Section 7).

Finally we observe that, with the same simplifying assumption  $\{1\} \in K^\circ$ , the discrete vector fields  $D_2$  and  $D_1$  have the same critical simplices:

- (1)  $(A_1, A_2, i)$  such that  $A_1 < i < A_2$  and
- (2)  $(\emptyset, \{1\}; \{2, 3, 4, \dots, n\})$ .

## 7. Proof of Theorem 4.1

The proof of Theorem 4.1 is based on the construction of a discrete Morse function  $D$  on the deleted join  $\mathcal{K}_\Delta^{*r}$ , where  $\mathcal{K} = \langle K_1, \dots, K_r \rangle$  is a collective  $r$ -unavoidable collection of complexes.

We will demonstrate that:

- If the  $r$ -tuple  $\mathcal{K}$  is collective  $r$ -unavoidable, then the critical simplices of the discrete Morse field  $D$  may appear only starting with dimension  $n - r$  (except for the unique 0-dimensional simplex). This observation immediately implies the connectivity bound in Theorem 4.1, part (1).
- Under the stronger hypothesis that  $\mathcal{K}$  is an Alexander  $r$ -tuple, the discrete vector field  $D$  has a single 0-dimensional critical simplex, while all other critical simplices have one and the same dimension  $n - r$ . Theorem 4.1 (part (2)) is an immediate consequence. Moreover, a direct dimension count will establish the purity of the complex  $\mathcal{K}_\Delta^{*r}$ .

As in Section 3, a simplex  $\beta = A_1 \uplus \dots \uplus A_r \in \mathcal{K}_\Delta^{*r}$  is in the ‘partition notation’ recorded as  $\beta = (A_1, \dots, A_r; B)$  where  $B = [n] \setminus \cup_{i=1}^r A_i$ . More explicitly, an ordered partition  $(A_1, A_2, \dots, A_r; B)$  of  $[n]$  into  $r+1$  parts, corresponds to a simplex in  $\mathcal{K}_\Delta^{*r}$  if and only if,

- (1)  $A_i \in K_i$  for each  $i = 1, \dots, r$ ;
- (2)  $\cup\{A_i\}_{i=1}^r \neq \emptyset$ , meaning that the partition  $(\emptyset, \dots, \emptyset, [n])$  is excluded.

Observe that the dimension of a simplex  $\beta = (A_1, \dots, A_r; B)$  is determined by the cardinality of  $B$ , indeed  $\dim(\beta) = n - |B| - 1$ .

Moreover, a facet of a simplex  $\beta = (A_1, A_2, \dots, A_r; B)$  is obtained by moving (we also say ‘migrating’) an element from one of the sets  $A_i$  to  $B$ . For example,  $(\{1, 2\}, \{6\}, \{5\}; \{3, 4, 7\})$  is a facet of  $(\{1, 2\}, \{6, 7\}, \{5\}; \{3, 4\})$  obtained by the migration of the element  $7 \in A_2$ .

**Construction of the discrete Morse function  $D$ .** The discrete vector field  $D$  is described by a step-by-step construction, generalizing the construction of the discrete vector field  $D_2$  from Section 6.2.

In the first step we match the simplices,

$$\alpha = (A_1, A_2, \dots, A_r; B \cup i_1) \text{ and } \beta = (A_1 \cup i_1, A_2, \dots, A_r; B)$$

iff the following holds:

- (1)  $i_1 < B, i_1 < A_1$ ;
- (2)  $A_1 \cup i_1 \in K_1$ .

In other words a simplex  $\alpha = (A_1, A_2, \dots, A_r; B')$  is matched (if possible) with the simplex  $\beta = (A'_1, A_2, \dots, A_r; B)$  obtained from  $\alpha$  by migrating the minimum  $i_1$  of the set  $B' = B \cup i$  into  $A'_1 = A_1 \cup i$  (provided  $i_1 < A_1$  and  $A'_1 \in K_1$ ).

Observe that many simplices are matched already in this step. Indeed, for  $\alpha = (A'_1, A_2, \dots, A_r; B')$  let  $i_1 = \min(A'_1 \cup B')$ . If  $i_1 \in B'$  then  $\alpha$  is matched with the simplex obtained by migrating  $i_1$  from  $B'$  to  $A'_1$ . If  $i_1 \in A'_1$ ,  $\alpha$  is obtained by the migration of  $i_1$  from its facet  $\gamma = (A_1, A_2, \dots, A_r; B)$ , where  $A_1 = A'_1 \setminus i_1$  and  $B = B' \cup i_1$ .

The remaining nonmatched simplices  $(A_1, A_2, \dots, A_r; B \cup i_1)$  fall into two types:

- (1) The first type:  
 $i_1 < B, i_1 < A_1$  and  $A_1 \cup i_1 \notin K_1$ .
- (2) The second type:  
 $B = \emptyset$  and  $A_1 = \emptyset$ .

Here we declare that the nonmatched simplices **of the second type** *will not participate* in matching in later steps of the construction, i.e., they will contribute to the critical simplices of  $D$ .

There is a single 0-dimensional nonmatched simplex,  $(\{1\}, \emptyset, \dots, \emptyset; \{2, \dots, n\})$ . Here (as in Section 6) we make a simplifying (nonessential) assumption that  $\{1\} \in K_1$ . (This condition can be easily satisfied by choosing a different linear order on  $\mathcal{K}$  and  $[n]$ , if necessary.)

We continue the construction by trying to migrate elements from  $B$  into  $A_2$  (in the second step), into  $A_3$  (in the third step), etc. Assume, inductively, that the first  $(k-1)$ -steps of the construction are completed.

In the  $k$ -th step of the construction we match the simplices,

$$\alpha = (\dots, A_k, \dots, A_r; B \cup i_1 \cdots \cup i_{k-1} \cup i_k) \text{ and}$$

$$\beta = (\dots, A_k \cup i_k, \dots, A_r; B \cup i_1 \cdots \cup i_{k-1})$$

iff the following holds:

- (1)  $\alpha$  and  $\beta$  are nonmatched simplices of **the first type** in all preceding steps,
- (2)  $i_k < B, i_k < A_k \setminus [1, i_{k-1}]$ ,
- (3)  $A_k \cup i_k \in K_k$ .

The remaining nonmatched simplices  $(A_1, A_2, \dots, A_k, \dots, A_r; B \cup i_1 \cdots \cup i_{k-1} \cup i_k)$  again fall into two types:

- (1) The first type:



- (a) the simplex is a first type nonmatched simplex on steps  $1, \dots, k-1$ ,
  - (b)  $i_k < B$ ,  $i_k < A_k \setminus [1, i_{k-1}]$ ,
  - (c)  $A_k \cup i_k \notin K_k$ .
- (2) The second type:
- (a) the simplex is a first type nonmatched simplex on steps  $1, \dots, k-1$ ,
  - (b)  $B = \emptyset$ ,  $A_k \subset [1, i_{k-1}]$ .

(As before we declare that the nonmatched simplices of the second type never participate in subsequent matchings.)

From the assumption that  $\mathcal{K}$  is a collective  $r$ -unavoidable collection of complexes we conclude that on the  $r$ -th step there are no nonmatched simplices of the first type. From here we deduce that the cardinality of  $B$  for critical simplices can vary from 0 to  $r-1$ , and in particular the dimension of any critical simplex is greater or equal than  $n-r$ . (The only exception being of course the 0-dimensional critical simplex  $(\{1\}, \emptyset, \dots, \emptyset; \{2, \dots, n\})$ ).

**An alternative description of the DMF.** It may be instructive to summarize the construction of the discrete Morse function  $D$  in the form of an ‘algorithm’ which describes the matching and lists the critical simplices.

For this purpose we introduce an operator  $\mathbf{a}$  which takes simplices

$$(A_1, \dots, A_r; B) \in K_1 *_{\Delta} \dots *_{\Delta} K_r$$

and maps them to strictly increasing  $r$ -tuples,

$$\mathbf{a} = (a_1 < a_2 < \dots < a_r) \in (\mathbb{N} \cup \{\infty\})^r,$$

by the following rule:

- 1:**  $a_1 := \min(B \cup A_1)$ ; if  $B \cup A_1 = \emptyset$  then  $a_1 = \dots = a_r := \infty$ .
- 2:**  $a_2 := \min((B \cup A_2) \setminus [1, a_1])$ ; if  $(B \cup A_2) \setminus [1, a_1] = \emptyset$  then  $a_k := \infty$  for all  $k \geq 2$ .
- ...
- i:**  $a_i := \min((B \cup A_i) \setminus [1, a_{i-1}])$ ; if  $((B \cup A_i) \setminus [1, a_{i-1}]) = \emptyset$  then  $a_k := \infty$  for all  $k \geq i$ .
- ...
- r:**  $a_r := \min\{(B \cup A_r) \setminus [1, a_{r-1}]\}$ ; if  $((B \cup A_r) \setminus [1, a_{r-1}]) = \emptyset$  then  $a_r := \infty$ .

We say that an element  $a_j$  of the  $r$ -tuple  $\mathbf{a}(A_1, \dots, A_r; B)$  is *potentially movable* if  $a_j \neq \infty$ . A potentially movable element  $a_j$  is *movable* if:

- (1) either  $a_j \in B$  and  $A_j \cup a_j \in K_j$ ,
- (2) or  $a_j \in A_j$ .

The *standard move* of a movable element  $a_j$  is the matching of:

- (1) either  $(A_1, \dots, A_j, \dots, A_r; B) \rightarrow (A_1, \dots, A_j \cup a_j, \dots, A_r; B \setminus a_j)$ ,
- (2) or  $(A_1, \dots, A_j \setminus a_j, \dots, A_r; B \cup a_j) \rightarrow (A_1, \dots, A_j, \dots, A_r; B)$ .

The following procedure finds the corresponding pair (if any) for each simplex. If the simplex is not matched, the algorithm reports that it is critical.

**Matching Algorithm.** A simplex  $(A_1, \dots, A_r; B) \in K_1 *_{\Delta} \dots *_{\Delta} K_r$  is paired with the simplex obtained by the standard move of the *minimal movable element* in  $\mathbf{a}(A_1, \dots, A_r; B)$ . If there are no movable elements, the simplex is critical.

Let us prove that the matching algorithm described above yields a pairing  $D'$  which equals  $D$ . Indeed, assume that  $(\alpha, \beta)$  is a pair in  $D$  with  $i_k$  moving from  $B$  to  $A_k$ . That is, this pair appears on the  $k$ -th step. Since  $i_k$  moves, it is a movable element in the sense of alternative description. Since  $(\alpha, \beta)$  was not paired on steps  $1, 2, \dots, k-1$ , it is a minimal movable element, and the same pair appears in  $D'$ . This argument certifies that the converse is also true: a pair from  $D'$  is also a pair from  $D$ .

PROPOSITION 7.1. *The “Matching Algorithm” describes a discrete Morse function  $D$ .*

It is clear that  $D$  is a discrete vector field. The proof of the acyclicity follows from the following lemmas.

LEMMA 7.1. *Assume the lexicographic order on the set  $(\mathbb{N} \cup \{\infty\})^r$ . Then the function  $\mathbf{a}$  decreases (nonstrictly) along any gradient path of the discrete vector field  $D$  described by the “Matching Algorithm”.*

PROOF. For any gradient path,  $\alpha_0^p \rightarrow \beta_0^{p+1} \rightarrow \alpha_1^p \rightarrow \beta_1^{p+1} \rightarrow \dots$ , we observe that  $\mathbf{a}(\alpha_i^p) = \mathbf{a}(\beta_i^{p+1})$  and  $\mathbf{a}(\beta_i^{p+1}) \leq \mathbf{a}(\alpha_{i+1}^p)$ .  $\square$

It immediately follows from Lemma 7.1 that the function  $\mathbf{a}$  must be constant along a cyclic gradient path (if it exists). In particular, along such a path the set of all potentially movable elements remains the same. The following lemma rules out this possibility.

LEMMA 7.2. *If the function  $\mathbf{a} = (a_1, a_2, \dots, a_r)$  is constant along a gradient path, then the path is acyclic.*

PROOF. Let us inspect a typical fragment of the gradient path,

$$(7.1) \quad \alpha_0^p \longrightarrow \beta_0^{p+1} \dashrightarrow \alpha_1^p,$$

which is more explicitly recorded as the path,

$$(\dots, A_r; B \cup a_k) \rightarrow (\dots, A_k \cup a_k, \dots; B) \dashrightarrow (\dots, A_k \cup a_k, \dots, A_m \setminus \nu, \dots; B \cup \nu).$$

If  $\nu < a_k$ , the value of  $a_j$  for some  $j \leq k$  would change, contrary to the assumption that  $\mathbf{a}$  is constant along the path. In the case  $\nu > a_k$  there are two possibilities.

The first possibility is that (by the matching algorithm) there is a matching,

$$(A_1 \dots, A_k, \dots, A_m \setminus \nu, \dots, A_r; B \cup a_k \cup \nu) \longrightarrow (A_1 \dots, A_k \cup a_k, \dots, A_m \setminus \nu, \dots; B \cup \nu)$$

or in other words a matching  $\gamma^{p-1} \longrightarrow \alpha_1^p$ , which would guarantee that the gradient path (7.1) terminates at  $\alpha_1^p$ .

This happens precisely if  $a_k$  is the minimal movable element in,

$$(7.2) \quad \mathbf{a}(A_1 \dots, A_k \cup \{a_k\}, \dots, A_m \setminus \nu, \dots, A_r; B \cup \nu).$$

The only possible scenario when  $a_k$  is not the minimal movable element in (7.2) is when  $m < k$  and the element  $a_m$  happens to be *movable* (as a consequence of  $\{a_m\} \cup (A_m \setminus \{j\}) \in K_m$ ).

Summarizing we observe that in the ‘worst case scenario’ the minimal movable element  $a_m$  of the simplex  $\alpha_1^p$  in (7.1) is strictly smaller than the minimal movable element  $a_k$  of the simplex  $\alpha_0^p$ . It follows that if this case persists, then the minimal movable element decreases along the path and the path must be acyclic.  $\square$

To establish the second statement in Theorem 4.1, we need the second half of Theorem 2.1. By Proposition 3.1 if  $\mathcal{K}$  is an Alexander  $r$ -tuple, then the complex  $\mathcal{K}_\Delta^{*r}$  is pure  $(n - r)$ -dimensional.

We end the proof with an efficient, combinatorial description of critical cells of the discrete Morse function  $D$ . The Corollary 4.1 is a consequence of the well known fact that the spheres in the wedge decomposition of  $\text{Bier}(\mathcal{K})$  are in one-to-one correspondence with the critical cells of  $D$ .

1. An  $(n - r)$ -dimensional simplex

$$(A_1, A_2, \dots, A_r; i_1 \cup i_2 \cup \dots \cup i_{r-1}) \quad \text{with} \quad i_1 < i_2 < \dots < i_{r-1}$$

is a critical simplex of the discrete Morse function  $D$  if and only if:

- (a)  $A_1$  avoids the segment  $[1, i_1]$ ,
- (b)  $A_k$  avoids the segment  $[i_{k-1}, i_k]$  for  $k \in [2, r - 1]$ ,
- (c)  $A_{r-1}$  avoids the segment  $[i_{r-2}, i_{r-1}]$ ,
- (d)  $A_k \cup i_k \notin K_k$  for  $k \in [1, r - 1]$ ,
- (e)  $A_r$  avoid the segment  $[i_{r-1}, n]$ .

2. There is a single 0-dimensional simplex:  $(\{1\}, \emptyset, \dots, \emptyset; \{2, 3, 4, \dots, n\})$ .

With this observation we complete the proof of Theorem 4.1 (Corollary 4.1).

EXAMPLE 7.1. The chessboard complex  $\Delta_{5,3}^{1,1,1;1}$  is the Bier complex associated to the Alexander 3-tuple  $\mathcal{K} = \langle K_1, K_2, K_3 \rangle$ , where  $K_1 = K_2 = K_3$  is the 0-dimensional skeleton of the 4-dimensional simplex  $\Delta([5])$ . Then the critical simplices  $(A_1, A_2, A_3; B)$  of the discrete Morse function  $D$  constructed in the proof of Theorem 4.1 are the following:

$$\begin{aligned} & (4, 5, 2; \{1, 3\}), (5, 4, 2; \{1, 3\}), (2, 5, 3; \{1, 4\}), (3, 5, 3; \{1, 4\}), (4, 5, 1; \{2, 3\}), \\ & (5, 4, 1; \{2, 3\}), (3, 5, 1; \{2, 4\}), (5, 1, 3; \{2, 4\}), (3, 1, 4; \{2, 5\}), (4, 1, 3; \{2, 5\}), \\ & (5, 2, 1; \{3, 4\}), (5, 1, 2; \{3, 4\}), (4, 2, 1; \{3, 5\}), (4, 1, 2; \{3, 5\}), (1, \emptyset, \emptyset; \{2, 3, 4, 5\}). \end{aligned}$$

## 8. Enumeration of critical simplices for optimal chessboard complexes

Optimal chessboard complexes (see Section 2.2 and Example 3.3 in Section 3) are our key examples of Alexander  $r$ -tuples for  $r \geq 3$ . In this section we enumerate critical simplices of  $\Delta_{n,r}^{m_1, \dots, m_r; 1}$  for  $n = m_1 + \dots + m_r + r - 1$ .

For a given simplex  $\beta \in \Delta_{n,r}^{m_1, \dots, m_r; 1}$ , let us encode the set of free columns  $1 \leq x_1 < \dots < x_{r-1} \leq n$  as an  $(r - 1)$ -tuple  $\mathbf{x} = (x_1, x_2, \dots, x_{r-1})$ . Observe that if (in the notation of Section 7)  $\beta = (A_1, \dots, A_n; B)$  then  $B = \{x_i\}_{i=1}^{r-1}$ .

Let  $\mathbf{b} = (b_1, b_2, \dots, b_r)$  denote the sequence that counts the number of rooks (in all rows) between consecutive free columns, i.e.,

$$b_1 = x_1 - 1, \quad b_2 = x_2 - x_1 - 1, \quad \dots, \quad b_r = r - x_r.$$

Let  $b_{i,j}$  denote the number of rooks in the  $i^{\text{th}}$  row between columns  $x_{j-1}$  and  $x_j$  (with the obvious interpretation of numbers  $b_{i1}$  and  $b_{ir}$ ). We know from the ‘‘critical simplices criterion’’ (found at the end of the proof of Theorem 4.1 in Section 7), that  $b_{ii} = 0$  for all  $i$ . For a given  $\mathbf{x}$ , all possible numbers of rooks between free columns in critical simplices (we ignore for a moment the order of rooks), corresponds to all nonnegative  $r \times r$  matrices,

$$B = \begin{pmatrix} b_{11} & b_{12} & \cdots & b_{1r} \\ b_{21} & b_{22} & \cdots & b_{2r} \\ \cdots & \cdots & \cdots & \cdots \\ b_{r1} & b_{r2} & \cdots & b_{rr} \end{pmatrix}$$

with nonnegative integers such that:

$$b_{11} = \cdots = b_{rr} = 0, \quad B \cdot \mathbf{1} = (m_1, \dots, m_r), \quad \mathbf{1}^t \cdot B = (b_1, b_2, \dots, b_r).$$

Denote the number of such matrices by  $R_{\mathbf{x}}$ . If the number of rooks between two consecutive free columns for each row is fixed, the number of all configurations is

$$\binom{b_1}{b_{11}, b_{21}, \dots, b_{r1}} \binom{b_2}{b_{12}, b_{22}, \dots, b_{r2}} \cdots \binom{b_r}{b_{1r}, b_{2r}, \dots, b_{rr}}.$$

Therefore, the number of all critical simplices of  $\Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}}$  is

$$\sum_{1 \leq x_1 < \cdots < x_{r-1} \leq r} R_{\mathbf{x}} \binom{b_1}{b_{11}, b_{21}, \dots, b_{r1}} \binom{b_2}{b_{12}, b_{22}, \dots, b_{r2}} \cdots \binom{b_r}{b_{1r}, b_{2r}, \dots, b_{rr}}$$

## 9. Discrete Morse function for a long chessboard complex

The ‘long’ chessboard complexes (described in Section 2.2) are not collective  $r$ -unavoidable complexes, let alone Alexander  $r$ -tuples. However, the construction of the discrete Morse function, described in Section 7, is sufficiently general and versatile to be applied in this case as well. This is very interesting since the existence of a perfect Morse function on this complex provides an alternative proof of the (critical case) of [8, Theorem 3.2]. Recall that this result paved the way for some new Tverberg–Van Kampen–Flores type results, including the [9, Theorem 1.2].

Recall that a multiple chessboard complex  $K = \Delta_{n,r}^{m_1, \dots, m_r; \mathbf{1}}$  is ‘long’ if  $n > m_1 + \cdots + m_r + r - 1$ . Following essentially the construction of the matching described in Section 7, one obtains the discrete Morse function  $D$  on  $K$  which has the following critical simplices.

With the exception of the unique 0-dimensional critical simplex, all other critical simplices are described as the configurations  $(A_1, A_2, \dots, A_r; B)$  for which there exist elements  $i_1 < i_2 < \cdots < i_r$  in  $B$  such that the following conditions are satisfied:

- (1)  $A_1$  avoids the segment  $[1, i_1]$ ,
- (2)  $A_k$  avoids the segment  $[i_{k-1}, i_k]$  for  $k \in [2, r]$ ,

- (3)  $A_k \cup i_k \notin K_k$  for  $k \in [1, r-1]$ ,  
(4)  $B \setminus \{i_1, \dots, i_r\} > i_r$ .

The condition (3) implies that all the critical simplices have one and the same dimension  $(m_1 + \dots + m_r - 1)$ .

EXAMPLE 9.1. The complex  $\Delta_{4,2}^{1,1;1}$  has 5 critical simplices of dimension 1:

$$(4, 3, \{1, 2\}) (3, 4, \{1, 2\}) (2, 4, \{1, 3\}) (4, 1, \{3, 2\}) (3, 1, \{4, 2\}).$$

The existence of a perfect discrete Morse function on the long, multiple chessboard complex  $\Delta_{n,r}^{m_1, \dots, m_r; 1}$  provides an alternative proof of the following theorem from [8]. (Two other proofs, both of them comparatively complex and nontrivial, relied respectively on a shelling construction, and the Nerve Lemma).

THEOREM 9.1. [8, Theorem 3.2] *The long chessboard complex is homotopy equivalent to a wedge of  $(m_1 + \dots + m_r - 1)$ -dimensional spheres.*

### 9.1. Enumeration of critical simplices for a long chessboard complex.

In this section we enumerate the critical simplices in the long chessboard complex  $\Delta_{n,r}^{m_1, \dots, m_r; 1}$ .

We use the notation as in Section 8. Recall that (for all  $i = 1, 2, \dots, r$ ) the integer  $b_{ij}$  evaluates the number of rooks in the  $i^{\text{th}}$  row between  $(j-1)^{\text{th}}$  and  $j^{\text{th}}$  free column (for  $j = 1, 2, \dots, r$ ). The distributions of rooks between columns (we again ignore for a moment the exact positions of the rooks) is encoded by the matrix  $B \in \text{Mat}_{r,r+1}(\mathbb{N}_0)$  where  $b_{11} = \dots = b_{rr} = 0$ ,  $B \cdot \mathbf{1} = (m_1, \dots, m_r)$ .

Also, in this case we have  $n - r - m_1 - m_2 - \dots - m_r$  free columns, and all of them are positioned behind the  $r^{\text{th}}$  free column.

Simply by counting all partitions of the corresponding multisets, we obtain the following formula for the number of all critical simplices in  $\Delta_{n,r}^{m_1, \dots, m_r; 1}$ ,

$$\sum_{\substack{B \in \text{Mat}_{r,r+1}(\mathbb{N}_0), \\ B \cdot \mathbf{1} = \mathbf{m}, b_{11} = \dots = b_{rr} = 0}} \binom{n - r - \sum_i m_i + \sum_j b_{jr}}{\sum_j b_{jr}} \prod_{i=1}^r \binom{b_{11} + b_{21} + \dots + b_{r1}}{b_{11}, b_{21}, \dots, b_{r1}}$$

For example, if  $r = 2$  we have that the number of critical simplices is

$$\sum_{b_{31}=1}^{m_1} \sum_{b_{32}=1}^{m_2} \binom{b_{13} + b_{23}}{b_{13}} \binom{n - 2 - m_1 - m_2 + b_{13} + b_{23}}{b_{13} + b_{23}}.$$

**Acknowledgements.** It is our pleasure to acknowledge the support and hospitality of the *Mathematisches Forschungsinstitut Oberwolfach*, where in the spring of 2016 this paper was initiated as a ‘research in pairs’ project. The construction from the proof of Theorem 4.1 and the constructions of Section 6 are supported by the Russian Science Foundation under grant 16-11-10039. R. Živaljević acknowledges the support of the Ministry of Education, Science and Technological Development of Republic of Serbia, Grant 174034. Supported by the Grants 174020 and 174034 of the Ministry of Education, Science and Technological Development of Serbia.

## References

1. A. Björner, *Topological methods*; in: R. Graham, M. Grötschel, and L. Lovász eds., *Handbook of Combinatorics*, 1819–1872. North-Holland, Amsterdam, 1995.
2. A. Björner, A. Paffenholz, J. Sjostrand, G. M. Ziegler, *Bier spheres and posets*, *Discrete Comput. Geom.* **34**(1) (2005), 71–86.
3. P. V. M. Blagojević, F. Frick, G. M. Ziegler, *Tverberg plus constraints*, *Bull. Lond. Math. Soc.* **46** (2014), 953–967.
4. M. K. Chari, *On discrete Morse functions and combinatorial decompositions*, *Discrete Math.* **217**(1–3) (2000), 101–113.
5. S. Lj. Čukić, E. Delucchi, *Simplicial shellable spheres via combinatorial blowups*, *Proc. Am. Math. Soc.* **135**(8) (2007), 2403–2414, (electronic).
6. R. Forman, *Morse theory for cell complexes*, *Adv. Math.* **134** (1998), 90–145.
7. ———, *A user’s guide to discrete Morse theory*, *Sémin. Lothar. Comb.* **48** (2002), Article B48c.
8. D. Jojić, S. T. Vrećica, R. T. Živaljević, *Multiple chessboard complexes and the colored Tverberg problem*, *J. Comb. Theory, Ser. A* **145** (2017), 400–425.
9. ———, *Symmetric multiple chessboard complexes and a new theorem of Tverberg type*, *J. Algebr. Comb.* **46** (2017), 15–31.
10. ———, *Topology and combinatorics of ‘unavoidable complexes’*, arXiv:1603.08472 [math.AT].
11. M. Jelić, D. Jojić, M. Timotijević, S. T. Vrećica, R. T. Živaljević, *Combinatorics of ‘unavoidable complexes’*, arXiv:1612.09487 [math.CO].
12. J. Matoušek, *Using the Borsuk–Ulam Theorem: Lectures on Topological Methods in Combinatorics and Geometry*, Springer, Heidelberg, 2003.
13. A. Schrijver, *On the history of combinatorial optimization (till 1960)*, in: K. Aardal, G. L. Nemhauser, R. Weismantel eds., *Handbook of Discrete Optimization*, Elsevier, Amsterdam, 2005, 1–68.
14. R. T. Živaljević, *Topological methods*, Chapter 14 in: J. E. Goodman, J. O’Rourke eds., *Handbook of Discrete and Computational Geometry*, Chapman and Hall / CRC 2004, 305–330.

Faculty of Science, University of Banja Luka  
 Republic of Srpska, Bosnia and Herzegovina  
 ducic68@teol.net

(Received 20 06 2017)

(Revised 30 10 2017)

Chebyshev Laboratory  
 St. Petersburg State University, Russia  
 geometr.nekrasov@yandex.ru

Mathematics & Mechanics Department  
 St. Petersburg State University  
 St. Petersburg Department of V. A. Steklov Institute of Mathematics  
 of the Russian Academy of Sciences, St. Petersburg, Russia  
 gaiane-panina@rambler.ru

Mathematical Institute of the Serbian Academy of Sciences and Arts  
 University of Belgrade, Serbia  
 rade@mi.sanu.ac.rs