

MEASURE AND INTEGRATION IN THE ALTERNATIVE SET THEORY

Miodrag Rašković

With the analysis point of view, there is a strong similarity between the saturated models of analysis (NA) and the Alternative Set Theory (AST). So, notions as for example “infinite external set”, “internal set” and “hyper-finite set” in NA have corresponding notions as “countable class”, “set” and “infinite” set in AST (“infinite” in AST meaning).

The internal definition principle is in relation with the comprehension schema in AST, and comprehension property in NA is in relation with the prolongation axiom.

This similarity between NA and AST becomes complete understandable, if one knows that the ultra-power of the set of hereditary-finite sets enriched with its subsets is a model for AST (see [5]).

On the other hand, AST allows us to make a natural fundation of analysis.

Our intention will be translating the notions and theorems from [2] to AST.

For basic motivations, notions, axioms, definitions and theorems for AST, one may consult [1]. We recall, that the class of natural numbers is N ,¹ finite natural number is FN ,² rational numbers is RN ,³ finite rational numbers is FRN ⁴ and bounded rational numbers is BRN .⁵ All of this classes are class-theoretically definable, as their relations and operations \leq , $+$ and \cdot are. The set of real numbers is $Real = BRN / \dot{=}$, where by $\dot{=}$ “infinitely near”⁶ relations is denoted.

$Real$ has all topological and algebraical property as the classical reals does. But for us, its selector $R \subseteq BRN$ ⁷ is more useful than the such $Real$ itself. R is

¹We model natural numbers in the manner of von Neumann.

² $n \in FN$ iff $(\forall X \subseteq n)$ (“ X is a set”).

³ $r \in RN$ iff $(\forall m, n \in Z)$ $(r = \langle m, n \rangle)$ and we write $r = \frac{m}{n}$, where $Z = N \cup \{ \langle 0, \alpha \rangle \mid \alpha \in N \}$.

⁴ $r \in FRN$ iff $(\exists m, n \in FZ)$ $(r = \frac{m}{n})$, where $FZ = FN \cup \{ \langle 0, \alpha \rangle \mid \alpha \in FN \}$.

⁵ $r \in BRN$ iff $(\exists n \in FN)$ $(|r| \leq n)$

⁶ $x = y$ iff $(\forall n \in FN)$ $(|x - y| \leq \frac{1}{n})$

⁷A class R is image of choice function on the class $Real$.

topologically isomorphic with *Real*, so that we can translate all interesting properties from *Real* to *R*.

For each $x \in R$, there is ${}^\circ x \in Real$ so that $x \in {}^\circ x$, and for each $y \in {}^\circ x$ we have $st(y) = x$. Observe that $st(x)$ and ${}^\circ x$ are not same.

For $A \subseteq R$, let ${}^\circ A = \{x \in Real \mid (\exists y) (y \in A \wedge y \in x)\}$.

The topological notions as an interval, open and closed set, convergence and so on, we can define in the usual manner (if we exchange a set with a class), but the role of N and RN play FN and FRN . The role of $*R$ play RN .

The set R omits some algebraic characteristic, which *Real* has. For example, in R , $(\exists x) (x = 2)$ is true, but not $(\exists x) (x = 2)$. However, this is not of an importance for us. R is more natural than *Real*, because $R \subseteq BRN$, while *Real* is the class of classes.

We can replace the saturation with the following theorems (see [1]).

THEOREM 1. *Each countable class is proper semiset.*

THEOREM 2. *Let X, Y be countable classes such that $\cap X \subseteq \cup Y$. Then there is a set u such that $\cup X \subseteq u \subseteq \cap Y$.*

THEOREM 3. *Let Z be set-theoretically definable class. Let X be a countable subsemiset of Z . If X is directed then there is a $u \in Z$ which is an upper bound of the elements of X ordered by inclusion. If X is dually directed then there is a $u \in Z$ which is a lower bound of the elements of X ordered by \subseteq .*

THEOREM 4. *Let $\{X_n, n \in FN\}$ be a sequence of revealed classes (for example, definable classes are revealed) such that for each $m \in FN$, $\cap\{X_n \mid n \leq m\}$ is non-empty. Then $\cap\{X_n \mid n \in FN\} \neq \emptyset$.*

1. Loeb and Lebesgue measure

Now, we will start to investigate Loeb measure in *AST*.

Let $x \approx n$ iff $(\exists f)(f: \frac{on}{1-\overline{f}} x)$.

Let Ω be a set, and $*\mathcal{P}(\Omega)$ the set of its subsets. It is easy to see that $*\mathcal{P}(\Omega)$ is a field of sets.

Let \bar{P} and P be functions so that for $\Omega \approx n$, $A \in *\mathcal{P}(\Omega)$ and $A \approx m$, we have $\bar{P}(A) = \frac{|A|}{|\Omega|} = \frac{m}{n}$ and $P(A) = st\bar{P}(A)$.

If $A \subseteq \Omega$, we can define inner measure so that $P_{\text{inner}}(A) = \sup\{P(B) \mid B \subseteq A, B \text{ set}\}$, and outer measure $P_{\text{outer}}(A) = \inf\{P(B) \mid A \subseteq B, B \text{ set}\}$.

DEF. 1. A class $A \subseteq \Omega$ is Loeb measurable iff $P_{\text{inner}}(A) = P_{\text{outer}}(A)$ and let us put $P(A) = P_{\text{inner}}(A) = P_{\text{outer}}(A)$.

Let $L(\Omega)$ be a class of Loeb measurable classes (which are sets or semisets).

LEMMA 1. *For each a class A we have:*

(1) $A \in L(\Omega)$ iff for each $\varepsilon > 0$ there are classes B and C so that $B \subseteq A \subseteq C$ and $P(C \setminus B) < \varepsilon$.

(2) $P_{\text{inner}}(A) = 1 - P_{\text{outer}}(\Omega \setminus A)$.

THEOREM 1. (Loeb) *The class $L(\Omega)$ is a σ -field and P is a σ -additive function.*

Proof. A proof that $L(\Omega)$ is a field and P is additive is as in (2).

To complete the proof, we must show that, if for each $i \in FN$, $A_i \supseteq A_{i+1}$, $A_i \in L(\Omega)$, $P(A_i) = r_i$, and $\lim_{i \rightarrow \infty} r_i = r$, then $\bigcap_{i \in FN} A_i \in L(\Omega)$ and $P(\bigcap_{i \in FN} A_i) = r$. We may apply the lemma 1 In this case is enough to choose sets B and C , so that for $A = \bigcap_{i \in FN} A_i$ and $\varepsilon > 0$, we have $B \subseteq A \subseteq C$, $P(B) \geq r - \varepsilon$ and $P(C) \leq r + \varepsilon$.

The sequence $\{r_i\}_{i \in FN}$ is descending and $\lim_{i \rightarrow \infty} r_i = r$, so we can choose $n \in FN^8$ and a set C , so that $A_n \subseteq C$ and $P(C) < r_n + \frac{\varepsilon}{2}$. But then $A \subseteq A_n \subseteq C$ and $P(C) < r_n + \frac{\varepsilon}{2} < r + \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = r + \varepsilon$, as required.

Let us find B . We can chose sets $B'_m \subseteq A_m$ with $P(B'_m) > r_m - 2^{-m}$.

Let $B_m = B'_1 \cap \dots \cap B'_m$. By induction, we can easily show that $P(B_m) > r_m - (1 - 2^{-m})\varepsilon$.

The class $X_m = \{B \in {}^* \mathcal{P}(\Omega) \mid \bar{P}(B) > r - \varepsilon \wedge B \subseteq B_m\}$ is revealed (and, more, a set). Also, we have $B_m \in X_m$, because $B_m \subseteq B_m$ and $\bar{P}(B_m) > r_m - (1 - 2^{-m}) > r - \varepsilon$. Therefore, $X_m \neq \emptyset$ and $X_1 \supseteq X_2 \supseteq \dots$. So, by Theorem 4. we have $\bigcap_{m \in FN} X_m \neq \emptyset$.

Let is $B \in \bigcap_{m \in FN} X_m$. Then, for each $m \in FN$, $B \subseteq B_m$ and $\bar{P}(B) \doteq P(B) > r - \varepsilon$. For each $m \in FN$, we have $B_m \subseteq A_m$. So $B \subseteq \bigcap_{m \in FN} A_m$, as required.

THEOREM 2. *For each $A \in L(\Omega)$, there is $B \in {}^* \mathcal{P}(\Omega)$ so that $P(A \Delta B) = 0$.*

The proof of the Theorem follows by the Theorem II.

THEOREM 3. *If $P(A) = 0$ and a semiset A is a countable union of sets, then there is a set $B \supseteq A$, so that $P(B) = 0$.*

The proof follows by the Theorem 3.

Our intention will be to define Lebesques measure on R and *Real*, and to show its conection with Loeb's measure.

Let $H \in N \setminus FN$ and $\Delta t = \frac{1}{H}$. Then, the class $T = \{0, \Delta t, 2\Delta t, \dots, 1\} = \{\langle k, H \rangle \mid k \in H\}$ is a set.

Let $[s, t] = \{x \in R \mid s \leq x \leq t\}$ and $st_T: T \rightarrow [0, 1]$, where $st_T = st \upharpoonright T$. Further on, we will write st instead of st_T , if it will not bring us to an ambiguity.

For $A \subseteq [0, 1]$ we have $st^{-1}(A) \subseteq T$ and $st(st^{-1}(A)) = A$, while for $B \subseteq T$ we have $st(B) \subseteq [0, 1]$ and $st^{-1}(st(B)) \supseteq B$.

Let $*[s, t] = \{x \in RN \mid s \leq x \leq t\}$ and $(s, t) = [s, t] \setminus \{s, t\}$.

⁸So that $r < r_n + \frac{\varepsilon}{2}$

Now, we give the definition of Lebesgue measure μ , on $[0, 1]$.

For $A = [s, t]$ we have $\mu([s, t]) = \mu((s, t)) = t - s$.

If $A \subseteq [0, 1]$, and A is open, then A can be written (uniquely) as a countable union of disjoint open intervals, $A = \bigcup_{n \in \mathbb{N}} A_n$, and then $\mu(A) = \sum_{n \in \mathbb{N}} \mu(A_n)$.

If $A \subseteq [0, 1]$ is closed, then $[0, 1] \setminus A$ is open, so $\mu(A) = 1 - \mu([0, 1] \setminus A)$.

For all other $A \subseteq [0, 1]$, we define $\mu_{\text{inner}}(A) = \sup\{\mu(B) \mid B \subseteq A, B \text{ is closed}\}$ and $\mu_{\text{outer}}(A) = \inf\{\mu(B) \mid A \subseteq B, B \text{ is open}\}$.

The class of Lebesgue measurable subclasses is $\text{Leb}[0, 1] = \{A \subseteq [0, 1] \mid \mu_{\text{inner}}(A) = \mu_{\text{outer}}(A)\}$, and for $A \in \text{Leb}[0, 1]$ we have $\mu(A) = \mu_{\text{outer}}(A) = \mu_{\text{inner}}(A)$.

Now, we give Fisher's theorem, which is connection between Loeb's and Lebesgue's measure. We need several lemmas for the proof of the theorem. However we omit their proofs, which are very similar to proofs in (2).

Only, we must use the Theorem 4. instead of ω -saturation.

LEMMA 2. *For each $r \in [0, 1]$, $st^{-1}(\{r\}) = \{t \in T \mid t \dot{=} r\} \in L(T)$ and has Loeb measure but any set containing it has a positive Loeb measure.*

LEMMA 3. *For all $s, t \in [0, 1]$, so that $s \leq t$ we have $P(T \cap^* [s, t]) = t - s$ and $P(st^{-1}[s, t]) = \mu([s, t]) = t - s$.*

LEMMA 4. *If $A \subseteq [0, 1]$ and A is closed, then $st^{-1}(A) \in L(T)$ and $\mu(A) = P(st^{-1}(A))$.*

LEMMA 5. *Let $B \subseteq T$ be set. Then $st(B)$ is closed.*

THEOREM 5. (Fisher) *for each $A \subseteq [0, 1]$, $A \in \text{Leb}[0, 1]$ iff $st^{-1}(A) \in L(T)$, and in this case, we have $\mu(A) = P(st^{-1}(A))$.*

The proof follows by the lemmas above.

We say ${}^\circ A \subseteq [0, 1]$ is Lebesgue measurable iff $A \in \text{Leb}[0, 1]$. Let us define ${}^\circ \mu$ with ${}^\circ \mu({}^\circ A) = \mu(A)$.

By Fishers theorem, we have ${}^\circ \mu(A) = P(st^{-1}(A))$.

2. Loeb measurable functions

First, will introduce two notions, and will give the theorem, which connects them.

DEF. 2. A class function F is Loeb measurable iff for each $r \in R$, we have $\{w \mid F(w) \leq r\} \in L(\Omega)$.

A set function f is lifting of a function $F: \Omega \rightarrow R$ iff $f: \Omega \rightarrow RN$ and $st f(w) = F(w)$ almost sure (on a set of measure 1).

THEOREM 5. *A class function F is Loeb measurable iff it has a lifting f . Moreover, if for each $w \in \Omega$, $|F(w)| \leq n$, then we can find a lifting f such that $|f(w)| \leq n$.*

PROOF: Let f lifts F on x , where $x = \{w \in \Omega \mid stf(w) = F(w)\}$ and $P(x) = 1$. For $r \in R$, a class $\{w \in x \mid F(w) \leq r\} = \cap \{w \in x \mid f(w) \leq r + \frac{1}{n}\}$ is measurable, as countable intersection of the measurable sets.

Let $Q = \{g_n \mid n \in FN\}$ be a sequence of finite rationals and F a Loeb measurable function.

The classes $A_n = \{w \mid F(W) \leq g_n\}$ are Loeb measurable, for each $n \in FN$. By the Theorem 2. there is a set B_n so that $P(B_n \Delta A_n) = 0$ and for $g_m \leq g_n$ we have $B_m \subseteq B_n$.

For each $n \in FN$, there is an function f_n such that:

$$(*) \quad (\forall m \leq n)(x \in B_m \leftrightarrow f_n(x) \leq g_m).$$

It is enough to take f_n , so that $f_n(B_0) = \{g_0\}$ and $f(B_{m+1} \setminus B_m) = \{g_m\}$, for each $1 \leq m \leq n$.

Let $X_n = \{f \mid \text{"}f \text{ is function"} \text{ and } (*)\}$. Then, we have $X_n \neq \emptyset$ and $X_1 \supseteq X_2 \dots$. According the Theorem 4., there is $f \in \cap_{n \in FN} X_n$. So we have $stf(w) = F(w)$ for we $\Omega \setminus \cup_{n \in FN} (A_n \Delta B_n)$ and

$$\begin{aligned} P\left(\Omega \setminus \bigcup_{n \in FN} (A_n \Delta B_n)\right) &= P(\Omega) - P\left(\bigcup_{n \in FN} (A_n \Delta B_n)\right) \\ &= 1 - \sum_{n \in FN} P(A_n \Delta B_n) = 1. \end{aligned}$$


So we have $st f(w) = F(w)$ almost surely.

The rest we can prove trivially, if we bound f in X_n by n .

DEF. 4. A set function f is lifting of $F_0: [0, 1] \rightarrow R$ iff $f: T \rightarrow RN$ and $st f(t) = F_0(st(t))$ almost surely on T .

LEMMA 6. A class function $F_0: [0, 1] \rightarrow R$ is Lebesgue measurable iff F_0 has a lifting $f: T \rightarrow RN$.

PROOF:



Let us define $F: T \rightarrow R$ by $F(t) = F_0(st^{-1}(t))$. By Fisher's theorem (Theorem 4) F_0 is Lebesgue measurable iff F is Loeb measurable iff (by Theorem 5.) F has a lifting f .

REMARK: If $f: T \rightarrow RN$ is almost surely finite set, then f is a lifting of some F , where is $F(w) = stf(w)$. Such F is Loeb measurable.

DEF. 5. A set function f is a uniform lifting of F iff for all $w \in T$, $stf(w) = F(w)$.

THEOREM 6. A class function F has a uniform lifting f iff $\{w \mid f(w) \leq r\}$, $\{w \mid f(w) \leq r\} \in \{x \mid \exists A(A: FN \rightarrow^* \mathcal{P}(T)^1 X = \bigcap_{n \in FN} A)\}$ for all $r \in R$.

The proof follows by Theorem 4.

Let $F_0: T \rightarrow Real$, so that ${}^\circ F(w) = F_0(w)$. The function F_0 is Loeb measurable iff for all $r \in R$, we have $\{w \in T \mid F(w) \leq r\} \in L(T)$.

3. Integration

Let Ω be infinite set, then there is $n \in N$ and a set function f so that $f: n \xrightarrow{\frac{on}{1-\gamma}} \Omega$. Then, we have $\sum_{w \in \Omega} F(w) = \sum_{m=1}^n F(f(m))$.

A bounded and Loeb measurable function $F: \Omega \rightarrow R$ is simple iff range (F) is finite.

DEF. 6. Let F be bounded and measurable function. If F is simple, then $\int_{\Omega} F(w)dw = \sum_{r \in \text{range}(F)} r \cdot P(F^{-1}(\{r\}))$. In general

$$\int_{\Omega} F(w)dw = \sup \left\{ \int_{\Omega} G(w)dw \mid G \text{ is simple and } G \leq F \right.$$

THEOREM 7. (Loeb) Let $F: \Omega \rightarrow R$ be bounded and Loeb measurable function and let $f: \Omega \rightarrow RN$ be a bounded lifting of F , then

$$\int_{\Omega} F(w)dw = st \sum_{w \in \Omega} f(w)\Delta w.$$

The proof is similar to the proof in (2).

THEOREM 8. Let $F: [0, 1] \rightarrow R$ be bounded Lebesgue measurable and let f be a lifting function of f . Then, we have $\int_{\Omega} F(t)dt = st \sum_{w \in \Omega} f(w)\Delta w$.

The proof follows easily by Theorem 4. and Theorem 7.

Let $F: \Omega \rightarrow RN$ be Loeb measurable and non-negative. Let us denote $\min(F(w), n)$ by $(F \wedge n)(w)$.

Then, we define $\int_{\Omega} F(w)dw = \lim_{n \rightarrow \infty} \int_{\Omega} (F \wedge n)(w)dw$.

A function F is Loeb integrable iff $\int_{\Omega} F(w)dw$ is finite.

We can denote $\max(F(w), 0)$ by $F^+(w)$ and $\min(F(w), 0)$ by $F^-(w)$. Then we have $F = F^+ + F^-$.

In general case, for unbounded F , we define

$$\int_{\Omega} F(w)dw = \int_{\Omega} F^+(w)dw - \int_{\Omega} -F^-(w)dw.$$

So, we see that a function F is Loeb integrable iff both F^+ and F^- are.

DEF. 7. Let f be non-negative set function so that $f: \Omega \rightarrow RN$. The function f is s -integrable iff $\sum_{w \in \Omega} f(w) \Delta w$ is finite, and

$$\lim_{\substack{n \rightarrow \infty \\ FN}} st \sum_{w \in \Omega} (f \wedge n)(w) \Delta w = st \sum_{w \in \Omega} f(w) \Delta w.$$

In general, a function f is s -integrable iff both f^+ and f^- are.

THEOREM 9. A function F is Loeb integrable iff F has an s -integrable lifting f .

The proof is similar to the proof in (2).

For $F_0: [0, 1] \rightarrow Real$ and $F_0(w) =^{\circ} F(w)$ we can define

$$\int_0^1 F(t)dt =^{\circ} \left(\int_0^1 F(t)dt \right).$$

REFERENCES

- [1] P. Vopěnka, *Mathematics in the Alternative Set Theory*, BSB B.G. Teubner Verlagsgesellschaft, Leipzig, 1979.
- [2] H. J. Keisler, *Hyperfinite Probability Theory and Probability Logic*, manuscript, Medison, 1979.
- [3] P. A. Loeb, *Conversion from Nonstandard to Standard Measure Spaces and Applications in Probability Theory*, TAMS, 1975.
- [4] K. D. Stroyan, W. A. J. Luxemburg, *Introduction to the Theory of Infinitesimals*, Academic Press, New York, 1976.
- [5] A. Sochor, *Diferential Calculus in the Alternative Set Theory*, Set Theory and Hierarchy Theory V, Lecture notes Math., 619, 1877.