

## ALMOST CONTINUITY AND NEARLY (ALMOST) PARACOMPACTNEES

Ilija Kovačević

**Abstract.** The purpose of the present paper is to investigate some properties of nearly (almost) paracompactness under almost continuous mappings.

Notation is standard except that  $\alpha(A)$  will be used to denote interior of the closure of  $A$ . The topology  $\tau^*$  is the semi-regularization of  $\tau$  and has as a base the regularly open sets from  $\tau$ .

### 1. Preliminaries

*Definition 1.1.* A subset of a space is said to be *regular open* iff it is the interior of some closed set or equivalently iff it is the interior of its own closure. A set is said to be *regularly closed* iff it is the closure of some open set or equivalently iff it is the closure of its own interior ([1]).

A subset is regularly open iff its complement is regularly closed.

*Definition 1.2.* A space  $X$  is said to be *almost regular* iff for any regularly closed set  $F$  and any point  $x \notin F$ , there exists disjoint open sets containing  $F$  and  $x$  respectively ([14]).

A space  $X$  is almost regular iff for each point  $x \in X$  and each regularly open set  $V$  containing  $x$  there exists a regularly open set  $U$  such that  $x \in U \subset \overline{U} \subset V$  ([14], Theorem 2.2).

*Definition 1.3.* A space  $X$  is *nearly paracompact* iff every regularly open cover of  $X$  has a locally finite open refinement ([18]).

*Definition 1.4.* A space  $X$  is *nearly strongly paracompact* iff every regularly open cover of  $X$  has a star finite open refinement ([8]).

*Definition 1.5.* A space  $X$  is said to be *almost paracompact* iff for every open covering  $\mathcal{U}$  of  $X$  there exists a locally finite family of open sets  $\mathcal{V}$  which refines  $\mathcal{U}$  and is such that the family of closures of members of  $\mathcal{V}$  forms a covering of  $X$  ([15]).

*Definition 1.6.* A space  $X$  is said to be *almost compact* iff each open covering of  $X$  has a finite subfamily the closures of whose members cover  $X$  ([15]).

*Definition 1.7.* A space  $X$  is said to be *almost strongly paracompact* iff for every open covering  $\mathcal{U}$  of  $X$  there exists a star finite family of open sets  $\mathcal{V}$  which refines  $\mathcal{U}$  and is such that the family of closures of members of  $\mathcal{V}$  forms a covering of  $X$  ([8]).

*Definition 1.8.* Let  $X$  be a topological space and  $A$  a subset of  $X$ . The set  $A$  is a  $\alpha$ -*almost paracompact* iff for every  $X$ -open covering  $\mathcal{U}$  of  $A$  there exists an  $X$ -locally finite family of  $X$ -open sets  $\mathcal{V}$  which refines  $\mathcal{U}$  and is such that the family of  $X$ -closures of members of  $\mathcal{V}$  forms a covering of  $A$  ([4]).

*Definition 1.9.* A space  $X$  is *locally almost paracompact* iff each point of  $X$  has an open neighbourhood  $U$ , such that  $\overline{U}$  is  $\alpha$ -almost paracompact ([4]).

*Definition 1.10.* Let  $X$  be a topological space, and  $A$  a subset of  $X$ . The set  $A$  is  $\alpha$ -*nearly paracompact* iff every  $X$ -regularly open cover of  $A$  has an  $X$ -open  $X$ -locally finite refinement which covers  $A$  ([6]).

*Definition 1.11.* A space  $X$  is *locally nearly paracompact* iff each point of  $X$  has an open neighbourhood  $U$  such that  $\overline{U}$  is  $\alpha$ -nearly paracompact ([5]).

*Definition 1.12.* Let  $X$  be a topological space and  $A$ , a subset of  $X$ . The set  $A$  is  $\alpha$ -*nearly strongly paracompact* iff every  $X$ -regularly open cover of  $A$  has an  $X$ -open star finite refinement which covers  $A$  ([7]).

*Definition 1.13.* A topological space  $X$  is called *locally nearly strongly paracompact* iff each point of  $X$  has an open neighbourhood  $U$  such that  $\overline{U}$  is an  $\alpha$ -nearly strongly paracompact subset of  $X$  ([7]).

*Definition 1.14.* A subset  $A$  of a space  $X$  is  $\alpha$ -*nearly compact (N-closed)* iff every  $X$ -regular open cover of  $A$  has a finite subcovering ([17]).

*Definition 1.15.* A topological space  $X$  is *locally nearly compact* iff each point has an open neighbourhood  $U$  such that  $\overline{U}$  is an  $\alpha$ -nearly compact subset of  $X$  ([2]).

*Definition 1.16.* A subset  $A$  of a space  $X$  is said to be *H-closed* iff for every  $X$ -open cover  $\{U_\alpha : \alpha \in I\}$  of  $A$ , there exists a finite subset  $I_0$  of  $I$  such that

$$A \subset \cup\{\overline{U}_\alpha : \alpha \in I_0\}.$$

*Definition 1.17.* A function  $f: X \rightarrow Y$  is said to be *almost continuous* iff for each point  $x \in X$  and each open neighbourhood  $V$  of  $f(x)$  in  $Y$  there exists an open neighbourhood  $U$  of  $x$  in  $X$  such that  $f(U) \subset \alpha(V)$  ([13]).

A function is almost continuous iff the inverse image of every regularly open set is open ([13], Theorem 2.2).

*Definition 1.18.* A function  $f: X \rightarrow Y$  is said to be *almost closed (resp. almost open)* iff for every regularly closed (resp. regularly open) set  $F$  of  $X$ ,  $f(F)$  is closed (resp. open) in  $Y$  ([13]).

*Definition 1.19.* A closed set  $F$  of  $(X, \tau)$  is said to be *star closed* iff  $F$  is closed in  $(X, \tau^*)$ . A function  $f: X \rightarrow Y$  is said to be *star closed* iff for every star closed set  $F$  of  $X$ ,  $f(F)$  is closed in  $Y$  ([11]).

LEMMA 1.1. *If a mapping  $f: X \rightarrow Y$  is almost continuous and almost closed, then*

- 1) *For each regularly closed set  $F$  of  $Y$ ,  $f^{-1}(F)$  is regularly closed in  $X$ ,*
- 2) *For each regularly open set  $V$  of  $Y$ ,  $f^{-1}(V)$  is regularly open in  $X$ .*

*Proof.* 1. Let  $F$  be any regularly closed subset of  $Y$ . Then  $f^{-1}(F)$  is closed, since  $f$  is almost continuous. Hence we have

$$\overline{[f^{-1}(F)]^0} \subset f^{-1}(F).$$

On the other hand, since  $f$  is almost continuous and  $F^0$  is a non empty regularly open subset of  $Y$ ,  $f^{-1}(F^0)$  is non empty open and hence we have

$$f^{-1}(F^0) \subset [f^{-1}(F)]^0 \subset \overline{[f^{-1}(F)]^0}.$$

Moreover, since  $f$  is almost closed and  $\overline{[f^{-1}(F)]^0}$  is a regularly closed subset of  $X$ ,  $f(\overline{[f^{-1}(F)]^0})$  is closed. Hence we have

$$F = \overline{F^0} \subset f(\overline{[f^{-1}(F)]^0}). \text{ Thus we obtain } f^{-1}(F) \subset \overline{[f^{-1}(F)]^0}.$$

This completes the proof of 1.

- 2) The proof of 2) follows easily from 1) and the following two facts:
  - a)  $f^{-1}(Y \setminus F) = X \setminus f^{-1}(F)$  for each subset  $F$  of  $Y$ .
  - b)  $F$  is regularly closed iff  $Y \setminus F$  is regularly open subset of  $Y$ .

## 2. Nearly (almost) paracompactness

THEOREM 2.1. *Let  $X$  be a nearly paracompact almost regular space. If  $f: X \rightarrow Y$  in an almost continuous, almost closed surjection, such that  $f^{-1}(y)$  is an  $\alpha$ -nearly compact subset of  $X$  for each point  $y \in Y$ , then  $Y$  is nearly paracompact almost regular.*

*Proof.* Since  $f: X \rightarrow Y$  is almost continuous and almost closed surjection such that  $f^{-1}(y)$  is  $\alpha$ -nearly compact for each  $y \in Y$ ,  $Y$  is almost regular ([5], Lemma 5).

Next, we shall shown that  $Y$  is nearly paracompact. Let  $\mathcal{U} = \{U_\alpha : \alpha \in I\}$  be any regularly open cover of  $Y$ . Then by Lemma 1.1  $f^{-1}(\mathcal{U}) = \{f^{-1}(U_\alpha) : \alpha \in I\}$  is a regularly open cover of  $X$ . Since  $X$  is nearly paracompact, there exists a locally finite regularly open refinement  $\mathcal{V} = \{V_\beta : \beta \in J\}$  or  $f^{-1}(\mathcal{U}) = \{f^{-1}(U_\alpha) : \alpha \in I\}$ . Since  $f$  is almost closed and  $f^{-1}(y)$  is  $\alpha$ -nearly compact for each  $y \in Y$ ,  $f(\mathcal{V}) = \{f(V_\beta) : \beta \in J\}$  is locally finite ([11], Lemma 2). Also,  $f(\mathcal{V})$  covers  $Y$  and is a refinement of  $\mathcal{U}$ .

Hence  $f(\mathcal{V})$  is a locally finite refinement of  $\mathcal{U}$  and thus  $Y$  is nearly paracompact ([18], Theorem 1.5).

**THEOREM 2.2.** *Let  $f$  be any almost closed, almost continuous and almost open mapping of a space  $X$  onto a space  $Y$  such that  $f^{-1}(y)$  is  $\alpha$ -nearly compact for each  $y \in Y$ . Then, the image of an  $\alpha$ -almost paracompact subset of  $X$  is an  $\alpha$ -almost paracompact subset of  $Y$ .*

*Proof.* Let  $A$  be any  $\alpha$ -almost paracompact subset of  $X$ . Let  $\mathcal{U} = \{U_\alpha : \alpha \in I\}$  be any  $Y$ -regularly open cover of a subset  $f(A)$ . Since  $f$  is almost continuous and almost open,

$$f^{-1}(\mathcal{U}) = \{f^{-1}(U_\alpha) : \alpha \in I\}$$

is an  $X$ -regularly open cover of  $A$ . Since  $A$  is  $\alpha$ -almost paracompact, then there exists an  $X$ -open  $X$ -locally finite family  $\mathcal{V} = \{V_\beta : \beta \in J\}$  which refines  $f^{-1}(\mathcal{U})$  and is such that  $\{\overline{V}_\beta : \beta \in J\}$  forms a covering of  $A$ . Consider the family

$$\mathcal{W} = \{\alpha(V_\beta) : \beta \in J\}.$$

Then  $\mathcal{W}$  is an  $X$ -locally finite  $X$ -regularly open family which refines  $f^{-1}(\mathcal{U})$  and is such that  $\{\overline{\alpha(V_\beta)} : \beta \in J\}$  forms a covering of  $A$ .

Since  $f$  is almost continuous, almost open and almost closed such that  $f^{-1}(y)$  is  $\alpha$ -nearly compact for each  $y \in Y$ ,  $\{f(\alpha(V_\beta)) : \beta \in J\}$  is a  $Y$ -locally finite family which refines  $\mathcal{U}$  ([11], Lemma 2). Since  $f$  is almost closed and almost continuous, therefore  $f(\overline{\alpha(V_\beta)}) = \overline{f(\alpha(V_\beta))}$ . Hence  $\{f(\alpha(V_\beta)) : \beta \in J\}$  is a  $Y$ -locally finite family of  $Y$ -open subsets refining  $\{U_\alpha : \alpha \in I\}$  and such that  $\{f(\alpha(V_\beta)) : \beta \in J\}$  is a covering of  $f(A)$ . Hence  $f(A)$  is  $\alpha$ -almost paracompact ([4] Lemma 1.3).

**COROLLARY 2.1.** *If  $f : X \rightarrow Y$  is any almost closed, almost continuous and almost open mapping of an almost paracompact space  $X$  onto a space  $Y$  such that  $f^{-1}(y)$  is  $\alpha$ -nearly compact for each  $y \in Y$ , then  $Y$  is almost paracompact.*

*Proof.*  $X$  is an  $\alpha$ -almost paracompact subset of  $X$ . Therefore  $f(X) = Y$  is an  $\alpha$ -almost paracompact subset of  $Y$ , i.e.  $Y$  is almost paracompact.

**COROLLARY 2.2.** ([15]. Theorem 6.3.1). *If  $f$  is a closed, continuous, open mapping of a space  $X$  onto a space  $Y$  such that  $f^{-1}(y)$  is compact for each  $y \in Y$ , then  $Y$  is almost paracompact if  $X$  is almost paracompact.*

**THEOREM 2.3.** *If  $f$  is an almost closed, almost continuous, almost open mapping of a locally almost paracompact space  $X$  onto a space  $Y$  such that  $f^{-1}(y)$  is  $\alpha$ -nearly compact for each  $y \in Y$ , then  $Y$  is locally almost paracompact.*

*Proof.* Let  $y \in Y$ . Then, there exists  $x \in X$  such that  $f(x) = y$ . Since  $X$  is locally almost paracompact, there exists an open neighbourhood  $U$  of  $x$  such that  $\overline{U}$  is  $\alpha$ -almost paracompact subset of  $X$ . Then  $\alpha(U)$  is a regularly open neighbourhood of  $x$  such that  $\overline{\alpha(U)} = \overline{U}$  is  $\alpha$ -almost paracompact subset of  $X$ .

Then,  $f(\alpha(U))$  is a  $Y$ -open neighbourhood of  $y$  such that  $f(\overline{\alpha(U)}) = \overline{f(\alpha(U))}$  is  $\alpha$ -almost paracompact subset of  $Y$ , therefore  $Y$  is locally almost paracompact.

**COROLLARY 2.3.** ([4], Theorem 2.2.) *If  $f$  is any closed, continuous, open mapping of a space  $X$  onto a space  $Y$  such that  $f^{-1}(y)$  is compact for each  $y \in Y$ , then  $Y$  is locally almost paracompact if  $X$  is locally almost paracompact.*

**LEMMA 2.1.** *Let  $f$  be any almost open and almost continuous mapping of a space  $X$  onto a space  $Y$  such that  $f^{-1}(G)$  is  $H$ -closed for each proper open subset  $G \subset Y$ . Then the image of any  $\alpha$ -nearly paracompact subset of  $X$  is  $\alpha$  nearly strongly paracompact subset of  $Y$ .*

*Proof.* Let  $A$  be any  $\alpha$ -nearly paracompact subset of  $X$ . Let  $\mathcal{U} = \{U_\alpha : \alpha \in I\}$  be any  $Y$ -regularly open cover of a subset  $f(A)$ . Since  $f$  is almost continuous and almost open,

$$f^{-1}(\mathcal{U}) = \{f^{-1}(U_\alpha) : \alpha \in I\}$$

is an  $X$ -regularly open cover of  $A$ . Since  $A$  is  $\alpha$ -nearly paracompact, then there exists an  $X$ -regularly open  $X$ -locally finite family  $\mathcal{V} = \{V_\beta : \beta \in J\}$  which refines  $f^{-1}(\mathcal{U})$  and is such that

$$A \subset \cup\{V_\beta : \beta \in J\}.$$

Since  $f$  is almost continuous, almost open that  $f^{-1}(G)$  is  $H$ -closed for each proper open subset  $G \subset Y$ ,  $\{f(V_\beta) : \beta \in J\}$  is a  $Y$ -open star finite family which refines  $\mathcal{U}$ , and is such that

$$f(A) \subset \cup\{f(U_\beta) : \beta \in J\}$$

([9]. Corollary 4.1). This implies that  $f(A)$  is  $\alpha$ -nearly strongly paracompact.

**THEOREM 2.4.** *If  $f$  is an almost open, almost closed and almost continuous mapping of a locally nearly paracompact space  $X$  onto a space  $Y$  such that  $f^{-1}(G)$  is  $H$ -closed for every proper open set  $G \subset Y$ , then  $Y$  is locally nearly strongly paracompact.*

*Proof.* Let  $y \in Y$  be any point. Then there exists  $x \in X$  such that  $f(x) = y$ . Since  $X$  is locally nearly paracompact, there exists an open neighbourhood  $U$  of  $x$  such that  $\overline{U}$  is  $\alpha$ -nearly paracompact subset of  $X$ . Then,  $\alpha(U)$  is a regularly open neighbourhood of  $x$  such that  $\overline{\alpha(U)} = \overline{U}$  is  $\alpha$ -nearly paracompact subset of  $X$ . Then,  $f(\alpha(U))$  is a  $Y$ -open neighbourhood of  $y$  such that  $\overline{f(\alpha(U))} = \overline{f(\alpha(U))}$  is  $\alpha$ -nearly strongly paracompact, therefore  $Y$  is locally strongly paracompact.

**LEMMA 2.2.** *A space  $X$  is almost strongly paracompact iff for every regularly open covering of  $X$  there exists a star finite family of open sets which refines it and the closures of whose members cover the space  $X$ .*

*Proof.* Only the "if" part needs to be proved.

Let  $\{U_\lambda : \lambda \in I\}$  be any open covering of  $X$ . Then  $\{\alpha(U_\lambda) : \lambda \in I\}$  is a regularly open covering of  $X$ . There exists an open star finite family  $\{H_\beta : \beta \in J\}$

which refines  $\{\alpha(U_\lambda) : \lambda \in I\}$  such that  $X \cup \{\overline{H}_\beta : \beta \in J\}$ . For each  $\beta \in J$  there exists  $\lambda(\beta) \in I$  such that  $H_\beta \subset \alpha(U_{\lambda(\beta)})$ . For each  $\beta \in J$ , let

$$M_\beta = H_\beta \setminus [\overline{U_{\lambda(\beta)}} \setminus U_{\lambda(\beta)}].$$

Since  $H_\beta \subset \alpha(U_{\lambda(\beta)}) \subset \overline{U_{\lambda(\beta)}}$ , therefore  $M_\beta = H_\beta \cap U_{\lambda(\beta)}$ .

Thus  $\{M_\beta : \beta \in J\}$  is a star finite family of open sets which refines  $\mathcal{U}$ . We shall prove that

$$X = \cup\{\overline{M}_\beta : \beta \in J\}.$$

Let  $x \in X$ . Then  $x \in \overline{H}_\beta$  for some  $\beta \in J$ . Now

$$\overline{M}_\beta = \overline{H_\beta \cap U_{\lambda(\beta)}} = \overline{H_\beta} \cap \overline{U_{\lambda(\beta)}} = \overline{H}_\beta.$$

Thus  $x \in \overline{M}_\beta$ . Hence  $\{M_\beta : \beta \in J\}$  is an open star finite family which refines  $\mathcal{U}$  and the closures of whose members cover the space  $X$ , therefore  $X$  is almost strongly paracompact.

**THEOREM 2.3.** *If  $f$  is an almost continuous, almost open mapping of an almost paracompact space  $X$  onto a space  $Y$  such that  $f^{-1}(G)$  is  $H$ -closed for each proper open set  $G \subset X$ , then  $Y$  is almost strongly paracompact.*

*Proof.* Let  $\{U_\alpha : \alpha \in I\}$  be any regularly open cover of  $Y$ . Then  $\{f^{-1}(U_\alpha) : \alpha \in I\}$  is a regularly open cover of  $X$ . There exists a locally finite family  $\{V_\beta : \beta \in J\}$  of open sets refining  $\{f^{-1}(U_\alpha) : \alpha \in I\}$  such that  $X = \cup\{\overline{V}_\beta : \beta \in J\}$ . Now,  $\{\alpha(V_\beta) : \beta \in J\}$  is a locally finite family of regularly open sets refining  $\{f^{-1}(U_\alpha) : \alpha \in I\}$  and is such that  $X = \cup\{\overline{\alpha(V_\beta)} : \beta \in J\}$ . Since  $f$  is almost open and  $f^{-1}(G)$  is  $H$ -closed for each proper open set  $G \subset Y$ ,  $\{f(\alpha(V_\beta)) : \beta \in J\}$  is a star finite family of open sets refining  $\{U_\alpha : \alpha \in I\}$  ([9], Lemma 4.2). Since  $f(\overline{\alpha(V_\beta)}) \subset \overline{f(\alpha(V_\beta))}$ , therefore  $\{f(\alpha(V_\beta)) : \beta \in J\}$  is a star finite family of open sets refining  $\{U_\alpha : \alpha \in I\}$  and is such that  $Y = \cup\{\overline{f(\alpha(V_\beta))} : \beta \in J\}$ .

Hence  $Y$  is almost strongly paracompact.

#### REFERENCES

- [1] Arya, S. P., *A note on nearly paracompact spaces*, Matematički vesnik **8** (23), (1971), 113–115.
- [2] Carnahan, D., *Locally nearly compact spaces*, Boll. Un Mat. Ital. (4)**6** (1972), 146–153.
- [3] Herrington, I., *Properties of nearly compact spaces*, Proc. Amer. Math. Soc. **45** (1974) 431–436.
- [4] Kovačević, I., *Locally almost paracompact spaces*, Zbornik radova PMF u Novom Sadu, **10**, (1980), 86–91.
- [5] Kovačević, I., *Locally nearly paracompact spaces*, Publ.Inst.Math. Belgrade **29** (43) (1981), 117–124.
- [6] Kovačević, I., *On nearly paracompact spaces*, Publ. Inst. Math. Belgrade, **25** (39), (1979), 63–69.
- [7] Kovačević, I., *On nearly strongly paracompact spaces*, Publ. Inst. Math. Belgrade, **27** (41), (1980), 125–134.

- [8] Kovačević, I., *On nearly strongly paracompact any almost strongly paracompact spaces*, Publ. Inst. Math. Belgrade, **23** (37), (1978), 109–116.
- [9] Mashhour, A. S. and I. A. Hasanein, *On  $S$ -closed any almost (nearly) strongly paracompactness* (to appear).
- [10] Noiri, T., *Almost continuity and some separation axioms*, Glasnik matematički Ser. III **9** (29) No 1, (1974), 131–135.
- [11] Noiri, T., *Completely continuous images of nearly paracompact spaces*, Matematički vesnik **1** (14) (24), (1977), 59–64.
- [12] Noiri, T.,  *$N$ -closed sets and almost closed mappings*, Glasnik matematički **4** (24), (1969), 89–99.
- [13] Singal, M. K. and A. R. Singal, *Almost continuous mappings*, Yokuhoma Math. J., **16** (1968), 63–73.
- [14] Singal, M. K. and S. P. Arya, *On almost regular spaces*, Glasnik matematički **4** (24) (1969), 89–99.
- [15] Singal, M. K. and S. P. Arya, *On  $m$ -paracompact spaces*, Math. Annal. **181** (1979), 119–133.
- [16] Singal, M. K. and A. Mathur, *On nearly compact spaces*, Boll. Un. Math. Ital. (4)**6** (1969) 702–710.
- [17] Singal, M. K. and A. Mathur, *On nearly compact spaces II*, Boll. Un. Math. Ital. (4)**9** (1974) 670–678.
- [18] Singal, M. K. and S. P. Aryas, *On nearly paracompact spaces*, Matematički vesnik **6** (21) (1969) 3–16.

University of Novi Sad  
Faculty of Technical Science  
Department of Mathematics  
Veljka Vlahovića 3  
21000 Novi Sad  
Yugoslavia