

A UNIFIED CLASS OF POLYNOMIALS*

Hukum Chand Agrawal**

Summary. In this paper we propose to study the polynomial set $\{f_n^{(\alpha)}\}(x)$ satisfying the functional relation

$$T(\Delta_\alpha)\{f_n^{(\alpha)}(x)\} = f_{n-1}^{(\alpha+1)}(x), \quad n = 1, 2, 3, \dots,$$

where $f(\alpha)_n(x)$ is the polynomial of degree n in x and T is the operator of infinite order defined by

$$T(\Delta_\alpha) = \sum_{k=0}^{\infty} h_k^{(\alpha)} \Delta_\alpha^{k+1}, \quad h_0^{(\alpha)} \neq 0,$$

in which $\Delta_\alpha\{f(\alpha)\} = f(\alpha+1) - f(\alpha)$.

1. Introduction. In his recent communication the author [1] studied the polynomial set $\{p_n^{(\alpha)}(x)\}$ satisfying the condition

$$\Delta_\alpha\{p_n^{(\alpha)}(x)\} = p_{(n-1)}^{(\alpha+1)}(x), \quad n = 1, 2, 3, \dots \quad (1.1)$$

A list of twelve polynomials is given which satisfy the above functional relation. In this paper we study another classification of polynomials which includes the class above as a particular case.

Consider the polynomial set $\{f_n^{(\alpha)}(x)\}$; $f_n^{(\alpha)}(x)$ are the polynomials of degree n in x , and the infinite operator

$$T(\Delta_\alpha) \equiv T = \sum_{k=0}^{\infty} h_k^{(\alpha)} \Delta_\alpha^{k+1}, \quad h_0^{(\alpha)} \neq 0, \quad (1.2)$$

in which $\Delta_\alpha\{f(\alpha)\} = f(\alpha+1) - f(\alpha)$.

*This work has been done under a fellowship of University Grants Commission, India.

**On study leave from Bundel Khand Post-graduate College, Jhansi (U. P.), India.

We are concerned here with those polynomials $f_n^{(\alpha)}(x)$ which satisfy the condition

$$T \left\{ f_n^{(\alpha)}(x) \right\} = f_{n-1}^{(\alpha+1)}(x), \quad n = 1, 2, 3, \dots \quad (1.3)$$

Obviously, for $h_0^{(\alpha)} = 1$ and $h_1^{(\alpha)} = h_2^{(\alpha)} = \dots = 0$ the condition (1.3) reduces to (1.1).

2. Certain Fundamental Properties of T -Operators.

THEOREM 1. *If $f_n^{(\alpha)}(x)$ is a simple set of polynomials in α , then there exists a unique difference operator of the form*

$$T = \sum_{k=0}^{\infty} h_k^{(\alpha)} \Delta_{\alpha}^{k+1}, \quad h^{(\alpha)} \neq 0 \quad (2.1)$$

where $h_k^{(\alpha)}$ is a polynomial of degree $\leq k$ in α , for which

$$T \left\{ f_n^{(\alpha)}(x) \right\} = f_{n-1}^{(\alpha+1)}(x), \quad n \geq 1, \quad (2.2)$$

Proof. From (2.1) and (2.2), we have

$$\sum_{k=0}^{n-1} h_k^{(\alpha)} \Delta_{\alpha}^{k+1} \left\{ f_n^{(\alpha)}(x) \right\} = f_{n-1}^{(\alpha+1)}(x). \quad (2.3)$$

The above equation shows that $h_k^{(\alpha)}$ is uniquely defined and is of degree $\leq k$, because $f_{n-1}^{(\alpha+1)}(x)$ is of degree $n-1$ for each $n (n \neq 0)$.

One can easily show that

THEOREM 2. *A necessary and sufficient condition that the simple sets of polynomials $f_n^{(\alpha)}(x)$ and $m_n^{(\alpha)}(x)$ belong to the same operator T is that there exist polynomial coefficients $b_k(x)$ of degree $\leq k$ in x and independent of α and n , such that*

$$f_n^{(\alpha)}(x) = \sum_{k=0}^n b_k(x) m_{n-k}^{(\alpha)}(x), \quad b_0(x) \neq 0. \quad (2.4)$$

Definition. Let $f_n^{(\alpha)}(x)$ be the simple set of polynomials belonging to the operator T defined by (1.2). If the maximum degree of the coefficient $h_k^{(\alpha)}$ in α is m , we say that the set $f_n^{(\alpha)}(x)$ is of α -type m . If the degree of $h_k^{(\alpha)}$ is unbounded, we say that $f_n^{(\alpha)}(x)$ is of α -type ∞ .

3. Some Properties of Sets of α -type Zero. According to the definition of α -type zero, any polynomial $f_n^{(\alpha)}(x)$ corresponding to the operator T is said to be of α -type zero, if

$$T = \sum_{k=0}^{\infty} h_k \Delta_{\alpha}^{k+1}, \quad h_0 \neq 0 \quad (3.1)$$

and

$$T \left\{ f_n^{(\alpha)}(x) \right\} = f_{n-1}^{(\alpha+1)}(x), \quad (3.2)$$

where h_k are independent of α .

Let $A(t)$ (independent of α) be a formal power series obtainable from the symbolic correspondence

$$T(A(t)) = t(1 + A(t)), \quad (3.3)$$

where $T(A(t))$ stands for $\sum_{k=0}^{\infty} h_k (A(t))^{k+1}$, ($h_0 \neq 0$),

Again, let

$$A(t) = \sum_{r=1}^{\infty} u_r t^r, \quad (3.4)$$

and denote

$$[A(t)]^k = \left[\sum_{r=1}^{\infty} u_r t^r \right]^k \quad \text{by} \quad \sum_{r=k}^{\infty} u_{rk} t^r.$$

Then, (3.3) on equating the coefficient of t^r on both sides, gives

$$u_r = \sum_{k=0}^r h_k u_{(r+1)(k+1)}, \quad r = 1, 2, \dots; \quad (3.5)$$

with $h_0 u_{11} = 1$.

THEOREM 3. *A necessary and sufficient condition for $f_n^{(\alpha)}(x)$ to be of α -type zero corresponding to the operator T is that $f_n^{(\alpha)}(x)$ possesses a generating function of the type*

$$(1 + A(t))^\alpha Q(x, t) = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n, \quad (3.6)$$

where¹ $A(t)$ and $Q(x, t)$ are independent of α and $A(t)$ is given by (3.3) and (3.4).

Proof. Transforming both sides of (3.6) by T , we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} t^n T \left\{ f_n^{(\alpha)}(x) \right\} &= (1 + A(t))^\alpha T(A(t)) Q(x, t) = \\ &= t(1 + A(t))^{\alpha+1} Q(x, t) = \sum_{n=0}^{\infty} f_n^{(\alpha+1)}(x) t^{n+1}, \end{aligned}$$

which gives $T \left\{ f_n^{(\alpha)}(x) \right\} = f_{n-1}^{(\alpha+1)}(x)$. Therefore, $f_n^{(\alpha)}(x)$ is of α -type zero.

Conversely, let $f_n^{(\alpha)}(x)$ be of α -type zero. Then from (3.2), we get

$$[(1-t)\Delta_\alpha - t] \sum_{n=0}^{\infty} t^n f_n^{(\alpha)}(x) = 0. \quad (3.7)$$

¹The generating function (3.6), in fact includes the generating functions given and studied by Appell [2], Sheffer [8], Brenke [4], Boas and Buck [3], and Rainville [6, §77].

Solving the above homogeneous-linear-difference equation, we get (3.6). Thus the theorem is proved.

COROLLARY 1. *A necessary and sufficient condition for $f_n^{(\alpha)}(x)$ to be of α -type zero and Sheffer A -type zero corresponding to the operator T and J^2 , respectively, is that $f_n^{(\alpha)}(x)$ possesses the generating function*

$$(1 + A(t))^\alpha \exp\{xH(t)\} = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x)t^n, \quad (3.8)$$

where $A(t)$ and $H(t)$ are independent of α and are given by (3.3) and $J(H(t)) = H(J(t)) = t$, respectively.

THEOREM 4. *Let $\{f_n^{(\alpha)}(x)\}$ be a set of α -type zero polynomials having the generating function*

$$(1 + A(t))^\alpha Q(x, t) = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x)t^n.$$

A necessary and sufficient condition for $f_n^{(\alpha)}(x)$ to satisfy the recurrence relation

$$nf_n^{(\alpha)}(x) = \sum_{r=0}^{n-1} (\alpha l_r + m_r(x)) f_{n-r-1}^{(\alpha)}(x), \quad n \geq 1 \quad (3.9)$$

is that there exist constants l_k and polynomial coefficients $m_k(x)$ of degree $\leq k$ in x , independent of α and n , given by

$$A'(t)/(1 + A(t)) = \sum_{r=0}^{\infty} l_r t^r \quad (3.10)$$

and

$$Q'(x, t)/Q(x, t) = \sum_{r=0}^{\infty} m_r(x)t^r, \quad (3.11)$$

respectively. Prime denotes differentiation with respect to t .

Proof. Differentiating both sides of (3.6), with respect to t , we get

$$\begin{aligned} \sum_{n=0}^{\infty} nt^n f_n^{(\alpha)}(x) &= t[\alpha A'(t)/(1 + A(t)) + Q'(x, t)/Q(x, t)](1 + A(t))^\alpha Q(x, t) \\ &= \sum_{n=0}^{\infty} \sum_r^n (\alpha l_r + m_r(x)) f_{n-r}^{(\alpha)}(x) t^{n+1}. \end{aligned}$$

²Here, as well as in what follows, J is defined by

$$J(D) \equiv J = \sum_{k=0}^{\infty} c_k D^{k+1}, \quad c_0 \neq 0, \quad D \equiv d/dx,$$

where the c_k 's are independent of α .

Equating the coefficients of t^n , we get (3.9). Thus, the sufficient part of the theorem is proved.

For, the necessary part, let

$$P = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x)t^n, \quad L = \sum_{n=0}^{\infty} \beta_n t^n \quad \text{and} \quad M = \sum_{n=0}^{\infty} \gamma_n t^n \quad (3.12)$$

where $r\beta_r = l_{r-1}1$, $\beta_0 = 0$, $r\gamma_r = m_{r-1}(x)$ and $\gamma_0 = 0$.

With these assumptions, (3.9) can be written as

$$\frac{dP}{dt} = \left[\alpha \frac{dL}{dt} + \frac{dM}{dt} \right] P,$$

which after some simplifications, gives

$$\alpha A'(t)/(1+A(t)) + Q'(x,t)/Q(x,t) = \alpha \sum_{r=0}^{\infty} \beta_r r t^{r-1} + \sum_{r=1}^{\infty} \gamma_r r t^{r-1}.$$

Since $A(t)$ and $Q(x,t)$ are independent of α , comparing the coefficient of α , we obtain

$$A'(t)/(1+A(t)) = \sum_{r=1}^{\infty} \beta_r r t^{r-1} = \sum_{r=1}^{\infty} l_{r-1} t^{r-1}$$

and

$$Q'(x,t)/Q(x,t) = \sum_{r=1}^{\infty} \gamma_r r t^{r-1} = \sum_{r=1}^{\infty} m_{r-1}(x) t^{r-1},$$

which are (3.10) and (3.11), respectively. Hence the theorem is proved.

Explicit form. The α -type zero polynomials satisfy the recurrence relation (3.9), viz.,

$$\begin{aligned} n f_n^{(\alpha)}(x) &= (\alpha l_0 + m_0(x)) f_{n-1}^{(\alpha)}(x) + (\alpha l_1 + m_1(x)) f_{n-2}^{(\alpha)}(x) + \dots \\ &\quad + (\alpha l_{n-1} + m_{n-1}(x)) f_0^{(\alpha)}(x). \end{aligned}$$

Eliminating $f_{n-1}^{(\alpha)}(x), f_{n-2}^{(\alpha)}(x), \dots, f_0^{(\alpha)}(x)$, we get the following explicit form for $f_n^{(\alpha)}(x)$

$$f_n^{(\alpha)}(x) = \sum \frac{s_1^{r_1} s_2^{r_2} \dots s_n^{r_n}}{r_1! r_2! \dots r_n!}, \quad (3.13)$$

where $\alpha l_k + m_k(x) = (k+1)s_{k+1}$, for $k = 0, 1, \dots, n-1$; $f_0^{(\alpha)}(x) = 1$ and the summation is taken over all positive integral values of r_1, r_2, \dots, r_n such that $r_1 + 2r_2 + \dots + nr_n = n$. (3.13) shows that $f_n^{(\alpha)}(x)$ is a polynomial of degree n in α .

THEOREM 5. *A necessary and sufficient condition for $f_n^{(\alpha)}(x)$ to be of α -type zero is that it satisfies a difference equation of the form*

$$\Delta_{\alpha} \left\{ f_n^{(\alpha)}(x) \right\} = \sum_{k=1}^{\infty} u_k f_{n-k}^{(\alpha)}(x), \quad (3.14)$$

where u_k is independent of n , α and is given by (3.3) and (3.4).

Proof. Applying Δ_α to both sides of (3.6), we obtain

$$\sum_{n=0}^{\infty} t_u \Delta_\alpha \left\{ f_n^{(\alpha)}(x) \right\} = A(t)(1 + A(t))^\alpha Q(x, t) = \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} u_k f_n^{(\alpha)}(x) t^{n+k}.$$

Equating the coefficients of t^n , we get (3.14).

Conversely, (3.14) can be written as

$$\Delta_\alpha \{P\} = A(t)P, \quad (3.15)$$

in the notation of (3.12).

The solution of the difference equation (3.15) is

$$P = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n = (1 + A(t))^\alpha Q(x, t).$$

Thus the theorem is proved.

The difference equation (3.15) can be generalized as $T(P) = t(1 + A(t))P$. Thus, $P = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n$ is also the solution of the difference equation $T(P) = t(1 + A(t))P$.

The following results can be proved easily.

COROLLARY 2. *If $f_n^{(\alpha)}(x)$ is of α -type zero and Sheffer A -type zero corresponding to the operator T and J , respectively, then so are the sets*

$$\{(\Delta_\alpha D) f_{n+1}^{(\alpha)}(x)\}, \{(\Delta_\alpha^2 D^2) f_{n+2}^{(\alpha)}(x)\}, \dots;$$

where $D \equiv d/dx$.

COROLLARY 3. *If $\{f_n^{(\alpha)}(x)\}$ is α -type and Sheffer A -type zero, then*

$$f_n^{(\alpha+\beta)}(x+y) = \sum_{r=0}^n f_{n-r}^{(\alpha)}(x) f_r^{(\beta)}(y). \quad (3.16)$$

(3.16) can also be written as

$$n! f_n^{(\alpha+\beta)}(x+y) = \sum_{r=0}^n \binom{n}{r} ((n-r)! f_{n-r}^{(\alpha)}(x)) (r! f_r^{(\beta)}(y)),$$

which shows that $\{n! f_n^{(\alpha)}(x)\}$ is a cross-sequence (for definition, see [7]).

The result (3.16) can be generalized as

$$f_n^{(\alpha_1 + \dots + \alpha_r)}(x_1 + \dots + x_r) = \sum_{m_1 + \dots + m_r = n} f_{m_1}^{(\alpha_1)}(x_1) \dots f_{m_r}^{(\alpha_r)}(x_r). \quad (3.17)$$

THEOREM 6. *If $\{f_n^{(\alpha)}(x)\}$ is of α -type zero corresponding to the operator Δ_α , then so is $\{f_n^{(\alpha+\beta_n)}(x)\}$ corresponding to the operator τ , defined by*

$$\tau(z) = z(1+z)^{-\beta} \quad (3.18)$$

where $z = u(t)/(1-u(t))$ and $t = u(t)(1-u(t))^\beta$.

Proof. With the help of the following generating relation [5]

$$\frac{1-u(t)]^{1-\alpha} F(x, u(t))}{1-(1+\beta)u(t)} = \sum_{n=0}^{\infty} f_n^{(\alpha+\beta_n)}(x) t^n \quad (3.19)$$

where $t = u(t)(1-u(t))^\beta$, the theorem can be proved easily.

4. A Characterization for α -type Zero Polynomials. Let us consider the set of polynomials $\{\psi_n^{(\alpha)}(x, A, Q)\}$ defined by

$$\psi_n^{(\alpha)}(x, A, Q) = \{E_\alpha^{-1}(1 + A(\nabla_\alpha))\}^\alpha Q(x, \nabla_\alpha) \frac{(\alpha)_n}{n!}, \quad (4.1)$$

where $A(t)$ and $Q(x, t)$ are formal power series in t independent of n .

(4.1), gives

$$\sum_{n=0}^{\infty} \psi_n^{(\alpha)}(x, A, Q) t^n = \{E_\alpha^{-1}(1 + A(\nabla_\alpha))\}^\alpha Q(x, \nabla_\alpha) (1-t)^{-\alpha}. \quad (4.2)$$

Now the application of the formula $\Phi(\nabla_\alpha)\{a^\alpha\} = a^\alpha \Phi(1-a^{-1})$, with $\Phi(x) = \sum_{r=0}^{\infty} b_r x^r$, reduces (4.2) to the form

$$(1 + A(t))^\alpha Q(x, t) = \sum_{n=0}^{\infty} \Psi_n^{(\alpha)}(x, A, Q) t^n. \quad (4.3)$$

Hence, we conclude that

THEOREM 7. *A necessary and sufficient condition for $f_n^{(\alpha)}(x)$ to be of α -type zero is that it is given by the operational formula*

$$f_n^{(\alpha)}(x) = \{E_\alpha^{-1}(1 + A(\nabla_\alpha))\}^\alpha Q(x, \nabla_\alpha) (\alpha)_n / n!, \quad (4.4)$$

and then the polynomial is defined by the generating function

$$(1 + A(t))^\alpha (x, t) = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n.$$

5. Algebraic Structure. Consider the set G_1 consisting of all α -type zero polynomials corresponding to the operator T as its elements, i. e.,

$$G_1 = \left\{ f_n^{(\alpha)}(x) : \tau(f_n^{(\alpha)}(x)) = f_{n-1}^{(\alpha+1)}(x) \right\}, \quad (5.1)$$

where τ is fixed and given by (3.1). For the sake of brevity, we denote the elements of G_1 by $f_n^{(\alpha)}, g_n^{(\alpha)}, \dots$.

THEOREM 8. *The set G_1 is an Abelian Group with respect to the operation $*$ defined by*

$$p_n^{(\alpha)} * q_n^{(\alpha)} = \sum_{k=0}^{\infty} p_{n-k}^{(0)} q_k^{(\alpha)}. \quad (5.2)$$

Before proving Theorem 8, we derive the following lemma:

$$\text{LEMMA 1.} \text{ If } (1 + A(t))^\alpha = \sum_{n=0}^{\infty} I_n^{(\alpha)} t^n, \text{ then} \quad (5.3)$$

$$(i) \ I_n^{(\alpha)} \in G_1,$$

$$(ii) \ I_r^{(0)} = \begin{cases} 0 & \text{for } r \neq 0 \\ 1 & \text{for } r = 0, \end{cases} \quad (5.4)$$

(iii) *the explicit form of any element $f_n^{(\alpha)} \in G_1$ is*

$$\sum I_r^{(\alpha)} f_{n-r}^{(0)}, \quad (5.5)$$

(iv) $I_n^{(\alpha)}$ *is the identity element for the set $(G_1, *)$.*

Proof of the Lemma 1. By Theorem 3, it is evident that $I_n^{(\alpha)} \in G_1$. Putting $\alpha = 0$ in (5.3), we obtain

$$1 = \sum_{n=0}^{\infty} I_n^{(0)} t^n.$$

On comparing the coefficients of various powers of t , we get (5.4).

For every $f_n^{(\alpha)} \in G_1$, by Theorem 3, we have

$$(1 + A(t))^\alpha Q(x, t) = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n,$$

in which the substitution $\alpha = 0$ gives

$$Q(x, t) = \sum_{n=0}^{\infty} f_n^{(0)}(x) t^n. \quad (5.6)$$

Now, putting the value of $(1 + A(t))^\alpha$, $Q(x, t)$ from (5.3) and (5.6), respectively, in (3.6), and equating the coefficients of t^n , we get the required result (5.5).

Since

$$f_n^{(\alpha)} * I_n^{(\alpha)} = \sum_{r=0}^n I_r^{(\alpha)} f_{n-r}^{(0)} = f_n^{(\alpha)}, \quad (\text{by (5.5)}),$$

and

$$I_n^{(\alpha)} * f_n^{(\alpha)} = \sum_{r=0}^n I_r^{(0)} f_{n-r}^{(\alpha)} = f_n^{(\alpha)}, \quad (\text{by (5.4)});$$

$I_n^{(\alpha)}$ is the identity element for the set $(G_1, *)$. Thus the lemma is proved.

Proof of Theorem 8. With the help of the lemma above, the theorem can be proved easily.

THEOREM 9. *The mapping $\rho: G_1 \rightarrow G_1$ such that*

$$\rho(f^{(\alpha)}) = T(f_n^{(\alpha)}), \quad \forall f_n^{(\alpha)} \in G_1 \quad (5.7)$$

is an isomorphism.

Proof. The proof is simple and hence omitted.

6. Polynomials of β -type m . Before defining β -type m polynomials consider the following example.

The classical Hermite polynomials are defined by means of the relation [6]

$$\exp(2xt - t^2) = \sum_{n=0}^{\infty} H_n(x)t^n/n!,$$

which gives

$$\Delta_x \{H_n(x)\} = 2H_{n-1}(x) + 4(x+1)H_{n-2}(x) + 8x(x+2)H_{n-3}(x) + \dots \quad (6.1)$$

The example above suggests the following extension of Theorem 5.

LEMMA 2. *For every polynomial $f_n^{(\alpha)}(x)$ there exist unique polynomial coefficients $u_k^{(\alpha)}$ of degree $\leq k$ in α and independent of n , such that*

$$\Delta_d \{f_n^{(\alpha)}(x)\} = u_1^{(\alpha)}f_{n-1}^{(\alpha)} + u_2f_{n-2}^{(\alpha)}(x) + \dots + u_n^{(\alpha)}f_0^{(\alpha)}(x), \quad (n \geq 1). \quad (6.2)$$

Definition. A set of polynomials $\{f_n^{(\alpha)}(x)\}$ is said to be of β -type m if in (6.2) the maximum degree of the coefficients $u_k^{(\alpha)}$ is m . If the degree of $u_k^{(\alpha)}$ is unbounded as $k \rightarrow \infty$ we say that the set $\{f_n^{(\alpha)}(x)\}$ is of β -type ∞ .

From Theorem 5 and Lemma 2, we conclude

THEOREM 10. *The set of polynomials $\{f_n^{(\alpha)}(x)\}$ is of β -type zero if, and only if, it is of α -type zero.*

caps Theorem 11. *A necessary and sufficient condition for the polynomial $f_n^{(\alpha)}(x)$ to be of β -type m is that*

$$\sum_{n=0}^{\infty} t^n f_n^{(\alpha)} = C \exp \left[\Delta_{\alpha}^{-1} \log \left(1 + \sum_{r=1}^{\infty} u_r^{(\alpha)} t^r \right) \right], \quad (6.3)$$

where C is an arbitrary periodic function of period unity in α .

Proof. From (6.2), we have

$$\Delta_\alpha \left\{ \sum_{n=0}^{\infty} f_n^{(\alpha)}(t^n) \right\} = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n \sum_{r=1}^{\infty} u_r^{(\alpha)} t^r$$

or

$$\left[\Delta_\alpha - \sum_{r=1}^{\infty} u_r^{(\alpha)} t^r \right] \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n = 0. \quad (6.4)$$

(6.4) is a homogeneous-linear-difference equation of order one, whose solution is (6.3). The converse part can be proved easily by transforming both sides of (6.3) by Δ_α , and hence the proof is omitted. Thus, the theorem is proved.

7. Polynomials of γ -type m . In this section we define another class of polynomials which are said to be of γ -type m , based on Lemma 3 (an extension of Theorem 4) given below. We also give a generating function for γ -type m polynomials.

LEMMA 3. *For every polynomial $f_n^{(\alpha)}(x)$ there exist unique polynomial coefficients $\nu_k^{(\alpha)}(x)$ of degree $\leq k$ in α and independent of n , such that*

$$n f_n^{(\alpha)}(x) = \nu_1^{(\alpha)}(x) f_{n-1}^{(\alpha)}(x) + \nu_2^{(\alpha)}(x) f_{n-2}^{(\alpha)}(x) + \cdots + \nu_n^{(\alpha)}(x) f_0^{(\alpha)}(x), \quad (n \geq 1). \quad (7.1)$$

Definition. A set of polynomials $\{f_n^{(\alpha)}(x)\}$ is said to be of γ -type m if in (7.1) the maximum degree of the coefficients $\nu_k^{(\alpha)}(x)$ is $(m+1)$ in α .

From the definition above and Theorem 4, it is evident that every α -type zero polynomial is also of γ -type zero.

THEOREM 12. *A necessary and sufficient condition for the polynomials $f_n^{(\alpha)}(x)$ to be of γ -type zero is that*

$$\sum_{n=0}^{\infty} t^n f_n^{(\alpha)}(x) = K \exp \left(\sum_{r=0}^{\infty} \nu_{r+1}^{(\alpha)}(x) t^{r+1} / (r+1) \right), \quad (7.2)$$

where K is an arbitrary constant (independent of t).

Proof. The sufficient part of the theorem can be proved easily by differentiating both sides of (7.2) with respect to t .

For the converse part write (7.1) as

$$\frac{\delta}{\delta t} \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n = \sum_{n=0}^{\infty} t^{n-1} \sum_{r=1}^n \nu_r^{(\alpha)}(x) f_{n-r}^{(\alpha)}(x) = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n \sum_{r=0}^{\infty} \nu_{r+1}^{(\alpha)}(x) t^r,$$

or

$$\left[\frac{\delta}{\delta t} - \sum_{r=0}^{\infty} \nu_{r+1}^{(\alpha)}(x) t^r \right] \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) t^n = 0. \quad (7.3)$$

On solving the difference equation (7.3), we get (7.2).

8. Generalized α -type Zero Polynomials. We conclude this paper by giving a generalization of α -type zero polynomials, introduced in § 3. We shall also give two characterizations for these polynomials.

Let us consider the following difference-operator of infinite order

$$T(\Delta_\alpha) \equiv T = \sum_{k=0}^{\infty} g_k \Delta_\alpha^{k+r}, \quad (8.1)$$

in which $g_0 \neq 0$, g_k ($k \geq 0$) are independent of α and r is some fixed positive integer.

Definition. Any polynomial $G_n^{(\alpha)}(x)$ for which there exists an operator T of the form (8.1), such that

$$T \left\{ G_n^{(\alpha)}(x) \right\} = G_{n-r}^{(\alpha+r)}(x), \quad (n = r, r+1, \dots) \quad (8.2)$$

where r is some fixed positive integer, we call a Generalized α -type zero polynomial. Obviously, for $r = 1$, (8.2) reduces to the condition required for $G_n^{(\alpha)}(x)$ to be of α -type zero.

THEOREM 13. *For any polynomial $G_n^{(\alpha)}(x)$ to be a Generalized α -type zero polynomial, the necessary and sufficient condition is that it satisfies a generating relation of the form*

$$\sum_{i=1}^r Q_i(x, t) (1 + B(\varepsilon_i t))^\alpha = \sum_{n=0}^{\infty} G_n^{(\alpha)}(x) t^n, \quad (8.3)$$

where, $B(t)$ is defined by the relation

$$T(B(t)) = t^r (1 + B(t))^r, \quad (8.4)$$

and $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r$ are the r roots of unity.

Proof. By operating both sides of (8.3), by T it can be shown easily with the help of (8.4) that $G_n^{(\alpha)}(x)$ satisfies the condition (8.2).

Conversely, let us write

$$\sum_{n=0}^{\infty} G_n^{(\alpha)}(x) t^n = G \quad (8.5)$$

Therefore, from (8.2) and (8.5), we have

$$[T - t^r E_\alpha^r] G = 0. \quad (8.6)$$

It is always possible to find out another difference-operator of the form

$$M(\Delta_\alpha) = \sum_{k=0}^{\infty} j_k \Delta_\alpha^{k+1}, \quad j_0 \neq 0 \quad (8.7)$$

such that

$$T(\Delta_\alpha) = (M(\Delta_\alpha))^r. \quad (8.8)$$

Hence from (8.6) and (8.8), we obtain

$$[(M(\Delta_\alpha))^r - (1 + \Delta_\alpha)^r t^r]G = 0.$$

Consequently, if $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r$, are the r roots of unity, we have

$$[M(\Delta_\alpha) - (1 + \Delta_\alpha)\varepsilon_i t]G = 0, \quad (i = 1, 2, \dots, r). \quad (8.9)$$

Solving the homogeneous-linear-difference equations above we get

$$G = \sum_{i=1}^r Q_i(x, t)(1 + B(\varepsilon_i t))^\alpha$$

where $B(t)$ is given by

$$M(B(t)) = t(1 + B(t)). \quad (8.10)$$

Therefore, the theorem is proved.

Like Theorem 2, one can show that

THEOREM 14. *If the set $\{G_n^{(\alpha)}(x)\}$ corresponds to the operator T , then a necessary and sufficient condition for the set $\{K_n^{(\alpha)}(x)\}$ to correspond also to the same operator T is that there exist polynomial coefficients $d_k(x)$ of degree $\leq h$ in x and independent of n and α , such that*

$$G_n^{(\alpha)}(x) = \sum_{i=0}^n d_i(x)K_n^{(\alpha)}(x), \quad d_0(x) \neq 0. \quad (8.11)$$

Finally, we give still another characterization for generalized α -type zero polynomials.

THEOREM 15. *Let $M(\Delta_\alpha)$ be the operator of type (8.7) and $u(\alpha)$ a function of bounded variation on $(0, \infty)$ such that $\int_0^\infty du(\alpha) \neq 0$. Then $G_n^{(\alpha)}(x)$ is a Generalized α -type zero polynomial if, and only if,*

$$\int_0^\infty \{M(\Delta_\alpha)\}^k G_n^{(\alpha-n)}(x) du(\alpha) = c_{n,k}, \quad (k = 0, 1, 2, \dots) \quad (8.12)$$

where $c_{n,k}$ are elements of an infinite triangular matrix, in which $c + n + r, k + r = c_{n,k}$.

Before proving the theorem above we first prove the following lemma:

LEMMA 4. *(8.12) is satisfied by one and only one $G_n^{(\alpha-n)}(x)$ for some given $M(\Delta_\alpha)$ and $u(\alpha)$ satisfying the conditions stated in the theorem above.*

Proof of Lemma 4. From Lemma 1 we know that the polynomials $I_n^{(\alpha)}$ defined by

$$(1 + B(t))^\alpha = \sum_{n=0}^{\infty} I_n^{(\alpha)} t^n$$

are of α -type zero corresponding to the operator $M(\Delta_\alpha)$, where $M(B(t)) = t(1 + B(t))$. We also have $I_0^{(\alpha)} = 1$.

Let $G_n^{(\alpha-n)}(x)$ satisfy (8.12); then we can write

$$G_n^{(\alpha-n)}(x) = \sum_{i=0}^n A(n, i, x) I_i^{(\alpha)}.$$

Therefore

$$\{M(\Delta_\alpha)\}^k G_n^{(\alpha-n)}(x) = \sum_{i=0}^{n-k} A(n, i+k, x) I_i^{(\alpha+k)}. \quad (8.13)$$

If we write

$$\int_0^{\infty} I_i^{(\alpha+k)} du(\alpha) = e_{i,k},$$

then from (8.12) and (8.13), we get

$$\sum_{i=0}^{n-k} A(n, i+k, x) e_{i,k} = c_{n,k}, \quad (k = 0, 1, 2, \dots, n). \quad (8.14)$$

Since $A(n, i, x) = 0$, if $i > n$, it follows that the determinant of the system (8.14) is $\prod_{k=0}^n e_{0,k} \neq 0$, and since $e_{0,k} = \int_0^{\infty} du(\alpha) \neq 0$, we conclude that $A(n, i, x)$ ($i = 0, 1, \dots, n$) are uniquely determined.

Proof of Theorem 15. Let $G_n^{(\alpha)}(x)$ be a Generalized α -type zero polynomial. Since $G_n^{(\alpha)}(x)$ is a polynomial of degree n in α , we have

$$[M(\Delta_\alpha)]^i G_n^{(\alpha-n)}(x) = 0, \quad \text{if } i > n$$

or

$$c_{n,i} = 0, \quad \text{if } i > n.$$

Again, since

$$M^k(\Delta_\alpha) G_n^{(\alpha-)}(x) = M^{k+r}(\Delta_\alpha) G_{n+r}^{(\alpha-n-r)}(x)$$

we get $c_{n,k} = c_{n+r,k+r}$.

Conversely, let $G_n^{(\alpha-n)}(x)$ satisfy (8.12). Then

$$\int_0^\infty M^{k+i}(\Delta_\alpha) G_{n+i}^{(\alpha-n-i)}(x) du(\alpha) = c_{n+i, k+i} = c_{n, k}.$$

The substitution

$$S_n^{(\alpha)}(x) = M^i(\Delta_\alpha) G_{n+i}^{(\alpha-n-i)}(x)$$

reduces (8.15) to

$$\int_0^\infty M^k(\Delta_k) S_n^{(\alpha)}(x) du(\alpha) = c_{n, k}.$$

But by Lemma 4, (8.12) is satisfied by the unique polynomial $G_n^{(\alpha-n)}$. Therefore $S_n^{(\alpha)}(x) = G_n^{(\alpha-n)}(x)$.

This completes the proof of the theorem.

I am grateful to Professor R. P. Agrawal for his kind guidance during the preparation of this paper.

REFERENCES

- [1] H. C. Agrawal, *On a unified class of polynomials*, (Under communication).
- [2] P. Appell, *Sur une classe de polynomes*, Ann. Sci. École Norm. Sup. **9** (1880), 119–144.
- [3] R. P. Boas, R. C. Buck, *Polynomials defined by generating relations*, Amer. Math Monthly, bf 63 (1956), 626–632.
- [4] W. C. Brenke, *On generating functions of polynomial systems*, Amer. Math Monthly, bf 52 (1945), 297–301.
- [5] J. W. Brown, *On the Sheffer A-type of certain modified polynomial sets*, Proc. Amer. Math. Soc., **23** (1969), 718–722.
- [6] E. D. Rainville, *Special Functions*, Macmillan, New York, 1965.
- [7] G. C. Rota, D. Kahaner, A. Odlyzko, *Finite operator calculus*, J. Math. Anal. Appl., **42** (1973), 685–760.
- [8] I. M. Sheller, *Some properties of polynomial sets of type zero*, Duke Math. J., **5** (1939), 590–622.

Department of Mathematics
Lucknow University
Lucknow, India

(Received 15 07 1980)