

A GENERALIZATION OF A THEOREM OF A. D. OTTO

Th. Exarchakos

Abstract. In this paper we prove that if G is a finite p -group of class c with G/G' of exponent p^r and L_i/L_{i+1} is cyclic of order p^r for $i = 1, 2, \dots, c-1$, where $L_i, i = 0, 1, \dots, c$ is the lower central series of G , then the order of G divides the order of the group $A(G)$ of automorphisms of G .

Introduction and notation. Let G be a finite p -group of class c . Let $G = L_0 \supset L_1 \supset \dots \supset L_c = 1$, $1 = Z_0 \subset Z_1 \subset \dots \subset Z_c = G$ be the lower and the upper central series of G respectively, where $L_1 = G' = [G, G]$ and $Z_1 = Z = Z(G)$. If G has no non-trivial abelian direct factor, then G is called a PN -group. A. D. Otto in [1] proved that if G is a PN -group with $|L_i/L_{i+1}| = p$ for all $i = 1, 2, \dots, c-1$ and $\exp(G/G') = p$, then the order of G divides the order of the group of automorphisms of G . We generalize this result by showing that if G is any finite p -group with L_i/L_{i+1} cyclic of order p' for all $i = 1, 2, \dots, c-1$ and $\exp(G/G') = p'$, then $|G|$ divides $A(G)$. We also show that the same result holds if Z_i/Z_{i-1} is cyclic of order p' , $i = 1, 2, \dots, c-1$, and $L_j = Z_{c-j}$ for some j , $1 \leq j \leq c-1$. Throughout this paper, G is a finite non-abelian p -group, $|G|$ is the order of G , $C(p^x)$ is the cyclic group of order p^x , $A(G)$, $I(G)$, $A_c(G)$ are the groups of automorphisms, inner automorphisms, central automorphisms of G .

* * *

We begin with

LEMMA 1. *Let G be a PN -group. If $\exp(G/G') \leq |Z|$, then $|A_c(G)| \geq |C/G'|$.*

Proof. Let $|G/G'| = p^m$ and $G/G' = C(p^{m_1}) \times C(p^{m_2}) \times \dots \times C(p^{m_t})$, where $m_1 \geq m_2 \geq \dots \geq m_t \geq 1$ and $\sum_{j=1}^t m_j = m$. Similarly let $|Z| = p^k$ and $Z = C(p^{k_1}) \times C(p^{k_2}) \times \dots \times C(p^{k_s})$ with $k_1 \geq k_2 \geq \dots \geq k_s \geq 1$ and $\sum_{i=1}^s k_i = k$. If

AMS Subject Classification (1980): Primary 1650, 1660; Secondary 0510.

a_x is the number of times p^x appears in the invariants of G/G' , then $\sum_{x \geq 1}^m xa_x = m$. Since G is a PN -group $|A_c(G)| = |\text{Hom}(G, Z)| = |\text{Hom}(G/G', Z)|$ [2]. So we have $|A_c(G)| = |\text{Hom}(G/G', Z)| = |\text{Hom}(\prod_{j=1}^t (C(p^{m_j}), \prod_{i=1}^s C(p^{k_i}))|$. Hence $|A_c(G)| = \prod_{j,i}^{t,s} |\text{Hom}(C(p^{m_j}), C(p^{k_i}))| \prod_{j,i}^{t,s} p^{\min(m_j, k_i)} = p^A$ for some A . Summing powers over $m_j = 1, 2, \dots, m_1$ for $k_i = k_s, \dots, k_1$ we have

$$\begin{aligned} A &= \left(\sum_{x \geq 1}^{k_s} xa_x + k_s \sum_{x > k_s}^{m_1} a_x \right) + \dots + \left(\sum_{x \geq 1}^{k_1} xa_x + k_1 \sum_{x > k_1}^{m_1} a_x \right) = \\ &= \sum_{i=2}^s \left(\sum_{x \geq 1}^{k_i} xa_x + k_i \sum_{x > k_i}^{m_1} a_x \right) + \sum_{i=1}^s \left(k_i \sum_{x > k_1}^{m_1} a_x \right) + \sum_{x \geq 1}^{k_1} xa_x = \\ &= \sum_{i=2}^s \theta_i + k \sum_{x > k_1}^{m_1} a_x + \sum_{x \geq 1}^{k_1} xa_x, \text{ where } \theta_i = \sum_{x \geq 1}^{k_i} xa_x + k_i \sum_{x > k_i}^{m_1} a_x. \end{aligned}$$

Since $k \geq m_1$ we have $k \sum_{x > k_1}^{m_1} a_x \geq \sum_{x > k_1}^{m_1} xa_x$ and so $A \geq k \sum_{i=2}^s \theta_i + \sum_{x > k_1}^{m_1} xa_x = \sum_{i=2}^s \theta_i + m$.

LEMMA 2. [4]. *Let G be a finite non-abelian p -group. Let $G = L_0 \supset L_1 \supset \dots \supset L_c = 1$, $1 = Z_0 \subset Z_1 \subset \dots \subset Z_c = G$ be the lower and the upper central series of G . If L_i/L_{i+1} is cyclic of order p^r for all $i = 1, 2, \dots, c-1$. then $L_i \cap Z_{c-i-1} = L_{i+1}$, $i = 1, 2, \dots, c-1$.*

LEMMA 3. [3]. *If G is a finite non-abelian group and Z_i/Z_{i-1} is cyclic of order p^r for all $i = 1, 2, \dots, c-1$, then $[G, Z_{i+1}] = Z_i$ for $i = 1, 2, \dots, c-1$.*

THEOREM 1. *Let G be a finite group of order p^n and class c . If L_i/L_{i+1} is cyclic of order p^r for all $i = 1, 2, \dots, c-1$ and $\exp(G/G') = p^r$, then $|G|$ divides, $|A(G)|$.*

Proof. Consider the following:

A: G is a PN -group. Since $L_i \subseteq Z_{c-i}$ and $L_i \not\subseteq Z_{c-i-1}$ we have $(Z_{c-i}/L_i) \supseteq (L_i Z_{c-i-1}/L_i) \simeq Z_{c-i-1}/L_i \cap Z_{c-i-1} = Z_{c-i-1}/L_{i+1}$ (by Lemma 2). Hence

$$|Z_{c-i}/Z_{c-i-1}| \geq |L_i/L_{i+1}| = p^r$$

for all $i = 1, 2, \dots, c-1$. But $|G/Z_{c-1}| = p^{2r}$ [4] and so $|G/Z_2| \geq p^{(c-1)r}$ which gives $|Z_2| \leq p^{n-(c-1)r}$. If $|Z| = p^k$ then $|I(G)| = |G/Z| = p^{n-k}$ and $|Z_2/Z| \leq p^{n-(c-1)r-k}$. Since $L_{c-1} \subseteq Z$ and $|L_{c-1}| = p^r$ we have $|Z| \geq p^r = \exp(G/G')$

and so by Lemma 1, $|A_c(G)| \geq |G/G'|$. But $|L_i/L_{i+1}| = p^r$ for $i = 1, 2, \dots, c-1$ which implies that $|L_{c-i}| = p^{ir}$ and so $|L_1| = p^{(c-1)r}$. Therefore we have $|G/G'| = |G/L_1| = p^{n-(c-1)r}$, and so $|A_c(G)| \geq p^{n-(c-1)r}$. Since $A_c(G)$ centralizes $I(G)$ in $A(G)$ we have $|I(G) \cap A_c(G)| = |Z(I(G))| = |Z(G/Z)| = |Z_2/Z| \leq p^{n-(c-1)r-k}$. Hence

$$\begin{aligned} |A(G)|_p &\geq |I(G)A_c(G)| = |I(G)| \cdot |A_c(G)|/|I(G) \cap A_c(G)| \geq \\ &\geq p^{n-k} p^{n-(c-1)r} / p^{n-(c-1)r-k} = p^n. \end{aligned}$$

B: $G = H \times K$, where H is abelian of order p^e and K is a PN -group. By [1], $|A(G)|_p \geq p^e |A(K)|_p$. Since $G = H \times K$, $|G'| = |K'|$, and by induction $|L_i(G)| = |L_i(K)|$ for all $i = 1, 2, \dots, c$. Hence $L_i(K)/L_{i+1}(K)$ is cyclic of order p^r for $i = 1, 2, \dots, c-1$. Moreover $G/G' = H \times K/K'$ and so $\exp(K/K') \leq p^r$. But $\exp(L_i(K)/L_{i+1}(K)) = p^r$ and so $\exp(K/K') = p^r$. Therefore by A, $|A(K)|_p \geq |K|$ and so $|A(G)|_p \geq p^e \cdot |K| = |G|$.

COROLLARY. *Let G be a PN -group. If $|L_i/L_{i+1}| = p$, $i = 1, 2, \dots, c-1$, and $\exp(G/G') \leq |Z|$, then $|G|$ divides $|A(G)|$.*

THEOREM 2. *Let G be a finite p -group of order p^n and class c . If Z_{i+1}/Z_i is cyclic of order p^r for all $i = 0, 1, \dots, c-2$, and $Z_{c-j} = L_j$ for some j , $1 \leq j \leq c-1$, then $|G|$ divides $|A(G)|$.*

Proof. By Lemma 3, $L_{j+1} = [L_j, G] = [Z_{c-j}, G] = Z_{c-j-1}$ and so $p^r = \exp(Z_{c-j}/Z_{c-j-1}) = \exp(L_j/L_{j+1}) \leq \exp(L_{j-1}/L_j) \leq |L_{j-1}/L_j| \leq |Z_{c-j-1}/Z_{c-j}| = p^r$. Hence $|L_{j-1}| = |Z_{c-j+1}|$ and since $L_{j-1} \subseteq Z_{c-j+1}$ we have $L_{j-1} = Z_{c-j+1}$. Therefore $L_j = Z_{c_j}$ for all $j = 1, 2, \dots, c$, and so L_j/L_{j+1} is cyclic of order p^r for all j . By [4], $G/Z_{c-1} = p^{2r}$ and so $|G/L_1| = |G/G'| = p^{2r}$. Let $p^{m_1} \geq p^{m_2} \geq \dots \geq p^{m_i}$ by the invariants of G/G' , if $m_2 < r$, then $\exp(L_1/L_2) \geq p^{m_2} < p^r$, which is a contradiction. Hence $m_2 \geq r$ and so $m_1 = m_2 = r$ and $\exp(G/G') = p^r$. The result follows from Theorem 1.

REFERENCES

- [1] A. D. Otto, *Central automorphisms of finite p -group*, Trans. Amer. Math. Soc. **125** (1966), 280–287.
- [2] J. E. Adney, J. Yen, *Automorphisms of a p -group*, Illinois J. Math. **9** (1965), 137–143.
- [3] G. Baumslag, N. Blackburn, *Groups with cyclic upper central factors*, Proc. London Math. Soc. **10** (1960), 531–544.
- [4] J. A. Gallian, *Finite p -groups with homocyclic central factors*, Canad. J. Math. **26** (1974), 636–643.
- [5] N. Blackburn, *On special class of p -groups*, Acta Math **100** (1958), 45–92.

Marasleios Academy of Athens
4. Marasli Street
Athens, GREECE

(Received 07 04 1980)