A NOTE ON EXTENSIONS OF BEAR AND P. P.-RINGS

N. J. Groenewald

Bear rings are rings in which the left (right) annihilator of each subset is generated by an idempotent [2]. Closely related to Bear rings are left P. P. -rings; these are the rings in which each principal left ideal is projective, or equivalently, ring in which the left annihilator of each element is generated by an idempotent. In [1] Armendariz showed that if R is a ring which has no nonzero nilpotent elements then R[X] is a Bear or P.P.-ring if and only if R is a Bear or P.P.-ring. In this note we generalize this result. A semigroup G is called au u.p. semigroup if, when A and B are nonempty finite subsets of G, then there always exists at least one $x \in G$ which has an unique representation in the form x = ab with $a \in A$ and $b \in B$. We prove that if R is a reduced ring and G a u. p. semigroup then the semigroup ring RG is a Bear or P.P.-ring if and only if R is a Bear or P.P.-ring.

We will assume throughout that rings have a unit. In a reduced ring left and right annihilators coicide for any subset U of R, hence we let $\operatorname{ann}_R(U) = l(U=r(U) = \{a \in R : aU = 0\}.$

The key lemma is the following characterization of zero divisors in RG when R is a reduced ring.

LEMMA 1. [3, COROLLARY 3.2] Let G be an u. p. semigroup and let R be a reduced ring. Let G be an u.p. semigroup and let $p, q \in RG$ such that pq = 0. Then for any $g, h \in G$ we have $p_q q_h = 0$.

COROLLARY 1. If R is a reduced ring and $f \in RG$, G an u.p. semigroup, such that $f^2 = f$ then $f \in R$.

Proof. Let $f = \sum_{i=1}^{n} a_i g_i$. It is easy to show that $g_i = e$ for at least one *i*.

Hence we may, without any loss in generality, put $f = a_1 e + a_2 g_2 + \ldots + a_n g_n$. Now f(f-1) = 0. From Lemma 1 we have $a_1(a_1-1) = 0$ and $a_i = 0$ for $i \ge 2$. Herce $f = a_1 = a_1^2 \in \mathbb{R}$.

If
$$f \in RG$$
 and $f = \sum_{i=1}^{n} a_i g_i$ let $S_f = \{a_1, a_2, \dots, a_n\}.$

N. J. Groenewald

COROLLARY 2. Let R be a reduced ring and $U \subseteq RG$. If $T = \bigcup_{f \in U} S_f$ then ann_{RG}U = ann_R(T)G.

Proof. This follows easily from Lemma 1.

THEOREM 1. Let R be a reduced ring and G an u.p. semigroup. Then RG is a P.P.-ring if and only if R is a P.P.-ring.

Proof. If RG is a P.P.-ring and $a \in R$ then $\operatorname{ann}_R(a) = R \cap \operatorname{ann}_{RG}(a) = R \cap (RG)e$ with $e^2 = e$. By Corollary 1, $e \in R$ and thus $R \cap RGe = Re$.

Now assume R is a P.P.-ring. Let $a, b \in R$ with $\operatorname{ann}_R(a) = Re_1$, $\operatorname{ann}_R(b) = Re_2$, where $e_1^2 = e_1$, $e_2^2 = e_2$. Put $e = e_1e_2$. Because the idempotents of R are central we have $e^2 = e$. We show that $\operatorname{ann}_R\{a,b\} = Re$. If xa = xb = 0 then $x = xe_1 = xe_2$ and $xe = xe_1e_2 = x$. Hence $\operatorname{ann}_R\{a,b\} \subseteq Re$. Further, let $t \in Re$, say $t = re_1e_2$. Now $ta = re_1e_2a = re_2e_1a = 0$ and $tb = re_1e_2b = 0$. Hence $Re \subseteq \operatorname{ann}_R\{a,b\}$. Therefore, $Re = \operatorname{ann}_R\{a,b\}$. Thus for any finite subset $U \subseteq R$, $\operatorname{ann}_R(U) = Re$ for some idempotent $e \in R$. If e RG then by Corollary 2, $\operatorname{ann}_{RG}(f) = \operatorname{ann}_R(S_f)G = (Re)G = (RG)e$ with $e^2 = e$, as Sf is finite. Thus RG is a P.P.-ring.

Similarly we can establish

THEOREM 2. Let R be a reduced ring and G an u. p. semigroup. Then RG is a Bear ring if and only if R is a Bear ring.

COROLLARY 3 [1, THEOREM A] Let R be a reduced ring. Then R[x] is a P.P.-ring if and only if R is a P.P.-ring.

Proof. It follows from the fact that the infinite cyclic semigroup $\langle X \rangle$ is an u.p. semigroup.

COROLLARY 4 [1, THEOREM B]. Let R be a reduced ring. Then R[x] is a Bear ring if and only if R is a Bear ring.

REFERENCES

- E. P. Armendariz, A note on extensions of Bear and P.P.-rings, J. Austral. Math. Soc. 18 (1974), 470-473.
- [2] Kaplansky, Rings of Operators, W. A. Benjamin, New York, 1968.
- [3] J. Krempa, D. Neiwieczerzal, Rings in which annihilators are ideals and their application to semigroup rings, Bull. Acad. Polon. Sci. Ser Sci., Math., Astronom. Phys. 25 (1977), 851-856.

University of Porth Elizabeth South Africa (Received 08 07 1982)

72