

ON THE AUTOMORPHISM GROUP OF AN INFINITE GRAPH

Aleksandar Torgašev

Abstract. In this paper, a specially defined automorphism group $\Gamma(G)$ of a connected countable simple infinite graph is considered. As the main result, we prove that $\Gamma(G)$ contains at most one non-trivial element. All infinite graphs with a non-trivial automorphism group are completely described.

Finally, for graphs with odd, or with a small even number (2 or 4) of non-zero eigenvalues, the corresponding automorphism groups are characterized.

1. Introduction. Throughout the paper, G is a connected infinite countable graph without loops or multiple edges, which we briefly call a graph. Its vertex set is $V(G) = N$, and its adjacency matrix $A = [a_{ij}]$ is an infinite $N \times N$ matrix, where

$$a_{ij} = \begin{cases} a^{i+j-2} & \text{if } i, j, \text{ are adjacent} \\ 0 & \text{otherwise} \end{cases}$$

(a is a fixed positive constant, $0 < a < 1$).

Hence, the whole graph G is labelled and the “weight” of vertex $v_i = i$ is a^{i-1} ($i \in N$).

For other definitions and results concerning spectra of infinite graphs, one can see [3, 4, 5].

2. Results. The automorphism group $\Gamma(G)$ of an infinite graph G defined here, depends on the matrix A , thus especially depends on the way of labelling of the vertex set $V(G)$.

Namely, we put $P \in \Gamma(G)$ if and only if

$$(1) \quad AP = PA,$$

where $P = [p_{ij}]$ is an infinite permutation matrix of the set $V(G)$,

$$p_{ij} = \begin{cases} 1, & j = \omega(i) \\ 0, & j \neq \omega(i) \end{cases}$$

and ω is the corresponding permutation of the set N .

In the sequel, we identify any automorphism $P \in \Gamma(G)$ with the corresponding permutation of the set N .

Obviously, each permutation $P \in \Gamma(G)$ is a unitary operator in the corresponding Hilbert space H , and

$$Pe_i = e_{\omega(i)} \quad (i \in N),$$

for a fixed orthonormal basis $\{e_i\}_1^\infty$ of H .

Relation (1) is equivalent to

$$(2) \quad a_{\omega(i)\omega(j)} = a_{ij} \quad (i, j \in N),$$

so that vertices i, j are adjacent if and only if $\omega(i), \omega(j)$ are adjacent. In this case (2) gives

$$(3) \quad \begin{aligned} a^{\omega(i)+\omega(j)-2} &= a^{i+j-2}, \text{ or} \\ \omega(i) - i &= -[\omega(j) - j] \quad (i, j, \text{-adjacent}). \end{aligned}$$

The last relation is very restrictive, and it is the main difference in comparison to the finite case.

LEMMA 1. (i) For every $\omega \in \Gamma(G)$ there is a unique integer $d = d(\omega)$ such that

$$(4) \quad |\omega(i) - i| = d \quad (i \in N).$$

(ii) If $\omega(i) = i$ for an $i \in N$, then $\omega = \text{id}$.

(iii) If G has at least one odd cycle, then $\Gamma(G)$ is trivial.


Proof. (i) For any two adjacent vertices $i, j \in V(G)$, relation (3) yields

$$|\omega(i) - i| = |\omega(j) - j|,$$

and the connectivity of G ends the proof.

The last two statements are then immediate by (i). \square


Examples. (1) The automorphism group of the one-way infinite path



is always trivial.

Indeed, for any $\omega \in \Gamma(G)$ we must have $\omega(i_1) = i_1$, thus $\omega = \text{id}$.

(2) The corresponding group of an infinite two-way infinite path



is either trivial or contains exactly one non-trivial element. If, for example, $i_p = 2p - 1$ and $j_p = 2p(p \in N)$, then it is non-trivial.

The following property is one of the most important properties of the groups considered.

THEOREM 1. *In each case $|\Gamma(G)| \leq 2$.*

Proof. Let $\omega \in \Gamma(G)$ and $d = \omega(1) - 1$. Then $|\omega(i) - i| = d$ ($i \in N$), and the only possibilities we have are

$$\begin{aligned} \omega(1) = d + 1, \quad 1 = \omega(d + 1), \dots, \quad \omega(d) = 2d, \quad d = \omega(2d), \quad \omega(2d + 1) = 3d + 1, \\ 2d + 1 = \omega(3d + 1), \dots \end{aligned}$$

Generally, we obtain

$$\omega(i) = i + (-1)^{\lfloor (i-1)/d \rfloor} d.$$

If now $\omega_1, \omega_2 \in \Gamma(G)$ are the automorphisms with the corresponding values $d_1 = d_2$, by the last relation we immediately find $\omega_1 = \omega_2$.

Next, let ω_1, ω_2 be the different automorphisms with $d_1 < d_2$. Then we get

$$\omega_2 \omega_1(1) = \omega_2(1 + d_1) = 1 + d_1 + d_2,$$

and also

$$\omega_2 \omega_1(d_1 + 1) = \omega_2(1) = 1 + d_2,$$

whence $d_1 = d_2 = d_2 - d_1$; thus $d_1 = 0$, $\omega_1 = \text{id}$, *q.e.d.*

Hence, $\Gamma(G)$ is always either trivial or contains at most one non-trivial element (which is then—involution). \square

So, the most important question concerning $\Gamma(G)$ is when it is non-trivial. In the next main theorem we completely describe all the infinite graphs which have a non-trivial automorphism group. It appears that the considered property depends only on the structure of the graph, and on the way of labelling of its vertex set.

First, let G be any bipartite graph. Its characteristic parts are denoted by N_1 and N_2 , assuming always that the minimal element is in N_1 . Note that N_1, N_2 are not the cardinals, but the corresponding sets of indices.

Next, we need the notion of *symmetric bipartite graphs* (briefly, SBGs). We call an infinite bipartite graph with the characteristic parts N_1, N_2 —*symmetric*, if there is a bijection $\pi : N_1 \rightarrow N_2$ such that two vertices $a \in N_1, \pi(b) \in N_2$ are adjacent if and only if the vertices $b \in N_1, \pi(a) \in N_2$ are.

If G is a SBG, then obviously N_1, N_2 are infinite.

If, additionally, we have that $\pi(a) - a = d = \pi(1) - 1$ for each $a \in N_1$, we say that N_1, N_2 are *good*. In this case, we obtain that

$$N_1 = \{(2s - 2)d + r \mid r \leq d, s \in N\}, \quad N_2 = \{(2s - 1)d + r \mid r \leq d, s \in N\}.$$

THEOREM 2. *Graph G has a non-trivial automorphism group if and only if it is a SBG with the good characteristic parts N_1 and N_2 .*

Proof. Let $\Gamma(G)$ be non-trivial, and let $\omega \in \Gamma(G)$ be the unique non-trivial automorphism (involution) of G . Then, by Lemma 1(iii), G can not have any odd cycle as an induced subgraph, thus it must be bipartite.

Let, next, the characteristic parts of G be N_1, N_2 with the minimal element in N_1 . Then, by the odd-path and the even-path characterizations of N_1, N_2 , we easily find that

$$\omega(a) = a + d(a \in N_1), \quad \omega(b) = b - d \quad (b \in N_2),$$

where $d = d(\omega) = \omega(1) - 1 > 0$.

Hence, N_1, N_2 are good, and ω is a needed bijection between N_1 and N_2 .

Since the converse statement is immediate, this completes the proof. \square

As examples, we consider the infinite graphs with a finite number $p(p \geq 2)$ of non-zero eigenvalues.

PROPOSITION 1. *Let G have an odd number of non-zero eigenvalues. Then its automorphism group is always trivial.*

Proof. As is known ([4]), each bipartite infinite graph, for every $a \in (0, 1)$, has the spectrum symmetric about the zero. Hence, if G has an odd number of non-zero eigenvalues, it cannot be bipartite, whence $\Gamma(G)$ is trivial. \square

Next, consider the infinite graphs with $p = 2$ or 4 non-zero eigenvalues.

We need the notion of characteristic subsets of G . The characteristic subsets N_1, N_2, \dots of an infinite graph are the equivalence classes related to the equivalence relation on the vertex set $N : x \sim y$ if and only if x, y are not adjacent and they have the same neighbors. Their number is finite or infinite and always greater than 1. If it is finite, G is said to be of finite type (type p , if this number is p) [5]. The corresponding quotient graph is denoted by g , and often called the canonical graph of G . If, for example, G is the complete m -partitive graph $K(N_1, \dots, N_m)(m \geq 2)$, then its characteristic subsets will be N_1, \dots, N_m , and its canonical graph is K_m .

LEMMA 2. (i) *If $\omega(x) \in N_i$ for an $x \in N_i$, then $\omega = \text{id}$.*

(ii) *If G is of finite type p and $\Gamma(G)$ is non-trivial, then p is even.*


Proof. (i) Assume, on the contrary, $\omega \neq \text{id}$, and denote by M_1, M_2 the characteristic parts of G . Since it follows easily that each N_i is contained either in M_1 or in M_2 , we get the statement.

(ii) Let ω be the non-trivial automorphism of $\Gamma(G)$. Since by (i), ω is an involution on the set $\{N_1, \dots, N_p\}$, without fixed elements, we have that p must be even. \square

PROPOSITION 2. *Let G have exactly two non-zero eigenvalues. Then $\Gamma(G)$ is non-trivial iff G is a complete bipartite graph with good characteristic parts.*

Proof. In [4], we proved that G has exactly two non-zero eigenvalues if and only if it is a complete bipartite graph. Hence, $\Gamma(G)$ is non-trivial iff N_1, N_2 are good (and consequently-infinite). \square

PROPOSITION 3. *The following graph*



where $N_4 = N_1 + d$, $N_3 = N_2 - d$ ($d \neq 0$) is the unique connected infinite graph with four non-zero eigenvalues and a non-trivial automorphism group.

Proof. In [5] we proved that G has exactly four non-zero eigenvalues if and only if its canonical graph is one of the eight particular graphs with 4, 5 or 6 vertices. Since six of them have a triangle as a subgraph, their automorphism groups must be trivial. Since next, the seventh of them is P_5 with 5 characteristic subsets, by Lemma 2 (ii), its automorphism group is trivial, too. Hence, only P_4 remains, and the remaining proof is easy. \square

The general problem for any even number p of non-zero eigenvalues ($p \geq 6$) is obviously equivalent to the determination of all finite connected canonical symmetric bipartite graphs with exactly p non-zero eigenvalues. The present author thinks it can be solved at least for $p = 6$, and may be for $p = 8$.

Acknowledgement. The author is very indebted to the anonymous referee for the final version of Theorem 1.

REFERENCES

- [1] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs—Theory and Application*, VEB Deutscher Verlag Wissen., Berlin, 1980; Academic Press, New York, 1980.
- [2] M. Petersdorf, H. Sachs, *Über spektrum, Automorphismengruppe und Teiler eines Graphen*, Wis. Z. THE Ilmenau **15** (1969), heft 4/5, 123–128.
- [3] M. Petrović, *The spectrum of an infinite labelled graph*, (in Serbian), Master's thesis, Fac. Sci., Beograd, 1981.
- [4] A. Torgašev, *On spectra of infinite graphs*, Publ. Inst. Math. (Beograd) **29** (**43**) (1981), 269–292.
- [5] A. Torgašev, *On infinite graphs with three and four non-zero eigenvalues*, Bull. Acad. Serbe Sci. et Arts LXXVI, Sci. Math. **11** (1981), 39–48.
- [6] A. Torgašev, *On infinite graphs with five non-zero eigenvalues*, Bull. Acad. Serbe Sci. et Arts LXXIX, Sci. Math. **12** (1982), 31–38.

Institut za matematiku
 Prirodno-matematički fakultet
 Beograd, Jugoslavija

(Received 21 04 1981)