

## ON THE $\mathcal{A}$ -COMPATIBILITY OF SUPPORTS OF DISTRIBUTIONS OF $\mathcal{K}'\{M_p\}$ -TYPE

Stevan Pilipović

**Abstract.** We determine relations between the notions of  $\mathcal{A}$ -compatibility and of  $M_p$ -convolution of distributions from  $\mathcal{K}'\{M_p\}$ .

**1. Introduction.** First we shall repeat two definitions. Let  $A$  and  $B$  be subsets of  $\mathbf{R}$ . If these sets satisfy the condition:

$$(1) \quad x_n \in A, \quad y_n \in B, \quad |x_n| + |y_n| \rightarrow \infty \Rightarrow |x_n + y_n| \rightarrow \infty, \quad (n \rightarrow \infty)$$

then they are called compatible.

The sets  $A$  and  $B$  are called polynomially compatible if there exists a polynomial  $P$  on  $\mathbf{R}$  such that

$$(2) \quad x \in A, \quad y \in B \Rightarrow |x| + |y| \leq P(|x + y|).$$

It is known that if  $f, g \in \mathcal{D}'$  ( $f, g \in \mathcal{S}'$ ) and the sets  $A = \text{supp } f$ ,  $B = \text{supp } g$  are compatible (polynomially compatible), then the convolution (tempered convolution) exists. The notion of compatibility of supports of distributions from  $\mathcal{D}'$  was investigated for example in [1] and the notion of tempered convolution and polynomial compatibility of supports of tempered distributions is introduced and investigated in [3], [4], [5].

In [5, Theorems 5.1 and 5.2] Kamiński proved that the notion of compatibility (polynomial compatibility) is essential for the convolution (tempered convolution) of distributions (tempered distributions).

KAMIŃSKI'S THEOREM [5]. *Let  $A$  and  $B$  be subsets of  $\mathbf{R}$  and let for every two non-negative measures (non-negative tempered measures)  $f$  and  $g$  with  $\text{supp } f \subset A$ ,  $\text{supp } g \subset B$ , the convolution (tempered convolution)  $f * g$  exist. Then the sets  $A$  and  $B$  are compatible (polynomially compatible) .*

The space  $\mathcal{S}'$  is an example of a space of  $\|\{M_p\}$ -type [2]. In [7] we generalize the notions of tempered convolution and of polynomial compatibility. We introduced the definition of  $M_p$ -convolution of elements from  $\|\{M_p\}$  and the definition of  $\mathcal{A}$ -compatibility.

In this paper we shall further investigate relations between the notions of  $\mathcal{A}$ -compatibility and of  $M_p$ -convolution. (We use the symbol  $\pm$  for this convolution). We shall give conditions on the sequence  $(M_p)$  such that the notion of  $\mathcal{A}$ -compatibility is essential for the  $M_p$ -convolution.

**2.  $\mathcal{A}\{M_p\}$ -compatible sets.** The space of  $\mathcal{K}\{M_p\}$ -type, where  $\{M_p(x)\}$  is a sequence of continuous functions on  $\mathbf{R}$  such that  $1 \leq M_1(x) \leq M_2(x) \leq \dots$ , was introduced in [2] as the dual space of the space  $\mathcal{K}\{M_p\}$ . The space  $\mathcal{K}\{M_p\}$  is a subspace of  $\mathbf{C}^\infty(\mathbf{R})$  defined in the following way:

$$\varphi \in \mathcal{K}\{M_p\} \text{ iff } \|\varphi\|_p := \sup\{M_p\}|\varphi^{(q)}(x)| : x \in \mathbf{R}, q \leq p\} < \infty \quad p = 1, 2, \dots$$

Topology in this space is defined by the sequence of norms  $(\|\cdot\|_p; \mathcal{K}\{M_p\})$  is an  $F$ -space, and if we suppose:

( $N$ ) for every  $p \in \mathbf{N}$  there is  $p' \in \mathbf{N}$  such that

$$M_p/M_{p'} \in L^1(\mathbf{R}) \text{ and } M_p(x)/M_{p'}(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty, \quad ([2])$$

then  $\mathcal{K}\{M_p\}$  is an  $FS$  space. ( $\mathbf{N}$  is the set of natural numbers.)

In this paper we shall suppose that  $M_p(x), p \in \mathbf{N}$ , are even functions which increase monotonically to infinity when  $x \rightarrow \infty$ . Also, we suppose that the sequence  $(M_p(x))$  satisfies:

( $N'$ ) ( $N$ ) holds and  $M_p(x)/M_{p'}(x) \rightarrow 0$  monotonically as  $x \rightarrow \infty$ .

Condition ( $N'$ ) implies that for every  $p' \in \mathbf{N}$  which correspond to some  $p \in \mathbf{N}$  in ( $N$ )

$$(3) \quad M_{p'}(x)/x \rightarrow \infty \text{ as } x \rightarrow \infty.$$

Namely, from the fact that  $M_p/M_{p'} \in L^1$  and  $M_p/M_{p'}(x) \rightarrow 0$  monotonically, it follows that for some  $\tilde{x}_{p,p'} > 0$

$$(3^*) \quad M_p(x)/M_{p'}(x) < 1/x \text{ if } x > \tilde{x}_{p,p'}.$$

If ( $3^*$ ) does not hold, then there exists a sequence  $(x_k)$  of positive numbers such that  $x_{k+1} > 1 + x_k^2$ ,  $k \in \mathbf{N}$ , and  $M_p(x_k)/M_{p'}(x_k) > x_k^{-1}$ . But then

$$\int_{x_1}^{\infty} (M_p(x)/M_{p'}(x)) dx \geq \sum_{k=1}^{\infty} x_{k+1}^{-1} (x_{k+1} - x_k) = \infty.$$

Condition ( $3^*$ ) implies that for a fixed  $m \geq 1$

$$M_p(mx) \leq M_{p'}(xm)/(xm) \text{ if } x > \tilde{x}_{p,p'}/m.$$

Since  $mM_p(x) \leq mxM_p(mx)$  if  $m \geq 1$  and  $x \geq 1$  we obtain that for every  $p \in \mathbf{N}$  there exists a  $p' \in \mathbf{N}$  and an  $x_{p,p'}$  such that

$$(C) \quad mM_p(x) \leq M_{p'}(mx) \quad \text{if } x > \tilde{x}_{p,p'}.$$

Without loss of generality we can suppose that (3) holds for every  $p \in \mathbf{N}$ .

For our investigations of  $M_p$ -convolutions the following condition on the sequence  $(M_p)$  is also needed:

$$(4) \quad \text{For every } p \in \mathbf{N} \text{ there is a } p' \in \mathbf{N} \text{ and a } C_{p,p'} > 0 \text{ such that}$$

$$M_p^2(x) \leq C_{p,p'} M_{p'}(x) \text{ for } x > C_{p,p'}.$$

Now we shall give a definition of the set  $\mathcal{A}$  that is somewhat different from the definition of this set in [7].

We denote by  $\mathcal{A}$  a set of non-negative functions defined on  $\mathbf{R}^+$ , bounded on bounded domains, directed according to the ordinary relation  $\leq$  (i.e. for every  $f$  and  $g$  from  $\mathcal{A}$  there is an  $h \in \mathcal{A}$  such  $\max\{f(x), g(x)\} \leq h(x), x \in \mathbf{R}$ ) such that:

- (A1) If a non-negative function  $\varphi$  defined on  $\mathbf{R}^+$  satisfies the inequality  $\varphi(x) \leq \psi(x), x \in \mathbf{R}^+$  for some  $\psi \in \mathcal{A}$ , then  $\varphi \in \mathcal{A}$ ;
- (A2) There are  $\varphi \in \mathcal{A}$  and  $x_0 \geq 0$  such that  $\varphi(x) \geq x$  for  $x \geq x_0$ ;
- (A3) For every  $\varphi \in \mathcal{A}$ ,  $m \in \mathbf{N}$  and  $n \in \mathbf{N}_0$  there is a  $\psi \in \mathcal{A}$  such that  $m\varphi(x+n) \leq \psi(x), x \in \mathbf{R}^+$ . ( $\mathbf{N}_0 = \mathbf{N} \cup \{0\}$ .)

Let us suppose that for a given sequence  $(M_p)$  and set  $\mathcal{A}$  the following condition holds:

$$(S) \quad \text{For every } p \in \mathbf{N} \text{ and } \varphi \in \mathcal{A} \text{ there is a } p' \in \mathbf{N} \text{ and an } x_{p,p'} > 0 \text{ such that}$$

$$M_p(\varphi(x)) \leq M_{p'}(x) \quad \text{if } x > x_{p,p'}.$$

In this case we shall denote the set  $\mathcal{A}$  by  $\mathcal{A}(M_p)$ .

Condition (S) implies some properties of the sequence  $(M_p(x))$ . For example:

$$(5) \quad \text{For every } p \in \mathbf{N} \text{ there are } p' \in \mathbf{N} \text{ and } x_{p,p'} > 0 \text{ such that}$$

$$M_p(px) \leq M_{p'}(x) \text{ if } x > x_{p,p'}.$$

Let us prove this. From (A2) and (A3) it follows that there exists a  $\varphi \in \mathcal{A}$  such that  $px \leq \varphi(x), x \in \mathbf{R}^+$ . Therefore (S) implies (5) because  $M_p(x)$  is monotonous for  $x \geq 0$ .

Sequences  $(M_p(x))$  which define spaces of exponential distributions quoted in [7, part 5], satisfy all the conditions above.

Let  $(M_p)$  be a sequence of even, monotonically increasing functions (when  $x \rightarrow \infty$ ) for which (N'), (4) and (5) hold. Then, we denote by  $\mathcal{B}(M_p)$  the set of all sets  $\mathcal{A}(M_p)$ .

PROPOSITION 1.  $\mathcal{B}(M_p) \neq \emptyset$ .

*Proof.* We denote by  $\mathcal{A}$  the set of all non-negative functions which are smaller or equal to some non-negative (on  $\mathbf{R}^+$ ) polynomial of order 1. It is easy to check that  $\mathcal{A}_0 \in \mathcal{B}(M_p)$ .

We denote by  $\mathcal{A}_{\max}(M_p)$  the set defined by  $\mathcal{A}_{\max}(M_p) = \bigcup_{A \in \mathcal{B}} \mathcal{A}(M_p)$ . It is easy to prove that  $\mathcal{A}_{\max}(M_p) \in \mathcal{B}(M_p)$ .

Let  $f, g \in \mathcal{K}'M_p$  and  $A = \text{supp } f$ ,  $B = \text{supp } g$ . As in [7] we say that  $A$  and  $B$  are compatible if there exists a  $\varphi \in \mathcal{A}(M_p)$  such that

$$x \in A, y \in B \Rightarrow |x| = |y| \leq \varphi(|x + y|).$$

We give now Theorem 9 from [7] in the following version:

THEOREM 2. *If  $A$  and  $B$  are  $\mathcal{A}_{\max}(M_p)$ -compatible, then the convolution  $f \sharp g$  exists. ( $\text{supp } f \in A, \text{supp } g \in B$ .)*

Consider now the precise characterization of the sets  $\mathcal{A}(M_p)$  for a given sequence  $(M_p(x))$  (which satisfies all the conditions mentioned).

THEOREM 3. *Let  $(M_p(x))$  satisfy the following condition:*

(B) *For every  $p \in \mathbf{N}$ ,  $r \in \mathbf{N}$  and  $\varepsilon > 0$  there exists  $p' \in \mathbf{N}$  and an  $x_{p,r,p',\varepsilon} > 0$*

$$\text{such that } M_p^{-1}(M_r(x)) \leq \varepsilon M_{p'}^{-1}(M_{p'}(x)) \text{ if } x > x_{p,r,p',\varepsilon}.$$

*Then  $\mathcal{A}_{\max}(M_p)$  is the set of all non-negative functions which are smaller or equal to some linear combinations of functions of the form  $x \rightarrow M_p^{-1}(M_q(x))$ ,  $(p, q) \in \mathbf{N}^2$ ,  $x > 0$ , and a constant function.*

*Proof.* We put  $\varphi_{p,q}(x) = M_p^{-1}(M_q(x))$ ,  $x \in \mathbf{R}^+$ ,  $(p, q) \in \mathbf{N}^2$  and denote by  $A$  the set of all non-negative functions which are smaller or equal to some linear combinations of functions of the form  $x \rightarrow \varphi_{p,q}(x)$ ,  $(p, q) \in \mathbf{R}^2$ ,  $x > 0$  and a constant function.

We have  $0 \leq \varphi_{p,p}(x)$ , for every  $p \in \mathbf{N}$  and

$$\max\{\varphi_{p_1,q_1}(x), \varphi_{p_2,q_2}(x)\} \leq \varphi_{p_0,q_0}(x), x \in \mathbf{R}^+,$$

where  $p_0 = \min\{p_1, p_2\}$ ,  $q_0 = \max\{q_1, q_2\}$ . From (5) and (B) it follows that for every  $p, q, m \in \mathbf{N}$  and  $n \in \mathbf{N}_0$  there exists a  $q' \in \mathbf{N}$  and  $\tilde{x}$  such that

$$m\varphi_{p,q}(x+n) \leq \varphi_{p,q'}(x) \text{ if } x > \tilde{x}.$$

Namely, for sufficiently large  $x > 0$  have

$$mM_p^{-1}(M_q(x+n)) \leq mM_p^{-1}(M_q(2x)) \leq mM_p^{-1}(M_{q_1}(x)) \leq M_p^{-1}(M_q(x)).$$

If for some non-negative function  $\varphi$  on  $\mathbf{R}^+$  we have the estimate  $\varphi(x) \leq M_p^{-1}(M_q(x))$  if  $x > x_{pq} > 0$  for some  $(p, q) \in \mathbf{N}^2$ , then  $\varphi(x) \in A$  because for suitable  $C > 0$

$$\varphi(x) \leq M_p^{-1}(M_q(x)) + C, \quad x \in \mathbf{R}^+.$$

Now it is clear that  $A \in \mathcal{B}(M_p)$ .

Condition (S) implies that if  $\varphi \in \mathcal{A}_{\max}(M_p)$ , then  $\varphi(x) \leq M_p^{-1}(M'_p(x))$  for  $x > x_{p,p'} > 0$ ; that is  $\varphi \in A$ . Since  $A \in \mathcal{B}(M_p)$ , the assertion is proved.

Let us remark that the sequences  $(M_p)$  given in [7, part 5] satisfy condition (B) and that the corresponding sets  $\mathcal{A}$  can be redefined to be  $\mathcal{A}_{\max}(M_p)$ .

**3. Conditions for the  $\mathcal{A}(M_p)$ -compatibility.** Let  $(M_p(x))$  be a sequence which satisfies all the conditions from Section 2 and let  $\mathcal{A}(M_p)$  be an element from  $\mathcal{B}(M_p)$  (we suppose that condition (S) is satisfied).

In Theorem 5, which will be stated later, the following condition concerning  $(M_p)$  and  $\mathcal{A}(M_p)$  will be used:

(B1) For every  $p \in \mathbf{N}$  and every  $q \in \mathbf{N}$  there exists  $\varphi \in \mathcal{A}(M_p)$   
and an  $x_\varphi > 0$  such that  $M_p(\varphi(x)) \geq M_q(x)$  if  $x > x_\varphi$ .

PROPOSITION 4. (i) If  $\mathcal{A}(M) = \mathcal{A}_{\max}(M_p)$ , then (B) implies (B1).

(ii) Conditions (S) and (B1), concerning the given set  $\mathcal{A} = \mathcal{A}(M_p)$ , imply that (B) holds and that  $\mathcal{A}(M_p) = \mathcal{A}_{\max}(M_p)$ .

*Proof.* (i) This follows easily from Theorem 3.

(ii) It follows from (B1) that for every  $p$  and every  $r \in \mathbf{N}$  there exists a  $\varphi \in \mathcal{A}(M_p)$  such that  $M_p^{-1}(M_r(x)) \leq \varphi(x)$  for sufficiently large  $x$ . From (A3) and (S) it follows that:

$$\begin{aligned} m\varphi(x) &\leq \psi(x), & \text{for some } \psi \in \mathcal{A}(M_p). \\ \psi(x) &\leq M_{p'}^{-1}(M_{p'}(x)) & \text{for some } p' \text{ and sufficiently large } x. \end{aligned}$$

Thus we obtain that for arbitrary  $m \in \mathbf{N}$ ,  $p \in \mathbf{N}$ ,  $r \in \mathbf{N}$ , there exists a  $p' \in \mathbf{N}$  and an  $\tilde{x} > 0$  such that

$$mM_p^{-1}(M_r(x)) \leq M_{p'}^{-1}(M_{p'}(x)) \text{ if } x > \tilde{x}.$$

that is, condition (B) holds. Since  $\mathcal{A}(M_p)$  contains functions of the form  $M_p^{-1} \cdot (M_q(x))$ ,  $(p, q) \in \mathbf{N}^2$ , and their non-negative linear combinations, it follows that  $\mathcal{A}(M_p) = \mathcal{A}_{\max}(M_p)$ .

**THEOREM 5.** *We suppose that  $(M_p(x))$  is a sequence of functions which satisfies all the conditions from Section 2 and that (S) and (B1) hold for a given set  $\mathcal{A}$ . Then the following assertion holds:*

(\*) *If  $A$  and  $B$  are subsets of  $\mathbf{R}$  such that for every two non-negative measures  $f$  and  $g$  from  $\mathcal{K}'\{M_p\}$  with supports in  $A$  and  $B$  respectively, the convolution  $f * g$  exists, then  $A$  and  $B$  are  $\mathcal{A}_{\max}(M_p)$ -compatible.*

*Proof.* We shall use the idea of the proof of Theorem 5.2. from [5]. Since tempered non-negative measures are non-negative measures from  $\mathcal{K}'\{M_p\}$ , we have that the sets  $A$  and  $B$  are compatible.

Let us suppose that  $A$  and  $B$  are not  $\mathcal{A}_{\max}\{M_p\}$ -compatible.

Let  $p \in \mathbf{N}$  be fixed. There are points  $x_i \in A$ ,  $y_i \in B$  such that

$$(6) \quad |x_i| + |y_i| \geq 2^i(M_p^{-1}(M_i(|x_i| + |y_i|)) + 1), \quad i \in \mathbf{N}.$$

This holds because functions of the form  $2^i(M_p^{-1}(M_i(x) + 1))$ ,  $x \in \mathbf{R}^+$ ,  $i \in \mathbf{N}$ , belong to  $\mathcal{A}_{\max}(M_p)$ . Condition (6) implies that  $|x_i| + |y_i| \rightarrow \infty$ , and therefore,  $|z_i| = |x_i + y_i| \rightarrow \infty$  as  $i \rightarrow \infty$ .

There are three possibilities:

- (i)  $|x_i| \rightarrow \infty$  and  $|y_i| \rightarrow \infty$ ;
- (ii)  $|x_i| \rightarrow \infty$  and  $|y_i| \not\rightarrow \infty$ ;
- (iii)  $|x_i| \not\rightarrow \infty$  and  $|y_i| \rightarrow \infty$ .

First we consider case (i). It is not a restriction if we suppose that  $|x_{i+1}| > |x_i|$ ,  $|y_{i+1}| > |y_i| + 1$  and  $|z_{i+1}| > |z_i| + 1$ ,  $i \in \mathbf{N}$ .

We put

$$f(t) = \sum_{i=1}^{\infty} M_{p'}(|x_i|)\delta(t - x_i); \quad g(t) = \sum_{i=1}^{\infty} M_{p'}(|y_i|)\delta(t - y_i),$$

where we shall choose  $p' \in \mathbf{N}$  later.

From (5) we obtain that for a given  $p \in \mathbf{N}$  there exists a  $p'$  and  $x_{pp'}$  such that

$$(7) \quad M_p(x) \leq M_{p'}(x/2) \leq M_{p'}(x - t)M_{p'}(t) \quad \text{if } x > x_{pp'} \text{ and } t \in \mathbf{R}.$$

Now we choose  $p'$  as an element from  $\mathbf{N}$  which corresponds to  $p$  ( $p$  was fixed earlier) in (7). From (7) we have

$$\begin{aligned} M_{p'}(|x_i|)M_{p'}(|y_i|) &= M_{p'}(|x_i| + |y_i| - |y_i|)M_{p'}(|y_i|) \\ &\geq M_{p'}(|x_i| + |y_i|/2) \geq M_p(|x_i| + |y_i|) \end{aligned}$$

if  $|x_i| > x_{pp'}$  (This is true for all  $i$  with  $i \geq i_0$  for some  $i_0$ .)

Since  $f$  and  $g$  belong to  $\mathcal{K}\{M_p\}$  and  $\text{supp } f \subset A$ ,  $\text{supp } g \subset B$ , then the convolutions  $f \sharp g$  and  $f * g$  exist and

$$(f \sharp g)(t) = (f * g)(t) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} M_{p'}(|x_i|)M_{p'}(|y_j|)\delta(t - x_i - y_j).$$

Using (7) and (6) we have

$$\begin{aligned} (f \sharp g)(t) &\geq \sum_{i=i_0} M_{p'}(|x_i|)M_{p'}(|y_i|)\delta(t - z_i) \geq \sum_{i=i_0} M_p(|x_i| + |y_i|)\delta(t - z_i) \geq \\ &\geq M_p(M_p^{-1}(M_i(|x_i| + |y_i|)))\delta(t - z_i) = \sum_{i=i_0} M_i(|z_i|)\delta(t - z_i). \end{aligned}$$

The last series is a distribution which does not belong to  $\mathcal{K}'\{M_p\}$ . Thus  $(f \sharp g)(t)$  is not in  $\mathcal{K}'\{M\}$  and so this is a contradiction.

Now we consider case (ii). It is not a restriction if we suppose that  $|x_{i+1}| > |x_i| + 1$ ,  $y_i \rightarrow y$  and  $|z_{i+1}| > |z_i| + 1$ ,  $i \in \mathbf{N}$ . From (C) and (5) it follows that there are sequences  $(p_i)$  and  $(L_i)$  such that  $2^i M_i(x) \leq M_{p_i}(x)$  if  $x > L_i$ .

We choose the sequence  $(x_i)$  such that  $M_{p_i}(|x_i + y_i|) \geq 2^i M_i(|x_i + y_i|)$ ,  $i \in \mathbf{N}$  (i.e.  $|x_i + y_i| > L_i$ ) and

$$(8) \quad |x_i| + |y_i| \geq M_p^{-1}(M_{p_i}(|x_i + y_i|)).$$

The existence of the sequence  $(x_i)$  for which (8) holds follows from the fact that the functions  $M_p^{-1}(M_{p_i}(x))$  are from  $\mathcal{A}_{\max}(M_p)$ .

Let  $f(t) = \sum_{i=1}^{\infty} M_{p'}(|x_i|)\delta(t - x_i)$  and  $g(t) = \sum_{i=1}^{\infty} 2^{-i} M_{p'}(|y_i|)\delta(t - y_i)$ . Clearly,  $f$  and  $g$  are from  $\mathcal{K}'\{M_p\}$ . Since  $f \sharp g$  and  $f * g$  exist, and

$$\begin{aligned} f \sharp g = f * g &= \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} M_{p'}(|x_i|) 2^{-j} M_{p'}(|y_j|) \delta(t - x_i - y_j) \geq \\ &\geq \sum_{i=i_0}^{\infty} 2^{-i} M_p(|x_i| + |y_i|) \delta(t - z_i) \geq \sum_{i=i_0}^{\infty} 2^{-i} M_p(M_p^{-1}(M_{p_i}(|x_i + y_i|))) \cdot \\ &\quad \cdot \delta(t - z_i) \geq \sum_{i=i_0}^{\infty} M_i(|z_i|) \delta(t - z_i), \end{aligned}$$

we obtain a contradiction as the last series is not an element from  $\mathcal{K}'\{M\}$ .

Case (iii) is symmetrical to case (ii), and the proof is complete.

From the preceding Theorem and Proposition 4 we directly obtain:

**THEOREM 6.** *If in Theorem 5 instead of (B1) we suppose that (B) holds, and in addition, if we suppose that  $\mathcal{A}(M_p) = \mathcal{A}_{\max}(M_p)$  (all the other conditions are the same as in Theorem 5), then the assertion (\*) holds.*

#### REFERENCES

- [1] P. Antosik, J. Mikusiński, R. Sikorski, *Theory of Distributions Sequential Approach*, PWN-Polish Sci. Publ., Warszawa. 1979.
- [2] I.M. Gel'fand, G.E. Shilov, *Generalized Functions, Vol. 2*, Academic Press, New York, 1968.
- [3] A. Kamiński, *On convolutions, products and Fourier transformations*, Bull. Acad. Polon. Sci. Mat. Astronom. Phys. **25**(1977), 369–374.
- [4] A. Kamiński, *Convolution, product and Fourier transformations of distributions*, Studia Math. **74**(1982), 369, 83–96.

- [5] A. Kamiński, *On the Rényi theory of conditional probability*, (to appear in *Studia Math.*).
- [6] S. Pilipović, A. Takači, *The space  $\mathcal{H}'\{M_p\}$  and convolutions*, Proc. Int. Conf., "on Generalized Functions and its Applications in Mathematical Physics" Moskow 1980., Math. Inst. M.V. Steklov, (1981) 415–426.
- [7] S. Pilipović, *On the convolution in the space of  $\mathcal{K}'\{M_p\}$ -type*, Math. Nachr. **120**(1985), 03–112.

Institut za Matematiku  
21000 Novi Sad  
Dr ilije Đuričića 4

(Received 13 08 1984)  
(Revised 30 04 1985)