

ALL REPRODUCTIVE SOLUTIONS OF FINITE EQUATIONS

Slaviša B. Prešić

Summary. An equation, the solution set of which is a subset of a given finite set, is called a *finite equation*. Applying some kind of algebraic structure we effectively determine all reproductive solutions of such equations (*Theorem 1* and *Theorem 2*).

1. Let E be a given non-empty set and $f: E \rightarrow E$ a given function. An x -equation

$$(1) \quad f(x) = x$$

is called *reproductive* [4] if the function f satisfies the identity

$$(2) \quad f(f(x)) = f(x)$$

All solutions of a reproductive equation can be found in a trivial way. Namely, the formula

$$(3) \quad x = f(p) \quad (p \text{ is an arbitrary element of } E)$$

determines all solution of (1) provided this equation is a reproductive one. The formula is an example of the so-called general reproductive solutions formula ([2], [3], [7]).

Next, any x -equation

$$(4) \quad eq(x) \quad (x \text{ is an unknown element of } E; eq \text{ is a given unary relation of } E)$$

which has at least one solution is equivalent to a reproductive equation [4].

Accordingly, to solve a given x -equation (4) it suffices to find any reproductive equation equivalent to (1). In this paper we are concerned with finding all such reproductive equations in case of a finite equation.

2. Let $B = \{\sigma_0, \sigma_1, \dots, \sigma_n\}$ be a given set of $n + 1$ elements and $S = \{0, 1\}$. Define the operation x^y by

$$(5) \quad x^y = \begin{cases} 1, & \text{if } x = y \\ 0, & \text{otherwise} \end{cases} \quad (x, y \in B \cup S)$$

The standard Boolean operations $+$ and \cdot ("or" and "and") are described by the following tables

$$\begin{array}{c|cc} + & 0 & 1 \\ \hline 0 & 0 & 1 \\ 1 & 1 & 1 \end{array} \quad \begin{array}{c|cc} \cdot & 0 & 1 \\ \hline 0 & 0 & 0 \\ 1 & 0 & 1 \end{array}$$

Extend these operations to partial operations of the set $B \cup S$ in the following way

$$(6) \quad x + 0 = x, \quad 0 + x = x, \quad x \cdot 0 = 0, \quad 0 \cdot x = 0, \quad x \cdot 1 = x, \quad 1 \cdot x = x \quad (x \in B \cup S)$$

We consider the following x -equation

$$(7) \quad s_0 \cdot x^{b_0} + s_1 \cdot x^{b_1} + \dots + s_n \cdot x^{b_n} = 0$$

where $s_i \in \{0, 1\}$ are given elements and $x \in B$ is unknown. Obviously the equation (7) is possible iff the condition

$$(8) \quad s_0 \cdot s_1 \cdots s_n = 0$$

holds.

In the sequel, assuming the condition (8), we are going to determine all reproductive solutions of the equation (7).

First we introduce the following definition.

Let $(\sigma_0, \sigma_1, \dots, \sigma_n) \in S^{n+1}$ be any element. Then the set $Z(\sigma_0, \dots, \sigma_n)$ ("the zero-set of $(\sigma_0, \dots, \sigma_n)$ ") is defined as follows

$$(9) \quad b_i \in Z(\sigma_0, \dots, \sigma_n) \Leftrightarrow \sigma_i = 0 \quad (i = 0, 1, \dots, n)$$

For instance, if $n = 3$ we have

$$Z(1, 0, 1, 0) = \{b_1 b_3\}, \quad Z(1, 1, 1, 1) = \emptyset, \quad Z(0, 0, 0, 0) = \{b_0, b_1, b_2, b_3\}$$

Let now s_0, \dots, s_n be any elements of S satisfying the condition (8). With respect to the sequence s_0, \dots, s_n we define the so-called *repro-function*¹ $A : B \rightarrow B$. This is any function defined by a certain formula of the form

$$(10) \quad A(x) = A_0(s_0, \dots, s_n)x^{b_0} \cdots + A_n(s_0, \dots, s_n)x^{b_n}$$

where each coefficient $A_k(s_0, \dots, s_n)$ is determined by some equality of the form

$$(11) \quad A_k(s_0, \dots, s_n) = b_k s_k^0 + \sum_{\sigma_k \neq 0, \sigma_0 \cdots \sigma_n = 0} F_k(\sigma_0, \dots, \sigma_n) s_0^{\sigma_0} \cdots s_n^{\sigma_n}$$

¹As a matter of fact, A is a function of the type $A : S^{n+1} \times B \rightarrow B$.

assuming that coefficients $F_k(\sigma_0, \dots, \sigma_n) \in B$ satisfy the condition

$$(12) \quad F_k(\sigma_0, \dots, \sigma_n) \in Z(\sigma_0, \dots, \sigma_n)$$

For instance, if $n = 2, k = 1$ the equality (11) reads

$$A_1(s_0, s_1, s_2) = b_1 s_1^0 + F_1(0, 1, 0) s_0^0 s_1^1 s_2^0 + F_1(0, 1, 1) s_0^0 s_1^1 s_2^1 + F_1(1, 1, 0) s_0^1 s_1^1 s_2^0,$$

where the coefficients $F_1(0, 1, 0), F_1(0, 1, 1), F_1(1, 1, 0)$ can be any elements of B satisfying the conditions (of type (12))

$$F_1(0, 1, 0) \in \{b_0, b_2\}, \quad F_1(0, 1, 1) = b_0, \quad F_1(1, 1, 0) = b_2$$

Note that generally, according to the condition $s_0 \cdots s_n = 0$, there exists at least one repro-function with respect to the sequence s_0, \dots, s_n .

THEOREM 1. *Let $s_0 \cdots s_n = 0$, then the equation of the form*

$$x = A(x)$$

is a reproductive equation and equivalent to the equation (7) if and only if A is a repro-function.

Proof. Denote the equation (7) by $g(x) = 0$. Firstly, we prove the following fact

(p) Let $A(x)$ be determined by a certain equality of type (10) assuming only that $A_0(s_0, \dots, s_n), \dots, A_n(s_0, \dots, s_n) \in B$. Then the implication $g(x) = 0 \Rightarrow x = A(x)$ holds if and only if for each coefficient $A_k(s_0, \dots, s_n)$ an equality of the form (11) is satisfied, where $F_k(\sigma_0, \dots, \sigma_n)$ can be any elements² of B .

The proof immediately follows from the following equivalences

$$\begin{aligned} (p) &\Leftrightarrow (\forall x \in B)[s_0 x^{b_0} + \dots + s_n x^{b_n} = 0 \Rightarrow x = A_0(s_0, \dots, s_n) x^{b_0} + \dots + \\ &\quad + A_n(s_0, \dots, s_n) x^{b_n}] \\ &\Leftrightarrow (\forall k \in \{0, \dots, n\})(s_k = 0 \Rightarrow A_k(s_0, \dots, s_n) = b_k) \\ &\Leftrightarrow A_k(s_0, \dots, s_n) \text{ is determined by means of a certain equality (11), where} \\ &\quad F_k(\sigma_0, \dots, \sigma_n) \text{ may be any elements of } B. \end{aligned}$$

Next we introduce the condition

$$(q) \quad (\forall x \in B) s_0 (A(x))^{b_0} + \dots + s_n (A(x))^{b_n} = 0.$$

Obviously the sentence " $x = A(x)$ is a reproductive equation, equivalent to the equation $f(x) = 0$ " is logically equivalent to the conjunction $(p) \wedge (q)$. Accordingly, the remaining part of the proof reads:

$x = A(x)$ is a reproductive equation, equivalent to the equation $f(x) = 0$

²Thus condition (8) is not assumed.

- $\Leftrightarrow (p) \wedge (q)$
 $\Leftrightarrow (\forall x \in B) s_0(A(x))^{b_0} + \dots + s_n(A(x))^{b_n} = 0$, and $A(x)$ is determined by means of some equalities of the form (10), (11), where $F_k(\delta_0, \dots, \delta_n)$ are certain elements of B
 $\Leftrightarrow (\forall x \in B) s_0(A_0(s_0, \dots, s_n)x^{b_0} + \dots + A_n(s_0, \dots, s_n)x^{b_n})^{b_0} + \dots + s_n(A_0(s_0, \dots, s_n)x^{b_0} + \dots + A_n(s_0, \dots, s_n)x^{b_n})^{b_n} = 0$
 and $A_k(s_0, \dots, s_n)$, with $k \in \{0, \dots, n\}$, are determined by some equalities of the form (11).
 $\Leftrightarrow (\forall i, k \in \{0, \dots, n\}) s_i A_k^{b_i}(s_0, \dots, s_n) = 0$ and $A_k(s_0, \dots, s_n)$, with $k \in \{0, \dots, n\}$, are determined by some equalities of the form (11). This part of the proof is based on the following general facts: If a_0, \dots, a_n, b are any elements of B then:
 $1^\circ (a_0x^{b_0} + \dots + a_nx^{b_n})^b = a_0^b x^{b_0} + \dots + a_n^b x^{b_n}$ (for all $x \in B$)
 $2^\circ (\forall x \in B) a_0x^{b_0} + \dots + a_nx^{b_n} = 0 \Leftrightarrow (\forall i \in \{0, \dots, n\}) a_i = 0$
 $\Leftrightarrow (\forall i, k \in \{0, \dots, n\}) s_i \cdot \left(\sum_{\sigma_k \neq 0, \sigma_0 \dots \sigma_n = 0} F_k^{b_i}(\sigma_0, \dots, \sigma_n) s_0^{\sigma_0} \dots s_n^{\sigma_n} \right) = 0$, where $F_k(s_0, \dots, s_n)$ are certain elements of B . We have used the equality of the form (11) and the identity $s_i \sigma_k^{b_i} s_k^0 = 0$
 $\Leftrightarrow (\forall i, k \in \{0, \dots, n\}) \left(\sum_{(\sigma_0, \dots, \sigma_n) \in S^{n+1}} \sigma_i s_0^{\sigma_0} \dots s_n^{\sigma_n} \cdot \sum_{\sigma_k \neq 0, \sigma_0 \dots \sigma_n = 0} F_k^{b_i}(\sigma_0, \dots, \sigma_n) s_0^{\sigma_0} \dots s_n^{\sigma_n} = 0 \right)$
 (For the identity $s_i = \sum_{(\sigma_0, \dots, \sigma_n) \in S^{n+1}} \sigma_i s_0^{\sigma_0} \dots s_n^{\sigma_n}$ holds).
 $\Leftrightarrow (\forall i, k \in \{0, \dots, n\}) \sum_{\sigma_k \neq 0, \sigma_0 \dots \sigma_n = 0} \sigma_i F_k^{b_i}(\sigma_0, \dots, \sigma_n) s_0^{\sigma_0} \dots s_n^{\sigma_n} = 0$
 $\Leftrightarrow (\forall i, k \in \{0, \dots, n\}) (\forall \sigma_0, \dots, \sigma_n \in S) (P \Rightarrow \sigma_i F_k^{b_i}(\sigma_0, \dots, \sigma_n) = 0)$
 where the condition $\sigma_k \neq 0, \sigma_0 \dots \sigma_n = 0$ is denoted by P .
 From $F_k \in B$ it follows that $(\forall k)(\exists j) F_k = b_j$. Hence we conclude the equality $F_k = b_{\varphi(k)}$ where $\varphi: \{0, \dots, n\} \rightarrow \{0, \dots, n\}$ is a certain function³.
 $\Leftrightarrow (\forall \sigma_0, \dots, \sigma_n \in S) (\forall i, k \in \{0, \dots, n\}) (P \Rightarrow \sigma_i b_{\varphi(k)}^{b_i} = 0)$
 $\Leftrightarrow (\forall \sigma_0, \dots, \sigma_n \in S) (\forall k \in \{0, \dots, n\}) (P \Rightarrow \sigma_{\varphi(k)} = 0)$
 For $x \neq y \Rightarrow x^y = 0$.
 $\Leftrightarrow (\forall \sigma_0, \dots, \sigma_n \in S) (\forall k \in \{0, \dots, n\}) (P \Rightarrow b_{\varphi(k)} \in Z(\sigma_0, \dots, \sigma_n))$
 Using definition (9).
 $\Leftrightarrow (\forall \sigma_0, \dots, \sigma_n \in S) (\forall k \in \{0, \dots, n\}) (P \Rightarrow F_k \in Z(\sigma_0, \dots, \sigma_n))$
 $\Leftrightarrow A$ is a repro-function

³ φ also depends on $\sigma_0, \dots, \sigma_n$.

The proof is complete.

From Theorem 1 immediately follows the following result on the reproductive solutions.

THEOREM 2. *If (7) is a possible equation then a formula*

$$x = A(p) \quad (p \text{ is any element of } B)$$

represents a general reproductive solution of the equation (7) if and only if the function A is a repro-function.

Example 1. Let $B = \{0, 1, 2\}$. Consider the x -equation

$$s_0 x^0 + s_1 x^1 + s_2 x^2 = 0 \quad (s_i \text{ are given and } x \text{ is unknown})$$

This equation is possible if and only if $s_0 s_1 s_2 = 0$. Any general reproductive solution has the following form

$$(14) \quad x = A_0 p^0 + A_1 p^1 + A_2 p^2$$

where A_i are defined by

$$A_0 = (1 \text{ or } 2)s_0^1 s_1^0 s_2^0 + 1s_0^1 s_1^0 s_2^1 + 2s_0^1 s_1^1 s_2^0, \quad A_1 = 1s_1^0 + (0 \text{ or } 2)s_0^0 s_1^1 s_2^0 + 2s_0^1 s_1^1 s_2^0$$

$$A_2 = 2s_2^0 + (0 \text{ or } 1)s_0^0 s_1^0 s_2^1 + 1s_0^1 s_1^0 s_2^1$$

In these equalities a symbol of the form $(p \text{ or } q)$ denotes an element which may be p or q . Consequently there are exactly 8 formulas of the form (14).

3. Now we state an application of Theorems 1 and 2. Let n be a given natural number and $B = S^n$. Then according to the definition (5) we have the following identity

$$(x_1, \dots, x_n)^{(i_1, \dots, i_n)} = x_1^{i_1} \dots x_n^{i_n}$$

In connection with it the equation of the type (7) may be written in the following form

$$(15) \quad \sum a_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} = 0$$

where $a_{i_1 \dots i_n} \in S$ are given, and $x_j \in S$ are unknown elements. In other words, (14) is the standard Boolean equation in x_1, \dots, x_n . Theorems 1 and 2 can be directly applied to the equation (15); consequently one effectively finds all general reproductive solutions of it.

REFERENCES

- [1] M. Božić, *A note on reproductive solutions*, Publ. Inst. Math. (Beograd) **19 (33)** (1975), 33-35.
- [2] L. Löwenheim, *Über das Auflösungsproblem im logischen Klassen-kalkul*. Sitzungsber, Berl. Math. Gesellschaft **7** (1980), 89-94.
- [3] L. Löwenheim, *Über die Auflösung von Gleichungen im logischen Gebietenkalkul*. Math. Ann. **68** (1910), 169-207.
- [4] S. B. Prešić, *Une classe d'équations matricielles et l'équation fonctionnelle $f^2 = f$* , Publ. Inst. Math. (Beograd) **8 (22)** (1968), 143-148.
- [5] S. B. Prešić, *Une méthode de resolution des équations dont toutes les solutions appartiennent à un ensemble fini*, C.R. Acad. Sci. Paris Ser. A **272** (1971), 654-657.
- [6] S. B. Prešić, *Ein Satz über reproductive Lösungen*, Publ. Inst. Math. Belgrade **14 (28)** (1973), 133-136.
- [7] S. Rudeanu, *Boolean Functions and Equations*, North-Holland, Amsterdam & Elsevier, New York, 1974.
- [8] S. Rudeanu, *On reproductive solutions of arbitrary equations*, Publ. Inst. Math. (Beograd) **24 (38)** (1978), 143-147.
- [9] D. Banković, *On general and reproductive solutions of arbitrary equations*, Publ. Inst. Math. (Beograd) **26 (40)**, (1979), 31-33.

Matematički fakultet
Studentski trg 16,
11000 Beograd
Jugoslavija

(Received 12 10 1987)