

INDEPENDENT VERTEX SETS IN SOME COMPOUND GRAPHS

Ivan Gutman

Abstract. Let G be an n -vertex graph and $R_1, R_2, \dots, R_n$ distinct rooted graphs. The compound graph $G[R_1, R_2, \dots, R_n]$ is obtained by identifying the root of R_i with the i -th vertex of G , $i = 1, 2, \dots, n$. We determine the number of independent vertex sets and the independence polynomial of $G[R_1, R_2, \dots, R_n]$. Several special cases of these results are pointed out.

1. Introduction

Consider finite graphs without loops. If G is such a graph, then $\mathbf{V}(G)$ will denote its vertex set. Any subset of $\mathbf{V}(G)$, such that no two elements of it are mutually adjacent, is called an independent vertex set of the graph G . Let $\text{Ind}(G)$ be the set of all independent vertex sets of G .

The number $\sigma(G)$ of independent vertex sets of the graph G , i.e. the cardinality of $\text{Ind}(G)$, has been examined in a number of recent papers [1–11]. In particular, Prodinger and Tichy [7, 11] called the quantity $\sigma(G)$ “the Fibonacci number of the graph G ”. The motivation for this was the fact that if P_n is the path-graph with n vertices, then $\sigma(P_n)$ is equal to the $(n+1)$ -th Fibonacci number.

In the present paper we determine the number of independent vertex sets of the compound graph $G[R_1, R_2, \dots, R_n]$ constructed in the following manner.

Let G be a graph with the vertex set $\mathbf{V}(G) = \{v_1, v_2, \dots, v_n\}$. Let further $R_1, R_2, \dots, R_n$ be distinct rooted graphs; the root of R_i is denoted by r_i , $i = 1, 2, \dots, n$. Then $G[R_1, R_2, \dots, R_n]$ is the graph obtained by identifying the vertex v_i of G with the root r_i of R_i , simultaneously for $i = 1, 2, \dots, n$ (see Fig. 1).

Denote by R_i° the graph obtained by deleting from R_i the root-vertex r_i and the edges incident to it. Denote by $R_i^\bullet$ the graph obtained by deleting from R_i the root-vertex r_i , the vertices adjacent to r_i and all the incident edges. Then the main result of our work can be formulated as follows.




Fig. 1

THEOREM 1. Let I be an independent vertex set of the graph G . Define

$$\sigma_i(I) = \begin{cases} \sigma(R_i^\circ) & \text{if } v_i \notin I, \\ \sigma(R_i^\bullet) & \text{if } v_i \in I. \end{cases}$$

Then

$$\sigma(G[R_1, R_2, \dots, R_n]) = \sum_{I \in \text{Ind}(G)} \prod_{i=1}^n \sigma_i(I).$$

Instead of Theorem 1 we prove a somewhat stronger result, namely Theorem 2. In order to do this we need some preparations.

2. The independence polynomial

Denote by $n(G, k)$ the number of distinct k -element independent vertex sets of the graph G . Then the polynomial

$$\omega(G) = \omega(G, x) = \sum_{k \geq 0} n(G, k) x^k \quad (1)$$

is called the independence polynomial of the graph G [3, 5, 6]. Evidently, $\omega(G, 1) = \sigma(G)$.

The basic properties of the independence polynomial have been determined by Gutman and Harary [5] and recently by Hoede and Li [6].

Two of these properties will be needed in the subsequent considerations:

- (a) If v is a vertex of the graph G and N_v is the set containing v and its first neighbors, then

$$\omega(G) = \omega(G - v) + x \omega(G - N_v). \quad (2)$$

- (b) If $G_1 \cup G_2$ is the graph composed of components G_1 and G_2 , then

$$\omega(G_1 \cup G_2) = \omega(G_1) \omega(G_2). \quad (3)$$

THEOREM 2. *Let I be an independent vertex set of the graph G . Define*

$$\omega_i(I) = \begin{cases} \omega(R_i^\circ) & \text{if } v_i \notin I, \\ \omega(R_i^\bullet) & \text{if } v_i \in I. \end{cases} \quad (4)$$

Then

$$\omega(G[R_1, R_2, \dots, R_n], x) = \sum_{I \in \text{Ind}(G)} x^{|I|} \prod_{i=1}^n \omega_i(I) \quad (5)$$

where $|I|$ stands for the number of elements of I .

Evidently, Theorem 1 is a special case of Theorem 2, obtained by setting $x = 1$ in formula (5).

3. Proof of Theorem 2

We demonstrate the validity of Theorem 2 by induction on the number n of vertices of the graph G . If $n = 1$, then $\mathbf{V}(G) = \{v_1\}$ and therefore $G[R_1, R_2, \dots, R_n]$ coincides with R_1 . On the other hand, for $n = 1$ the set $\text{Ind}(G)$ consists of only two elements, namely $I_1 = \emptyset$ and $I_2 = \{v_1\}$. Bearing in mind (4) we have $\omega_1(I_1) = \omega(R_1^\circ)$ and $\omega_1(I_2) = \omega(R_1^\bullet)$. Consequently, the right-hand side of (5) is equal to $\omega(R_1^\circ) + x\omega(R_1^\bullet)$. Because of (2) this latter expression is equal to $\omega(G)$.

Thus the statement of Theorem 2 is true for $n = 1$. In a similar manner one can check that Theorem 2 is satisfied for $n = 2$ and $n = 3$.

Assume now that Theorem 2 holds for all graphs G with less than n vertices. In order to accomplish the inductive proof we have to show that this assumption implies the validity of Theorem 2 for the graphs G having n vertices.

Suppose that $n \geq 3$ and apply formula (2) to the vertex v_n of the graph G . Without loss of generality we may label the vertices of G so that v_n is adjacent to $v_{n-1}, \dots, v_{n-d}$. Then by using (3),

$$\begin{aligned} \omega(G[R_1, R_2, \dots, R_n]) &= \omega(R_n^\circ) \omega((G - v_n)[R_1, R_2, \dots, R_n]) \\ &\quad + \omega(R_n^\bullet) \omega(R_{n-1}^\circ) \cdot \dots \cdot \omega(R_{n-d}^\circ) \omega((G - N_{v_n})[R_1, R_2, \dots, R_n]). \end{aligned} \quad (6)$$

The subgraphs $G - v_n$ and $G - N_{v_n}$ have $n - 1$ and $n - 1 - d$ vertices, respectively. Therefore according to the induction hypothesis:

$$\omega((G - v_n)[R_1, R_2, \dots, R_n]) = \sum_{I \in \text{Ind}(G - v_n)} x^{|I|} \prod_{i=1}^{n-1} \omega_i(I) \quad (7)$$

$$\omega((G - N_{v_n})[R_1, R_2, \dots, R_n]) = \sum_{I \in \text{Ind}(G - N_{v_n})} x^{|I|} \prod_{i=1}^{n-d-1} \omega_i(I). \quad (8)$$

The set $\text{Ind}(G)$ can be partitioned into two disjoint subsets $\text{Ind}^\circ(G)$ and $\text{Ind}^\bullet(G)$, such that $\text{Ind}^\circ(G)$ is the set of independent vertex sets of G which do not contain

the vertex v_n whereas $\text{Ind}^\bullet(G)$ is the set of those independent vertex sets of G which do contain v_n . It is easy to see that

$$\text{Ind}^\circ(G) = \text{Ind}(G - v_n) \quad (9)$$

$$\text{Ind}^\bullet(G) = \{I \cup \{v_n\} \mid I \in \text{Ind}(G - N_{v_n})\}. \quad (10)$$

Bearing in mind (7)–(10), the relation (6) is transformed into

$$\begin{aligned} \omega(G[R_1, R_2, \dots, R_n]) &= \omega(R_n^\circ) \sum_{I \in \text{Ind}^\circ(G)} x^{|I|} \prod_{i=1}^{n-1} \omega_i(I) \\ &\quad + x \omega(R_n^\bullet) \omega(R_{n-1}^\circ) \cdots \omega(R_{n-d}^\circ) \sum_{I \in \text{Ind}^\bullet(G)} x^{|I|-1} \prod_{i=1}^{n-d-1} \omega_i(I). \end{aligned} \quad (11)$$

For all $I \in \text{Ind}^\circ(G)$, $v_n \notin I$ and therefore $\omega_n(I) = \omega(R_n^\circ)$. For similar reasons, the relations $\omega_n(I) = \omega(R_n^\bullet)$ and $\omega_j(I) = \omega(R_j^\circ)$, $j = 1, \dots, d$, are satisfied for all $I \in \text{Ind}^\bullet(G)$. Consequently, equation (11) becomes

$$\omega(G[R_1, R_2, \dots, R_n]) = \sum_{I \in \text{Ind}^\circ(G)} x^{|I|} \prod_{i=1}^n \omega_i(I) + \sum_{I \in \text{Ind}^\bullet(G)} x^{|I|} \prod_{i=1}^n \omega_i(I)$$

and formula (5) follows from the fact that $\text{Ind}^\circ(G) \cup \text{Ind}^\bullet(G) = \text{Ind}(G)$. This completes the proof of Theorem 2.

4. Special cases

4.1. All R_i are isomorphic. The graph $G[R_1, R_2, \dots, R_n]$ in which all R_i , $i = 1, 2, \dots, n$ are isomorphic to the rooted graph R is denoted by $G[R]$. For the compound graphs $G[R]$ formula (5) is much simplified by the fact that the product $\prod_{i=1}^n \omega_i(I)$ depends only on the cardinality k of the independent vertex set I and is equal to $\omega(R - r)^{n-k} \omega(R - N_r)^k$ where r stands for the root of R . Since the number of k -element independent vertex sets of the graph G is equal to $n(G, k)$ we further have

$$\omega(G[R]) = \sum_{k \geq 0} x^k n(G, k) \omega(R - r)^{n-k} \omega(R - N_r)^k. \quad (12)$$

This, bearing in mind the definition of $\omega(G)$, immediately leads to Corollary 2.1.

COROLLARY 2.1. $\omega(G[R], x) = \omega(R - r, x)^n \omega(G, \omega(R - N_r) / \omega(R - r))$.

COROLLARY 2.2. *If k^* is the maximum cardinality of an independent vertex set of the graph G , then the polynomial $\omega(R - r)^{n-k^*}$ divides the polynomial $\omega(G[R])$.*

4.2. The corona. The corona $G \circ Q$ of the graphs G and Q is obtained from G and n copies of Q , so that each vertex of G is joined to all vertices of a copy of Q . Whence, $G \circ Q$ is a special case of $G[R]$ when the root r of R is adjacent to all other vertices of R . In this notation, $Q = R - r$.

COROLLARY 2.3. $\omega(G \circ Q, x) = \omega(Q, x)^n \omega(G, 1/\omega(Q))$.

COROLLARY 2.4. *If k^* is the maximum cardinality of an independent vertex set of the graph G , then the polynomial $\omega(Q)^{n-k^*}$ divides the polynomial $\omega(G \circ Q)$.*

4.3. Some more special cases. If G is the complete graph K_n then $\text{Ind}(G)$ consists of $n + 1$ elements: the empty set and n one-element sets, each containing one vertex of G . Formula (5) gives then

$$\omega(K_n[R_1, R_2, \dots, R_n]) = x^0 \prod_{i=1}^n \omega_i(\emptyset) + \sum_{j=1}^n \prod_{i=1}^n \omega_i(\{v_j\}).$$

Bearing in mind (4) we arrive at

$$\text{COROLLARY 2.5. } \omega(K_n[R_1, R_2, \dots, R_n]) = \left[1 + x \sum_{j=1}^n \frac{\omega(R_j^\bullet)}{\omega(R_j^\circ)} \right] \prod_{i=1}^n \omega(R_i^\circ).$$

It is easy to deduce combinatorial formulas for the $n(G, k)$ -numbers of the path P_n and the circuit C_n [2, 5]. Then equations (12) and (2) lead to

COROLLARY 2.6.

$$\begin{aligned} \omega(P_n(R)) &= \sum_{k \geq 0} \binom{n-1-k}{k} \omega(R-r)^{n-k} [\omega(R) - \omega(R-r)]^k, \\ \omega(C_n(R)) &= \sum_{k \geq 0} \frac{n}{n-k} \binom{n-k}{k} \omega(R-r)^{n-k} [\omega(R) - \omega(R-r)]^k. \end{aligned}$$

REFERENCES

- [1] K. Engel, *On the Fibonacci number of an $M \times N$ lattice*, Fibonacci Quart. **28** (1990), 72–78.
- [2] I. Gutman, *On independent vertices and edges of a graph*, in: R. Bodendiek, R. Henn (Eds.), *Topics in Combinatorics and Graph Theory*, Physica-Verlag, heidelberg 1990, 291–296.
- [3] I. Gutman, *An identity for the independence polynomials of trees*, Publ. Inst. Math. (Beograd) **50** (1991), 19–23.
- [4] I. Gutman, *Numbers of independent vertex and edge sets of a graph: some analogies*, Graph Theory Notes New York **22** (1992), 18–22.
- [5] I. Gutman, F. Harary, *Generalization of the matching polynomial*, Utilitas Math. **24** (1983), 97–106.
- [6] C. Hoede, X. L. Li, *Clique polynomials and independent set polynomials of graphs*, Memor. Fac. Appl. Math. Univ. Twente **982** (1991), 1–14.
- [7] P. Kirschenhofer, H. Prodinger, R. F. Tichy, *Fibonacci numbers of graphs. II*, Fibonacci Quart. **21** (1983), 219–229.
- [8] V. Linek, *Bipartite graphs can have any number of independent sets*, Discrete Math. **76** (1989), 131–136.
- [9] R. E. Merrifield, H. E. Simmons, *Enumeration of structure-sensitive graphical subsets: theory*, Proc. Nat. Acad. Sci. USA **78** (1981), 692–695.
- [10] R. E. Merrifield, H. E. Simmons, *Enumeration of structure-sensitive graphical subsets: calculations*, Proc. Nat. Acad. Sci. USA **78** (1981), 1329–1332.
- [11] H. Prodinger, R. F. Tichy, *Fibonacci numbers of graphs*, Fibonacci Quart. **20** (1982), 16–21.

Prirodno-matematički fakultet
34000 Kragujevac, p.p.60
Jugoslavija

(Received 20 11 1991)