

LOGICS WITH TWO TYPES OF INTEGRAL OPERATORS

Radosav S. Dorđević

Abstract. We prove completeness theorems for absolutely continuous and singular biprobability models of a logic with integrals. Also in both cases, we prove the finite compactness theorem for a set of sentences of the form $\tau \in [r, s]$.

We assume throughout the paper that \mathcal{A} is a countable admissible set with $\omega \in \mathcal{A}$. In [2], Keisler introduced a logic $L_{\mathcal{A}f}$ which has an integral operator which builds terms with bound variables. In our case two types of integral operators $\int_1 \dots dx$ and $\int_2 \dots dx$ are allowed.

A biprobability model for $L_{\mathcal{A}f_1f_2}$ logic is a model $\mathfrak{A} = \langle A, R_i, c_j, \mu_1, \mu_2 \rangle_{i \in I, j \in J}$, where $\langle A, R_i, c_j \rangle$ is a first-order model without operations and μ_1, μ_2 are probability measures on A . We shall see a difference in semantics for $L_{\mathcal{A}f_1f_2}^a$ and $L_{\mathcal{A}f_1f_2}^s$ by means of the following definition.

Definition 1. (a) An absolutely continuous biprobability model for $L_{\mathcal{A}f_1f_2}^a$ is a biprobability model \mathfrak{A} such that μ_1 is absolutely continuous with respect to μ_2 , i. e. $\mu_1 \ll \mu_2$.

(b) A singular biprobability model for $L_{\mathcal{A}f_1f_2}^s$ is a biprobability model \mathfrak{A} such that μ_1 is singular with respect to μ_2 , i. e. $\mu_1 \perp \mu_2$. \square

In both cases, quantifiers are interpreted by

$$\left(\int_k \tau(x, \vec{a}) dx \right)^{\mathfrak{A}} = \int \tau(b, \vec{a})^{\mathfrak{A}} d\mu_k(b) \quad \text{for } k = 1, 2,$$

where $\tau(x, \vec{y})$ is a term and $\vec{a} \in A^n$.

Diagonal products $\mu_k^{(n)}$, which are the corresponding restrictions of completions of μ_k^n 's ($k = 1, 2$) to σ -algebras generated by the measurable rectangles and the diagonal sets $\{ \vec{x} \in A^n : x_i = x_j \}$, can be replaced by sequences of probability measures on A^n 's which satisfy the Fubini theorem. That generalization of a probability structure is relevant for us.

Mathematics Subject Classification (1991): Primary 03C70

This research was supported by Government of Serbia grant number 0401A, through Matematički institut.

Definition 2. A graded biprobability model for $L_{\mathcal{A} f_1 f_2}$ is a model $\mathfrak{A} = \langle A, R_i, c_j, \mu_n^1, \mu_n^2 \rangle_{i \in I, j \in J, n \geq 1}$ such that:

- (1) Each μ_n^k is a countably additive probability measure on A^n .
- (2) Each n -ary relation R_i is μ_n^k -measurable and the identity relation is μ_2^k -measurable.
- (3) $\mu_n^k \times \mu_m^k \subseteq \mu_{n+m}^k$.
- (4) Each μ_n^k is preserved under permutation of $\{1, 2, \dots, n\}$.
- (5) $\langle \mu_n^k : n \in \mathbb{N} \rangle$ has the Fubini property: If B is μ_{m+n}^k -measurable, then
 - (a) For each $\vec{x} \in A^m$, the section $B_{\vec{x}} = \{ \vec{y} : B(\vec{x}, \vec{y}) \}$ is μ_n^k -measurable.
 - (b) The function $f(\vec{x}) = \mu_n^k(B_{\vec{x}})$ is μ_m^k -measurable.
 - (c) $\int f(\vec{x}) d\mu_m^k = \mu_{m+n}^k(B)$. \square

Definition 3. (a) A graded biprobability model for $L_{\mathcal{A} f_1 f_2}^a$ is a graded biprobability model \mathfrak{A} such that $\mu_n^1 \ll \mu_n^2$ for each $n \in \mathbb{N}$.

(b) A graded biprobability model for $L_{\mathcal{A} f_1 f_2}^s$ is a graded biprobability model \mathfrak{A} such that $\mu_n^1 \perp \mu_n^2$ for each $n \in \mathbb{N}$. \square

1. The logic $L_{\mathcal{A} f_1 f_2}^a$. Axioms and rules of inference for $L_{\mathcal{A} f_1 f_2}^a$ are those for $L_{\mathcal{A} f}$, as listed in [3] with both \int_1 and \int_2 playing the role of \int , together with the following axioms:

(A₁) *Axioms of continuity of integral operators:* ($i, j = 1, 2$)

$$(a) \bigwedge_n \bigvee_m \bigvee_k \int_i \mathbb{F}_k \left(\int_j \tau(\vec{x}, \vec{y}) d\vec{x} \right) d\vec{y} < \frac{1}{n},$$

$$\text{where } F_k(s) = \begin{cases} 1, & \text{if } r - 1/m + 1/k \leq s \leq r - 2/k \\ 0, & \text{if } s \leq r - 1/m \text{ or } s \geq r - 1/k \\ \text{linear,} & \text{for other cases} \end{cases}$$

is a continuous real function such that $F_k \upharpoonright \mathbb{Q} \in \mathcal{A}$.

$$(b) \bigwedge_n \bigvee_m \bigvee_k \int_i \mathbb{G}_k \left(\int_j \tau(\vec{x}, \vec{y}) d\vec{x} \right) d\vec{y} < \frac{1}{n},$$

$$\text{where } G_k(s) = \begin{cases} 1, & \text{if } r + 2/k \leq s \leq r + 1/m - 1/k \\ 0, & \text{if } s \leq r + 1/k \text{ or } s \geq r + 1/m \\ \text{linear,} & \text{for other cases.} \end{cases}$$

(A₂) *Axiom of absolute continuity:*

$$\bigwedge_{\varepsilon \in \mathbb{Q}_+} \bigvee_{\delta \in \mathbb{Q}_+} \bigwedge_n \bigwedge_{\tau \in T_n} (|\int_2 \tau(\vec{x}) d\vec{x}| < \delta \implies |\int_1 \tau(\vec{x}) d\vec{x}| < \varepsilon),$$

where $T = \bigcup_n T_n$, T_n is a set of terms with n free variables and $T, T_n \in \mathcal{A}$.

(A₃) $\int_1 (\int_2 \tau dy) dx = \int_2 (\int_1 \tau dx) dy$.

Now we introduce two sorts of auxiliary models.

Definition 4. (a) A weak model for $L_{f_1 f_2}^a$ is a model $\langle \mathfrak{A}, I_1, I_2 \rangle$ where \mathfrak{A} is a first-order model and I_k is what may be called an \mathcal{A} -Daniell integral on A , that is, I_k is a positive linear real function on the set of terms with at most one free

variable x and parameters from A , i. e.

$$\begin{aligned} I_k(r) &= r, & k &= 1, 2 \\ I_k(r \cdot \sigma + s \cdot \tau) &= r \cdot I_k(\sigma) + s \cdot I_k(\tau), \\ \text{if } \tau(b, \vec{a})^{\mathfrak{A}} &\geq 0 \text{ for all } b \in A, \text{ then } I_k(\tau(x, \vec{a})) &\geq 0. \end{aligned}$$

(b) A middle model for $L_{\mathcal{A} f_1 f_2}^{\alpha}$ is a weak model \mathfrak{A} such that for each $\varepsilon > 0$ there is $\delta > 0$ such that for each term $\tau(x, \vec{y})$ and $\vec{a} \in A^n$, if $|I_2(\tau(x, \vec{a}))| < \delta$ then $|I_1(\tau(x, \vec{a}))| < \varepsilon$. \square

In both cases, for τ a term, define $\tau^{\mathfrak{A}}$ inductively as for biprobability models, except that at the integral step, we define

$$(\int_k \tau(x, \vec{a}) dx)^{\mathfrak{A}} = I_k(\tau(x, \vec{a})) .$$

LEMMA 1. (Middle Completeness Theorem for $L_{\mathcal{A} f_1 f_2}^{\alpha}$) *Let T be a set of sentences of $L_{\mathcal{A} f_1 f_2}^{\alpha}$ such that T is Σ_1 -definable over \mathcal{A} . Then T is consistent with the axioms of this logic iff it has a middle model in which each theorem of $L_{\mathcal{A} f_1 f_2}^{\alpha}$ is true.*

Proof. The soundness is easy to prove because all the axioms represent known properties of integrals (the Generalized Radon-Nikodym Theorem and the Fubini Theorem prove that each function $\tau(x, y)^{\mathfrak{A}}: A \times A \rightarrow \mathbb{R}$ is compatible with absolutely continuous measures μ_1 and μ_2 , i. e.

$$\iint \tau(x, y)^{\mathfrak{A}} d\mu_1(x) d\mu_2(y) = \iint \tau(x, y)^{\mathfrak{A}} d\mu_2(y) d\mu_1(x) .$$

A Henkin argument is used to construct a weak model $\langle \mathfrak{A}, I_1, I_2 \rangle$ of T in which each theorem of $L_{\mathcal{A} f_1 f_2}^{\alpha}$ is true. Let $K = L \cup C$ be the language introduced in this construction, where C is a set of new constant symbols and $C \in \mathcal{A}$. We wish the axiom A_2 to hold for all the terms and that is done by the following construction (see [9]).

Let K' be a language with four kinds of variables: X, Y, Z, \dots are variables for sets, x, y, z, \dots are variables for urelements, r, s, t, \dots are variables for reals from $[0, 1] \cap \mathcal{A}$, and U, V, W, \dots are variables for functions $A^n \mapsto \mathbb{R}$, $n \geq 0$. Predicates are: $E_n^s(\vec{x}, X)$ for sets, $n \geq 1$; $E_{n+1}^t(\vec{x}, r, U)$ for terms, $n \geq 0$; $I_k(U, r)$ for $U: A^0 \rightarrow \mathbb{R}$ or $U: A^1 \rightarrow \mathbb{R}$, $k = 1, 2$; and \leq for reals. Function symbols are f, g, h, \dots for each continuous real function $F: \mathbb{R}^n \rightarrow \mathbb{R}$ such that $F \upharpoonright \mathbb{Q}^n \in \mathcal{A}$. Constant symbols are: X_{φ} for each formula φ ; U_{τ} for each term τ ; and \bar{r} for each real number $r \in [0, 1] \cap \mathcal{A}$.

Let S be the following theory of $K'_{\mathcal{A}}$:

1. *Axioms of validity:*

- 1.1 $(\forall X) \bigwedge_{n < m} \neg(\exists \vec{x}, \vec{y})(E_m^s(\vec{x}, \vec{y}, X) \wedge E_n^s(\vec{x}, X)),$ where $\{\vec{x}\} \cap \{\vec{y}\} = \emptyset$;
- 1.2 $(\forall U) \bigwedge_{n < m} \neg(\exists \vec{x}, \vec{y}, r, s)(E_{m+1}^t(\vec{x}, \vec{y}, r, U) \wedge E_{n+1}^t(\vec{x}, s, U));$
- 1.3 $(\forall U)(\forall \vec{x}, r, s)((E_{n+1}^t(\vec{x}, r, U) \wedge E_{n+1}^t(\vec{x}, s, U)) \implies r = s);$

2. Axioms of extensionality:

- 2.1 $(\forall \vec{x})(E_n^s(\vec{x}, X) \iff E_n^s(\vec{x}, Y)) \iff X = Y;$
 2.2 $(\forall \vec{x}, r)(E_{n+1}^t(\vec{x}, r, U) \iff E_{n+1}^t(\vec{x}, r, V)) \iff U = V;$

3. Axioms of terms:

- 3.1 $(\forall \vec{x})(E_{n+1}^t(\vec{x}, 0, U_\tau) \vee E_{n+1}^t(\vec{x}, 1, U_\tau))$ if τ is $\mathbf{1}(R(\vec{x}))$;
 3.2 $(\forall x, y)(E_{2+1}^t(x, y, 0, U_\tau) \vee E_{2+1}^t(x, y, 1, U_\tau))$ if τ is $\mathbf{1}(x = y)$;
 3.3 $E_{0+1}^t(\vec{r}, U_\tau)$ if τ is r ;
 3.4 $(\forall \vec{x}, r)(E_{n+1}^t(\vec{x}, r, U_\tau) \iff (\exists \vec{s})(\bigwedge_{i=1}^k E_{n+1}^t(\vec{x}, s_i, U_{\tau_i}) \wedge$
 $\wedge f(s_1, \dots, s_k) = r))$ if τ is $\mathbb{F}(\tau_1, \dots, \tau_k)$;
 3.5 $(\forall \vec{x}, r)(E_{n+1}^t(\vec{x}, r, U_\tau) \iff (\exists V)((\forall y, s)(E_{1+1}^t(y, s, V) \iff$
 $\iff E_{n+1+1}^t(\vec{x}, y, s, U_\sigma)) \wedge I_k(V, r)))$ if τ is $\int_k \sigma(\vec{v}, v_0) dv_0$, $k = 1, 2$;

4. Axioms of satisfaction:

- 4.1 $(\forall \vec{x})(E_n^s(\vec{x}, X_\varphi) \iff (\exists r \geq 0)E_{n+1}^t(\vec{x}, r, U_\tau))$ if φ is $\tau \geq 0$;
 4.2 $(\forall \vec{x})(E_n^s(\vec{x}, X_{\neg\varphi}) \iff \neg E_n^s(\vec{x}, X_\varphi))$;
 4.3 $(\forall \vec{x})(E_n^s(\vec{x}, X_{\wedge\Phi}) \iff \bigwedge_{\varphi \in \Phi} E_n^s(\vec{x}, X_\varphi))$;

5. Axioms of integral operators:

- 5.1 $(\forall U)((\bigwedge_{n \geq 2} \neg(\exists \vec{x}, r)E_{n+1}^t(\vec{x}, r, U)) \iff (\exists_1 s)I_k(U, s))$, $k = 1, 2$;
 5.2 $(\forall r)I_k(U_r, r)$;
 5.3 $(\forall U, V, r, s)I_k(r \cdot U + s \cdot V) = r \cdot I_k(U) + s \cdot I_k(V)$, where $I_k(U) = r$ iff $I_k(U, r)$;
 5.4 $(\forall U)((\forall x)(\exists r \geq 0)E_{1+1}^t(x, r, U) \implies (\exists s \geq 0)I_k(U, s))$;

6. Axiom of absolute continuity:

$$(\forall \varepsilon > 0)(\exists \delta > 0)(\forall U)(|I_2(U)| < \delta \implies |I_1(U)| < \varepsilon);$$

7. Axioms for an Archimedean field;

8. Transformations of axioms of $K_{\mathcal{A} f_1 f_2}^a$:

$$(\forall \vec{x})E_n^s(\vec{x}, X_\varphi), \text{ where } \varphi \text{ is an axiom of this logic};$$

9. Axioms of realizability of all sentences φ of T :

$$(\forall x_0)E_1^s(x_0, X_\varphi).$$

A weak model $\langle \mathfrak{A}, I_1, I_2 \rangle$ for $K_{\mathcal{A} f_1 f_2}^a$ can be transformed to a standard model \mathfrak{B} for $K_{\mathcal{A}}'$ by taking: $B_\varphi^{\mathfrak{B}} = \{ \vec{a} \in A^n : \mathfrak{A} \models \varphi[\vec{a}] \}$, $U_\tau^{\mathfrak{B}}(\vec{a}) = \tau^{\mathfrak{A}}(\vec{a})$ for $\vec{a} \in A^n$ and $I_k^{\mathfrak{B}}(U_\tau^{\mathfrak{B}}) = I_k(\tau)$ for each term τ with at most one free variable. By the Barwise Compactness Theorem (see [1]), it can be shown that S has a standard model \mathfrak{D} , because S is Σ -definable over \mathfrak{A} and A_2 holds in \mathfrak{A} . \mathfrak{D} can be transformed to a middle model \mathfrak{C} of T by taking:

$$R^{\mathfrak{C}} = \{ \vec{x} \in D^n : E_n^s(\vec{x}, X_{\mathbf{1}(R(\vec{x})=1)}) \} \text{ and}$$

$$I_k^{\mathfrak{C}}(\tau(x, \vec{a})) = I_k^{\mathfrak{D}}(U_{\tau(x, \vec{a})}) \text{ for } \vec{a} \in D^n \text{ and } k = 1, 2.$$

This completes the proof of the Middle Completeness Theorem. \square

In order to construct an absolutely continuous biprobability model, we need the following lemma.

LEMMA 2. (Loeb [4]) *In an ω_1 -saturated nonstandard universe, let M be an internal vector lattice of functions from an internal set A into ${}^*\mathbb{R}$ (the set of hyperreal numbers), and let I be an internal positive linear functional on M , such that $\mathbf{1} \in M$ and $I(\mathbf{1}) = 1$. Then there is a complete probability measure μ on A such that for each finitely bounded $\varphi \in M$, the standard part of φ is integrable with respect to μ and its integral is equal to the standard part of $I(\varphi)$.*

THEOREM 1. (Completeness Theorem for $L_{\mathcal{A}f_1f_2}^a$) *Let T be a set of sentences of $L_{\mathcal{A}f_1f_2}^a$ such that T is Σ_1 on \mathcal{A} and consistent. Then there is an absolutely continuous biprobability model of T .*

Proof. Let $\langle \mathfrak{A}, I_1, I_2 \rangle$ be a middle model of T in which each theorem of $L_{\mathcal{A}f_1f_2}^a$ is true. The Daniell integral construction from Lemma 2 produces probability measures μ_1, μ_2 on *A such that for each $*$ -term $\tau(x)$, the standard part of ${}^*I_k(\tau)$ is the integral $\int \text{st}(\tau(b))^{\mathfrak{A}} d\mu_k(b)$ (we define measures μ_n^k on ${}^*A^n$ by using iterated integrals). The absolute continuity in the middle model \mathfrak{A} implies the absolute continuity for all measurable sets. Also, using axiom A_3 , it can be shown that $\mu_n^1 \ll \mu_n^2$ for each $n \in \mathbb{N}$. This graded biprobability model $\hat{\mathfrak{A}} = \langle {}^*\mathfrak{A}, \mu_n^1, \mu_n^2 \rangle$ can be used to produce an absolutely continuous biprobability model of T (see [3]). \square

We can look only for a part of $L_{\mathcal{A}f_1f_2}^a$ which satisfies the finite compactness property, because this logic cannot satisfy the full compactness (for example, each finite subset of $T = \{ \int_1 \mathbf{1}(R(x)) dx > 0 \} \cup \{ \int_1 \mathbf{1}(R(x)) dx \leq \frac{1}{n} : n \in \mathbb{N} \}$, where R is a unary predicate, has a probability model, but not T itself).

THEOREM 2. *Let T be a set of sentences of $L_{\mathcal{A}f_1f_2}^a$ of the form $\tau \in [r, s]$. If every finite subset of T has a graded biprobability model, then T has a graded biprobability model.*

Proof. Let us suppose that each finite subset $\Psi \subseteq T$ has a model \mathfrak{A}_Ψ . By Lemma 1 we can suppose that \mathfrak{A}_Ψ is a middle model. Take an ultraproduct ${}^*\mathfrak{A}$ such that, for each $\varphi \in T$, almost every \mathfrak{A}_Ψ satisfies φ . Then form a graded biprobability model $\hat{\mathfrak{A}}$ from ${}^*\mathfrak{A}$ by the Daniell integral construction (Lemma 2). It can be shown by induction that every sentence of $L_{\mathcal{A}f_1f_2}^a$ of the form $\tau \in [r, s]$ which is true in almost all \mathfrak{A}_Ψ holds in $\hat{\mathfrak{A}}$, too. The absolute continuity condition can be expressed in the middle model by the first-order sentence

$$(\forall \varepsilon > 0)(\exists \delta > 0)(\forall U)(|I_2(U)| < \delta \implies |I_1(U)| < \varepsilon) .$$

By Łos's Theorem and Loeb construction the sentence

$$(\forall \varepsilon > 0)(\exists \delta > 0)(\forall X)(\mu_2(X) < \delta \implies \mu_1(X) < \varepsilon)$$

holds in $\hat{\mathfrak{A}}$. \square

2. The logic $L_{\mathcal{A} f_1 f_2}^s$. Axioms and rules of inference for the logic $L_{\mathcal{A} f_1 f_2}^s$ are those of $L_{\mathcal{A} f}$ (with both f_1 and f_2 in place of f , see [3]) together with the axioms of continuity A_1 and

(A₄) *Axiom of singularity:*

$$\bigvee_k \int_i \mathbb{H}_k \left(\int_1 \mathbf{1}(x=y) dy, \int_2 \mathbf{1}(x=y) dy \right) dx = 0, \quad i = 1, 2,$$

$$\text{where } H_k(s, t) = \begin{cases} 1, & \text{if } s \geq \frac{2}{k} \text{ and } t \geq \frac{2}{k} \\ 0, & \text{if } s \leq \frac{1}{k} \text{ or } t \leq \frac{1}{k} \\ \text{linear,} & \text{for other cases} \end{cases}.$$

THEOREM 3. (Completeness Theorem for $L_{\mathcal{A} f_1 f_2}^s$) *A theory T of $L_{\mathcal{A} f_1 f_2}^s$ is consistent iff T has a singular biprobability model.*

Proof. The proof of soundness is easy. Let $\langle \mathfrak{A}, I_1, I_2 \rangle$ be a weak model of T in which each theorem of $L_{\mathcal{A} f_1 f_2}^s$ is true. Let $\mathcal{F} = \{ B \subseteq A : \chi_B \in \text{dom}(I_1) = \text{dom}(I_2) \}$ be an algebra of subsets of A , where $\chi_B(x) = \begin{cases} 1, & \text{if } x \in B \\ 0, & \text{if } x \notin B \end{cases}$. Define finitely additive probability measures ν_1, ν_2 on \mathcal{F} by $\nu_k(B) = I_k(\chi_B)$, $B \in \mathcal{F}$ and $k = 1, 2$.

Then, for $a \in A$, the singleton $\{a\}$ belongs to \mathcal{F} because $\chi_{\{a\}} = (x=a)^{\mathfrak{A}}$, the set $B = \{a \in A : \nu_1\{a\} > 0 \ \& \ \nu_2\{a\} > 0\}$ belongs to \mathcal{F} and $\nu_k(B) = 0$, $k = 1, 2$ by A_4 .

By construction from [7], the measures ν_1, ν_2 can be extended so that $\nu_1 \subseteq \bar{\nu}_1$, $\nu_2 \subseteq \bar{\nu}_2$ and the measures $\bar{\nu}_1, \bar{\nu}_2$ are singular. Then construct a middle biprobability model $\langle \mathfrak{A}, \bar{I}_1, \bar{I}_2 \rangle$ of T by

$$\text{dom}(\bar{I}_k) = \text{dom}(I_k) \cup \{ \chi_C : C \in \bar{\mathcal{F}} \setminus \mathcal{F} \} \quad \text{and} \quad \bar{I}_k(\chi_C) = \bar{\nu}_k(C),$$

for each C from the extension $\bar{\mathcal{F}}$ of \mathcal{F} .

By Loeb's construction (Lemma 2) and the construction of the biprobability model from a graded biprobability model (see [3]), the singularity of finitely additive measures in the middle model will be preserved in the biprobability model. \square

Finally, we prove Finite Compactness Theorem for the singular case.

THEOREM 4. *Let T be a set of sentences of $L_{\mathcal{A} f_1 f_2}^s$ of the form $\tau \in [r, s]$. If every finite subset of T has a graded biprobability model, then T has a graded biprobability model.*

Proof. As in Theorem 2, our proof is based on the ultraproduct and Daniell integral construction. Now, we can suppose that \mathfrak{A}_Ψ is a weak model for each finite subset $\Psi \subseteq T$. Let $\bar{\mathfrak{A}}_\Psi$ be a middle model as in Theorem 3. Take an ultraproducts $^*\mathfrak{A} = \prod \bar{\mathfrak{A}}_\Psi$ such that, for each $\varphi \in T$, almost every $\bar{\mathfrak{A}}_\Psi$ satisfies φ . The condition of singularity can be express in the middle model by the first-order sentence $(\exists f)(\bar{I}_1(f) = 1 \wedge \bar{I}_2 = 0)$. By Los's Theorem and Loeb's construction the sentence $(\exists X)(\mu_1(X) = 1 \wedge \mu_2(X) = 0)$ holds in $^*\mathfrak{A}$. \square

REFERENCES

- [1] J. Barwise, *Admissible Sets and Structures*, Springer-Verlag, Berlin, 1975.
- [2] H.J. Keisler, *Hyperfinite model theory*; in Logic Colloquium '76, North-Holland, Amsterdam, 1977, pp.5-110.
- [3] H.J. Keisler, *Probability quantifiers*; in Model Theoretic Languages (J. Barwise and S. Feferman, eds.), Springer-Verlag, Berlin, 1985, pp. 509-556.
- [4] P.A. Loeb, *A functional approach to nonstandard measure theory*, Contemporary Math. **26** (1984), 251-261.
- [5] J. Los, E. Marczewski, *Extension of measures*, Fund. Math. **36** (1949), 267-276.
- [6] M.D. Rašković, *Completeness theorem for biprobability models*, J. Symbolic Logic **51**, (1986), 586-590.
- [7] M.D. Rašković, *Completeness theorem for singular biprobability models*, Proc. Amer. Math. Soc. **102** (1988), 389-391.
- [8] M.D. Rašković, R.S. Dordević, *Finite compactness theorem for biprobability logics*, Math. Balkanica **5** (1991), 12-14.
- [9] R.S. Dordević, *Analytic completeness theorem for absolutely continuous biprobability models*, Zeitschr. Math. Logik Grundlagen Math. **38** (1992), 241-246.

Prirodno-matematički fakultet
Radoja Domanovića 12
34001 Kragujevac, p.p. 60
Jugoslavija

(Received 16 11 1992)