

ON TOPOLOGICAL SPACES WITH DENSE COMPLETELY METRIZABLE SUBSPACES

T. Nagamizu

Dedicated to Professor Akihiro Okuyama on his 60th birthday

Abstract. We obtain some characterizations for the spaces which have a dense completely metrizable subspace and some results related with these spaces.

1. Introduction. In 1991, Michael [7] gave some characterizations for the spaces called *almost Čech-complete* which are simply called *almost complete*.

We know that for a metrizable space X , the following statements are equivalent [7, Proposition 4.4]:

- (1) X is an almost complete space,
- (2) X has a dense completely metrizable subspace.

The first purpose of this paper is to obtain some characterizations for the spaces which have a dense completely metrizable subspace and some results related with these spaces.

Arhangel'skiĭ and Kočinac asked several questions on weakly perfect spaces and spaces with dense G_δ -diagonal [1]. The second purpose of this paper is to give answers to their Questions 8 and 9.

2. Definitions and notations. All considered spaces are completely regular. A sequence $\{U_n \mid n \in \mathbf{N}\}$ of subsets of a space X is said to be *complete* if every filter base \mathcal{F} on X which is controlled* by $\{U_n \mid n \in \mathbf{N}\}$ clusters at some $x \in X$.

A sequence $\{\mathcal{U}_n \mid n \in \mathbf{N}\}$ of collections of subsets of X is said to be *complete* if $\{U_n \mid n \in \mathbf{N}\}$ is a complete sequence whenever $U_n \in \mathcal{U}_n$ for all $n \in \mathbf{N}$.

A collection \mathcal{U} of subsets of a space X is said to be an *almost cover* if $\bigcup \mathcal{U}$ is dense in X . Let \mathcal{U} and \mathcal{V} be collections of subsets of X . We say that \mathcal{V} is a *strong*

AMS Subject Classification (1991): Primary 54C10, 54E99.

* \mathcal{F} is controlled by $\{U_n\}$ if each U_n contains some $F \in \mathcal{F}$.

refinement of \mathcal{U} if \mathcal{V} is a refinement of \mathcal{U} and for each element $V \in \mathcal{V}$ there exists an element $U \in \mathcal{U}$ with $\text{Cl}(V) \subset U$.

The following lemma is proved in [7, Lemma 4.6].

LEMMA 2.1. *If X has a complete sequence $\{\mathcal{U}_n \mid n \in \mathbf{N}\}$ of open almost covers, then there exists a complete sequence $\{\mathcal{V}_n \mid n \in \mathbf{N}\}$ of disjoint open almost covers of X such that \mathcal{V}_{n+1} is a strong refinement of \mathcal{V}_n for each $n \in \mathbf{N}$.*

Let \mathcal{U} be a collection of subsets of X . \mathcal{U} is said to *separate points of X* if x and y are distinct points of X , then there exists different elements U_x and U_y in \mathcal{U} such that $x \in U_x$ and $y \in U_y$. \mathcal{U} is said to have a *finite intersection property* (f.i.p.) if every finite subcollection of \mathcal{U} have a nonempty intersection.

3. Characterizations. The main purpose in this section is to prove Theorem 3.4.

LEMMA 3.1. *Let X has a complete sequence $\{\mathcal{U}_n \mid n \in \mathbf{N}\}$ of open almost covers such that for each sequence $\{U_n \mid U_n \in \mathcal{U}_n, n \in \mathbf{N}\}$ with f.i.p., the set $\bigcap \{\text{Cl}(U_n) \mid n \in \mathbf{N}\}$ is a singleton. Then there exists a complete sequence $\{\mathcal{V}_n \mid n \in \mathbf{N}\}$ of disjoint open almost covers of X such that*

- (i) \mathcal{V}_{n+1} is a strong refinement of \mathcal{V}_n for each $n \in \mathbf{N}$, and
- (ii) for each decreasing sequence $\{V_n \mid V_n \in \mathcal{V}_n, n \in \mathbf{N}\}$, the set $\bigcap \{\text{Cl}(V_n) \mid n \in \mathbf{N}\}$ is a singleton.

Proof. From Lemma 2.1, there exists a complete sequence $\{\mathcal{V}_n \mid n \in \mathbf{N}\}$ of disjoint open almost covers of X such that \mathcal{V}_{n+1} is a strong refinement of \mathcal{V}_n and \mathcal{U}_n for each $n \in \mathbf{N}$.

Let $\{V_n \mid n \in \mathbf{N}\}$ is a decreasing sequence where $V_n \in \mathcal{V}_n$ for each $n \in \mathbf{N}$. By the construction of \mathcal{V}_n , for each $n \in \mathbf{N}$, there exists $U_n \in \mathcal{U}_n$ such that $\text{Cl}(V_{n+1}) \subset V_n \cap U_n$. Since $\{V_n \mid n \in \mathbf{N}\}$ is decreasing, $\{U_n \mid n \in \mathbf{N}\}$ has f.i.p.

For each $n \in \mathbf{N}$, we put $F_n = \text{Cl}(V_{n+1})$. By completeness of $\{\mathcal{V}_n \mid n \in \mathbf{N}\}$, we have:

$$\emptyset \neq \bigcap_{n \in \mathbf{N}} F_n = \bigcap_{n \in \mathbf{N}} V_n = \bigcap_{n \in \mathbf{N}} \text{Cl}(V_n) = \bigcap_{n \in \mathbf{N}} \text{Cl}(U_n).$$

Since $\bigcap_{n \in \mathbf{N}} \text{Cl}(U_n)$ is a singleton, $\bigcap_{n \in \mathbf{N}} \text{Cl}(V_n)$ is also a singleton. \square

THEOREM 3.2. *Let X have a complete sequence $\{\mathcal{U}_n \mid n \in \mathbf{N}\}$ of disjoint open almost covers such that*

- (i) \mathcal{U}_{n+1} is a strong refinement of \mathcal{U}_n for each $n \in \mathbf{N}$, and
- (ii) the set $\bigcap_{n \in \mathbf{N}} \text{Cl}(U_n)$ is a singleton for each decreasing sequence $\{U_n \mid U_n \in \mathcal{U}, n \in \mathbf{N}\}$.

Then X has a dense G_δ completely metrizable subspace.

Proof. By [7, Proposition 4.5], X is a Baire space. Since $G_n = \bigcup \mathcal{U}_n$ is an open dense subset in X for each $n \in \mathbf{N}$, then $M = \bigcap_{n \in \mathbf{N}} G_n$ is a dense G_δ set in X .

By the condition (ii), if x and y are two distinct points of M , then there exists $n \in \mathbf{N}$ such that \mathcal{U}_n separates x and y .

Let us define the metric ρ on M by

$$\rho(x, y) = \begin{cases} 0, & x = y \\ \min\{n \mid \mathcal{U}_n \text{ separates } x \text{ and } y\}^{-1}, & \text{otherwise.} \end{cases}$$

It is easy to check that ρ is a complete metric on M , by the condition (i) and (ii). Moreover, $U_n \cap M$ is a $1/n$ -open ball at x for each $U_n \in \mathcal{U}_n$ and $x \in U_n \cap M$. Hence the original topology on M is stronger than ρ -topology.

Now we show the next claim.

CLAIM. *Let F be a closed subset of M and $x \in M \setminus F$. Then there exist $n \in \mathbf{N}$ and $U_n \in \mathcal{U}_n$ such that $x \in U_n$ and $U_n \cap F = \emptyset$.*

Proof of the claim. Suppose that $U_n \cap F \neq \emptyset$ whenever $x \in U_n$ for each $n \in \mathbf{N}$. Pick a point x_n in $U_n \cap F$, and put $F_n = \text{Cl}\{x_m \mid m \geq n+1\}$ for each $n \in \mathbf{N}$. Then by the condition (i), $\{U_n \mid n \in \mathbf{N}\}$ is a decreasing sequence, and $\{x\} = \bigcap_{n \in \mathbf{N}} U_n$, by the condition (ii). It follows that

$$\emptyset \neq \bigcap_{n \in \mathbf{N}} F_n \subset \bigcap_{n \in \mathbf{N}} U_n = \{x\}.$$

Hence $x \in F$. This is a contradiction, and the claim is proved.

By the claim, ρ -topology is stronger than the original topology. It follows that M is a dense G_δ completely metrizable subspace. The proof is complete. \square

THEOREM 3.3. *Let X be a space with a dense completely metrizable subspace. Then there exists a complete sequence $\{\mathcal{U}_n \mid n \in \mathbf{N}\}$ of open almost covers of X such that for each sequence $\{U_n \mid U_n \in \mathcal{U}_n, n \in \mathbf{N}\}$ with the f.i.p., the set $\bigcap_{n \in \mathbf{N}} \text{Cl}(U_n)$ is a singleton.*

Proof. Let M be a dense completely metrizable subspace of X and ρ a compatible metric on M . Let $U(x, n)$ be an open subset of X such that $B(x, 1/n) = U(x, n) \cap M$ for each $x \in M$ and $n \in \mathbf{N}$, where $B(x, 1/n) = \{y \in M \mid \rho(x, y) < 1/n\}$ be a $1/n$ -open ball in M . Then for each $n \in \mathbf{N}$, $\mathcal{U}_n = \{U(x, n) \mid x \in M\}$ is an open almost cover of X .

Now we show that $\{\mathcal{U}_n \mid n \in \mathbf{N}\}$ is a complete sequence. Let $\{U_n \mid U_n \in \mathcal{U}_n, n \in \mathbf{N}\}$ be a sequence and \mathcal{F} a filter base on X which is controlled by $\{U_n \mid n \in \mathbf{N}\}$. Then for each $n \in \mathbf{N}$, there exists $F_n \in \mathcal{F}$ such that $F_n \subset U_n$. By the construction of \mathcal{U}_n , there exists x_n such that $x_n \in M$, $U_n = U(x_n, n)$ for each $n \in \mathbf{N}$. Since $\{U_n \mid n \in \mathbf{N}\}$ has the f.i.p., it follows that $\{x_n \mid n \in \mathbf{N}\}$ is a ρ -Cauchy sequence. Then there exists $x_0 \in M$ such that $\{x_n \mid n \in \mathbf{N}\}$ converges to x_0 . Therefore we have that $x_0 \in \bigcap \{\text{Cl}(F) \mid F \in \mathcal{F}\}$. Hence $\{U_n \mid n \in \mathbf{N}\}$ is a complete sequence.

In the same way, it is easy to see that $\bigcap_{n \in \mathbf{N}} \text{Cl}(U_n) = \{x_0\}$. The proof is complete. \square

These results lead to the following theorem.

THEOREM 3.4. *For the space X , the following conditions are equivalent.*

- (1) X has a complete sequence $\{\mathcal{U}_n \mid n \in \mathbb{N}\}$ of open almost covers such that for each sequence $\{U_n \mid U_n \in \mathcal{U}_n, n \in \mathbb{N}\}$ with f.i.p., the set $\bigcap_{n \in \mathbb{N}} \text{Cl}(U_n)$ is a singleton.
- (2) X has a complete sequence $\{\mathcal{U}_n \mid n \in \mathbb{N}\}$ of disjoint open almost covers such that
 - (i) \mathcal{U}_{n+1} is a strong refinement of \mathcal{U}_n for each $n \in \mathbb{N}$, and
 - (ii) for each decreasing sequence $\{U_n \mid U_n \in \mathcal{U}_n, n \in \mathbb{N}\}$, the set $\bigcap_{n \in \mathbb{N}} \text{Cl}(U_n)$ is a singleton.
- (3) X has a dense G_δ completely metrizable subspace.
- (4) X has a dense completely metrizable subspace.

A space X is said to be a *Namioka space* if the following condition is satisfied:

- (*) for any compact space Y and any separately continuous function $f: X \times Y \rightarrow \mathbb{R}$, there exists a dense G_δ subset $A \subset X$ such that f is jointly continuous at each point of $A \times Y$.

Next we consider the following game. Let α and β be two players with β the first to move. β starts by choosing a nonempty open subset $U_1 \subset X$. Then α chooses an open subset $V_1 \subset U_1$ and a point $x_1 \in V_1$. β then chooses a nonempty open subset $U_2 \subset V_1$ (he may choose as he wishes but is expected to escape from x_1). Next α chooses an open subset $V_2 \subset U_2$ and a point $x_2 \in V_2$, and so on. α wins if any subsequence $\{x_{n_p} \mid p \in \mathbb{N}\}$ of the sequence $\{x_n \mid n \in \mathbb{N}\}$ accumulates to at least one point of the set $\bigcap_{i=1}^{\infty} V_i = \bigcap_{i=1}^{\infty} U_i$. Then X is said to be σ -well α -favorable if α has a winning strategy in the game above.

It is well known that σ -well α -favorable spaces are Namioka [9, Theorem 6.3].

THEOREM 3.5. *Let X be a space with a dense completely metrizable subspace M . Then X is a σ -well α -favorable space. Hence X is a Namioka space.*

Proof. Let U_1 be a nonempty open subset of X . Since M is a dense subspace, we can pick a point x_1 in $M \cap U_1$. Then there exists a nonempty open subset V_1 of X such that $x_1 \in V_1 \subset \text{Cl}(V_1) \subset U_1$ and $d_M - \text{diam}(V_1 \cap M) \leq 1/2$, where d_M is a compatible metric on M . By induction, there exists a sequence $\{x_n \mid n \in \mathbb{N}\}$ in X and sequences $\{V_n \mid n \in \mathbb{N}\}$, $\{U_n \mid n \in \mathbb{N}\}$ of subsets of X such that

$$x_n \in V_n \cap M, \quad U_{n+1} \subset V_n \subset \text{Cl}(V_n) \subset U_n, \quad \text{and} \quad d_M - \text{diam}(V_n \cap M) \leq 1/n + 1$$

for each $n \in \mathbb{N}$. Since $\{x_n \mid n \in \mathbb{N}\}$ is a d_M -Cauchy sequence in M , there exists x_0 in M such that $\{x_n \mid n \in \mathbb{N}\}$ converges to x_0 . By the construction of $\{V_n \mid n \in \mathbb{N}\}$, we have $x_0 \in \bigcap_{n \in \mathbb{N}} V_n = \bigcap_{n \in \mathbb{N}} \text{Cl}(V_n)$. The proof is complete. \square

THEOREM 3.6. *Let X be a space with a dense completely metrizable subspace, Y a space and $f: X \rightarrow Y$ an irreducible, closed, continuous and onto map. Then Y has a dense completely metrizable subspace.*

Proof. By Theorem 3.4, there exists a complete sequence $\{\mathcal{U}_n \mid n \in \mathbb{N}\}$ of disjoint open almost covers of X , which satisfies the conditions (i) and (ii) of (2). For each $U \in \mathcal{U}_n$, put $W(U) = Y \setminus f(X \setminus U)$. Then each $W(U)$ is a nonempty open subset of Y . Now put $\mathcal{V}_n = \{W(U) \mid U \in \mathcal{U}_n\}$ for each $n \in \mathbb{N}$. It is easy to see that $\{\mathcal{V}_n \mid n \in \mathbb{N}\}$ is a complete sequence of open almost covers of Y , which satisfies the condition (1) of Theorem 3.4. The proof is complete. \square

4. countable dense Δ -base. Here $\Delta_X = \{(x, x) \mid x \in X\}$ is the diagonal in $X \times X$. Arhangel'skii and Kočinac [1] asked the following questions:

Question 1. When there exist a countable family \mathcal{U} of open sets in $X \times X$ such that $\bigcap \mathcal{U} \cap \Delta_X$ is dense in Δ_X and for each open neighborhood V of Δ_X in $X \times X$ one can find $U \in \mathcal{U}$ such that $U \subset V$? Such \mathcal{U} will be called a *dense Δ -base* of X .

Question 2. Let X be a compact space with a countable dense Δ -base. Does there exist a dense open metrizable subspace $Y \subset X$? A dense separable subspace $Z \subset X$?

It is clear that if X has a dense discrete subspace, then X has a countable dense Δ -base. Now we prove the following theorem.

THEOREM 4.1. *Let X be a compact space. If X has a dense completely metrizable subspace, then X has a countable dense Δ -base.*

Proof. Let M be a completely metrizable subspace of X and ρ a compatible metric on M . For each $n \in \mathbb{N}$, put $V_n = \{(x, y) \in M \times M \mid \rho(x, y) < 1/n\}$. Since each V_n is open set in $M \times M$, there exists an open set U_n in $X \times X$ such that $V_n = U_n \cap (M \times M)$. We show that $\mathcal{U} = \{U_n \mid n \in \mathbb{N}\}$ is a countable dense Δ -base of X .

Let V be an open neighborhood of Δ_X in $X \times X$. Then we prove that there exists $n \in \mathbb{N}$ such that $U_n \subset V$. By normality of $X \times X$, it is enough to show that $U_n \subset \text{Cl}(V)$.

Indeed, suppose that $U_n \not\subset \text{Cl}(V)$ for each $n \in \mathbb{N}$. Then there exists $(x_n, y_n) \in V_n \setminus \text{Cl}(V)$ for each $n \in \mathbb{N}$. By the definition of V_n , $\rho(x_n, y_n) < 1/n$ and $\{(x_n, y_n) \mid n \in \mathbb{N}\} \subset (X \times X) \setminus \text{Cl}(V) \subset (X \times X) \setminus V$. Since $(X \times X) \setminus V$ is compact, there exists a cluster point (x_0, y_0) of $\{(x_n, y_n) \mid n \in \mathbb{N}\}$ such that $(x_0, y_0) \in (X \times X) \setminus V$. Hence $x_0 \neq y_0$. Then there exist open subsets V_{x_0} and V_{y_0} such that $x_0 \in V_{x_0}$, $y_0 \in V_{y_0}$ and $\text{Cl}(V_{x_0}) \cap \text{Cl}(V_{y_0}) = \emptyset$. By the completeness, it follows that $\text{dist}(\text{Cl}(V_{x_0}) \cap M, \text{Cl}(V_{y_0}) \cap M) > 0$. But $\rho(x_n, y_n) < 1/n$ for each $n \in \mathbb{N}$, a contradiction.

Finally, since $\Delta_M \subset (\bigcap \mathcal{U}) \cap \Delta_X$, the set $(\bigcap \mathcal{U}) \cap \Delta_X$ is dense in Δ_X . The proof is complete. \square

Next we consider Question 2. We remark the following proposition.

PROPOSITION 4.2. *Let X be a space and M a dense completely metrizable subspace of X . Then the following conditions are equivalent.*

- (1) X is separable.

- (2) M is separable.
- (3) X satisfies the countable chain condition.

We have the negative answer of the second part of Question 2.

Example 4.3. Let X be the closed ordinal space $[0, \Omega]$, where Ω is the first uncountable ordinal. Since X is a compact scattered space, it has a dense uncountable discrete subspace. Therefore X has a countable dense Δ -base. But it is clear that X does not have any dense separable metrizable subspaces.

Let us note that if M is a dense open metrizable subspace of a compact space X , then M is a completely metrizable.

REFERENCES

- [1] A.V. Arhangel'skii and Lj.D. Kočinac, *On a dense G_δ -diagonal*, Publ. Inst. Math. **47** (61) (1990), 121–126.
- [2] R. Engelking, *General Topology*, PWN, Warszawa, (1977).
- [3] Z. Frolík, *Generalizations of the G_δ -property of complete metric spaces*, Czech. Math. J. **10** (1960), 359–379.
- [4] Z. Frolík, *Baire spaces and some generalizations of complete metric spaces*, Czech. Math. J. **11** (1961), 237–247.
- [5] R.W. Hansell, J.E. Jayne and M. Talagrand, *First class selector for weakly upper semi-continuous multi-valued maps in Banach spaces*, J. Reine. Angew. Mathematik **361** (1985), 201–220.
- [6] E. Michael, *A note on completely metrizable spaces*, Proc. Amer. Math. Soc. **96** (1985), 513–522.
- [7] E. Michael, *Almost complete spaces, hypercomplete spaces and related mapping theorems*, Topology Appl. **41** (1991), 113–130.
- [8] I. Namioka, *Radon-Nikodým compact spaces and fragmentability*, Mathematika **34** (1987), 258–281.
- [9] Z. Piotrowski, *Separately and jointly continuity*, Real Anal. Exchange **11** (1985/86), no. 2, 293–322.
- [10] N.K. Ribarska, *Internal characterization of fragmentable spaces*, Mathematika **34** (1987), 243–257.
- [11] D.B. Shakhmatov, *Compact spaces and their generalizations*, (preprint).
- [12] C. Stegall, *Topological spaces with dense subspaces that are homeomorphic to complete metric spaces and the classification of $C(K)$ Banach spaces*, Mathematika **34** (1987), 101–107.
- [13] C. Stegall, *The topology of certain spaces of measures*, Topology Appl. **41** (1991), 73–112.

Fukui National College of Technology
 Geshi Sabae
 Fukui 916
 JAPAN

(Received 12 11 1993)
 (Revised 19 04 1994)