

ON GAPS BETWEEN BOUNDED OPERATORS

Dragana Cvetković

Communicated by Mirosljub Jevtić

ABSTRACT. We consider the distance between an arbitrary bounded operator on a Banach space X and the null operator. The distance is computed in terms of various gaps. Thus, we generalize the Habibi's result from [4].

1. Introduction

In [4] Habibi considered the spherical gap between a linear operator and the null operator on finite dimensional Hilbert spaces. We extend (see also [1]) his result to various gaps on arbitrary Banach spaces.

Let X, Y be arbitrary Banach spaces. We denote by $\mathcal{G}(X)$, and $\mathcal{B}(X, Y)$, respectively, the set of all closed subspaces of X and the set of all bounded linear operators from X to Y . The closed unit sphere of a Banach space X is denoted by $S(X)$. If x is a vector of a Banach space X and M is a subset of X we put $\text{dist}(x, M) = \inf_{m \in M} \|x - m\|$. If M, N are subspaces of X , the set of all invertible operators C on X such that $C(M) = N$ is denoted by $\mathcal{B}(X; M, N)^{-1}$. Let $X \times Y$ be the space with the norm

$$\|(x, y)\| = (\|x\|^p + \|y\|^p)^{1/p}, \quad x \in X, y \in Y, p \geq 1.$$

Let $M, N \in \mathcal{G}(X)$. The spherical gap between M and N is defined by [2]

$$\tilde{\Theta}(M, N) = \max \left\{ \tilde{\Theta}_0(M, N), \tilde{\Theta}_0(N, M) \right\},$$

where

$$\tilde{\Theta}_0(M, N) = \sup_{m \in S(M)} d(m, S(N)).$$

2000 *Mathematics Subject Classification*. Primary: 46B20; Secondary 15A60.
Key words and phrases. spherical gap; geometric gap.

The geometric gap between M and N is defined by [6]

$$\Theta(M, N) = \max \{ \Theta_0(M, N), \Theta_0(N, M) \},$$

where

$$\Theta_0(M, N) = \sup_{m \in S(M)} d(m, N).$$

If $A, B \in \mathcal{B}(X, Y)$, then their graphs $G(A), G(B)$ are closed subspaces of $X \times Y$, so the gap between the operators A and B is defined as the gap between their graphs, i.e.,

$$\tilde{\Theta}(A, B) = \tilde{\Theta}(G(A), G(B)) \text{ and } \Theta(A, B) = \Theta(G(A), G(B)).$$

Also,

$$\tilde{\Theta}_0(A, B) = \tilde{\Theta}_0(G(A), G(B)) \text{ and } \Theta_0(A, B) = \Theta_0(G(A), G(B)).$$

Obviously,

$$\tilde{\Theta}(A, B) = \max \{ \tilde{\Theta}_0(A, B), \tilde{\Theta}_0(B, A) \},$$

$$\Theta(A, B) = \max \{ \Theta_0(A, B), \Theta_0(B, A) \}.$$

In [4] Habibi considered the spherical gap between operators on finite dimensional Hilbert spaces, assuming $p = 2$. We shall consider spherical and geometric gap of bounded operators on Banach spaces, where $p \geq 1$ is arbitrary. Thus, a generalization of Habibi's result is obtained.

2. Results

First we compute the spherical gap between a bounded operator and the null operator.

THEOREM 2.1. *If X, Y are Banach spaces and $A \in \mathcal{B}(X, Y)$, then*

$$\tilde{\Theta}(A) = \tilde{\Theta}(A, 0) = \left(\left(1 - \frac{1}{(1 + \|A\|^p)^{1/p}} \right)^p + \frac{\|A\|^p}{1 + \|A\|^p} \right)^{1/p}.$$

Proof. Since

$$G(A) = \{(x, Ax) : x \in X\} \text{ and } G(0) = \{(x, 0) : x \in X\},$$

it follows that

$$\begin{aligned} \tilde{\Theta}_0(A, 0) &= \sup_{(y, Ay) \in S(G(A))} \inf_{(x, 0) \in S(G(0))} \|(x, 0) - (y, Ay)\| \\ &= \sup_{\substack{y \in X \\ \|y\|^p + \|Ay\|^p = 1}} \inf_{\substack{x \in X \\ \|x\| = 1}} (\|x - y\|^p + \|Ay\|^p)^{1/p}. \end{aligned}$$

Consider the function

$$f(x, y) = (\|x - y\|^p + \|Ay\|^p)^{1/p}.$$

Suppose that $x \in X$, $y \in Y$ satisfy

$$(1) \quad \|x\| = 1, \quad \|y\|^p + \|Ay\|^p = 1.$$

Then

$$f(x, y) \geq ((\|x\| - \|y\|)^p + \|Ay\|^p)^{1/p} = ((1 - \|y\|)^p + \|Ay\|^p)^{1/p}.$$

Hence,

$$(2) \quad \inf_{\substack{x \in X \\ \|x\|=1}} f(x, y) \geq ((1 - \|y\|)^p + \|Ay\|^p)^{1/p}.$$

Also,

$$(3) \quad \inf_{\substack{x \in X \\ \|x\|=1}} f(x, y) \leq f\left(\frac{y}{\|y\|}, y\right) = ((1 - \|y\|)^p + \|Ay\|^p)^{1/p}.$$

So, from (2) and (3) it follows that

$$(4) \quad \inf_{\substack{x \in X \\ \|x\|=1}} f(x, y) = ((1 - \|y\|)^p + \|Ay\|^p)^{1/p}.$$

According to the condition (1) we have

$$\|y\| \geq \frac{1}{(1 + \|A\|^p)^{1/p}}.$$

Since the function $\phi(t) = (1 - t)^p + 1 - t^p$ is decreasing for $0 \leq t \leq 1$, we have

$$(5) \quad \begin{aligned} \tilde{\Theta}_0(A, 0) &= \sup_{\substack{y \in X \\ \|y\|^p + \|Ay\|^p = 1}} ((1 - \|y\|)^p + 1 - \|y\|^p)^{1/p} \\ &= \left(\left(1 - \frac{1}{(1 + \|A\|^p)^{1/p}}\right)^p + \frac{\|A\|^p}{1 + \|A\|^p} \right)^{1/p}. \end{aligned}$$

Let $y = cx$, where $c = \frac{1}{(1 + \|Ax\|^p)^{1/p}}$. It is obvious that y satisfy the condition (1)

$$\text{and } f(x, y) = \left(\left(1 - \frac{1}{(1 + \|Ax\|^p)^{1/p}}\right)^p + \frac{\|Ax\|^p}{1 + \|Ax\|^p} \right)^{1/p}.$$

So,

$$(6) \quad \inf_{\substack{y \in Y \\ \|y\|^p + \|Ay\|^p = 1}} f(x, y) \leq \left(\left(1 - \frac{1}{(1 + \|Ax\|^p)^{1/p}}\right)^p + \frac{\|Ax\|^p}{1 + \|Ax\|^p} \right)^{1/p}.$$

The function $\psi(t) = (1 - \frac{1}{(1+t)^{1/p}})^p + \frac{t}{1+t}$ is increasing, so from (6) it follows that

$$(7) \quad \begin{aligned} \tilde{\Theta}_0(0, A) &\leq \sup_{\substack{x \in X \\ \|x\|=1}} \left(\left(1 - \frac{1}{(1 + \|Ax\|^p)^{1/p}} \right)^p + \frac{\|Ax\|^p}{1 + \|Ax\|^p} \right)^{1/p} \\ &= \left(\left(1 - \frac{1}{(1 + \|A\|^p)^{1/p}} \right)^p + \frac{\|A\|^p}{1 + \|A\|^p} \right)^{1/p}. \end{aligned}$$

Finally, from (5) and (7) we have that

$$\begin{aligned} \tilde{\Theta}(A) &= \max \{ \tilde{\Theta}_0(A, 0), \tilde{\Theta}_0(0, A) \} = \tilde{\Theta}_0(A, 0) \\ &= \left(\left(1 - \frac{1}{(1 + \|A\|^p)^{1/p}} \right)^p + \frac{\|A\|^p}{1 + \|A\|^p} \right)^{1/p} \quad \square \end{aligned}$$

From the proof of Theorem 2.1 it follows that

$$\tilde{\Theta}_0(0, A) \leq \tilde{\Theta}_0(A, 0) = \left(\left(1 - \frac{1}{(1 + \|A\|^p)^{1/p}} \right)^p + \frac{\|A\|^p}{1 + \|A\|^p} \right)^{1/p}.$$

Notice that Habibi's main result in [4] is a special case of our Theorem 2.1 for $p = 2$.

Now, we consider the geometric gap.

THEOREM 2.2. *If X, Y are Banach spaces and $A \in \mathcal{B}(X, Y)$, then*

$$\Theta_0(A, 0) = \frac{\|A\|}{(1 + \|A\|^p)^{1/p}}.$$

Proof. It follows that

$$\begin{aligned} \Theta_0(A, 0) &= \sup_{\substack{y \in X \\ \|y\|^p + \|Ay\|^p = 1}} \inf_{x \in X} (\|x - y\|^p + \|Ay\|^p)^{1/p} \\ &= \sup_{\substack{y \in X \\ \|y\|^p + \|Ay\|^p = 1}} \|Ay\| = \sup_{z \in X} \frac{\|Az\|}{(\|z\|^p + \|Az\|^p)^{1/p}} = \frac{\|A\|}{(1 + \|A\|^p)^{1/p}} \quad \square \end{aligned}$$

In the case $p = 2$ we can obtain the following Habibi's result.

THEOREM 2.3. *If X and Y are Banach spaces, $A \in \mathcal{B}(X, Y)$ and $p = 2$, then the following holds:*

$$\Theta(A, 0) = \frac{\|A\|}{(1 + \|A\|^2)^{1/2}}.$$

Proof. By Theorem 2.2 it is sufficient to prove

$$\Theta_0(0, A) \leq \frac{\|A\|}{(1 + \|A\|^2)^{1/2}}.$$

Actually, this is a Habibi's result from [3]. For reader's convenience, we give a complete proof. Consider the function $f(x, y) = (\|x - y\|^2 + \|Ay\|^2)^{1/2}$. Let $x \in S(X)$ and $y = \frac{x}{1 + \|Ax\|^2}$. Then we have

$$\text{dist}((x, 0), G(A)) \leq f(x, y) = \frac{\|Ax\|}{(1 + \|Ax\|^2)^{1/2}}.$$

Since the function $\mu(t) = \frac{t}{1+t}$ is increasing, it follows that

$$\begin{aligned} \Theta_0(0, A) &= \sup_{\|x\|=1} \text{dist}((x, 0), G(A)) \leq \sup_{\|x\|=1} \frac{\|Ax\|}{(1 + \|Ax\|^2)^{1/2}} \\ &= \frac{\|A\|}{(1 + \|A\|^2)^{1/2}}. \quad \square \end{aligned}$$

Finally, we can prove the following result.

THEOREM 2.4. *Let X, Y be Banach spaces and $X \times Y$ is the space with the norm*

$$\|(x, y)\| = (\|x\|^2 + \|y\|^2)^{1/2}, \quad x \in X, \quad y \in Y.$$

If $A, B \in B(X, Y)$, then

$$\Theta(A, B) \leq 2(1 + \min\{\|A\|, \|B\|\}^2) \frac{\|A - B\|}{(1 + \|A - B\|^2)^{1/2}}.$$

Proof. From [5], Theorem 2.17, page 204, it follows that

$$\Theta(A, B) = \Theta((A - B) + B, 0 + B) \leq 2(1 + \|B\|^2)\Theta(A - B, 0),$$

and

$$\Theta(A, B) = \Theta(0 + A, (B - A) + A) \leq 2(1 + \|A\|^2)\Theta(B - A, 0).$$

Now, by Theorem 2.3 we have that

$$\Theta(A, B) \leq 2(1 + \|B\|^2) \frac{\|A - B\|}{(1 + \|A - B\|^2)^{1/2}},$$

and

$$\Theta(A, B) \leq 2(1 + \|A\|^2) \frac{\|A - B\|}{(1 + \|A - B\|^2)^{1/2}},$$

i.e.,

$$\Theta(A, B) \leq 2(1 + \min\{\|A\|, \|B\|\}^2) \frac{\|A - B\|}{(1 + \|A - B\|^2)^{1/2}}. \quad \square$$

The author wishes to thank referee for suggestions which greatly improved several proofs.

References

- [1] D. Cvetković, *Gaps between Banach spaces(in Serbian)*, Master Thesis, Faculty of Science and Mathematics, University of Nis (2000), 1–81.
- [2] I. C. Gohberg and A. S. Markus, *Two theorems on the opening between subspaces of a Banach space*, Uspekhi Mat. Nauk. **14 (5)** (1959), 135–140 (Russian).
- [3] J. F. Habibi, *The gap of the graph of a matrix*, Linear Algebra Appl. **186** (1993), 55–57.
- [4] J. F. Habibi, *The spherical gap of the graph of a linear transformation*, Linear Algebra Appl. **212** (1994), 501–503.
- [5] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [6] M. G. Krein, M. A. Krasnoselskii and D. P. Milman, *On the defect numbers of linear operators in a Banach space and on some geometric questions*, Sbornik Trudov Inst. Mat. AN Ukrainian SSR **11** (1948), 97–112 (Russian).

Prirodno-atematički fakultet
18000 Niš
Serbia and Montenegro
dragana@pmf.pmf.ni.ac.yu

(Received 17 01 2001)
(Revised 28 02 2002)